

**NUTRIENTS AND PHYTO-CHEMICAL ASSAY OF AFRICAN STAR APPLE KERNEL
(*Chrysophyllum africanum*) AS POTENTIAL FEEDSTUFF IN FISH AND
LIVESTOCK PRODUCTION**

AGBABIAKA, LA

Department of Animal Science and Fisheries, Imo State University , Owerri, Nigeria.

e-mail : adebokson2@yahoo.com

ABSTRACT

Nutrients assay of raw and fermented African Star Apple seeds were evaluated. The proximate analyses differ ($p < 0.05$) significantly between the raw (RSA) and processed (FSA) *Chrysophyllum africanum* seed in crude protein, lipid, dietary fibre and soluble carbohydrate. Protein value was $10.13 \pm 0.02\%$, Ash $7.25 \pm 0.01\%$, Crude fat $9.72 \pm 0.02\%$ and Soluble carbohydrate $69.45 \pm 0.84\%$ for RSA. However, the values of $5.55 \pm 0.01\%$, $14.49 \pm 0.76\%$, $4.94 \pm 0.07\%$, $1.19 \pm 0.01\%$ and $51.04 \pm 0.08\%$ were recorded for Ash, Crude protein, Crude fat, Crude fibre and Soluble carbohydrate in FSA respectively. Nevertheless, similar observations were recorded in the concentration of phyto-chemicals which decreased in all the parameters in FSA ($p < 0.05$) compared with RSA. The FSA contained relatively higher concentration of minerals than RSA ($p > 0.05$). Generally, there was a significant loss of nutrients in processed seed except crude protein, however, the reduction in the phyto-toxin in FSA may be an encouragement coupled with rich nutrients as a component of animal feed to replace the expensive orthodox energy feedstuff.

INTRODUCTION

The nutritional value of any ingredient or feed can be evaluated by biological, chemical and physical scores (Eddy and Udo, 2004). Chemical score has proven to be a vital tool in food chemistry because it tends to assess the nutritional value based on the proximate composition of the food/feed which includes the protein, carbohydrate, lipid, moisture, ash and dietary fibre contents respectively.

Plants are primary sources of medicine, food and shelter used by human on daily basis, their roots, leaves, fruits and seeds often provide food for humans (Amaechi, 2009). The fruits are main source of minerals, fibre and vitamins which are inevitable for human health (Berounmand and Deokule, 2009).

African star apple (*C. africanum*) popularly called “Agbalumo” among the Yoruba tribe of Western Nigeria is also known as “Agwaluma” or “Udara” in Hausa and Igbo languages respectively. It is primarily cultivated for its sweet fleshy fruits which had been reported as an excellent source for vitamin C, Iron, thickener or jam and flavours to diets and raw materials to some manufacturing industries such as resin (Adisa *et al.*, 2000). Star apple is believed to be

originated from low-lands of Central America and West Indian. It is common in both urban and rural centre in Nigeria especially during the months of December through April, when most orthodox cereals for feed formulation are usually scarce. The ripe fruit is highly perishable, and deteriorate within 5 days of harvest (Adisa *et al.*, 2000).

There has been a lot of work on the potential of the seeds/kernels which are often discarded after leaving the fleshy fruit. This experiment was conceived at investigating the nutritional potential(s) of the African star apple seeds with the aim of providing baseline information on its suitability or otherwise as animal feedstuff.

MATERIALS AND METHODS

Sample collection and preparation:- The fruits of African star apple used for this research were collected within the college premises of Federal Polytechnic, Nekede, Owerri, Nigeria. The fruits were washed thoroughly under tap, skin peeled and the pulp (edible portion) was removed to expose the seeds. The seeds were dehulled to expose the mesocarp (kernel). The mesocarp was divided into 2 batches; one portion was sundried as raw star apple (RSA), while the other portion was added water (3 times the volume of the seed) and allowed to ferment for 3 days. Thereafter, the supernatant water was decanted and sundried for 4 days until crispy.

The sundried samples were passed through a metal blade blender to produce star apple meals, packed in air tight bottles and labeled as RSA and FSA for raw and fermented star apple meals respectively. These samples were subjected to laboratory analyses to determine the chemical composition and phyto-toxins according to AOAC (2000) and Enujiugha and Olubunmi (2003) respectively.

RESULTS AND DISCUSSION

The results of the proximate and mineral composition of raw and fermented star apple kernel is shown in tables 1 and 2 respectively, while table 3 presents the anti nutrients of the test ingredients.

The percentage moisture content of the star apple kernel (RSA and FSA) ranged between 10.69 – 14.66±0.01%. These values were higher than (9.5%) reported for sweet maize (Cortez and Altamirano, 1999) and 6% for mango seed kernel (Zein *et al.*; 2005) but lower than the value of 68.42% reported for cassava (Bressani and Ortiz, 2000). The moisture content of any food is an index of its water activity which is used to measure or predict the stability and susceptibility to microbial contamination (Scott, 1995; Frazier and Wurstoff, 2010).

The Crude fat (Ether extract) concentrations were 9.72±0.02% and 4.94 ±0.07% for RSA and FSA respectively. These values were higher than 3.9% recorded for sweet maize (Cortez and Altamirano, 1999) and 5.08% recorded for raw mango seed kernel (Agbabiaka *et al.*; 2010). This

is an indication that African star apple seed / kernel contained a moderately high level of calorie comparable to some conventional feedstuff.

The Crude protein content of RSA was $10.13 \pm 0.12\%$ which is lower than the value of $14.49 \pm 0.76\%$ found in FSA. The higher Crude protein content of the FSA is attributed to the increase in build-up and effect of single celled microbes responsible for fermentation. Similar result was observed when cassava peel was fermented (Ofuya and Obilor, 1993; Ezeronye, 2001; Ubalua and Ezeronye, 2008).

However, the crude protein values recorded in this experiment also showed that African star apple seed is endowed with crude protein comparable to some orthodox energy feedstuffs such as maize and millet (CP = 9 – 10%) and tigernut (7 – 9.5%) according to Aduku (1993); Oladele *et al.* (2009) and Agbabiaka *et al.* (2012) respectively. Nevertheless, the Nitrogen free extract (soluble carbohydrate) decreased from 69.45% in RSA to $51.04 \pm 0.08\%$ in FSA. The decline was due to the effect of fermentation process which allowed the microbes such as yeast and bacteria to use up the carbohydrate as source of energy (Ekpe *et al.*, 2007).

There were noticeable decline in all the values of the anti-nutritional factors found in FSA compared with RSA (table 3), this may be due to the effect of fermentation and or leaching of some poly-phenols (Agbabiaka *et al.*, 2011). This experiment also revealed high concentration of macro and micro elements than RSA ($p > 0.05$), which is attributed to the role of micro-organisms of fermentation (Brock *et al.*, 1994).

Table 1: Proximate composition of African star apple (*Chrysophyllum africanum*).

Nutrients (%)	RSA	FSA
Moisture	14.66 ± 0.01	10.69 ± 0.01
Ash	7.25 ± 0.01	5.55 ± 0.01
Crude protein	10.13 ± 0.12	14.49 ± 0.76
Crude fat	9.72 ± 0.02	4.94 ± 0.07
Crude fibre	1.22 ± 0.02	1.19 ± 0.01
Nitrogen free extracts	69.45 ± 0.84	51.04 ± 0.08

Table 2: Mineral Composition of African star apple (*Chrysophyllum africanum*)

Constituents	Sample Concentration (%)	
	RSA	FSA
Sodium	12.54 ± 0.02	17.83 ± 0.01
Potassium	31.32 ± 0.01	12.17 ± 0.03
Calcium	0.54 ± 0.01	0.86 ± 0.01
Magnesium	0.39 ± 0.02	0.71 ± 0.01
Zinc	0.25 ± 0.01	0.44 ± 0.01
Iron	0.16 ± 0.01	0.15 ± 0.02
Copper	0.02 ± 0.01	0.02 ± 0.01
Manganese	0.05 ± 0.01	0.12 ± 0.01
Phosphorus	0.44 ± 0.01	1.09 ± 0.01

Table 3: Anti-nutrients of African star apple (*Chrysophyllum africanum*) .

Chemical Constituents	Sample concentration	
	FSA	RSA
Tannin (mg/100g)	4.87± 0.02	10.42± 0.01
Phytate (mg/g)	9.88± 0.12	12.37± 0.04
Phytin phosphorus (mg/g)	2.78± 0.01	3.48± 0.02
Oxalate (mg/g)	2.25± 0.06	4.10± 0.16
Alkaloid (%)	0.85± 0.07	0.48± 0.03
Saponin (%)	1.25± 0.13	0.74± 0.02

REFERENCES

- Adebisi A.A. (1997). Preliminary Survey of post-harvest and Marketing constraints of *Chrysophyllum aibidum* (African Star Apple) in Nigeria in: proceedings of a National workshop on the potentials of the star apple in Nigeria (eds) Denton OA, Ladipo DO, Adetoro MA, Sarumi MB, pp 84-102.
- Adisa V.A., Fajola A.O. (2000). Post Harvest seed rots of three species of citrus in Southwestern Nigeria. *Indian Phytopathol.* 35:595-603.
- Aduku A.O. (1993). *Tropical feedstuffs analysis table*. Department of Animal Science, ABU, Samaru, Zaria, Nigeria.
- Agbabiaka, L. A , Ekeocha. C. A and Amadi .S.A (2010). Performance of catfish (*Clarias gariepinus* Burchell, 1822) fed mango seed kernel based diets as substitute for maize. *Int. J. Trop. Agric. Food Syst.*, 4(4): 293-296.
- Agbabiaka L. A, F. N. Madubuike and B. O. Esonu, (2011). Effect of Processing on Nutrients and Anti-nutrients of Castor Oil Bean (*Ricinus communis*) Seeds and by products. *Pakistan Journal of Nutrition* 10(6): 561-563.
- Agbabiaka, L.A., Madubuko .C.U and Anyanwu .C.N (2012). Replacement value of tigernut meal (*Cyperus esculentus*) with maize in catfish (*Clarias gariepinus*) diets. *Science Research Reporter*. 2(2):160-164.
- Amaechi, N.C. (2009). Nutritive and Anti-Nutritive Evaluation of wonderful Kola (*Bucchozia coricea*) seeds *Pakistan Journal of Nutrition*, 8(8): 1120 – 1122.
- Ali Berounmand and Deokule, S.S. (2009). Studies on Nutritional values of some wild plants from Iran *Indian. Pak. J., Nutri.* 8(1):26-31
- A.O.A.C.(2000) *Official methods of Analysis*. Association of Official Analytical Chemists (ed. K. Herrick) 17th ed. Washington D.C. U.S.A.
- Brock T.D., Madigan, M.T., Martinko, J.M. and Parker J. (2004). *Biology of Micro-organisms* 7th Ed. Prentice Hall Int, USA, pp 282

- Bressani and Ortiz (2000). Nutritional content of cassava. "In praise food" UTNE reader http://www.utne.com/food_cullinary-Ethis.aspx pp 8. Retrieved 24-09-2010.
- Cortez and Altamirano (1999). Nutritional content of sweet maize. <http://www.Nutr.gov.uk/pdfs> (Accessed 12-07-2012).
- Eddy, N.O. and Udo, C.L. (2004). The energy value of some Nigeria soup. *Pakistan Journal of Nutrition*, 3(5):101-132
- Ekpe O.O , Umoh I.B and Eka O.U (2008). Effect of a typical rural processing method on the proximate composition and amino acid profile of bush mango seeds (*Irvingia gabonensis*). *African journal of Food, Agric. Nutrition and development* 7(1):1-12 .
- Enujiugha, V.N. and A.O. Olubunmi, 2003. Evaluation of nutrients and some anti-nutrients in lesser-known under-utilized oil seeds. *Int. J. Food Sci. Technol.*, 38: 1-4.
- Ezeronye, O.U., 2001. Fermentation and protein enrichment of cassava peel and rumen digesta and their evaluation as diets in growing swiss rats. *J. Agric. Biotechnol. Environ.*, 3 (1/2): 27-31
- Frazier, W.S. and Wwstoff, D.C. (2010). *Food Microbiology* 3rd edition, Mc Graw Hill, New York.
- Ofuya, C.O. and S.N. Obilor, 1993. The suitability of fermented cassava peel as a poultry feedstuff. *Bioresour. Technol.*, 44: 101-104.
- Oladele A.K., Oshundahunsi F.O. and Adebawale A.Y. (2009). Influence of processing techniques on the nutrients and anti-nutrients of tiger nut (*Cyperus esculentus*). *World Journal of Dairy and Food Sciences*, 4(2): 88-93.
- Scott, W.S. (1995). Water relations of food spoilage microorganisms. *Adv. Food res* 7:84-0127
- Ubalua AO and Ezeronye.O.U (2008). Growth Responses and Nutritional Evaluation of Cassava Peel Based Diet on Tilapia (*Oreochromis niloticus*) Fish Fingerlings. *Journal of Food Technology* 6 (5): 207-213

Zein, R. E., El-Bagoury, A. A. and Kassab, H. E. (2005). Chemical and nutritional studies on mango seed kernel. *Journal of Agric. Sci. Mansoura Univ.*, 30(6): 3285 – 3299.