

Biopesticide Activities of Extracts from *Calotropis procera* and *Annona squamosa*: A Potential Alternative to Chemical Pesticides

Nighat Begum¹, B. Sharma² and Ravi S. Pandey^{1*}
Departments of Zoology¹ and Biochemistry², University of Allahabad;
Allahabad, 211002, India;
*Corresponding author; E-mail:rspandey2004@yahoo.com

ABSTRACT

The control of harmful insects in agriculture and forestry, stores, cattle-breeding production, keeping of domestic animals and in human hygienic hygiene sector is still considered a serious problem. Widespread use of chemical pesticides represents a potential risk to human and environmental health. Therefore, the search for alternative strategies in pest control is the need of hour timely to overcome these problems. Desirable are preparations that exhibit new modes of actions and impair processes that are rather specific for the pests to be combated that are targetted. In the last twenty five years, much attention has been devoted to natural pest control agents. One of the most important groups among them are plant based active substances or mixtures of substances commonly known as 'botanicals'. Such natural products typically occur as cocktails of metabolically related compounds with differing activity/spectrum towards different insects. This-The present paper is a mini review presenting an updated account of biopesticide properties of extracts from two different plant species, which could be developed as a potential substitute of the for chemical pesticides.

Running title: Plant extracts as biopesticides

Key words: Biopesticides, plant extracts, chemical pesticides, toxicity, insects

1. INTRODUCTION

The ~~uncontrolled-unregulated~~ use of ~~chemical-and~~ synthetic insecticides has resulted in an increase ~~of-in~~ resistant strains of insect populations to conventional chemicals. Public awareness ~~seeking-clear-of~~ environmental ~~al~~ and food contamination from pesticides ~~lead~~ ~~has led~~ environmental protection agencies ~~(EPA) rulings to be formulated which to~~ ~~banned~~ the use of some of these chemicals like chlorinated, organophosphorus and carbamate insecticides. The synthetic/chemical pesticides have inherent drawbacks like (i) ~~biomagnifications~~, (ii) ~~their~~ loss of efficacy due to resistance development in insects, (iii) persistence of ~~some~~ active compounds in soil, ground water and lakes, (iv) ~~effects~~ effects on non-target organisms, (v) ~~disruption of biological control by natural enemies~~ (vi) resurgence of stored product insect pests and (vii) ~~fostered environmental and~~ human health concerns [1-4].

For the last two decades scientists ~~had been engaged in finding~~ ~~have searched for~~ botanical insecticides based on naturally occurring substances as a substitute for synthetic insecticides, ~~with an-~~ ~~They emphasized~~ on their ~~use in practice of~~ integrated pest management (IPM) rather than insect control [5]. The use of botanicals in pest management is useful in suppressing pest population as well as maintaining the sound ecological balance as the non target organisms are ~~not-less~~ affected.

The ~~reservoir~~ of ~~the~~ plant ~~species~~ possessing insecticidal substances is enormous. More than 2000 species ~~of the plants~~ are known ~~that to~~ possess some insecticidal activity [5].

Though pyrethrum, nicotine and rotenone were recognized as effective insect control agents since the middle of the 17th century, the most economically important and natural plant compounds are the pyrethrins obtained from the flower heads of pyrethrum ~~Chrysanthemum-Tanacetum~~ *cinerariaefolium* which are in use for ~~the~~ commercial insect

control. Despite the relative safety of ~~the some~~ well-known botanical insecticides, most of these substances have their drawback hindering large-scale application. Pyrethrins are unstable in ~~the sun~~light and are rapidly metabolized thus limiting their potency and application [6]. These limitations gave the impetus for synthesis of active analogues, termed ~~as~~ pyrethroids. Nicotine isolated from a number of Nicotiana species ~~of Nicotiana~~ is insecticidal, but its use in insect control has dropped steadily because of the high cost of production, disagreeable odour, extreme mammalian toxicity, instability in the environment and limited insecticidal activity [6].

Rotenone is ~~unstable and highly~~ toxic to fish. Further, ~~several insects many pest species~~ have ~~exhibited developed~~ resistance to pyrethroids [6]. For these reasons, the search for new, safer and more effective insecticides from ~~the~~ plants is desirable. Indeed the research in this area has led to the discovery of substances with increasing insecticidal activities. The substances include insect growth regulators / inhibitors and antifeedants. Keeping the importance of application of some environmentally ~~sound friendly~~ plant based molecules as potential substitutes ~~of the for~~ synthetic pesticides; ~~an~~ an endeavour has been made in this paper to present an updated account of the information available on the ~~biopesticidal biopesticide efficacies effects~~ of different ~~plant~~ species in general and *Calotropis procera* and *Annona squamosa* in particular.

2. PARTS OF THE PLANTS AND THEIR EXTRACTS USED AS BIOPESTICIDES

It was estimated that nearly 2400 species of plants in India possess insecticidal properties [7]. Botanical insecticides break down readily in soil and are not stored in animal and plant tissues. Often their effects are not as long lasting as those of synthetic insecticides and some of these products may be very difficult to find. ~~The p~~Plant parts used for extraction or assay ~~were the~~ have included leaves, roots, tubers, fruits, seeds,

flowers, the whole plant, bark, sap, pods and wood. The most commonly utilized parts were the leaves (62 species) followed by roots (16 species) and tubers (12 species). The plant families Asteraceae, Annonaceae, Asclepiadaceae, Fabaceae and Euphorbiaceae contain most the majority of the insecticidal plant species reported [8].

Recently several other plants viz. Neem, Pongamia, Indian privet, *Adathoda*, *Chrysanthemum*, Turmeric, Onion, Garlic, *Ocimum*, *Cedrus deodara*, *Nicotiana tabacum*, Custard apple, Ginger, *Citrus* fruits and some other plants have been reported as to have insecticidal plants-properties which-and therefore can be used in insecticide preparation [9,10]. Spinosad, a secondary metabolite produced by the fermentation from-of the fungus *Saccharopolyspora spinosa* mushroom and the active principle of the commercial products of the *Naturalyte*TM class and the azadirachtines-azadirachtins - a group of limonoids, obtained from the seeds of the Neem-neem tree (*Azadirachta indica*), have shown the efficacy to-in the control of fall webworm (*Hyphantria cunea*) [11].

Garlic acts as a repellent against-to various pests and is grown as a border intercrop to prevent pests from going-nearattacking the main crop. Extracts and powder preparations of garlic and onion bulbs are used to check pests in the-fields and grain ariges. Similarly, plants like Nochi-nochi (*Vitex negundo*), Pongamia-pongamia (*Pongamia glabra*), Adathoda-adathoda (*Adathoda vasica*) and Sweet-sweet flag (*Acorus calamus*) are-have been found to be effective against various pests of field crops during storage [12].

Extracts of *Pomoea cornea fistulosa*, *Calotropis gigantea* and *Datura strumarium* contain active principles toxic to many crop pests. Similarly an ethyl acetate extract of *Leucas aspera* leaves was found to be quite-quite effective against the early third instar larvae of the malaria mosquito *Anopheles stephensi* [13].

The data presented by a recent study showed that plant extracts cited by TRAMIL ethnopharmacological surveys have the potential to control the leaf-cutting ant,

Formatted: Font: Italic

Acromyrmex octospinosus. In particular, ~~the a~~ *Mammea americana* extract, with its natural low repellent effect and its high toxicity by ingestion, and *Nerium oleander* extracts, with their natural delayed action, are possibly the best extracts for the control of these ants [14].

The extract of flowers of champak (*Michelia champaca*) is potent against mosquito larvae. ~~Whereas the leaf~~ *Leaf* extracts of *Strychnos nux-vomica* had been shown to possess larvicidal efficacy against the filarial vector *Culex quinquefasciatus* [15]. The leaf extracts of lantana (*Lantana camara*), ~~Cit~~ *citrus* oil, tulsi (*Ocimum basilicum*, *O. sanctum*) and vetiver (*Vetivera zizanoides*) are useful in controlling leaf miners in potato, beans, brinjal, tomato and chillies, etc. Crushed roots of marigold (*Tagetes erecta*) provide good control of root-knot nematodes when applied to soil in mulberry gardens [16]. The seed extract of custard apple (~~Annona~~ *squamosa*) and citrus fruit (*Citrus paradisi*) are effective against ~~the~~ diamond back moth and ~~Colarado~~ *Colorado* potato beetle, respectively. Bark extract of *Melia azadiarach* acts as ~~a~~ potential antifeedant against ~~the~~ tobacco caterpillar (*Spodoptera litura*) and ~~the~~ gram pod borer (*Heliothis armigera*) [17,18]. Leaf extracts of lemon grass (*Cymbopogon citratus*), argemone (*Argemone mexicana*), cassia (*Cassia occidentalis*), artemesia (*Artemesia absinthium*) and sigesbekia (*Sieges beekii* *orientalis*) are strong antifeedants ~~of to~~ caterpillar pests like *Crocidolomia binotalis* [19]. ~~A Root-root~~ extract of drumstick (*Moringa oleifera*) inhibits growth of bacteria [20]. Plant extracts of *Azadirachta indica*, *Garcinia kola*, *Zingiber officinale* and *Allium sativum* have been used for the control of bacterial leaf spot of two varieties of *Solanum* (*S. gilo* and *S. torvum*) caused by *Xanthomonas campestris* [21]. These plant ~~extracts in harmoniouswhen integration-integrated~~ with other safe methods of pest control like biological control, trap crops and cultural practices etc. can provide eco-friendly and economically viable solutions for pest problems in near future.

Formatted: Font: Italic

4. PROPERTIES OF AN IDEAL INSECTICIDAL PLANT AND THEIR EXTRACTS

An ideal insecticidal plant should be ~~perennial~~ with wide distribution and abundantly present in nature. The plant parts to be used should be removable: ~~harvest of~~ leaves, flowers or fruit ~~and harvesting~~ should not ~~mean destruction of~~ damage the plant. The plants should require ~~small a modest footprint~~ space, ~~reduced minimal~~ management and little water and fertilization ~~and~~ should ~~not have a high~~ be required economic value. The active ingredient should be effective at low rates.

The crude plant extracts are advantageous in terms of efficacy and pest resistance management as the active substances present in them act ~~synergistically~~ [22,23]. Furthermore, they are decomposed in the environment much faster and easier than ~~most~~ synthetic compounds [24]. In the light of differences in geo-climatic zones and biodiversity, the plant kingdom still remains an untapped vast reservoir of new molecules endowed with massive biopesticidal potential. ~~Over 2000 plants belonging to some 60 plant families are known to exhibit insecticidal activities~~ [25,26]. Their crude preparations are applied as powders or dusts (for example neem leaf dust, pyrethrum flower dusts etc.) and aqueous or organic solvent extracts [27].

However, deriving new biopesticidal principle(s) from plants remains a complex and time consuming task, because it needs interdisciplinary skills ~~of for~~ isolation, purification, characterization, synthesis of standards (new/standard chemicals) and ~~their~~ screening for biological effect(s). While plant extracts may afford additive/synergistic action of several weak and strong biopesticidal activities, their purification and structure determination is essential for standardization, ~~as also~~ and for bioefficacy improvement. In the grim scenario of mounting hazards and cost of synthetic chemical pesticides, natural chemistry of plants shows a ray of hope for ~~environment and human friendly and~~

Comment [M1]: I would argue the opposite – an annual plant that could be readily propagated and cultivated would be a better choice. Most perennials do not produce enough biomass on an annual basis for continuous harvest.

Comment [M2]: Cite Miresmailli et al. (2006), Jiang et al. (2009) for synergy; Feng et al. (1995) for resistance delay to neem.

Formatted: Highlight

Comment [M3]: You said this in the introductory section; no need to repeat it.

sustainable pest management with minimal environmental and health impacts in future.

In this regard, leaf and seed extracts of *Calotropis procera* and *Annona squamosa* have shown enormous-considerable potential to be as promising-biopesticides [28-30].

5. THE BIOPESTICIDE ACTIVITIES OF PLANTS

The biopesticide activities of two known plant species are described as following:

5.1. *Calotropis procera*

Calotropis procera (Ait.) known as Aak and Madar, is a member of the plant family Asclepiadaceae, a shrub widely distributed in West Africa, Asia and other parts of the tropics [31]. The plant is erect, tall, large, much-multi-branched and-perennial with a milky latex throughout. A large quantity of latex can be easily collected from its green parts [31]. The abundance of latex in the green parts of the plant indicates that it is probably produced and accumulated as a defense strategy against organisms such as virus, fungi, and-insects and larger herbivores [32]. The presence of plant defense related proteins such as hevein, an alpha-amylase inhibitor, has been described to occur in from the latex secretion of other plants [33]. Hence-Thus it has been found to be used by local indigenous people to successfully combat some cutaneous fungal infections successfully.

Despite some reports of toxicity associated with *Calotropis* feeding-toingestion in animals, its use in ethnoveterinary medicine is increasing based on empirical evidence in the successful treatment of different ailments. Different plant parts as well as latex of *C. procera* have been reported to have emetic, purgative and anthelmintic effects in traditional medicine. *C. procera* flowers are mostly used as an anthelmintic in small ruminants in the form of decoction and/or crude powder mixed with jaggeryjaggery (?) and administered as physic drench/balls [32].

5.1.1 Chemical constituents of *C. procera* extract

The active ingredients of *C. procera* are a number of alkaloids, enzymes and other inorganic elements. Cardenolides, the principal steroidal toxins isolated from *C. procera*, are cardiac poisons have been reported to inhibit the ubiquitous and essential animal enzyme Na^+/K^+ -ATPase. Moreover, only some special sorts of insects are known to feed on cardenolide-containing plants [34]. Coagulum contains resins and caoutchouc. The latex contains caoutchouc, calotropin, uscharin 0.45%, calotoxin 0.15%, calactin (composed of calotropagenin and hexose) 0.15%, trypsin, voruscharin, uzarigenin, syriogenin and proceroside. Leaves and stalks bears calotropin and calotropagenin [35]. Bark of the rootRoot bark possesses the phenolics benzoylneolone, benzoyl isolineolone, madaralban and madar fluavil. Flowers contains the anthocyanin cyanidin-3-rhamnoglucoside. The whole plant contains various enzymes such as trypsin, α -calotropeol, β -calotropeol and β -amyrin. Inorganic components such as calcium oxalate, nitrogen and sulphur are also found. The isolated fatty acid composition in the extract of *C. procera* has 7 saturated fatty acids and 11 unsaturated fatty acids. The essential elements such as Al, As, Cu, Ca, Cr, Cd, Fe, K, Mn, Na, Pb, and Zn have been analyzed from the medicinal plant in variable range. The total protein in *C. procera* was 27-32% [36]. The chemical structures of some phytochemicals with biopesticide activities are shown in the Fig.1.

Formatted: Font: Italic




Fig.1(a): Chemical Structure of Calotropagenin




Fig.1(b): Chemical Structure of Uscharidin




Fig.1(c): Chemical Structure of Uscharin




Fig.1(d): Chemical Structure of Calotropin

Source: Hanna et al. [35]

Comment [M14]: This structure should follow calotropagenin

5.1.2 Impact of phytochemicals isolated from *C. procera* showing biopesticide activities against non-target systems

Calotropin found in latex causes slowing of heart beat and gastroenteritis in frogs. [Latex](#) [The latex](#) is [an](#) irritant to the skin and mucous membranes and may cause blindness. It may rupture the muscles of [the](#) intestine and colon and death may occur. The plant may cause severe bullous dermatitis, slowed but stronger heart beat, laboured respiration, increased blood pressure, convulsions and death [37]. A recent finding indicates that the root [part](#) of *C. procera* possesses *in vitro* cytotoxicity against oral and CNS human cancer cell lines [38]. Further investigations are required to obtain the clinically important lead molecules for [the](#) drug development.

The antimicrobial activities of the organic solvent extracts of stem, leaves and flowers of *C. procera* against *Alternaria alternate*, *Aspergillus flavus*, *Asperigellus niger*, *Bipolaris bicolor*, *Curvularia lunata*, *Pencillium expansum*, *Pseudomonas marginales*, *Rhizoctonia solani* and *Ustilago* have been reported [by](#) [39]. In Unani and Ayurvedic medical system, various parts of this plant have been used in curing a number of ailments [34].

5.2 *Annona squamosa*

The Annonaceae (custard-apple family) is a large family of almost exclusively tropical trees and shrubs comprising about 130 genera and 2300 species. Plant parts of some species of this family have been used traditionally as insecticides. For example, the powdered seeds and leaf juices of *Annona* spp. are used to kill head and body lice, and bark of *Goniiothalamus macrophyllus* is used to repel mosquitoes [40].

Annona squamosa L., commonly known as Sitaphal and Custard [Apple](#), is a native of West Indies and is cultivated throughout India, mainly for its edible fruit. The young leaves of *A. squamosa* are used extensively for [its-their](#) antidiabetic activity. The plant contains aporphine alkaloids, carvone, linalool, limonene [41], squamosin [42] and

Comment [M15]: Commonly known as sweetsop in southeast Asia

quercetin [43]. Acetogenins, ~~another a characteristic group of compounds~~ isolated from *Annona squamosa* ~~seeds~~ have been suggested to act as potential anti-neoplastic agents [44]. ~~These are also the principal insecticidal constituents of Annona seed extracts.~~

5.2.1 Chemical constituents of *A. squamosa*

The leaf extracts of this plant are known to contain different types of flavonoids some of which can act as phytoalexins [45]. These are mainly involved with the defense mechanisms of the plant and some are known to possess several antimicrobial and insecticidal properties [46]. Annotemoyin, annotemoyin, squamocin and cholesteryl glucopyranosides are isolated from the seeds of *A. squamosa* [47].

~~Acetogenins a different class of secondary metabolites was found to occur in various parts of A. squamosa~~ [48]. More than 13 different alkaloids, several terpenes, kauranes were isolated. Antibacterial activity was attributed to terpenes and kauranes. Seeds yielded fixed oil containing hydroxyacids and found to contain anti-inflammatory cyclic peptides. Many pharmacological activities were experimentally reported ~~on for~~ extracts of *A. squamosa* L. ~~These~~ included antitumour, cytotoxic, anti-inflammatory, analgesic, antidiabetic, antioxidant, larvicidal, insecticidal, molluscicidal, licial, antibacterial, nutritive and ~~antithyroid~~ ~~antithyroid~~ properties [49].

The seeds are acrid and poisonous. Bark, leaves and seeds contain the alkaloid, anonaine. Six other aporphine alkaloids have been isolated from the leaves and stems: corydine, roemerine, norcorydine, norisocarydine, isocorydine and glaucine. Aporphine, norlaureline and dienone may be present also. A paste of the seed powder has been ~~applied to the head used~~ to kill ~~head~~ lice but care must be taken to avoid eye contact. If applied to the uterus, it induces abortion. Heat-extracted oil from the seeds has been employed against agricultural pests. Studies have shown the ether extract of the seeds

to have no residual toxicity after two days. In Mexico, the leaves are rubbed on floors and put in hen's nests to repel lice [40].




Figure 2. Chemical structure of squamosin

5.2.2 Impact of phytochemicals from *A. squamosa* on non-target systems

Mehra and Hiradher [50] reported larvicidal action of *A. squamosa* against larvae and pupae of *Culex quinquefasciatus*. ~~Its~~ The seed oil is larvicidal against [the rusty grain beetle](#) *Tribolium castaneum* (Herbst) and mosquitoes [51].

Annonaceous acetogenins extracted from tree leaves, bark and seeds have pesticidal and/or insect antifeedant properties [52]. This group of $C_{32/34}$ fatty-acid-derived natural products is among the most potent inhibitors of complex I in the mitochondrial electron transport system [53], [which is consistent with the mode-of-action of rotenone](#). To date, nearly 400 of these compounds have been isolated from the genera *Annona*, *Asimina*, *Goniothalamus*, *Rollinia* and *Uvaria* [54]. Their biological activities include cytotoxicity, and *in vivo* antitumor, antimalarial, parasitocidal and pesticidal effects [55].

A summary of some important Indian traditional plants exhibiting varied biological activities are displayed in Table 1.

Table 1. Some Indian traditional plants exhibiting ~~varied~~varius medicinal properties

Botanical Name	Family	Parts Used	Medicinal Use
----------------	--------	------------	---------------

<i>Emblica officinalis</i>	Euphorbiaceae	Fruit	Vitamin - C, cough, diabetes, cold, laxative, hyper acidity
<i>Withania somnifera</i>	Solanaceae	Root, Leaves	Restorative tonic, stress, nerves disorder, aphrodisiac
<i>Solanum nigrum</i>	Solanaceae	Fruit/ whole plant	Dropsy, general debility, diuretic, antidyenteric.
<i>Rauwolfia serpentina</i>		Root	Hyper tension, insomnia
<i>Santalum album</i>	Santalinaceae	Heart wood oil	Skin disorder, burning, sensation, Jaundice, cough
<i>Asparagus racemosus</i>	Liliaceae	Tuber, root	Enhance lactation, general weakness, fatigue, cough
<i>Ocimum sanctum</i>	Lamiaceae	Leaves/ Seed	Cough, cold, bronchitis, expectorant
<i>Mentha pipertia</i>	Lamiaceae	Leaves, Flower, Oil	Digestive, pain killer
<i>Aloe vera</i>	Liliaceae	Leaves	Laxative, wound healing, skin burns, ulcer
<i>Adhatoda vesica</i>	Sacanthaceae	Whole plant	Antispasmodic, respiratory stimulant

Antimicrobial and insecticidal properties of partially purified flavonoids from [an](#) aqueous extract of *A. squamosa* have been reported against *Callosobruchus chinensis* [56].

Ethanollic seed extracts of *A. squamosa* from Maluku (Indonesia) were highly inhibitory to larval growth of *Spodoptera litura* [57].

Many plants have been reported for their potential insecticidal actions on larvae and/or adults of house flies [58-60]. They also affect their metamorphosis, ~~of~~ emergence, ~~of~~ fecundity ~~and/or longevity~~ ~~or life span~~ [61].

6. CONCLUSION

The reports cited above ~~reports~~ clearly indicate the potential of the ~~aforesaid~~ aforementioned ~~two plants as panacea for the pest population control~~ management. Some of the phytochemicals isolated from them are also useful in management of certain diseases. Further validation of the extracts from these plants through multidimensional biochemical and molecular approaches is required and ~~their~~ field trials may be useful in evaluating their suitability as a safer, economic and ecofriendly biopesticides.

7. REFERENCES

1. Denholm I, Pickett JA, Devonshire AL. (Eds) Insecticide resistance: from mechanisms to management. IACR-Rothamsted, Harpenden;1999.
2. Nivsarkar M, Cherian B, Padh H. Alpha-terthienyl: A plant-derived new generation insecticide. Current Science. 2001;81:667-72.
3. Taskin V, Kence M, Göçmen B. Determination of malathion and diazinon resistance by sequencing the MdE7 gene from Guatemala, Colombia, Manhattan, and Thailand housefly (*Musca domestica* L.). Russian J Genetics. 2004;40:377-80.
4. Ramoutar D, Alm SR, Cowles RS. Pyrethroid resistance in populations of *Listronotus maculicollis* (Coleoptera: Curculionidae) from southern New England golf courses. J Econ Entomol. 2009;102(1):388-92.
5. Etebari K, Matindoost L, Singh RN. Decision tools for mulberry thrips *Pseudodendrothrips mori* (Niwa, 1908) management in sericultural regions; An overview. Entomologia Sinica. 2004;11:243-58.
6. Isman MB. Botanical insecticides, deterrents and repellents in modern agriculture and in increasingly regulated world. Ann Rev Ent. 2006; 51:44-56.
7. Baskaran V, Narayanasamy P. Traditional Pest Control. Caterpillar Publications, Mariyappa Nagar, Tamil Nadu, India; 1995.

Formatted: Font: Italic

8. Boulogne I, Petit P. Insecticidal and antifungal chemicals produced by plants: a review. Environ Chem Lett, 2012; 10(4): 325-47.
9. Rahuman AA, Bagavan A, Kamaraj C, Saravanan E, Zahir AA, Elango G. Efficacy of larvicidal botanical extracts against *Culex quinquefasciatus* Say (Diptera: Culicidae). Parasitol Res. 2009;104:1365–372.
10. Osipitan AA, Oseyemi AE. Evaluation of the Bio-insecticidal Potential of Some Tropical Plant Extracts Against Termite (Termitidae:Isoptera) in Ogun State, Nigeria. J Entomol. 2012;9:257-65.
11. Brudea V, Rîșca I, Enea C, Tomescu C. Efficacy of some biopesticides and plant secondary metabolites against fall webworm *Hyphantria Cunea* Drury (F. Arctiidae-Lepidoptera) in the Lab Conditions. Cercetari Agronomice in Moldova. 2012;45(1):73–80.
12. Sadek MM. Antifeedant and toxic activity of *Adhatoda vasica* leaf extract against *Spodoptera littoralis* (Lep., Noctuidae). J Appl Ent, 2003;127:396–404.
13. Arivoli S, Ravindran KJ, Tennyson S. Larvicidal efficacy of some plant extracts against the malarial vector *Anopheles stephensi* Liston (Diptera: Culicidae). World J Med Sci. 2012;7(2):77-80.
14. Boulogne I, Germosen-Robineau L, Ozier-Lafontaine H, Jacoby-Koaly C, Aurela L, Loranger-Merciris G. *Acromyrmex octospinosus* (Hymenoptera: Formicidae) management. Part 1: Effects of TRAMIL's insecticidal plant extracts. Pest Manag Sci. 2012;68(2):313–20.
15. Arivoli S, Samuel T. Larvicidal efficacy of *Strychnos nux-vomica* Linn. (Loganiaceae) leaf extracts against the filarial vector *Culex quinquefasciatus* Say. (Diptera: Culicidae). World J Med Sci. 2012; 7(1): 6-11.
16. Chitwood DJ. Phytochemical based strategies for nematode control. Annu Rev Phytopathol, 2002;40: 221–49.
17. Wheeler DA, Isman MB, Sanchez-Vindas PE, Arnason JT. Screening of Costa Rican *Trichilia* species for biological activity against the larvae of *Spodoptera litura* (Lepidoptera: Noctuidae). Biochem Syst Ecol. 2001; 29:347–58.
18. Nathan SS. Effects of *Melia azedarach* on nutritional physiology and enzyme activities of the rice leaffolder *Chaphalocrocis medinalis* (Guenée) (Lepidoptera: Pyralidae), Pestic Biochem Physiol. 2006; 84:98–108.

Formatted: Font: Italic

19. Abdelgaleil SAM, Abbassy MA, Belal AH, Abdel Rasoul MAA. Bioactivity of two major constituents isolated from the essential oil of *Artemisia judaica* L. [Biores Tech.](#) 2008; [99 \(13\)](#): 5947-950.
20. Fahey JW. *Moringa oleifera*: A Review of the Medical Evidence for Its Nutritional, Therapeutic, and Prophylactic Properties. Part1. Trees for Life J. 2005;1:5-10.
21. Opara EU, Obani ET. Performance of Some Plant Extracts and Pesticides in the Control of Bacterial Spot Diseases of *Solanum*. Agri Journal.2010;5(2):45-49.
22. Schmutterer H. Properties and potential of natural pesticides from the neem tree, *Azadirachta indica*. Ann Rev Entomol.1990;35:271-97.
23. Voßlinger M, Schmutterer H. Development of resistance to [Azadirachtin-azadirachtin](#) and other neem ingredients. In: Schmutterer, H. (Ed) The neem tree, 2nd edn. Neem Foundation, Mumbai. 2002; 598–606.
24. Ujvary I. Transforming natural products into natural pesticides- experience and expectations. Phytoparasitica. 2002;30:1–4.
25. Copping LG, Menn, JJ. Biopesticides: a review of their action, applications and efficacy. Pest Manag Sci. 2000; 56:651–76.
26. Walia S, Koul O. Exploring plant biodiversity for botanical insecticides. In: Sustainable crop protection, Biopesticide strategies, Kalyani Publishers, New Delhi. 2008;191-206.
27. George DR, Guy JH, Arkle S, Harrington D, De Luna C, Okello EJ, Shiel RS, Port G, Sparagano O.A. Use of plant-derived products to control arthropods of veterinary importance: a review. Ann N Y Acad Sci. 2008;1149:23-6.
28. Begum N, Sharma B, Pandey RS. Evaluation of Insecticidal efficacy of *Calotropis procera* and *Annona squamosa* ethanol extracts against *Musca domestica*. J Biofertil Biopestici. 2010; 1:1,.doi.org/10.4172/2155-6202.1000101.
29. Begum N, Sharma B, Pandey RS. Toxicity potential and anti AChE activity of some plant extracts in *Musca domestica*. J Biofertil Biopestici. 2011; 2:2,.doi.org/10.4172/2155-6202.1000108.

30. Begum N, Sharma B, Pandey RS. Insecticidal potential of *Calotropis procera* and *Annona squamosa* ethanol extracts against *Musca domestica*. Natural Products – An Indian journal. 2012;7:5.
31. Irvine FR. Woody plants of Ghana. Oxford University Press, London. 1961;48-50.
32. Larhsini M, Bousad M, Lazrek, JM, Amarouch H. Evaluation of antifungal and molluscicidal properties of extracts of *Calotropis procera*. Fitoterapia. 1997;68:371-73.
33. Wititsuwannakul D, Sakulborirug C, Wititsuwannakul RA. Lectin from the bark of the rubber tree (*Hevea brasiliensis*). Phytochem. 1998;47:183-87.
34. Agrawal AA, Petschenka G, Bingham RA, Weber MG, Rasmann S. Toxic cardenolides: chemical ecology and coevolution of specialized plant–herbivore interactions. New Phytologist, 2012;194: 28–45.
35. Hanna AG, Shalaby NMM, Morsy NAM, Simon A, Tóth G, Malik S, Duddeck, H. Structure of a calotropagenin-derived artifact from *Calotropis procera*. Magnetic Resonance in Chemistry. 2002;40:599–602.
36. Khanzada SK, Shaikh W, Kazi TG, Sofia S, Kabir A, Kandhro AA. (Analysis of fatty acid, elemental and total protein of *Calotropis procera* medicinal plant from Sindh, Pakistan. Pak J Bot. 2008; 40(5):1913-921.
37. Duke JA. CRC Handbook of Medicinal Herbs, CRC Press, Inc. Raton Florida, USA; 1986;292-95.
38. Bhagat M, Arora JS, Saxena, AK. *In vitro* cytotoxicity of extracts and fractions of *Calotropis procera* (Ait.) roots against human cancer cell lines. Int J Green Pharmacy. 2010;4(1):36-40.
39. Varahalarao V, Chandrashekar KN. *In vitro* bioactivity of Indian medicinal plant *Calotropis procera* (Ait). J Glob Pharma Technol. 2010;2(2):43-45.
40. Morton J. (1987). Sugar Apple. In: Fruits of warm climates. Julia FM, Miami FL (Ed) p. 69-72.
41. Ekundayo O. A review of the volatiles of the Annonaceae. J Essent Oil Res. 1989;1: 223.
42. Yu JG, Luo, XZ, Sun L, Li DY, Huang WH, Liu CY. Chemical constituents from the seeds of *Annona squamosa*. Yao Xue Xue Bao. 2005;40(2):153-58.

43. Panda S, Kar A. *Annona squamosa* seed extract in the regulation of hyperthyroidism and lipidperoxidation in mice: Possible involvement of quercetin. *J Plant Biochem.* 2007;14(12):799-805.

Formatted: Font: Not Italic

44. Yuan SSF, Chang HL, Chen HW, Yeh YT, Kao YH, Lin KH, Wu YC, Su JH. Annonacin, a mono-tetrahydrofuran acetogenin, arrests cancer cells at the G1 phase and causes cytotoxicity in a Bax and caspase-3-related pathway. *Life Sci.* 2003;72(25):2853-861.

45. Chatterjee A, Pakrashi SC, editors. *Annona squamosa* in the treatise of Indian medicinal plants Publication and Information Directorate, New Delhi, 1995;130.

46. Padmavati M, Reddy AR. Flavonoid biosynthetic pathway and cereal defence response: An emerging trend in biotechnology. *J Plant Biochem Biotechnol.* 1999;8:15-20.

47. Mukhlesur RM. Antimicrobial and cytotoxic constituents from the seeds of *Annona squamosa*. *Fitoterapia.* 2005;7:484-89.

48. Yang H, Zhang N, Li X, Chen J, Cai B. Structure–activity relationships of diverse annonaceous acetogenins against human tumor cells. *Bioorg. Med. Chem. Lett.* 2009;19(8):2199-202.

49. Jagtap NS, Nalamwar VP, Khadabadi SS, Pratapwar AS. Phytochemical and Pharmacological Profile of *Annona squamosa* Linn: A Review. *Res J Pharmac phytochem.* 2009;1(3):139.

50. Mehra BK, Hiradhar PK. Effect of crude acetone extract of seeds of *Annona squamosa* Linn. (Family: Annonaceae) on possible control potential against larvae of *Culex quinquefasciatus* Say. *J Entomol Res.* 2000;24(2):141–46.

51. Malek MA, Wilkins RM. Effects of *Annona squamosa* L. seed oil on the pupae and adults of *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *Bangladesh. J Entomol.* 1994;4:25-31.

52. González MC, Lavaud C, Gallardo T, Zafra-pollo MC, Cortes D. New Method for the determination of the absolute Stereochemistry in-of Antitumoral Annonaceous Acetogenins. *Tetrahedron.* 1998;54(22):6079-88.

53. Zafra-Polo MC, Gonzales MC, Estornell E, Sahpaz S, Cortes D. Acetogenins from Annonaceae: inhibitors of mitochondrial complex 1. *Phytochem.* 1996;42:253-71.

54. Johnson HA, Oberlies NH, Alali FQ, McLaughlin JL. Thwarting resistance: annonaceous acetogenins as new pesticidal and antitumor agents, *in*: Cutler, S.J. and Cutler, H.G. [Eds.] Biologically Active Natural Products. Pharmaceuticals. CRC Press, Washington, DC. 2000;173-83.
55. Asmanizar AD, Idris AB. Evaluation of *Jatropha curcas* and *Annona muricata* seed crude extracts against *Sitophilus zeamais* infesting stored rice. *J Entomol.* 2012; 9:13-22.
56. Kotkar HM, Mendki PS, Sadan SVG, Jha SR, Upasani SM, Maheshwari VL. Antimicrobial and pesticidal activity of partially purified flavonoids of *Annona squamosa*. *Pest Manag Sci.* 2002;58: 33-37.
57. Leatemia, J.A., Isman, M.B. Insecticidal activity of crude seed extracts of *Annona* spp. (Annonaceae), *Lansium domesticum* and *Sandoricum koetjape* (Meliaceae) against lepidopteran larvae. *Phytoparasitica.* 2004;32:32-37.
58. Morsy TA, Rahem MM, Allam KA. Control of *Musca domestica* third instar larvae by the latex of *Calotropis procera* (Family: Asclepiadaceae). *J Egypt Soc of Parasitol.* 2001;31:107-10.
59. Sukontason KL, Boonchu N, Sukontason K, Choochote W. Effects of eucalyptol on housefly (Diptera: Muscidae) and blow fly (Diptera: Calliphoridae). *Rev Inst Med Trop S Paulo.* 2004;46: 97-101.
60. Abdel Halim AS, Morsy TA. Efficacy of *Trigonella foenum-graecum* (fenugreek) on third stage larvae and adult fecundity of *Musca domestica*. *J Egypt Soc Parasitol.* 2006; 36: 329-34.
61. Abdel Halim AS, Morsy TA. The insecticidal activity of *Eucalyptus globulus* oil on the development of *Musca domestica* third stage larvae. *J Egypt Soc Parasitol.* 2005; 35: 631-36.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic