

Genetic and Environmental Attributes of Chronic Pancreatitis

Shweta Singh, Gourdas Choudhuri* and Sarita Agarwal#

Department of Genetics, *Department of Gastroenterology, Sanjay Gandhi Post
Graduate Institute of Medical Sciences, Lucknow-226014; #Corresponding author:
E-mail: saritasgpgi@gmail.com,

ABSTRACT

Chronic pancreatitis (CP) is a chronic progressive inflammatory disease of the pancreas. Tropical chronic pancreatitis is a severe form of CP with multifactorial aetiology. In addition to the growing evidence of attributes of genetic factors, the environmental factors may also play crucial roles in the progression and pathogenesis of CP as these factors may interact with the genetic components of individuals and cause aberrations. Oxidative stress related genes may modify the effects of ambient tobacco smoking and alcohol on CP. This review presents an updated account of the contribution of gene-environment interactions on the onset and progression of CP.

Keywords: Environmental factors, chronic pancreatitis, oxidative stress, smoking, alcohol

Running Title: Factors influencing chronic pancreatitis

1. INTRODUCTION

25 CP is a progressive inflammatory disorder ultimately leading to irreversible structural
26 changes resulting in functional impairment of exocrine and/or endocrine physiology (1, 2).
27 The existing reports indicate that despite the involvement of some genetic factors, the
28 environmental factors also influence the onset as well as progression of the disease. Alcohol
29 is recognised as the major cause of CP. In developed countries, 60–70% of patients with CP
30 have a long history of heavy alcohol consumption; whereas smoking has been described as
31 an additional risk factor (3-4). Since 1982 (5) smoking has been recognized as one of the
32 major risk factors for CP, particularly in alcohol-related CP compared to CP arising from
33 other causes (6). Alcohol is used by a far greater number of people than those who actually
34 go on to develop CP, indicating variable genetic susceptibility.

35 Environmental factors such as cigarette smoking, alcohol, high-fat diets, fossil fuel
36 products and high temperature cooked meat sources may expose the human pancreas to
37 toxins released by these factors (7,8). Induction of phase I xenobiotics metabolising enzymes
38 in the pancreas, principally cytochrome monooxygenases in response to a number of these
39 agents, have been reported to produce cytotoxic lipophilic intermediates that can potentially
40 overcome their abilities to detoxify such products (9). Initially in 1994, Cavallini and
41 coworkers had described the effect of tobacco and alcohol on pancreatic lithogenesis during
42 the course of CP. They found that medium to heavy smokers had a significantly increased
43 risk of developing calcifications in pancreas but found no effects in those using alcohol (10).

44 The results of recent studies suggest that CP has a strong genetic basis. The
45 increasing knowledge of gene-environment interaction has provided new insight in the
46 pathogenesis of CP (11). Tobacco smokings are known to contain free radicals and to
47 induce their direct formation at the tissue level causing damage of cell membranes, proteins,
48 DNA, chronic tissue inflammation and their remodelling in the long run (12,13). Upon
49 exposure, different protein systems including those scavenging reactive oxygen species
50 (ROS) are up-regulated, and the level of response is influenced by variation in expressions

51 of underlying genes. Likewise, polymorphisms in oxidative stress related candidate genes
52 like glutathione s-transferases (GSTs), microsomal epoxide hydroxylase or heme-
53 oxygenase 1 , have been associated with decline in CP function. The other environmental
54 factors such as ethanol, smoking and some putative etiologic factors may contribute to
55 chronic pancreatic injury. In this review, an endeavour has been made to present an updated
56 account of the contribution of gene-environment interactions on the onset, progression and
57 severity of chronic pancreatitis.



58 **2. ALCOHOL CONSUMPTION MAY ADVERSELY AFFECT THE DIGESTIVE SYSTEM**

59 **AND CAUSE PANCREATITIS:** Alcohol abuse was proposed as a causative agent of
60 pancreatitis in 1878 by Freidrich (14). In 1970s Sarles proposed that alcohol causes
61 pancreatitis via precipitation of the secreted proteins (protein plug formation) within small
62 pancreatic ducts leading to acinar atrophy and fibrosis (14, 15, 16). Some animal
63 experiments have shown that chronic alcohol administration produces changes in the acinar
64 cells. It has been shown that the pancreatic content of the digestive enzymes such as
65 trypsinogen, chymotrypsinogen, lipase and the lysosomal enzyme such as cathepsin B is
66 increased in alcohol fed rats (17).The risk for developing pancreatic disease occurs when
67 there is a daily consumption of greater than 100 g of ethanol for men and 80 g for women for
68 more than five years. Therefore, there is variability in the susceptibility of males and females
69 to the alcohol in disease progression.

70 **3. SMOKING INDUCED GENERATION OF FREE RADICALS AND OXIDATIVE STRESS**

71 **DURING CP:** Smoking has been proved to be independently associated with CP (18). The
72 pancreatic disease is a complex disorder resulting from multiple defects, which, when
73 combined, lead to failure of control systems and metabolic homeostasis. The developmental
74 factors like protein, genes and other pathways that govern interaction between
75 environmental factors and their impacts on the cellular level lead to make the pathogenesis
76 more complicated. The central role of free radicals in tobacco smoke mediated

77 carcinogenesis and oxidative stress has been established by a series of studies (19-20). An
78 important finding by Pryor and co-workers was that cigarette tar has high concentrations of
79 stable free radicals, identified as a semiquinone (QH•) and carbon-centered radicals (-C•) as
80 detected by **electron paramagnetic resonance** (EPR) (21). The most interesting is a
81 quinone/semiquinone/hydroquinone (Q/QH•/QH₂) system in the tar polymeric matrix (22).
82 The stable free radicals were identified as o- and p-benzosemiquinone radicals and their role
83 in the DNA damage through the formation of HO• were detected by EPR spin-trapping (23).
84 The following mechanisms have been identified: the QH• radicals reduce O₂ into O₂•-, which
85 can dismutate to form H₂O₂ and then with Fe²⁺ (cigarette tar itself contains high
86 concentrations of iron) can generate highly oxidizing hydroxyl radicals through the Fenton
87 reaction shown as following:



90 **4. FREE RADICALS ADVERSELY INFLUENCE MULTIPLE FACTORS IN PANCREATIC**

91 **CELLS:** The reactions of free radicals especially OH• with these **membrane** constituents
92 lead to lipid peroxidation within the membranes followed by disintegration of the cells and
93 ultimately to cell death. ROS can also disrupt mitochondrial membrane potential (øm),
94 leading to release of cytochrome c and subsequent DNA fragmentation due to H₂O₂-induced
95 mitochondrial damage (24). In other pathways, switching on proinflammatory cascades not
96 only exerts direct effects on the pancreatic cells, but also initiates the migration, adhesion,
97 and infiltration of inflammatory cells into the exocrine pancreas. Genes to be involved in
98 migration and adhesion process are chemokines. **The intercellular adhesion molecules** are
99 under the regulation of the redox-sensitive kinases or transcription factors such as
100 MAPKs, **(Mitogen-activated protein kinase)**, NF-κB **(Nuclear factor kappa-light chain**
101 **enhancer of activated B cells)** and **activator protein-1** (25, 26, 27) .So the mechanism by

which smoking contributes to pancreatic injury or by which smoking accelerates the pancreatic inflammatory process, is associated to the activation of multiple signal transduction pathways due to nicotine exposure explained in figure.1. This event results in high levels of intracellular calcium release and may be responsible for cell cytotoxicity and cell injury (28).

5. SMOKING AND DEVELOPMENT OF ACUTE AND CP: The large effects of tobacco smoking were caused by long-term cumulative exposure in current smokers as well as the remaining effects in former smokers, especially in men. The recent advances in basic research have revealed that stimulus or upstream signals regulating the ROS/RNS-generating enzymes might also play a role in the pathogenesis of pancreatic inflammation, thus opening a possible new therapeutic avenue against these diseases (29). Oxidative stress plays a critical role in both acute and chronic pancreatic inflammation. Generation of ROS/RNS from numerous enzymatic systems, including xanthine oxidase, nitric oxide synthase, CYP2E1, and NADPH(Nicotinamide adenine dinucleotide phosphate) oxidase, not only directly oxidizes a wide range of biomolecules, but also switches on several stress-activated pathways like MAPK, NF-kb, cytokines and other pro-inflammatory pathways (30, 31). Apart from the epidemiologic evidence of an association between smoking and development of acute and CP, a biological effect of smoking seems plausible because both animal studies and human studies have demonstrated changes into the pancreas and in pancreatic functioning after exposure to tobacco smoke.

6. INTERACTION OF CP WITH VARIOUS OTHER ENVIRONMENTAL FACTORS: A review article presented by Morinville and Whitcomb (32) has indicated that some environmental factors can cause injury to the pancreas. The different environmental factors, metabolic and oxidative stress can lead to pancreatic injury via premature activation of zymogens within the pancreas. Certain environmental factors such as infections by bacteria, viruses and parasites, medications such as aminosalicylic acid derivatives, anticonvulsants,

antineoplastics, antimicrobials, estrogens, diuretics and immunomodulators etc.; toxins such as alcohol, organophosphates, scorpion bite toxins, methylene chloride; or trauma including blunt trauma (motorcycle accidents etc.) and child abuse; and metabolic factors such as biochemical disturbance including hyperglycemia, hypertriglyceridemia, inborn errors of metabolism, diabetic ketoacidosis and uremia can cause damage to the pancreas to an irreparable extent. However, the reports available indicate that environmental factors alone do not account for a significant percentage of acute pancreatitis episodes (32). In addition, other factors such as obesity, diets containing high animal protein, fat and antioxidant deficiencies etc. may contribute to pancreatitis. The diets containing highly processed or red meat or deficient in fruits and vegetables or phytochemicals (lycopene and flavonols) may also act as risk factors (33).

7. ASSOCIATION OF GENETIC FACTORS IN CP: The sequencing of numerous genes have suggested that the genetic polymorphisms may act as predisposing risk factors for pancreatitis disorders. It has been observed that not all individuals exposed to a known environmental factor develop pancreatitis, and also that not all individuals possessing a genetic mutation associated with pancreatitis express it phenotypically. It suggests involvement of a complex gene-gene and gene-environmental interactions (32). The mutations in the genes such as **PRSS1** with chromosomal location at 7q35 (which encodes cationic trypsinogen) with mutations such as R122H, N29I, A16V, K23R, E79K etc.; **SPINK1** located at 5q32 (which encodes serine protease inhibitor Kazal type 1) contain mutations N34S, P55S, -253C alleles etc. and cystic fibrosis transmembrane conductance regulator (**CFTR**) having chromosomal location at 7q31.2 (which encodes cystic fibrosis transmembrane conductance regulator) with multiple mutations are associated with chronic pancreatitis. The mutations in **PRSS1** and **CFTR** manifest within the acinar cells and ductular system, respectively. **SPINK1** mutations appear to have a disease-modifying effect (eg, it has a synergistic effect in the presence of **CFTR**) (32, 33). Mutations causing loss of

function in other genes such as chymotrypsinogen C (**CTRC**) and calcium-sensing receptor (**CASR**) are also reported to contribute in chronic pancreatitis (34). The detailed roles of these factors are described in couple of recently published reviews (35, 36).

8. STONE FORMATION, DUCT OBSTRUCTION AND NECROSIS-FIBROSIS IN CP

The stone formation and duct obstruction due to excessive alcohol consumption has been found to increase the lithogenicity of pancreatic juice which finally culminates into stone formation. Upon chronic contact with the stones, the epithelial cells of pancreatic duct produce ulceration and scarring. The chronic obstruction of the acini leads to atrophy and fibrosis (37). The underlying cellular mechanisms of pancreatic fibrogenesis involve pancreatic stellate cells. The alcohol and oxidative stress are the key players which stimulate the stellate cells which upon activation migrate to the periacinar areas and deposit collagen and fibronectin. In addition, some specific cytokines emitted during the inflammatory phase of acute pancreatitis have been shown to stimulate the pancreatic stellate cells (38). Also, the transforming growth factor beta 1 has been shown to mediate onset of pancreatic fibrosis and its pathogenesis (37, 38).

9. CONCLUSION

Among the numerous environmental factors reported so far playing crucial role in influencing the health of humans, consumption of alcohol and smoking has been found not only to generate toxic free radicals but also adversely influence the oxidative pathway and transcription processes into pancreatic cells resulting into further aggravation of CP. These factors may exert synergistic effect with those of genetic factors making the disease conditions even worse as they interact with the genetic components of individuals.

177 **ACKNOWLEDGEMENTS:** Authors are grateful to Sanjay Gandhi Postgraduate Institute of
178 Medical Sciences (SGPGIMS), Lucknow-India for providing infrastructural facilities. SS is
179 thankful to DST for financial support in the form of a INSPIRE fellowship.

180 REFERENCES

- 181 1. Tandon RK, Sato N, Garg PK. Chronic pancreatitis: Asia Pacific consensus report. J
182 Gastroenterol Hepatol 2002; 17:508e18.
- 183 2. Steer ML, Waxman I, Freedman S. Chronic pancreatitis. N Engl J Med. 1995; 332:1482-
184 490.
- 185 3. Talamini G, Bassi C, Falconi M, Sartori N, Salvia R, Rigo L, Castagnini A, Di Francesco V,
186 Frulloni L, Bovo P, Vaona B, Angelini G, Vantini I, Cavallini G, Pederzoli P. Alcohol and
187 smoking as risk factors in chronic pancreatitis and pancreatic cancer. Dig Dis Sci. 1999;
188 44:1303–311.
- 189 4. Chowdhury P, Rayford PL. Smoking and pancreatic disorders. Eur J Gastroenterol
190 Hepatol. 2000; 12:869–77.
- 191 5. Yen S, Hsieh CC, MacMahon B. Consumption of alcohol and tobacco and other risk
192 factors for pancreatitis. Am J Epidemiol. 1982; 116: 407–14.
- 193 6. Morton C, Klatsky AL, Udaltsova N: Smoking, coffee, and pancreatitis. Am J
194 Gastroenterol. 2004; 99: 731–38.
- 195 7. Maisonneuve P, Lowenfals AB. Chronic pancreatitis and pancreatic cancer. Dig. Dis.2002;
196 20:32-37.
- 197 8. Bourliere M, Barthet M, Berthezene P, Durbec JP, Sarles H. Is tobacco a risk factor for
198 chronic pancreatitis and alcoholic cirrhosis? Gut 1991; 32:1392-395.
- 199 9. Rahman SH, Nanny C, Ibrahim K. Genetic polymorphisms of GSTT1, GSTM1, GSTP1,
200 MnSOD, and catalase in nonhereditary chronic pancreatitis: evidence of xenobiotic stress
201 and impaired antioxidant capacity. Dig Dis Sci. 2005; 50: 1376-383.

- 202 10. Cavallini G, Talamini G, Vaona B, Bovo P, Filippini M, Rigo L, Angelini G, Vantini I, Riela
203 A, Frulloni L. Effect of alcohol and smoking on pancreatic lithogenesis in the course of
204 chronic pancreatitis. *Pancreas* 1994; 9: 42–46.
- 205 11. Muddana V, Lamb J, Greer JB, Elinoff B, Hawes RH, Cotton PB, et al. Association
206 between calcium sensing receptor gene polymorphisms and chronic pancreatitis in a US
207 population: role of serine protease inhibitor Kazal 1type and alcohol. *World J Gastroenterol*
208 2008; 14: 4486–491.
- 209 12. Rahman I, Adcock IM. Oxidative stress and redox regulation of lung inflammation in
210 COPD. *Eur Resp J*. 2006; 28: 219–42.
- 211 13. MacNee W. Pulmonary and Systemic Oxidant/Antioxidant Imbalance in Chronic
212 Obstructive Pulmonary Disease. *Proc Am Thoracic Soc*. 2005; 2: 50–60.
- 213 14. Friedrich N. Disease of the Pancreas. New York: William Wood, 1878
- 214 15. Sarles H. Alcoholism and pancreatitis. *Scand J Gastroenterol* 1971; 6:193-98.
- 215 16. Nakamura K, Sarles H, Payan H. Three dimensional reconstructions of the pancreatic
216 ducts in chronic pancreatitis. *Gastroenterology* 1972; 62: 942-49.
- 217 17. Apte MV, Wilson JS, Korsten MA, McCaughan GW, Haber PS, Pirola RC. Effects of
218 ethanol and protein deficiency on pancreatic digestive and lysosomal enzymes. *Gut*.
219 1995;36: 287-93.
- 220 18. Yadav D, Whitcomb DC. The role of alcohol and smoking in pancreatitis. *Nat Rev*
221 *Gastroenterol. Hepatol*. 2010; 7: 131-45.
- 222 19. Leaderson P, Tagesson C. Cigarette smoke-induced DNA damage in cultured human
223 cells: role of hydroquinone and catechol in the formation of oxidative DNA-adduct, 8-
224 hydroxydeoxyguanosine. *Chem.-Biol Interact*. 1990; 75: 71–81.
- 225 20. Asami S, Manabe H, Miyake J, Tsurudone Y, Hirano T. Cigarette smoking induces an
226 increase in oxidative DNA damage, 8-hydroxydeoxyguanosine, in a central site of human
227 lung. *Carcinogenesis*.1997; 18: 1763–766.

- 228 21. Huang MF, Lin WL, Ma YC. A study of reactive oxygen species in mainstream of
229 cigarette. *Indoor Air*. 2005; 15: 135–40.
- 230 22. Pryor WA. Biological effects of cigarette smoke, wood smoke, and the smoke from
231 plastics: the use of Electron spin resonance. *Free Radic Biol Med*. 1992; 13: 659–76.
- 232 23. Pryor WA, Hales BJ, Premovic PI, Church DF. The radicals in cigarette tar: their nature
233 and suggested physiological implications. *Science* 1983; 220:425–27.
- 234 24. Satoh T, Enokido Y, Aoshima H, Uchiyama Y, Hatanaka H. Changes in mitochondrial
235 membrane potential during oxidative stress-induced apoptosis in PC12 cells.
236 *J Neurosci Res*. 1997; 50: 413–20.
- 237 25. Chipitsyna G, Gong Q, Gray CF, Haroon Y, Kamer E, Arafat HA. Induction of monocyte
238 chemoattractant protein- 1 expression by angiotensin II in the pancreatic islets and beta-
239 cells. *Endocrinology* 2007; 148: 2198–208.
- 240 26. Deng Y, Ren X, Yang L, Lin Y, Wu X. A JNK-dependent pathway is required for TNF
241 alpha-induced apoptosis. *Cell* 2003; 115: 61–70.
- 242 27. Zaninovic V, Gukovskaya AS, Gukovsky I, Mouria M, Pandol SJ. Cerulein upregulates
243 ICAM-1 in pancreatic acinar cells, which mediate neutrophil adhesion to these cells. *Am J*
244 *Physiol Gastrointest Liver Physiol*. 2000; 279: G666–G676.
- 245 28. Chowdhury P, MacLeod S, Udupa KB, et al. Pathophysiological effects of nicotine on the
246 pancreas: An update. *Exp Biol Med*. 2002; 227:445–54.
- 247 29. Chvanov M, Petersen OH, Tepikin A. Free radicals and the pancreatic acinar cells: role
248 in physiology and pathology. *Phil Trans R Soc Lond B Biol Sci*. 2005; 360:2273–284.
- 249 30. Burim RV, Canalle R, Martinelli Ade L, Takahashi CS. Polymorphisms in glutathione S-
250 transferases GSTM1, GSTT1 and GSTP1 and cytochromes P450 CYP2E1 and CYP1A1
251 and susceptibility to cirrhosis or pancreatitis in alcoholics. *Mutagenesis* 2004;19: 291–98.

31. Cave AC, Brewer AC, Narayanapanicker A, Ray R, Grieve DJ, Walker S, Shah AM. NADPH oxidases in cardiovascular health and disease. *Antioxid Redox Signal.* 2006; 8: 691–728.
32. Morinville V, Whitcomb DC. Recurrent Acute and Chronic Pancreatitis: Complex Disorders With a Genetic Basis. *Gastroenterol Hepatol.* 2005; 1: 195-205.
33. Nitsche C, Simon P, Weiss FU, Fluhr G, Weber E, Gärtner S, Behn CO, Kraft M, Ringel J, Aghdassi A, Mayerle J, Lerch MM. Environmental risk factors for chronic pancreatitis and pancreatic cancer. *Dig Dis.* 2011; 29: 235-42.
34. Chen J-M, Férec C. Chronic Pancreatitis: Genetics and Pathogenesis. *Annual Review of Genomics and Human Genetics* 2009; 10: 63-87.
35. Chen JM, Férec C. Chronic pancreatitis: genetics and pathogenesis. *Annu. Rev. Genomics Hum. Genet.* 2009; 10:63–87
36. Braganza JM, Lee SH, McCloy RF, McMahon MJ. Chronic pancreatitis. *Lancet.* 2011; 377: 1184-197.
37. Stevens T, Conwell DL, Zuccaro G. Pathogenesis of chronic pancreatitis: An evidence-based review of past theories and recent developments. *Am J Gastroenterol.* 99: 2004; 2256-270.
38. Affronti J. Chronic Pancreatitis and Exocrine Insufficiency. *Primary Care-Clinics in Office Practice*, 2011; 38: 515–37.




Fig.1. Direct and indirect effects of oxygen free radicals