

Rootstocks influence the Growth, Biochemical Contents and Disease Incidence in Thompson Seedless Grapevines

Ramhari G. Somkuwar*, Satisha Jogaiah, Sanjay D. Sawant, Pravin B. Taware, Devanand D. Bondage and Perna Itrotwar

National Research Centre for Grapes, P.O. Box 03, Manjri Farm Post, Pune 412307 India.

ABSTRACT

The variations in growth, powdery mildew and anthracnose incidence and biochemical changes in Thompson seedless grapevines grafted onto Dog Ridge and 110 R rootstocks in comparison to the own rooted grape vines was investigated. Bud sprouting after pruning was achieved earlier in case of own-rooted vines. However, the vine growth pattern studied in terms of the shoot length, inter nodal length, shoot diameter and leaf area was highest in case of vines grafted onto Dog Ridge. Foliar powdery mildew and anthracnose incidence was highest in vines grafted onto Dog Ridge and own-rooted vines, respectively. The disease incidence was recorded least in vines grafted onto 110-R. The biochemical analysis was also influenced by the rootstocks. The leaves of vines grafted onto Dog Ridge recorded the highest reducing sugars and protein contents and the least total sugars and phenolic contents. The vines grafted onto 110-R topped in total phenolic contents and other phenolic derivatives.

Keywords: grape, rootstocks, canopy architecture, diseases, phenolics.

1. INTRODUCTION

Grape (*Vitis vinifera* L.) is one of the major important commercial fruit crops of India with 1,11,000 ha area with an annual production of 12,35,000 tones [1]. Table grapes occupy more than 90% of the area under grape cultivation in India. Amongst the table grape varieties, Thompson Seedless is a popular cultivar for both fresh consumption (local and export market) and raisin making. Traditional grape cultivation in India involves growing commercial varieties of grapes on own roots. Due to increasing problems of soil salinity, bad quality water and drought, the decline in the productivity of own rooted vines in Maharashtra and neighboring states has been reported. Owing to the adverse conditions use of rootstock for the establishment of grape vineyards became important. As a result, the use of rootstock increased for enhancing grape production in India. The primary use of rootstocks is for pest resistance [2, 3]. The major functions of the grapevine root system are vine water relations, the uptake and translocation of nutrients, the synthesis and metabolism of plant growth substances and the storage of carbohydrates [4]. Reynolds and Wardle [5] outlined seven major criteria for choosing rootstocks in the order of importance. Numerous reports have also proved that rootstock affect vine growth, yield and fruit quality. These effects take place due to the interaction between environmental factors and the physiology of scion and rootstock cultivars. Grape rootstocks have a primary effect on vine size measured in terms of pruning weights, as indicated by [2, 3, 6, 7]. An increase in vine size when the canopy length

*Author for correspondence: Phone +912026956060; Fax +912026956099

Email: rgsgrapes@gmail.com

is fixed, results in crowding of shoots and internal canopy shading (8). Most secondary effects of rootstocks are mediated through their influence on vine size and internal canopy shading. The difference in performance of vines grafted on rootstock and own rooted vines is experienced by the grape growers. In the tropical parts of the country where the grapes are being grown for table as well as wine purpose, canopy development is an important aspect for the production of quality grapes and also to minimize the disease incidence.

For sustainable viticulture, it is important to know the interactions among rootstocks, different soil characters and scion productivity [9]. A rootstock found to be beneficial for one cultivar may not be universally advantageous for others, as the interaction of stock and scion influences the vine performance more than the stock or scion alone [10]. With increased awareness about the use of rootstocks in overcoming the adverse effects of drought and salinity, growers started using Dog Ridge rootstock for the cultivation of Thompson Seedless grapes. However, in the tropical and subtropical climate of India, Dog Ridge induced more vegetative vigour in the scions, which reduced the bud fruitfulness of Thompson Seedless in the long run. In this context, there was an increased demand for an alternative rootstock that would be suitable for Thompson Seedless in the tropical and subtropical climate of the Indian subcontinent. Another grape rootstock i.e., 110-R which is alternative rootstock to the present Dog Ridge [11] was introduced to overcome these problems.

Use of Dog Ridge and 110-R rootstocks is gaining importance in Indian viticulture and the majority of vineyards are being established on these rootstocks to overcome the adverse effects of abiotic stresses such as drought and salinity. In addition, rootstocks are known to influence the growth pattern of scions after grafting and some may even induce resistance to diseases by various physiological and biochemical changes in grafted vines [12]. Grape species and their cultivars differ in their susceptibility to powdery mildew infection [13]. The rootstocks alter the canopy architecture of a grapevine thereby creating favourable or unfavourable microclimate for the development of various diseases. This also alters the biochemical composition of a grapevine as a whole and of final harvest. Considering this, the present study was initiated with the objective to evaluate the changes in growth behaviour, biochemical composition and disease incidence of Thompson Seedless grafted onto Dog Ridge and 110-R rootstocks, with own rooted Thompson Seedless as a control.

2. MATERIALS AND METHODS

The trial was conducted at the farm of National Research Centre for Grapes, Pune during 2008-2010. The grape rootstock Dog Ridge and 110-R along with own-rooted vines of Thompson Seedless were planted during the year 1999 with 3.0 m distance between the rows and 1.8 m between the vines and the grafting of table grape variety Thompson seedless was performed during October 1999. The experimental site is situated in Mid-West Maharashtra at an altitude of 559 m above mean sea level; it lies on 18.32°N latitude and 73.51°E longitudes. The climate in this region is mild to slightly dry. The soil is heavy black with pH 7.75 and EC 0.46 dS m⁻¹. The vines were trained on 'extended Y' system of training with four cordons (H shape – Height = 1.20m from surface, Cross arm width = 0.60m) developed horizontally with vertical shoot positioning. The distance from the fruiting wire to the top of foliage support wire was 0.60 m.

2.1 Growth parameters

The investigation was carried out after foundation pruning on nine year old Thompson Seedless vines grafted onto Dog Ridge and 110-R rootstock and was compared with own rooted Thompson Seedless. The canes were pruned during mid-April leaving only a single visible bud at the base on a cordon. The days taken for sprouting were recorded on day to

**Author for correspondence: Phone +912026956060; Fax +912026956099*

Email: rgsgrapes@gmail.com

day basis by visual observations. The shoot length was measured at 30, 60 and 90 days after pruning (DAP) with the help of measuring tape. Inter nodal length, shoot diameter and leaf area was also measured after 90 DAP with the help of measuring scale, vernier calliper and leaf area meter (CID, Inc), respectively. Days taken for cane maturity were also recorded as the time taken for complete browning of cane from the date of foundation pruning inspecting visually of proper lignifications.

2.2 Disease assessment

Powdery mildew and anthracnose incidence were assessed in-situ by visual observations as the percent number of the leaves and canes infected by each disease at 90 days after pruning. The infection of each disease on leaf as well as on cane was recorded using the 0-5 scale and was then mentioned as percent infection.

2.3 Biochemical analysis of leaves

The total sugar content was measured by hydrolyzing the polysaccharides into simple sugars by acid hydrolysis and estimating the resultant monosaccharides by Anthrone method [14]. Quantity of the carbohydrate was expressed as glucose equivalent as determined from linear regression obtained by plotting absorption against known standard glucose concentration. Reducing sugars were estimated by using Dinitrosalicylic Acid (DNSA) method [15]. The results were expressed as glucose equivalent as determined from linear regression obtained by plotting the absorption against known standard glucose concentration.

Proteins were estimated by using standard procedure as suggested by Lowry et al. [16] method. The blue colour developed by the reduction of the phosphomolybdic-phosphotungstic components in the Folin-Ciocalteu reagent by the amino acids tyrosine and tryptophan present in the protein plus the colour developed by the biuret reaction of the protein with the alkaline cupric tartarate are measured in Lowry's method. The blue colour developed was read at 660nm on UV-visible spectrophotometer (Shimadzu- 1601). The quantity of protein was determined from linear regression obtained by plotting the absorption against known standard BSA (Bovine serum albumin-Fraction V) concentration. The proteins were expressed as mg g^{-1} fresh weight of sample.

Total phenolic content: One gram fresh samples of leaf were homogenized in 80% ethanol repetitively and final volume made up to 10.0ml. The mixtures were sonicated for 15min for complete extraction and centrifuged at 9000rpm for 10min. The supernatants were utilized for analysis of total phenolic contents and other phenolic derivatives. The total phenolic contents were determined by Folin-Ciocalteu method [17], using gallic acid as the standard. The total polyphenolic concentration was calculated from a calibration curve using gallic acid as a standard ($0-10 \text{ mg L}^{-1}$). Data were expressed in mg g^{-1} GAE. Total flavan-3-ol content was estimated using DMACA (*p*-dimethylaminocinnamaldehyde) method [18]. The concentration of total flavan-3-ols was expressed as the CE (mg g^{-1}) of the fresh sample. Flavonol content was estimated by measuring the absorbance of the extract at 360 nm after the addition of 2% HCl [19] and was expressed as the quercetin equivalent (QE mg g^{-1}). Total flavonoids were determined calorimetrically, following the procedures proposed by Kim et al. [20]. The amount of total flavonoids was expressed as the catechin equivalent (CE, mg g^{-1}).

2.4 Statistical Analysis

The experiment was conducted in randomized block design consisting of three treatment grafted onto Dog Ridge, 110R and own root vines of Thompson Seedless. Each treatment

*Author for correspondence: Phone +912026956060; Fax +912026956099

Email: rgsgrapes@gmail.com

consisted of 20 plants and was replicated five times totaling 45 plants under each treatment. The data was analysed statistically using Sigma Stat version 3.5.

3. RESULTS AND DISCUSSION

The results on days taken for sprouting (Table 1) revealed late sprouting in grafted Thompson Seedless vines (13-14 days) after pruning while the early sprouting was reported in own-rooted vines (10 days) after pruning. However, the shoot growth lead in case of grafted vines and the vines grafted onto Dog Ridge exhibit more vigour. The results on shoot length measurements made it clear that at all the stages of investigation i.e. 30, 60 and 90 DAP, the shoot lengths were in the sequence of own-rooted Thompson seedless vines < vines grafted on 110R < vines grafted onto Dog Ridge. The internode length, shoot diameter and leaf area measurements at 90 DAP (Table 1) exhibited the same pattern. The days taken for cane maturity were significantly more in vines grafted onto Dog Ridge (84.51 days) followed by 110-R (79.80 days) and was least in case of own rooted vines (74.59 days). The effect of rootstocks on days taken for sprouting was highly significant. The results of the present investigation are in accordance with the reports of several workers in the past, who established the influence of rootstocks on bud burst. Satisha et al. [21] reported that Thompson Seedless grafted onto Dog Ridge take significantly more time to sprout after pruning. Prakash and Reddy [22] reported the effect of different rootstocks on bud break in the grape cultivar Anab-e-Shahi, with a significant effect of rootstocks on bud burst. These results are similar to the current findings of delayed bud sprouting on Dog Ridge rootstock. However, Tangolar and Ergenoglu [23] reported that time required for bud break was not significantly affected by rootstocks. The rootstock Dog Ridge is imparting more vigour compared to other rootstock under tropical condition. This has also expressed in terms of increase in shoot length in grafted vines than that of own rooted vines. Sommer et al [24] also reported the increase in shoot length on grafted vines. Satisha et al. [21] while working on performance of Thompson Seedless grafted onto different rootstock also reported increase in shoot length and cane diameter in the vines grafted onto Dog Ridge rootstock compared to own rooted vines. Bica et al. [25] found significant effect of use of rootstock on higher leaf area. The rootstock Dog Ridge is performing better under Indian condition considering the yield and quality requirement. However, 110-R is another addition which is an alternative to present Dog Ridge, looking into the soil and water problem in grape cultivation [11]. The vines grafted onto 110-R rootstock are early to sprout than Dog Ridge grafted vines in addition to uniform sprouts is also achieved. The cane maturity is an indication of proper lignifications in the shoot. The early cane maturity helps to store enough reserve food material in shoot in addition to the healthy vine. The vines grafted onto 110-R rootstock were early to mature indicating proper vine health.

190
191

Table 1. Effect of Dog Ridge and 110-R rootstocks on growth, powdery mildew and anthracnose disease incidence in Thompson Seedless grapes

<i>Parameters</i>	<i>Vines grafted on Dog Ridge</i>	<i>Vines grafted on 110R</i>	<i>Own-rooted Thompson Seedless</i>	<i>Significance</i>
Days taken for sprouting	13.48 ±1.12 a	12.64 ±0.93 a	10.08 ±0.59 b	**
Shoot length at 30 DAP (cm)	39.71 ±0.74 a	35.03 ±1.27 b	31.44 ±1.66 c	**
Shoot length at 60 DAP (cm)	72.01 ±1.78 a	66.50 ±1.74 b	59.30 ±1.27 c	**
Shoot length at 90 DAP (cm)	104.21 ±2.78 a	93.34 ±2.40 b	85.01 ±2.25 c	**
Internode length (cm)	6.60 ±0.20 a	5.28 ±0.22 b	4.90 ±0.24 c	**
Shoot diameter (mm)	8.98 ±0.25 a	8.00 ±0.16 b	6.71 ±0.12 c	**
Leaf area (cm ²)	8411.14 ±424.29 a	6836.35 ±348.92 b	4727.22 ±252.50 c	**
Days required for cane maturity	84.51 ±0.86 a	79.80 ±0.71 b	74.59 ±1.02 c	**
Foliar powdery mildew incidence (%)	3.19 ±0.62 a	1.57 ±0.41 b	2.08 ±0.65 b	*
Foliar anthracnose incidence (%)	1.30 ±0.54 a	1.14 ±1.22 a	1.96 ±0.55 a	ns
Anthracnose incidence on canes (%)	0.80 ±0.12 c	1.20 ±0.13 b	2.00 ±0.39 a	**


192
193
194
195
196

The data are the means of five replicates ±standard deviation. ‘’, ‘**’ and ‘ns’ represent significance P<0.05, p<0.01 and non-significance, respectively by one-way-ANOVA statistics. Values followed by different letters differ significantly by DMRT at P=0.05.*

197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220

The biochemical composition of the leaves at 90 DAP (Fig. 1) revealed that the total sugars were significantly more in case of own-rooted and 110-R grafted vines than the vines grafted onto Dog Ridge. Reducing sugars and protein contents were significantly higher in case of the vines grafted onto Dog Ridge. The results on phenolic compounds in the leaves (Fig. 2) indicated that the total phenols, flavan-3-ols, flavonols and flavonoids contents were significantly high in the vines grafted onto 110-R rootstock. Own-rooted vines were at second position and the vines grafted onto Dog Ridge had the least phenolic contents. The results on biochemical composition exhibited significant difference due to grafting of Thompson Seedless onto Dog Ridge and 110-R rootstocks. This might be due to the alterations in the growth pattern of the vines by rootstocks as well as the differences in their uptake of nutrients and water from soil solution, as root development patterns vary with the rootstocks. Most secondary effects of rootstocks are mediated through their influence on vine size and internal canopy shading. Previous research has found a relationship between canopy structure and sunlight exposure and subsequent phenolic contents [26]. In addition, a relationship between variations in vine growth and differences in total phenolic levels (measured as absorbance at 280 nm) has been reported [27, 28]. Influence of rootstocks on changing pattern of phenolic compounds in Thompson Seedless reported by [29] confirms the results of the present investigation. Grape berries act as typical sink organ which rely on the use of available carbohydrate resource produced by photosynthesis to support their growth and development. In addition to the number of factors assessing to monitor ripening, berry sugar concentration is mainly due to water loss from the berries at maturity [30]. The effect of rootstock on fruit composition has been reported earlier by several workers, especially in wine grapes, with a close link between fruit quality and wine prepared from those grapes. Soluble solids, organic acids, pH, phenolic and anthocyanins, monoterpenes

221 and other components [31] are fruit quality parameters that affect the quality of wine. Hale
 222 and Brien [32] first investigated the influence of Salt Creek rootstock on the composition and
 223 quality of Shiraz grapes and wine. Their results showed that grafted Shiraz had larger
 224 berries with lower soluble solids and higher pH, titratable acidity, malate and potassium.
 225 Kubota et al. [33] on his study on Fujimori grapes grafted onto seven different rootstocks
 226 found that the glucose and fructose content of the pulp was higher in berries grafted onto
 227 3309 C, 3306C and 8B. However, some researchers reported no significant effects of
 228 rootstock on fruit composition in grapes. Reynolds and Wardle [34] studied nine grape
 229 cultivars and four rootstocks (3309C, 5BB, 5C, SO4) and they reported non-significant
 230 differences in titratable acidity as well as pH.



231
 232 **Fig.1. Effect of Dog Ridge and 110-R rootstocks on biochemical composition of**
 233 **Thompson Seedless grapes as compared to own-rooted vines**
 234

235 The data on disease incidence on Thompson Seedless grapevines (Table 1) revealed that
 236 the powdery mildew incidence in case of vines grafted onto Dog Ridge (3.19%) was
 237 significantly higher than the vines grafted onto 110-R (1.57%) and own-rooted Thompson
 238 Seedless vines (2.08%). There was no significant difference in anthracnose infection on
 239 leaves. However, the anthracnose infection of the canes was the highest in own rooted
 240 (2.00%) Thompson Seedless followed by vines grafted onto 110-R (1.20%) and the least in
 241 case of the vines grafted onto Dog Ridge (0.80%). Rootstocks not only perform a critical
 242 function in water and nutrient assimilation [35], but also provide increased tolerance to insect
 243 pest and diseases of grapes [36]. In the present investigation, use of Dog Ridge rootstock
 244 found increase in foliar powdery mildew disease susceptibility. However, the vines grafted
 245 onto 110-R rootstock exhibited significantly low powdery mildew incidence. The finding of the
 246 present investigation confirms the results of [29] who reported less powdery mildew infection
 247 in vines grafted onto 110-R rootstocks. The species of genus *Vitis* that contain high
 248 polyphenols content (*V. rotundifolia*) are more resistant to infection caused by *Uncinula*
 249 *neator*, and when infected they produce larger amounts of polyphenols than varieties that
 250 are more susceptible [37]. Rootstocks may influence the biochemical composition of the
 251 scion leaves grafted onto them, which in turn affects the degree of resistance or
 252 susceptibility to disease [29]. Variation in the incidence of disease in Thompson Seedless
 253 grafted onto different rootstocks may have been due to the genetic makeup of the rootstock,
 254 which have indirectly contributed to the synthesis of various secondary metabolites. The
 255 results on reduced anthracnose incidence on Thompson Seedless grafted vines were similar

*Author for correspondence: Phone +912026956060; Fax +912026956099
 Email: rgsgrapes@gmail.com

256 with the finding of Ren and Lu [36] where they reported increased resistance of Florida
 257 hybrid grapes to anthracnose disease. Pospisilova [38] rated the severity of mildew on both
 258 the fruit and leaves for 926 cultivars of *V. vinifera* and found that disease severity on the fruit
 259 generally corresponded to that on the leaves for a given cultivar.
 260




Fig.2. Effect of Dog Ridge and 110-R rootstocks on phenolic contents of Thompson Seedless leaves as compared to own-rooted vines

261
 262
 263
 264
 265
 266 The correlation coefficients (Table 2) between the growth parameters, biochemical
 267 composition and disease incidence of the vines revealed that the days taken for sprouting
 268 after pruning had no significant impact on further growth, biochemical composition as well as
 269 disease incidence of the vines. The vine vigour measured in terms of shoot length, inter
 270 nodal length and leaf area were significantly inter-related and had positive correlation with
 271 foliar powdery mildew incidence and anthracnose infection of canes with mixed influence on
 272 biochemical composition. Disease incidence of Thompson Seedless vines had no significant
 273 correlation with biochemical composition of the leaves at 90 DAP.
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289

290
291

Table 2. Correlation between vine growth, disease incidence and biochemical composition in own-rooted and grafted Thompson Seedless grape

<i>Parameters</i>	<i>Shoot length</i>	<i>Inter nodal length</i>	<i>Leaf area</i>	<i>Foliar PM</i>	<i>Foliar Anthracnose</i>	<i>Cane Anthracnose</i>	<i>Total phenolics</i>	<i>Reducing sugar</i>	<i>Total sugars</i>	<i>Protein</i>
Days taken for sprouting	-0.19ns	-0.13ns	-0.29ns	-0.19ns	0.15ns	0.41ns	0.10ns	-0.05ns	0.31 ns	0.03ns
Shoot length	0.89**	0.89**	0.88**	0.70**	-0.29ns	-0.84**	-0.72**	0.57*	-0.86**	0.26ns
Inter nodal length			0.86**	0.80**	-0.24ns	-0.79**	-0.83**	0.68**	-0.74**	0.50*
Foliar PM incidence					-0.39ns	-0.63**	-0.66**	0.79**	-0.41ns	0.47*
Foliar Anthracnose						0.33ns	-0.02ns	-0.38ns	0.05 ns	0.10ns
Cane Anthracnose							0.50*	-0.51*	0.80**	-0.08ns
Total phenolic								-0.38ns	0.69**	-0.75**
Reducing sugars									-0.19ns	0.38 ns
Total sugars										-0.10ns

292
293
294
295
296

‘*’, ‘**’ and ‘ns’ represent significance of the correlation coefficients at $P < 0.05$, $p < 0.01$ and non-significance, respectively.

297
298

4. CONCLUSION

299
300
301
302
303
304
305

The rootstock Dog Ridge and 110-R were found to alter the growth of the Thompson Seedless grapevines as compared to the own-rooted vines. The rootstock Dog Ridge imparted more vigour to the scion followed by 110-R. Although the own rooted vines sprouted earlier, they were less vigorous. The results on biochemical composition, powdery mildew and anthracnose disease incidence of vines were also influenced due to use of rootstocks. These traits may have significance in selection of rootstock considering the local soil and climatic conditions.

306
307

ACKNOWLEDGEMENTS

308
309
310

The authors are thankful to the Director, NRC Grapes, Pune for providing the facilities to carry out the research and motivation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

AUTHORS' CONTRIBUTIONS

The research work was carried out at the Research and developmental plot of NRC Grapes, Pune. Authors RGS, SJ and SDS designed the experiment, wrote the protocol and recorded the observations. Authors PBT and DDB done the field work and collected samples for study and author PI performed biochemical and statistical analysis. Author RGS prepared the manuscript. All authors read and approved the final manuscript.

REFERENCES

1. Anonymous. In: Indian Horticulture Database (Eds.: B. Kumar, N. C. Mistry, B. Singh and C. P. Gandhi). National Horticulture Board, Gurgaon, India: 68-75; 2012.
2. Pongracz DP. Rootstocks for grapevines. David Philip Publisher, Cape Town; 1983.
3. Howell GS. Vitis rootstocks. In: Rom RC and Carlson RF (Eds.). Rootstocks for food crops. Wiley and Sons, New York, pp. 451-474; 1987.
4. Richard D. The grape root system. Hort. Rev. 1983; 5: 127-168.
5. Reynolds AG, Wardle DA. Rootstocks impact vine performance and fruit composition of grapes in British Columbia. Hort. Technol. 2001; 11: 419-427.
6. Carbonneau A, Casteran P. Optimization of vine performance by the lyre training system. In: Lee, T. (ed.) Proc. 6th Aust. Wine Ind. Tech. Conf., Australian Industrial Publishers, Adelaide, Australia, pp. 194-204; 1987.
7. Pouget P. Usefulness of rootstocks for controlling vine vigour and improving wine quality. Acta. Hort. 1987; 206: 109-118.
8. Shaulis NJ. Responses of grapevines and grapes to spacing of and within canopies. In: Webb, D. (ed.) Grape and Wine Centennial Symposium Proceedings, University of California, Davis, pp. 353-360; 1982.
9. Keller M, Kummer M, Vasconcelos MC. Soil nitrogen utilization for growth and gas exchange by grapevines in response to nitrogen supply and rootstock. Aust. J. Grape Wine Res. 2001; 7: 2-11.
10. Hartmann HT, Kester DE, Davies FT. Plant pathogen- principles and practices. Prentice Hall, New Delhi, India; 1993.
11. Somkuwar RG, Satisha J, Ramteke SD. Effect of different rootstocks on fruitfulness in Thompson Seedless grapes (*Vitis vinifera* L.) Journal of Plant Sciences. 2006; 1(2): 176-181.
12. Satisha J, Prakash GS, Murti GSR, Upreti KK. Response of grape genotypes for water deficit: Root shoots growth and endogenous hormones. Indian. Pl. Physiol. 2005; 10: 225-230.
13. Peros JP, Nguyen TH, Troulet C, Michel-Romiti C, Notteghem JL. Assessment of powdery mildew resistance of grape and Erysiphe necator pathogenicity using a laboratory assay. Vitis. 2006; 45: 29-36.
14. Hedge JE, Hofreiter BT. In: Whistler, R. L. and Be Miller, J. N. (eds.) Carbohydrate Chemistry, Academic Press, New York: 17; 1962.
15. Miller GL. Use of dinitrosalicylic acid reagent for determination of reducing sugars. Anal. Chem. 31: 426-428; 1972.
16. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with folin phenol reagent. J. Biol. Chem. 1951; 193:265.

*Author for correspondence: Phone +912026956060; Fax +912026956099

Email: rgsgrapes@gmail.com

- 363 17. Singleton VL, Rossi JA. Colorimetry of total phenolics with phosphomolybdic-
364 phosphotungstic acid reagent. *Am. J. Enol. Vitic.* 1965; 16: 144-158.
- 365 18. Arnous A, Makris DP, Kefalas P. Effect of principle polyphenolic composition in
366 relation to antioxidant characteristics of aged red wines. *J. Agric. Food Chem.* 2001;
367 49: 5736-5742.
- 368 19. Mazza G. Anthocyanins in grape and grape products. *Critical Rev. Food Sci.*
369 *Nutrition.* 1999; 35:341-371.
- 370 20. Kim D, Chun OK, Kim YJ, Moon H, Lee CY. Quantification of polyphenols and their
371 antioxidants capacity in fresh plums. *J. Agric. Food Chem.* 2003; 51: 6509-6515.
- 372 21. Satisha J, Somkuwar RG, Sharma J, Upadhyay AK, Adsule PG. Influence of
373 rootstocks on growth yield and fruit composition of Thompson seedless grapes
374 grown in the Pune region of India. *S. Afr. J. Enol. Vitic.* 2010; 31: 1-8.
- 375 22. Prakash GS, Reddy NN. Effect of different rootstocks on budbreak in grape cv.
376 Anab-e-shahi. *Crop Res.* 1990; 3: 51-55.
- 377 23. Tangolar S, Ergenoglu F. The effects of different rootstocks on the levels of mineral
378 elements in the leaves and the carbohydrate contents of the canes of some early
379 maturing grape cultivars. *Doga, Turk Tarim ve Ormancilik Dergisi.* 1989; 13: 1267-
380 1283.
- 381 24. Sommer KJ, Clingeleffer PR, Olat N. Effect of minimal pruning on grapevine canopy
382 development, physiology and cropping level in both cool and warm climates. *Vitic.*
383 *Enol. Sci.* 1993; 48: 135-139.
- 384 25. Bica D, Gay G, Morando A, Soave E, Bravdo, BA. Effects of rootstock and *Vitis*
385 *vinifera* genotype on photosynthetic parameters. *Acta. Hort.* 2000; 526: 373-379.
- 386 26. Mabrouk H, Sinoquet H. Indices of light microclimate and canopy structure of
387 grapevines determined by 3D digitizing and image analysis. *Aust. J. Grape Wine*
388 *Res.* 1998; 4: 2-13.
- 389 27. Lamb DW, Weedon MM, Bramley RGV. Using remote sensing to predict grape
390 phenolics and colour at harvest in a Cabernet Sauvignon vineyard: Timing
391 observations against vine physiology and optimizing image resolution. *Aust. J.*
392 *Grape Wine Res.* 2004; 10: 46-54.
- 393 28. Cortell JM, Halbleib M, Gallagher AV, Righetti TL, Kennedy JA. Influence of vine
394 vigour on grape (*Vitis vinifera* L. cv. Pinot Noir) and wine proanthocyanidins. *J.*
395 *Agric. Food Chem.* 2005; 53: 5798-5808.
- 396 29. Satisha J, Doshi P, Adsule PG. Influence of rootstocks on changing the pattern of
397 phenolic compounds in Thompson seedless grapes and its relationship to the
398 incidence of powdery mildew. *Turk. J. Agric.* 2008; 32: 1-9.
- 399 30. Ali K, Frederica Maltese, Ana Margarida Forts, Maria Salome Pais, Young Hae Choi
400 and Robert Verpoorte. Monitoring biochemical changes during grape berry
401 development in Portuguese cultivars by NMR spectroscopy. *Food Chemistry.* 2011;
402 124: 1760-1769.
- 403 31. Jackson DI, Lombard PB. Environmental and management practices affecting grape
404 composition and wine quality - a review. *Am. J. Enol. Vitic.* 1993; 44, 409-430.
- 405 32. Hale CR, Brien CJ. Influence of Salt Creek rootstock on composition and quality of
406 Shiraz grapes and wine. *Vitis.* 1978; 17, 139-146.
- 407 33. Kubota N, Li xG, Yasui K. Effects of rootstocks on sugar, organic acid, amino acid,
408 and anthocyanin contents in berries of potted 'Fujiminori' grapes. *J. Japan. Soc.*
409 *Hort. Sci.* 1993; 62, 363-370.
- 410 34. Reynolds AG, Wardle DA, Cliff MA, King M. Impact of training system and vine
411 spacing on vine performance, berry composition, and wine sensory attributes of
412 Riesling. *Am. J. Enol. Vitic.* 2004; 55:96-103.
- 413 35. Rom CR. Coordination of root and shoot growth: roots and rootstocks. In: *Tree Fruit*
414 *Physiology*, K. M. Maib (ed.), pp53-58; 1996.

- 415 36. Ren Z, Lu J. Muscadine rootstock increased the resistance of Florida hybrid bunch
416 grape cv. 'Blanc Du Bois' to pierce's and anthracnose diseases. Proc. Fla. State
417 Hort. Soc. 2002; 115: 108-110.
- 418 37. Dai GH, Andary C, Cosson Mondolot L, Boublas D. Polyphenols and resistance of
419 grape vines to downy mildew. Acta Horticulturae. 1994; 381: 763-766.
- 420 38. Pospisilova D. Susceptibility of cultivars of Vitis vinifera to powdery mildew (Uncinula
421 necator Schw. Burr.). In Grapevine Genetics and Breeding. pp 251-7. Institut
422 National de la Recherche Agronomique. Paris (1978).
- 423

424 **ABBREVIATIONS**

425

426 DNSA - Dinitrosalicylic Acid ; BSA - Bovine serum albumin-Fraction V ; DMACA - *p*-
427 dimethylaminocinnamaldehyde