

Original Research Article

Characterization of the Chemical Properties of some Selected Refined Vegetable Oils Commonly sold in Nigeria.

ABSTRACT

SAMPLE ABSTRACT:

Aims: To ascertain the suitability for consumption of some selected refined vegetable oils commonly found and sold in most of the markets located in Nigeria.

Study design: Five different brands of commonly found vegetable oils in Nigeria markets, were subjected to Chemical analysis via determination of their iodine value, acid value, peroxide value, saponification value and ester value.

Place and Duration of Study: The chemical analysis was carried out in the Chemistry laboratory of the National Open University of Nigeria suited on Ahmadu Bello Way, Victoria Island, Lagos State between May – June, 2014.

Methodology: The iodine value, acid value, Peroxide value and saponification value were determined by standard procedures described by AOAC (1980) while the ester value was determined by baltes, 1964 method.

Results: The analysed chemical parameters shows that the iodine value of the oils are in the range of 21.83 g/100g to 79.95 g/100g, while the Saponification values of the oils are in the range of 235.62 mg/KOH/g to 274.19mg. The acid values and peroxide values of the oils are in the range of 0.39 mgKOH/g to 1.54 mgKOHg and 0.26 meq/kg – 1.17 meq/kg respectively while the ester value range from 206.70 mgKOH/g to 272.65 mgKOH/g.

Conclusion: The result of the study shows that the oils are non drying oils of low saturation, slow to oxidation and rancidity, will remain liquid for a long time and are suitable for consumption

Keywords: *iodine value, acid value, non drying oil, oxidation and low saturation.*[Put four to eight keywords } (Arial, inclined, 10 font, justified)

1. INTRODUCTION

Vegetable oils and fats are composed predominantly of triglycerides, which are long-chain fatty acid esters of glycerol (1,2). The predominant fatty acids present in vegetable oils and fats are saturated and unsaturated compounds with straight aliphatic chains. An even number of other fatty acids may be present in same Vegetable sources, including a small

amount of branched chain, cyclic and odd number straight chain acids (3). The (minor) non ester portion of vegetable oils and fats include; phospholipids (or phosphatides), sterols, vitamins and their precursors. The non-ester portion is usually less than 2% of the total oil. The glycerides themselves contain about 95% fatty acids and 5% glycerol (2). The amount of glycerol is the same in all vegetable oils, hence, the differences on properties of different oils are largely determined by the variations on the fatty acid structure (2,3). Glycerides composed mostly of unsaturated fatty acids are oils (liquids) at room temperature, while those composed mostly of saturated fatty acids are fats (solid) at room temperature (2). An important feature common to most plant origin oils and fats is the high percentage of unsaturated fatty acids in the glyceride (3).

Vegetable oils are one of the major components of human diets comprising as much as 25% of average caloric intake (3). While high levels of saturated fatty acids is desirable to increase oil stability, on the other hand, nutritionally they become undesirable. At room temperature (25°C), the saturated fatty acids from 12 to 24 carbon length have a waxy consistency, whereas unsaturated fatty acids of these lengths are only liquids (4). This difference is due to different degrees of packing of the fatty acids molecules. In the fatty saturated compounds, free rotation around each carbon-carbon bond gives the hydrocarbon chain great flexibility, the most stable conformation is the fully extended form in which the steric hindrance of neighbouring atom is minimized. These molecules can pack together tightly in nearly crystalline arrays, with atoms all along their length. Van der Waals contacts with atoms of neighbouring molecules. In unsaturated fatty acids, a cis double bond (the double bond in most naturally occurring unsaturated fatty acids are in the cis configuration) forces a kink in the hydrocarbon chain. Fatty acids with one or several such kink cannot pack together as tightly as fully saturated fatty acids, and their interactions with each other are therefore weak (4,5). Because of the waxy consistency of the saturated fatty acids, oils rich in saturated fatty acids tend to form diets, which result in the commonly known coronary diseases. When this oils are constantly consumed, they form solid fats deposit at both the cardiac region, the arteries and the veins. These fats tends to block the free flow of blood in these arteries and veins and thus increases the pressure to the blood flow and this eventually result to high blood pressure in which case if not treated carefully can lead to stroke and finally kills the victim (4).

Though the presence of unsaturated fatty acids in oils is preferred to saturated fatty acids in terms of consumption, high amount of unsaturated fatty acids in an oil makes it susceptible to polymerization and gum formation caused by oxidation during storage (6, 7). The polymerization product is a dry or hardened (solid) film. This imparts on the oil a level of industrial utility than edibility (2, 7).

The quality and purity of oils are assessed by determining the number of physical and chemical constants. The chemical constants include acid value, iodine value, saponification value etc (2). Of particular interest is the iodine value, because depending on its value, oils can be classified into drying, semi drying and non drying. The ability of an oil to be converted into a solid film in the presence of oxygen is known as drying.

There are various brands of vegetable oils sold in the markets located in Asaba, some are produced in the country while some are imported. Despite the strict regulations and enforcement by relevant regulatory agencies of sale of standard commodities to consumers, a times, manufacturers and importers do not comply to standards. Again, despite the tight security measures at the Nigerian sea ports, airports and borders, banned and substandard products still find their way into the country and eventually the markets (8).

In the light of the possible consequences of consuming oils high in saturated and unsaturated fatty acids, this study was undertaken, so as to access the quality of oils sold in the city (Asaba), if they are good for consumption, through examination of their chemical properties.

2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

Five different brands of vegetable oils found in almost all the market places in Nigeria, were purchased from the market. The five different brands of vegetable oils are Baron vegetable oil, Gino, Turkey, Ideal and Grand vegetable oils. Gino and Turkey are imported vegetable oils into the country while Ideal, Baron and Grand are produced in the country (Nigeria). The chemical properties of the oil namely the iodine value, saponification value, peroxide value and acid value were determined by standard procedures described by AOAC (1980) while the ester value was obtained by subtracting the acid value from the saponification value (Baltes, 1964). The iodine value, saponification value, peroxide value and acid value were determined in triplicate then their mean and standard deviation were calculated.

3. RESULTS AND DISCUSSION

The iodine value is a measure of the unsaturation of an oil. The higher the iodine value, the more the double bonds (unsaturation) present, which consequently reflects the reactivity of the oil; i.e the oil becomes more susceptible to oxidation and rancidification (11). Oleic acid containing one double bond absorbs 90% of iodine, linoleic acid (2 double bonds) absorbs 181% iodine and linoleic acid (3 double bonds) absorbs 274% iodine (12). Non drying oils, such as olive oil, used for soap making and in the food products have 1 double bond and have relatively low iodine values below 90. Semi drying oils such as soya bean oil contain some proportion of double bonds and have intermediate iodine values about 130 (below 140). Drying oils used in the paint and varnish industry have relatively high iodine values (about 190) (12). Tables 1 shows the iodine values of the oils, the result classifies/suggest that these oils are non-drying oil of low unsaturation. Some commonly used drying oils include linseed oil, tung oil etc, these oils are susceptible to drying and are often unsuitable for cooking (13), while Non drying oils are slow to oxidize and so remain liquid for a long time (14). Since the oils of this study contains one double bond on the basis of their classification as non-drying oils, they are expected to be less reactive, more stable, slow to oxidation and remain liquid for a long time. This suggest or gives an insight that they are suitable for consumption

Table 1. Iodine value in mg/100g of the five selected refined vegetable oils

Brands of refined vegetable oils	Iodine value
----------------------------------	--------------

Baron vegetable oil	57.70±0.92
Gino vegetable oil	79.95±0.12
Ideal vegetable oil	67.89±0.61
Grand vegetable oil	50.13±1.35
Turkey vegetable oil	21.83±0.07

The results are mean ± standard deviation of triplicate determination.

According to benardin (1973), the characteristics that are necessary for the confirmation of identity and edibility include acid value (15). High acid value are poisonous for livestock and consumption (16). Acid value is a measure of the free fatty acid present in fats and oils and it is a measure of hydrolytic rancidity (17). The higher the acid value, the higher the level of free fatty acid which translates into hydrolysis of triglycerides and decreased oil quality (18). According to Codex Standards for Edible Fats and Oils (1999), the maximum level of acid value for refined fats and oils is 0.6mgKOH/g fat or oil (19). From the scoring table (table 2), except for Gino vegetable oil, the other brands of vegetable oils had acid value a little above the maximum level for acid value quoted for refined oils. This suggests that Gino vegetable oil is suitable for consumption. An increment in the amount of free fatty acid in an oil sample indicates inadequate processing and storage conditions (18), probably poor storage conditions brought about oxidative and chemical changes in the oil leading to increase in free fatty acid. The acid value of the other brands may still be considered low, which implies that the oils may be considered edible.

163
164

Table 2. Acid value in mgKOH/g of the five selected refined vegetable oils

Brands of refined vegetable oils	Acid value
Baron vegetable oil	1.49±0.14
Gino vegetable oil	0.39±0.02
Ideal vegetable oil	1.17±0.01
Grand vegetable oil	1.00±0.35
Turkey vegetable oil	1.54±0.33

165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195

The results are mean ± standard deviation of triplicate determination.

The saponification value of most edible oils are in the range of 180-200mgKOH/g (20). Most of the oils namely sun flower oil, coconut oil, cotton seed oil, palm oil, sesame oil, soybean oil used as salad and cooking oils have saponification values ranging from 186mgKOH/g to 265mgKOH/g(21, 22). Also, saponification values of 200mgKOH/g and above had been reported to possess low molecular weight fatty acids (23). Therefore, it reveals that the oils studied possessed low molecular weight fatty acids and are edible.

196 **Table 3. Saponification value in mgKOH/g of the five selected refined vegetable oils**
197

Brands of refined vegetable oils	Saponification value
Baron vegetable oil	235.62 ± 0.8
Gino vegetable oil	234.92 ±0.3
Ideal vegetable oil	206.87±0.07
Grand vegetable oil	246.14±0.42
Turkey vegetable oi	274.19±1.70

198 *The results are mean ± standard deviation of triplicate determination.*
199

200 The peroxide value gives a measure of the extent to which an oil sample has undergone
201 primary oxidation (24). Oils with a high degree of unsaturation are most susceptible to
202 oxidation (24, 25). The maximum level of peroxide value for fats and oils is 10 milliequivalent
203 of active oxygen /kg oil (19). From table 4, the peroxide values of the different brands of oils
204 can be said to be very low indicating very low oxidation/ rancidity. These peroxide values
205 also suggest that these oils contain low molecular weight fatty acids of low unsaturation
206 hence, it further reveals that the oils are slow to polymerization and will remain liquid for a
207 long time. Summarily, they are suitable for consumption.

208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227

228
229
230

Table 4. Peroxide value in meq/kg of the five selected refined vegetable oils

Brands of refined vegetable oils	Peroxide value
Baron vegetable oil	0.81±0.06
Gino vegetable oil	1.17±0.06
Ideal vegetable oil	0.26±0.03
Grand vegetable oil	0.88±0.02
Turkey vegetable oil	0.74±0.01

231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259

The results are mean ± standard deviation of triplicate determination.

Ester value of the oil is very high compared with the acid value which is a very essential characteristics of edible oil (26).

260

Table 5. Ester value in mgKOH/g of the five selected refined vegetable oils

Brands of refined vegetable oils	Ester value
Baron vegetable oil	234.13
Gino vegetable oil	234.53
Ideal vegetable oil	206.70
Grand vegetable oil	245.14
Turkey vegetable oil	272.65

261

262

263

264

265

4. CONCLUSION

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

REFERENCES

285

286

287

288

289

290

291

292

293

1. Vegetable oils. Accessed 3 June 2014. Available: http://en.wikipedia.org/wiki/Vegetable_oil.
2. Ibemesi J A. From studies in polymers and vegetable oils to sanitization of the academic system. Accessed 11 July 2014. Available: <http://www.unn.edu.ng/files/Inaugural%20Lecture%20Documents/Physical%20science/19th%20Inaugural%20Lecture.pdf>.

- 294 3. Zambiasi R C, Przybylski R, Zambiasi M W and Mendonca C B. Fatty acid composition
295 of vegetable oils and fats. Accessed 6 June 2014. Available:
296 <http://ojs.c3s.ufpr.br/ojs/index.php/alimentos/article/download/8399/5848>.
- 297 4. David L N, Michael M C. A practical method for determining the oil content of avocado.
298 Citrus Subtropical Journal. 2005: 84-92
- 299 5. The kinds of fats. Accessed 6 June 2014. Available:
300 <http://www.indiana.edu/~oso/Fat/trans.html>.
- 301 6. Ma F, Hanna M A. Biodiesel production: A review. Bioresource technology. 1999; 70(1).
302 1-15
- 303 7. SLTC, the society of leather traders chemists official method of analysis. 1963.
- 304 8. Jide O. policing our porous borders. Accessed 6 April 2013. Available:
305 www.nigeriaintel.com/2013/04/03/policing-our-porous-borders.
- 306 9. AOAC, Official Methods of Analysis. Association of Official Analysis Chemist. 13th ed.
307 Washington DC. 1980.
- 308 10. Baltes J. 1964 Classical methods in fat analysis. In: Bockenoogen H A. Analysis and
309 characterization of oils, fats and fat products. London: Interscience publishers.
- 310 11. Iodine value. Accessed 10 June 2014. Available:
311 <http://www.britannica.com/EBchecked/topic/292677/iodine-value>.
- 312 12. Estimation of iodine value of oil. Accessed 10 June 2014. Available:
313 <http://www.classle.net/book/estimation-iodine-value-oil>.
- 314 13. Dry oils. Accessed 10 June 2014. Available: http://en.wikipedia.org/wiki/Drying_oil.
- 315 14. Alternative vegetable oils perennials. Accessed 10 June 2014. Available:
316 <http://www.pfaf.org/user/cmspage.aspx?pageid=41>.
- 317 15. Benerdini E. (1973). Oil and fat technology. In: Abitogun A S, Borokini F B.
318 Physicochemical parameters and fatty acid composition of cashew nut(anacardium
319 occidentale) oil. Journal of Research in National Development. 2009;7(2):1-3.
- 320 16. Umerie S C, Okonkwo I F, Nwadiolor N A, Okonkwo J C. Studies on the oil and
321 nutritive value of seeds of crotalaria retusa L.(fabaceae). Pakistan Journal of Nutrition.
322 2010;9(9):912-914.

- 323 17. Sampling and Analysis of Commercial Fats and Oils.AOCS. American Oil Chemists
324 Society Official Method Cd 8 – 53. 2003.
- 325 18. Acid value(AV) and the quality of fats and oils. Accessed 23 June 2014. Available:
326 [http://wiki.chemprime.chemeddl.org/index.php/Acid_Value_\(AV\)_and_the_quality_of_fats_and_oils](http://wiki.chemprime.chemeddl.org/index.php/Acid_Value_(AV)_and_the_quality_of_fats_and_oils).
327
- 328 19. CODEX STAN. 1999. Codex General Standard for Fats and Oils. Accessed 23 June
329 2014. Available: <http://www.fao.org/docrep/004/y2774e/y2774e03.htm>.
- 330 20. Glasser A N. 2008. Analysis of fixed oils, fats and waxes. In: Pharmaceutical chemistry
331 applications. Vol 1.New Delhi. Leslie G. Chattern.
- 332 21. CODEX Standard for Named Vegetable Oils, 1999. Accessed 23 June 2014. Available:
333 <http://www.codexalimentarius.org/input/download/standards/336/CXS210e.pdf>.
- 334 22. List of vegetable oils. Accessed 26 June 2014
- 335 23. Afolabi IS. Chemical qualities of oils from some fresh and market vegetable crops within
336 kwara state of Nigeria. Biokemistri. 2008; 20(2): 71-75
- 337 24. Peroxide value. Accessed 14 June 2014. Available:
338 http://en.wikipedia.org/wiki/Peroxide_value.
- 339 25. Milk placers for wildlife rancidity. Accessed 15 June 2014. Available:
340 <http://www.ewildagain.org/Milk%20Replacers/rancidity.htm>.
- 341 26. Gregory IO. 2005. In: Salau AB, Mada SB, Bola MA. Physicochemical properties of oil
342 ectracted from winged termite macrotermis bellicosus.Int. J. of Modern Biochemistry.
343 2012;1(1):36-40.

344

345