

TOXICITY AND HISTOPATHOLOGICAL EFFECT OF CYPERMETHRIN ON JUVENILES OF *CLARIAS GARIEPINUS*

²R.O OJUTIKU ¹F.P ASUWAJ U *, ²R.J. KOLO ¹R.A OBANDE ¹O.O AGBELEGE

¹ FEDERAL COLLEGE OF FRESHWATER FISHERIES TECHNOLOGY, BAGA,
MAIDUGURI,

² DEPARTMENT OF WATER RESOURCES, AQUACULTURE AND FISHERIES
TECHNOLOGY, FEDERAL UNIVERSITY OF TECHNOLOGY, MINNA.

³ CONVENANT UNIVERSITY, DEPARTMENT OF BIOLOGICAL SCIENCE, OGUN STATE.

rasheedojutiku@yahoo.com

ABSTRACT

*The toxicity and histopathological effects of Cypermethrin on the gill and liver tissues of the *Clarias gariepinus* were determined in this study. The fish were exposed to six (6) acute concentrations (0.025mg/l, 0.050mg/l, 0.075mg/l, 0.100mg/l, 0.125mg/l and 0.000mg/l) for 96 hours. The 96 hours LC50 of the toxicant to the test fish was 0.060mg/L . The most common gill changes at all doses of cypermethrin in solution were destruction of gill lamella, epithelial hyperplasia and epithelial hypertrophy. Hepatic lesions in the liver tissues of fish exposed to Cypermethrin in solution were characterized by degeneration of hepatocyte, vasculization of cell cytoplasm, fatty degeneration and hypertrophy of hepatocytes. Histological comparison of tissues indicated that most damage occurred in the gill rather than in the liver.*

INTRODUCTION

The application of environmental toxicology studies on non-mammalian vertebrates is rapidly expanding, and for aquatic system, fish have become indication for the evaluation of the effects of noxious compounds. (Evrnest, 2004). Pesticides occupy a unique position among many chemicals which are encountered daily by man. They are deliberately added to the environment for the purpose of killing, injuring or at times enhancing the development of some forms of life. Pesticides represent one group of agro-chemicals that are used in large quantities by agro-farmers.

The use of pesticides has increased rapidly since World War II, and several recent studies show that pesticide residues frequently occur in surface water in agricultural areas (Høysæter, 1994; Larson et al., 1999; Ule'n et al., 2002). Many aquatic species are taxonomically related to the target organisms of pesticides. Cypermethrin is a synthetic pyrethroid used for the control of ectoparasites which infest cattle, sheep, poultry and some companion animals. Recently, the compound has been used as a chemotherapeutic agent for the control of ectoparasite infestations (*Lepeophtheirus salmonis* and *Caligus elongatus*) in marine cage culture of Atlantic salmon, *Salmo salar* (Richards, 1983; Roth et al., 1993; Hart et al., 1997; Boxaspen and Holm, 2001; Treasurer and Wadsworth, 2004).

Fish sensitivity to pyrethroids may be explained by their relatively slow metabolism and elimination of these compounds. The half-lives for elimination of several pyrethroids by rainbow trout are all longer than 48 h, while elimination half-lives for birds and mammals range from 6 to 12 h (Bradbury and Coats, 1989b).

The present study aimed to determine the effect of acute toxicity and histopathological effect of Cypermethrin on juveniles of *Clarias gariepinus* in a static bioassay.

MATERIAL AND METHODS

180 healthy juveniles of *Clarias gariepinus* of the same cohort with average weight (16.62 ± 4.36)g, standard length (12.64 ± 1.03) cm and total length (14.97 ± 8.94) cm were sourced from the hatchery unit of the Federal college of Freshwater Fisheries Technology, Baga Maiduguri, Borno State. They were acclimatized for seven (7) days during which they were fed 5% of their body weight with commercial Coppens (2mm). Feeding was stopped 24hrs prior to the commencement of the toxicity test experiment

TOXICITY TEST

A preliminary range finding test was carried out based on the concentration of the active ingredient in the test chemical. The range finding was done using the following concentrations; 0.1mg/l, 10mg/l and 100mg/l of Cypermethrin for 24hrs in triplicates. The concentration was done using a serial dilution formula $C_1V_1 = C_2V_2$.

The result obtained from the range finding test provided a guide for the definitive test. Following this, the definitive test was carried out using; 0.025mg/l, 0.050mg/l, 0.075mg/l, 0.100mg/l, 0.125mg/l and 0.000mg/l of Cypermethrin. The result obtained was used to determine the median lethal concentration (LC_{50}) using Probit analysis.

EXPERIMENTAL PROCEDURE

A total of eighteen (18) glass aquaria were used for the definitive toxicity test. Ten juveniles of *Clarias gariepinus* were introduced into each aquarium with 20 litres of water with; 0.025mg/l, 0.050mg/l, 0.075mg/l, 0.100mg/l, 0.125mg/l and 0.000mg/l concentration of Cypermethrin at the same time. Each of the toxicant concentration was replicated three times each. The experiment was carried out using a static non-renewal bioassay for 96 hours. Mortality and general behavior of fish were noted 24 hourly.

HISTOPATHOLOGICAL EXAMINATION

The histology of the gills and liver of *Clarias gariepinus* was carried out after 96 hours exposure period to the various concentration of the toxicant (Cypermethrin).

TISSUE PROCESSING

Organs of gills and liver were collected and fixed in 10% formal saline for one week. They were then processed for routine paraffin histological sectioning. The tissues were dehydrated through graded concentration of ethanol (30%, 50%, 70%, and 90%), absolute ethanol and cleared in xylene. The tissues were pre-impregnated in xylene paraffin wax in the oven and embedded in pure paraffin wax. The organs were sectioned at 7µm thickness and tissues were stained with Haematoxylin and Eosin (H&E) for light microscopic examinations (Drury et al. 1967).

RESULTS

FISH BEHAVIOUR AND MORTALITY

During exposure, the test fish exhibited the following behavioural patterns before death occurred; restlessness, respiratory distress, loss of balance, gasping for air, vertical movement, excessive accumulation of mucus on the skin and death. The reaction to the toxicant was more pronounced in the aquaria containing the highest three (3) of 0.075mg/l, 0.1mg/l and 0.1mg/l concentrations of the toxicant. The mean cumulative mortality value, log 10 values of the concentrations, the total numbers of exposed fish and the probit kills' values are presented on Table 4.1. The observation made with control group were however different, they had stable swimming and no respiratory stress.

LETHAL CONCENTRATION (LC50)

The result obtained from this research work showed that the 96 hour LC50 was determined to be 0.060mg/l by plotting the graph of probit kill values against log of concentration and finding the values of 50% probit kill as against the antilog value on log concentration value (FIG.1).

The stressful behaviour exhibited by the fish may be attributed to the effects of the toxicant on the gill. This was clearly noted from the result of histopathological examination of the gill. The observed increase in erratic swimming, instability and subsequent immobilization before death which is directly proportional to the concentration of the toxicant could be attributed to the

respiratory impairment of the gills filaments by the insecticide. Several authors have reported work on similar patterns of abnormal behavioural responses in fish exposed to toxicant (Wade, et al., 2002, Oti, 2002) .Previous histopathological studies of fish exposed to pollutant have shown that fish gills are efficient indicators of water quality. Fish gills are vulnerable to pollutants in water because of their large surface area and external location. However, the gills perform numerous functions, which include respiration, excretion of nitrogenous waste products and acid-base balance.




Fig. 1: 96 hour LC50 of *Clarias gariepinus* exposed to different concentration of cypermethrin in static system

LIVER TISSUE

The liver tissue (plate I) of fish in the control set up showed a normal hepatocyte radiating from the central vein and sinusoids. It exhibited the typical parenchymatous appearance. The histomorphology of liver specimens exposed to cypermethrin insecticides showed a significant variation from the control group. Liver specimens (plate II) exposed to 0.025mg/l of cypermethrin shows a wide spread sinusoids haemorrhage, mild area of congestion, necrosis and vacuolation. A focal area of mononuclear cell infiltration was also evident. Specimen of liver

exposed to 0.05mg/l (plate III) of cypermethrin showed an enlarged vascular channel of congestion, cloudy swelling of hepatocyte and a focal area of inflammatory cells. Liver of fish exposed to 0.075mg/l (plate IV) showed a wide spread of fatty tissue degeneration thus, leading to a fatty tissue, cloudy swelling of hepatocyte with a severe thick fibrous connective tissue observed in the liver of fish exposed to 0.1mg/l (plate V a and b) of Cypermethrin. Plate VI indicates inflammatory cells, vacuolation enlarge fatty tissue





Plate. I. Photomicrograph of fish liver of control showing normal hepatocytes (arrows), radiating away from central vein (CV) and sinusoids (arrow head) H&E x400.

135

136




137

138 Plate II. Photomicrograph of fish liver treated with 0.025mg/L of Cypermethrin showing wide
139 spread sinusoidal haemorrhages (S) and mild areas of coagulative necrosis (CN) H&E x400.

140

141




142

143 Plate : III .Photomicrograph of fish liver treated with 0.050mg/L of Cypermethrin showing
144 enlarged vascular channel (arrow), congestion (CG) and cloudy swelling hepatocytes (CW) H&E
145 x400.


146

147 .



148

149 Plate: III Photomicrograph of fish liver treated with 0.050mg/L of Cypermethrin showing wide
150 spread cloudy swelling of hepatocytes (CH) and focal area of inflammatory cells (arrow) H&E
151 x400.



152

Plate:VI Photomicrograph of fish liver treated with 0.125mg/L of Cypermethrin showing thick fibrous connective tissue (FB), inflammatory cells (thick arrows) and vacuolation (arrows) H&E x400.




Plate.V Photomicrograph of fish liver treated with 0.10mg/L of Cypermethrin showing enlarged vascular channel (arrow), congestion (CG) and cloudy swelling (CW) H&E x400

GILL TISSUE

The degree of damages and variation noticed are shown in photomicrograph. Photomicrograph of fish gill treated with 0.025mg/l of cypermethrin shows gill necrosis, congestion, cartilage and interstitial haemorrhages (Plate VII). This was more evident in higher concentration of the insecticide. Gills exposed to 0.05mg/l showed a multi focal area of interstitial haemorrhages. The gills exhibits a marked alteration in their epithelia (plate VIII). Plate VIII b further shows a moderate gill necrosis with multi focal haemorrhages. Plate IX showed the gills of fish treated with 0.075mg/l of cypermethrin with severe gill necrosis. (Plate X) also showed gills of fish exposed to 0.100mg/l of cypermethrin with extensive vacuolation within the epithelia of the gills. Infiltration, cartilage and interstitial haemorrhages were variation noticed in the gills of fish treated with 0.125mg/l of cypermethrin.(Plate X). The gill arches of *Clarias gariepinus* in the control fish showed normal arrangement pattern. The arches contain primary lamella (Plate XI) projecting on the lateral sides of the primary and secondary lamella (respiratory lamella) .This ultimately showed a normal gill without any alteration or damages. Varied morphological changes occurred in the gill tissue of fish in the aquaria treated with cypermethrin pesticide.




Plate :VII . Photomicrograph of fish gills treated with 0.025mg/L of Cypermethrin showing gill necrosis (GN), congestion (CG), cartilage (CT) and interstitial haemorrhage (arrows) H&E x40.




Plate: VIII . Photomicrograph of fish gills treated with 0.050mg/L of Cypermethrin showing (IH) interstitial haemorrhage (arrows) H&E x400




Plate: IX Photomicrograph of fish gills treated with 0.075mg/L of Cypermethrin showing severe gill necrosis H&E x400




Plate:X . Photomicrograph of fish gill treated with 0.100mg/L of Cypermethrin showing necrosis of the gill filament (thin arrows), interstitial haemorrhage (thick arrows) cartilage (C) H&E x400




Plate:XI . Photomicrograph of control fish gills showing normal gills H&E x400

DISCUSSION

Clarias gariepinus juveniles were stressed progressively with time before death. The stressful behaviour of respiratory impairment due to the toxic effect of cypermethrin on the gills was similar with the reports of Omitoyin et. al., (2006) and Aguigwo (2002) that pesticide impairs respiratory organs. Death could therefore have occurred either by direct poisoning or indirectly by making the medium uncondusive or even by both, whichever is the case, the source of death was cypermethrin. Several abnormal behaviour such as incessant jumping and gulping of air, restlessness, loss of equilibrium, increase opercula activities, surface to bottom movement, sudden quick movement, resting at the bottom were similar to the observations of Omoniyi et al., (2002), Rahman et al., (2002) and Aguigwo (2002). The stressful and erratic behaviour of *Clarias gariepinus* juvenile in the experiment indicates respiratory impairment; probably due to the effect of the toxicant cypermethrin on the gills. The fishes became inactive at higher concentrations with increased time of exposure to toxicant.

The result of the light microscopic studies showed that the morphologic changes were more evidence in the liver of exposed fish as changes were not observed in the control fish. The liver of the exposed fish compared to the control showed varying degree of degeneration of cells, which include hypertrophy of hepatocytes, Fatty degeneration, vascular channel congestion and vascuolization of cell cytoplasm; these alterations were dose-dependent. Cengiz et al., (2001) observed similar histological alterations when varied concentrations of thiodanR were applied to mosquito fish (*Gambusia affinis*). Velmurugan et al. (2003) reported congestion, cloudy swelling of hepatocyte and focal necrosis in liver of *Cirrhinus mrigala* exposed to fenvalerate.

Similar result was reported by Adamu et al (2008), when *Oreochromis niloticus* was exposed to solution of Portland cement. Changes such as hyperplasia, disintegration of hepatic mass and focal coagulative necrosis were found in *Labeo rohita* exposed to cypermethrin (Jee et al., 2005). Liver is the detoxification place of toxicants. The hepatic changes suggested mobilization of same kind of defensive mechanism in an endeavour to detoxify the toxicant-cypermethrin.

CONCLUSION

Juveniles of *Clarias gariepinus* exposed to various concentration of Cypermethrin showed that this synthetic pyrethroid was highly toxic to the *Clarias gariepinus* fingerlings with lethal concentration (LC50)of 0.060mg/l. Evidences of toxicity manifested significantly in the damages caused to the gills and liver studied. The severity of these damages to some selected organs in the body of the fish is proportional to the concentration of the pesticides.

REFERENCES

- Boxaspen K., Holm J.C. (2001): The development of pyrethrum-based treatments against the ectoparasitic salmon lice *Lepeophtheirus salmonis* in sea cage rearing of Atlantic salmon *Salmo salar* L. *Aquaculture Research*, 32, 701–707.
- Bradbury S.P., Coats J.R. (1989a): Comparative toxicology of the pyrethroid insecticides. *Bulletin of Environmental Contamination and Toxicology*, 108, 134–177.
- Drury, R. A., B., wallington, e. A. And camerron, s. R. (1967).Carleton, *Histological techniques* 4th edition.Oxford university press.P. 114 – 35.
- Ervnest, H. (2004): *A Textbook of Modern Toxicology* (3rd ed). John Wiley and Sons, Inc., Hoboken, New Jersey. ISBN 0-471-26508-X, pp 557.
- Høysæter, T., 1994. Pesticides in surface water and in sediments in Norway. In: Helweg, A. (Ed.), *Pesticides in Precipitation and Surface water*. TemaNord 1995:558. Nordic Council of Ministers, Copenhagen, pp. 103—/118.
- Larson, S.J., Gilliom, R.J., Capel, P.D., 1999. Pesticides in streams of the United States-Initial results from the national water-quality assessment program. US Geological Survey, Water-Resource, Investigation Report 98-4222.

246 Richards R.H. (1983): Diseases of farmed fish: Salmonids. Veterinary Records, 112, 124–126.

247 Roth M., Richards R.H., Sommerville C. (1993): Current practices in the chemotherapeutic

248 control of sea lice infestations in aquaculture: a review. Journal of Fish Diseases,

249 16, 1–26.

250 Hart J.L., Thacker J.R.M., Braidwood J.C., Fraser N.R., Matthews J.E. (1997): Novel

251 cypermethrin formulation for the control of sea lice on salmon (*Salmo salar*).

252 Veterinary Records, 140, 179–181

Conc.	Log ₁₀ of	Total	Numb.of	%	Probit kill value
-------	----------------------	-------	---------	---	-------------------

253 Ule'n, B., Kreuger, J., Sundin, P., 2002. Undersökning av bekämpningsmedel i vatten från

254

255 Treasurer J.W., Wadsworth S.L. (2004): Interspecific comparison of experimental and natural

256 routes of *Lepeophtheirus salmonis* and *Caligus elongatus* challenge and

257 consequences for distribution of chalimus on salmonids and therapeutic

258 screening. Aquaculture Research, 35, 773–783.

259

260

261

262

Mg/l	conc.	numb. Of exposed fish	dead fish	mortality	
0.000	0.00	10	0.0	0.0	00000
0.025	- 1.60	10	1.3	13	3.8736
0.050	- 1.30	10	2.7	27	4.3872
0.075	- 1.12	10	3.3	33	4.5603
0.100	- 1.00	10	5.0	50	5.0000
0.125	- 0.90	10	7.3	73	5.6128

Table 4.1: Mortality rate, percentage mortality and Probit value of *Clarias gariepinus* juveniles exposed to acute dose of Cypermethrin for 96 hour in a static non-renewal bioassay

263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279

280