

Original Research Article

COMPARATIVE ANALYSIS OF THE NUTRITIONAL QUALITIES OF SEEDS OF SHEANUT (*Vitellaria paradoxa*) AND COCOA (*Theobroma cacao*)

Abstract

*This research work involves a study of the chemical, nutritional and anti-nutritional compositions of sheanut (*Vitellaria paradoxa*), cocoa (*Theobroma cacao*), and the effect of roasting on these properties. The ash, crude fibre, fat and carbohydrate contents of the two seeds showed some  of similarities, with sheanut having lower crude protein content of 6.74 - 6.84% when compared with that of cocoa, 13.26 -13.56%. The theobromine content of the sheanut was higher (2.34 - 4.50%) than that of cocoa (1.98- 2.11%). The mineral compositions of the two seeds were very low but the effect of roasting on the mineral content followed the same trend of increasing both the macro and micro elements contents of the samples. The anti-nutrient compositions of the two seeds were similar, with roasting having an increasing effect on the anti-nutritional factors (phytic acid, oxalate and tannin) when compared with the raw samples. The results revealed that sheanut will be a suitable substitute for cocoa in  industrial applications.*

Keywords: Cocoa, sheanut, theobromine, cocoa nibs.

INTRODUCTION

The tropical crops under study are cocoa (*Theobroma cacao*) and sheanut (*Butyrospermum parkii*). Cocoa belongs to the genus *Theobroma* in

the family of *Sterculiaceae*, while sheanut belongs to the genus *Butyrospermum* in the family of *Sapotaceae* (Samba-Murty and Subrahmanyam, 1989). The tree crop, cocoa (*Theobroma cacao*) has played an important role in the economies of various regions of the world. Cocoa which is harvested from the tree had been a major foreign-exchange earner in West Africa and South America. Cocoa beans had served as money in South American civilizations (Are & Gwynne-Jones, 1974). Sheanut tree is a medium-sized deciduous tree which occurs mostly in West Africa, Northern Uganda and Southern Sudan. The seed contains 45 – 60% of the oil called shea butter, which is solid at room temperature (Samba-Murty and Subrahmanyam, 1989; Eagle, 2009). According to Eagle (2009), shea nut butter is one of the Africa's most sustainable natural resources and the shea tree is an important source of income; it's the third largest cash crop in Burkina Faso and Ghana, where it's surpassed only by cocoa and coffee. The shea trees grow naturally in the grasslands of West and Central Africa and do not need irrigation, fertilizer, or pesticides. The shea trees are native to the savanna and are difficult to grow in plantations. It takes at least 25 years for a shea nut butter tree to produce large numbers of fruits. Shea butter has been used for centuries in Africa for its moisturizing and healing properties, where it has been used to protect and condition

skin, which have been damaged by the sun and wind (Maranz et al, 2004).

According to Dogbevi (2008), the economic importance of the shea tree cannot be overemphasized, especially in the face of the unstable world market price for cocoa and the need to find suitable substitutes for cocoa in the beverage, confectionery and cocoa butter industries. In order to determine the suitability of sheanut as a substitute for cocoa and as a potential industrial raw material, there is need for comprehensive study of the sheanut seeds.

Thus, this paper compares the chemical, nutritional and anti-nutritional compositions of sheanut (*Butyrospermum parkii*) with cocoa (*Theobroma cacao*) and the effect of roasting on these properties.

MATERIALS AND METHODS

Sample collection and preparation

The samples used for this study were raw cocoa nibs, roasted cocoa nibs, raw shea nuts and roasted shea nuts. The dried cocoa beans were obtained from Idanre Town, in Idanre Local Government Area of Ondo State, while the dried shea nuts were obtained from Bida Town, in Bida Local Government Area of Niger State; both states are located in Nigeria. The samples obtained were sun-dried, cleaned thoroughly to remove extraneous materials (stones, pebbles and twigs) and then separated into two portions. One portion was roasted in the oven at 120°C for 2 hours,

while the other portion was used in its raw state. The raw and roasted cocoa beans were shelled to obtain the cocoa nibs. The different samples were pounded separately with a pestle in a clean wooden mortar and thereafter ground with a laboratory Waring blender to obtain suitable particle sizes for laboratory analysis. The ground samples were stored in an air-tight plastic container and kept in the refrigerator, prior to analysis.

Determination of proximate and mineral composition

The proximate composition was determined by official methods of Analysis of the Association of Official Analytical Chemists (AOAC, 1990). For the minerals, sodium, calcium and potassium contents were determined by Flame Photometer (Hachway PFP 7 model), while phosphorus content of the samples was determined by Colorimetric method using Vanado Molybdate reagent (Ibitoye, 2005). Other minerals were determined after wet digestion with aqua regia using Atomic Absorption Spectrophotometer (Buck 205 model).

Theobromine Determination

Theobromine content was determined by the method of Pearson (1976).

Determination of Phytic acid, oxalate and tannin

The method of Young and Greaves (1940) was employed for phytic acid determination; the

oxalate content of the samples was determined by the method of Day and Underwood (1986), while tannin was quantified as described by Padmaja (1989).

RESULTS AND DISCUSSION

The chemical composition of the samples under study is shown in figure 1. The percentage moisture contents of the raw cocoa nib, roasted cocoa nib, raw sheanut and roasted sheanut are within the 7.5% maximum moisture content recommended by the International Cocoa Organization, Standard Organization of Nigeria and Cocoa Research Institute for commercial cocoa (Oyebode, 2000; NIS, 2003).

The moisture content determination gives an idea of the amount of water content in a given food; low moisture content prevents deterioration and spoilage in foods and is an indication of good shelf life of food samples (Yasgajau and Clifton, 1978).

Although, the moisture content of shea nut is higher than that of cocoa nib, the result showed that the moisture content of the roasted samples are generally lower than that of the raw samples. This implies that roasting has a reducing effect on the moisture content of the samples.

The ash content of all the samples is in agreement with the average total ash content of 3.3% specified for cocoa nib by Pearson (1976). The importance of ash content is that it gives an

amount of mineral elements and inorganic matter present in a food sample. These values for the ash content indicate that all the samples have low inorganic matter. These values for the ash content of the samples are a little bit lower than 3.49 to 3.70 % obtained for cocoa nib by Oyeboode (2000).

The fat content ranges from 60.44 to 67.23% for the samples. This shows that both cocoa and sheanut can be used as an alternative source of vegetable oil. The values are in agreement with the percentage fat composition of about 60% reported for cocoa and sheanut by previous researchers (Samba Murty and Subrahmanyam, 1989; Oyeboode, 2000; Dei *et al.*, 2007; Ogunlola, 2007). Roasting had a slight increasing effect on the fat content of the samples. Oils and fats are a vital component of the human diet because they are an important source of energy and also act as a carrier of fat soluble vitamins (Kochhar, 1986).

The crude fibre content ranges between 3.26 to 4.88% for the ground samples. These satisfy the maximum crude fibre content of 5.21% specified for commercial cocoa by Pearson, 1976. Roasting also increased the crude fibre content of the samples. The samples fibre contents are normal and this shows that the food samples when eaten can serve to prevent diseases associated with diets low or high in fibre content. Lack of fibre in the diet usually makes it difficult to eliminate food waste from the body. Consistent shortage of fibre

in the diet could lead to constipation and other disturbance of the colon (Okon, 1983; Ibitoye, 2005).

The crude protein content ranges from 6.74 to 13.56% for the samples. Protein serves as a source of energy and amino acids required by the body and they also play a part in the organoleptic properties of foods (Okon, 1983).

The recommended daily allowance (RDA) of protein for children ranges from 23.0 – 36.0g and for adult, 44-56g (NRC, 1989). The mean crude protein values of the samples are too low to be recommended as a protein source for children and adults.

But the crude protein content of cocoa samples compare well with the results 13.56- 13.94 % obtained by Oyeboode (2000). Results showed that roasting reduced the crude protein content of both samples.

The carbohydrate content ranges from 12.36 to 15.09%. This shows that consumption of cocoa and sheanut products will provide reasonable amount of energy for the body and they will also be a good source of feed for livestock (Ogunlola, 2007). The carbohydrate content of these samples are lower than 21.07- 21.17% reported for cocoa nibs by Oyeboode (2000).

The theobromine content of the cocoa samples ranges from 1.98 to 2.11%, while the sheanut samples has between 2.34 to 4.50% theobromine content. According to Ghilleen and

Mark (2004), the theobromine content of cocoa powder may reach up to 10%. Roasting of the samples increased the theobromine content. The stimulating properties of cocoa and chocolate are due to the presence of theobromine and small quantities of caffeine (Kochhar, 1986; Matissek, 1997; Smit *et al.*, 2004).

Figure 2 shows the results for the mineral composition of the samples.

All forms of living things require different minerals for their life processes.

All the samples were low in essential mineral elements such as Ca, K, P and Mg. Calcium, an essential element is required for growth and maintenance of healthy teeth and bones. Potassium is required for enzyme activities. Magnesium is an essential element present in ionic form in all tissues where it plays a part in many reactions involved in energy utilization (Olafe *et al.*, 1987).

Although, sodium is one of the major (essential) mineral elements in foods, the recommended daily allowance (RDA) by NAS (1974) and NRC (1989) is 2500 mg/kg per day. Sodium is responsible for liquid membrane transport in the body and the values obtained for sodium in all the samples are generally low, its values range from 600 to 870 mg/kg in the samples.

All the samples contain Mn, Fe, Cu and Zn at trace levels.

Iron is a trace element which is a component of haemoglobin (the pigment of red blood cells). Manganese is used in the process of respiration and copper is responsible for the formation of haemoglobin, healthy bones, blood vessel and nerves (Olafe *et al.*, 1987). Zinc is essential for maintaining a good skin and its deficiency leads to skin lesions and testicular atrophy (Robert and Daryl, 2000).

According to Nieman *et al.* (1992), Na/K ratio in the body is of great importance in the prevention of high blood pressure. Na/k ratio less than one is recommended. From the results, all the four samples analysed have Na/K ratio less than one, this indicated that their consumption would probably reduce high blood pressure.

The observed low essential elements may be due to the presence of the anti-nutritional factors such as phytic acid, oxalate and tannin. It has been reported that the presence of these anti-nutritional factors in foods inhibit the bioavailability of essential minerals (Aletor, 1993; Tacon, 1995; NRC, 1993; Eradman, 1979; Hassan *et al.*, 2005).

Figure 3 shows the anti-nutrients content of the samples. The anti-nutritional nature of phytic acid lies in its ability to chelate certain important minerals such as calcium, magnesium, iron and zinc in humans and animals, forming mineral-phytic acid complexes. The net result of this is reduced mineral bioavailability in animals

or humans consuming diets high in phytic acid (NRC, 1993). The phytic acid content of these samples was relatively low when compared to phytic acid content of other plants such as legumes (Balogun and Fetuga, 1980; Aletor, 1990 & 1993; Tacon, 1995).

The charge on oxalate allows it to act as a chelator of various positively charged metal ions such as Ca^{2+} , Fe^{2+} and Mg^{2+} , to deposit crystals of the corresponding oxalates in the body, which irritate the gut and kidneys. Methods to reduce the oxalate content in food are of current interest to scientist (Betsche and Frezdorff, 2005). The oxalate content of the examined samples was relatively low when compared to oxalate content of some tropical leafy vegetables (Aletor and Adeogun, 1995; Balogun and Fetuga, 1980).

According to existing literature, natural tannins are widely distributed in the plant kingdom. They occur in small amounts in most plant tissues (Kochhar, 1986; Pearson, 1976; Ihekoronye and Ngoddy, 1985). The percentage composition of natural tannin in cocoa nib is about 5.8 % (Samba-Murty and Subrahmanyam 1989).

The bitterness and astringency of the cocoa liquor is due to the presence of tannin. Tannic acid has been reported to bind with the proteins of saliva and the mucosal membrane of the mouth during the mastication of food and, consequently inhibit the digestive enzymes of the mouth and bind proline-rich proteins in the saliva (Aletor,

1993; Tacon, 1995). Among the anti-nutritional effects attributed to the tannins is a decrease in the digestibility of proteins and carbohydrate as a result of the formation of insoluble enzyme-resistant complexes with tannins. Other anti-nutritional effects of tannins include: damage to the gastrointestinal tract, interference with the absorption of iron and possibly, a carcinogenic effect (Ford and Hewitt, 1979).

It was observed that roasting increased the anti- nutrient contents of the two seeds when compared with the raw samples. According to Aletor and Adeogun (1995), the variation can largely be attributed to the lower moisture content of the roasted seeds.

However, these results were in agreement with the earlier work of Makkar *et al.* (1997) on *Jatropha curcas* where it was reported that concentration of anti-nutritional factors such as phytate, tannin were either similar or higher in processed seeds.

CONCLUSION

This research work compared the chemical, nutritional and anti- nutritional composition of the seeds of cocoa and sheanut, and most importantly the effect of roasting at improving or enhancing the desirable properties.

The results obtained from this research work revealed that sheanut, have similar chemical, nutritional and anti-nutritional compositions as cocoa. The results also revealed that roasting

significantly reduced moisture and crude protein contents of the seeds, while roasting significantly increased the minerals, ash, crude fibre, fat, theobromine, phytic acid, oxalate and tannin contents of the seeds.

It can be conveniently concluded that sheanut and its products will be a suitable substitutes for cocoa and its products in the beverage, confectionery, chocolate and other industries where cocoa is being widely used. As a result, sheanut can serve the same household and industrial purposes as cocoa.




Figure 1: The chemical composition (%) of samples




Figure 2: The mineral composition (mg/kg) of samples




Figure 3: The antinutrients content of samples

REFERENCES

- Aletor, V.A. (1990).** ‘Analytical report on anti - nutritional factors in some Nigerian feedstuffs, herbage by products, crop residues and browse plants’. A monograph prepared for the presidential task force on alternative formulation of livestock feeds. Product development, quality evaluation and health implications, Cabinet Office, Lagos, Nigeria.
- Aletor, V.A. (1993).** Allelochemicals in plant food and feeding stuffs (1): Nutritional, Biochemical and Physiopathological aspects in animal production. *Journal of Veterinary and Human Toxicology*, 35(1) PP 57-65.
- Aletor, V.A. & Adeogun O.A (1995).** Nutrient and anti- nutrient components of some tropical leafy vegetables. *Journal of Food chemistry* 53, PP 375-379, Elsevier science Ltd.
- AOAC (1990).** Official methods of Analysis of the Association of Official Analytical Chemists, 12th edition, Washington DC, USA.
- Are, L.A. & Gwynne- Jones, O.R.G. (1974).** Cacao in West Africa. 1st edition, Oxford University Press London. PP 1-10.
- Balogun, A.M. & Fetuga, B.L (1980).** Tannin, Phytin and oxalate content of some wild underutilized crop seeds in Nigeria. *Journal of Food Chemistry* 30, PP 37-43.
- Bernardini, E. (1973).** Oil and fat Technology, Publishing House Technologies S.R.L. Rome. PP 69-101, 709-719
- Betsche, T. & Fretzdorff, B. (2005).** Biodegradation of oxalic acid from spinach using cereal radicles. *Journal of Food Chemistry* 53(25): PP 9751-9758
- Cocks, L.V & Rede, V.C. (1973).** Laboratory handbook for oil and fat analysis. 2nd edition. Academic Press Inc. New York Pp 105-130.
- Day, R.A. & Underwood, A.L. (1986).** Quantification Analysis. 5th edition.

Prentice Hall Publication, New York. Pp 701.

Dei, H.K; Rose, S.P. & Mackenzie, A.M.

(2007). Sheanut (*Vitellaria paradoxa*) meal as a feed ingredient for poultry. World's poultry science journal 63: pp 611-624. Cambridge University press on behalf of world's poultry science association.

Deman, J.M. (1992). Chemical and physical

properties of fatty acids, fatty acids in food and their health implications. Marcel Dekker Inc. New York pp 18-46.

Dogbevi, E.K. (2008). Ghana's scientific

breakthrough and the sheanut industry. An article on www.ghananewstoday.com

Eagle, William (2009). Economic and Health

potential of Africa's Shea Nut

Eka, O.U. (1980). Proximate composition of

bush mango tree and some properties of dika fat. *Nigeria Journal of Nutritional Science*, v(i) PP 33-36.

Eradman, J.W. (1979). Oil seeds phytate:

Nutritional implications. *Journal of Oil Chemistry*, 56: PP 738-741

Ford, J.E. & Hewitt, D. (1979). Protein quality

in cereals and pulses. *British Journal of Nutrition*, 41, PP 341-352.

Ghilleen, P. & Mark, N. (2004). The cultural

history of plants. New York. Routledge, PP 137, 175, 178-180.

Hassan, A.B., Osman, G.A. & Babiker, E.E.

(2005). Effect of domestic processing on antinutrients and availability of protein and mineral of Lupin (*lupinus termis*) seeds. *Journal of Food Technology* 3:PP 263-268

Herbaut, C. (2006): Omega – 3 and health. (In

French). *Rev Med Brue* 27(4): PP 5355-5360

Ibitoye, A.A. (2005). Laboratory Manual on

Basic methods in plant analysis. 1st edition published by concept IT and Educational consults, Akure, Nigeria.

Ihekoronye, A.I. and Ngoddy, P.O. (1985).

Integrated Food Science and Technology for the tropics. 1st edition, Macmillan press Ltd, London

James, C.S. (1995). Analytical Chemistry of

foods, published by Blackie Academic and Professional, Glasgow, United Kingdom. PP 140 – 141.

Kochhar, S.L. (1986). Tropical crops: A

textbook of economic botany. 2nd edition, Macmillan publishers Ltd, London. PP 310-328.

Makkar, H. P. S., Becker, K. & Schmook, B.

(1997). Edible provenances of *Jatropha curcas* from Quintna Roo state of Mexico and effect of roasting on antinutrient and toxic factors in seeds

Maranz, S, Wiesman, Z., Bisgard, J. and

Bianchi, G. (2004). Germplasm resources of *Vitellaria paradoxa* based on variations in fat composition across the species distribution range. *Agro forestry System 60: PP 71-76*

Matissek, R. (1997). Evaluation of xanthine

derivatives in chocolate: Nutritional and chemical aspects. *European Food Research and Technology 205(3):PP 175-184.*

Meyer, L. H. (1987). Food Chemistry. CBS

publishers, India, PP 10-50

National Academy of Science (NAS) (1974).

Recommended dietary allowances. National Research Council, Washington D.C. 8th Edition. PP 25-26

National Research Council (NRC) (1989).

Recommended dietary allowances 10th edition. National academic press, Washington D.C., U.S.A.

National Research Council (NRC) (1993).

Nutrient requirements of fish. Committee on animal nutrition, board of agriculture, national research council, National Academy Press, Washington DC, USA. PP 114.

Nieman, D.C., Butterworth, D.E and Nieman,

C.N. (1992). Nutrition.

W.M.C. Brown Publisher, Dubugue, PP 9-540

Nigeria Industrial Standard (NIS) (2003):

Standard organization of Nigeria (SON)
468.

Ogunlola, O.G (2007). Proximate analysis of cocoa products and comparison of the mineral content. A post graduate diploma (PGD) thesis submitted to the Chemistry Department, Federal University of Technology, Akure, Nigeria.

Okon, B.D. (1983). Studies on the chemical composition and nutritive value of the fruits of African star apple. M.Sc. Thesis, University of Calabar pp 67.

Olaofe, O., Oladeji, F.O. and Ayodeji, O.I. (1987). Metal content of some cocoa bean produced in Ondo state. Nigerian Journal of Science, *Food and Agriculture* 41: PP 241-244.

Oyebode, E.T. (2000). Chemical, Mineral and Microbiological analysis of cocoa beans. PGD thesis submitted to the Chemistry

Department, Federal University of Technology, Akure, Nigeria.

Padmaja, G. (1989). Evaluation of techniques to reduce assayable tannin and cyanide in cassava leaves. *Journal of Agriculture and Food Chemistry* 37, PP 712-716.

Pearson, D. (1976). The chemical analysis of foods. 7th edition, Churchill Living Stone, London, PP 6-14, 200-222.

Reiffel, J.A and McDonald, A. (2006): Antiarrhythmic effects of Omega -3 fatty acids. *American Journal of Cardiology* 98 (4A): PP 501-601.

Robert, K.M. and Daryl, K.G. (2000). Harper's Biochemistry. 25th edition, McGraw Hills companies, USA PP 658-660.

Rossel, J.B. (1991). Vegetable oils and fats, analysis of oil seeds, fats and fatty foods. Elsevier science publishers Ltd, London. PP 261-325.

Samba-Murty, A.V. S.S. and Subrahmanyam, N.S. (1989). A textbook of Economic

Botany. Wiley Eastern Ltd, New Delhi, India. PP 501-509

Smit, H.J., Gaffan, E.A., Rogers, P.J. (2004):

Methylxanthine are the psycho pharmacologically active constituents of chocolate. *Psychopharmacology* 176(3-4): PP 412-419.

Tacon, A.G. J. (1995). Fishmeal replacers:

Review of anti nutrients within oil seeds and pulses – a limiting factor for the aqua feed green revolution. Paper originally presented at feed ingredient, Asia 1995, PP 153-181.

The Goldbook (1997): International Union of Pure and Applied Chemistry (IUPAC) compendium of chemical Terminology. 2nd edition.

Vermani, O. P. & Narula, A. K. (1989).

Applied chemistry, Theory and Practice. Witey Eastern Ltd, India, PP 94-97

Yasgajau, P. and Clifton, E.M. (1978): Food

Analysis, Theory & Practice. AVI Publisher. New York

Young, S.M. & Greaves, J.S. (1940). Influence

of variety and treatment on phytic acid content of wheat. *Food Research* 5, PP 103-105.