

Original Research Article

Multifactorial Analyses of Morphological Traits of Extensively Reared Helmeted Guinea Fowls *Numidia meleagris* in Kaduna and Katsina States of Nigeria

ABSTRACT

An experiment was conducted, using a total of 2005 sexually matured randomly-selected helmeted guinea fowl comprising of 5 varieties, to assess the magnitude of genetic diversity and the interdependence of morphological traits with the aim of reducing the number of characters for genetic and breeding purposes. Data was collected on some morphological traits from selected locations in Kaduna and Katsina states using a cross-sectional design and were analyzed using Principal Component Analysis (PCA) procedure and cluster analysis. The PCA showed that the first two principal components accounted for 53.10% of the total variation. Feather, ear lobe and beak colours, age, body and neck lengths were found to be responsible for most of the variations among populations of helmeted guinea fowl. Body, shank and neck lengths, helmet height and wing span were found to have a direct and positive relationship with body weight, while feather, breast and beak colours had a negative relationship. Cluster analysis revealed that the Ash variety was more closely related to the Pearl Black variety while Pearl Ash variety was more similar to the white. The Black variety was found to be more distinct from all the other four varieties. ~~It could be concluded that there are t~~Two main components accounted for variation in helmeted guinea fowl population in Kaduna and Katsina states. Furthermore, since variation in body weight was associated with body, shank and neck lengths, wing spans, helmet height and feather colour (varieties), these characters could be useful at the genomic level or as markers for improving body weight.

Keywords: *helmeted guinea fowl, breeding, markers, selection*

1. INTRODUCTION

Numida pitlorhycha and *Numida meleagris* are the two types of guinea fowl species in Nigeria. The former is indigenous to the rain forest zone of southern Nigeria while the latter is domiciled in the Northern part of Nigeria [1]. Guinea fowl play a vital role in socio-economic life of the rural populace. Apart from being a valued source of meat, eggs and income, they are also valued for their ability to serve as a watch animal and pest control [2]. The advantages of guinea fowl when compared with local chickens include: lower production cost, better meat quality, scavenging and survival abilities and resistance to common poultry parasites and diseases such as, Newcastle Disease and Fowl Pox [3]. Therefore, improvement in its production performance will increase household protein supply, combat protein-energy malnutrition and increase income.

Comment [SJN1]: Write out the aim of the study properly followed by the methodology

Comment [SJN2]: Arrange in alphabetical order

Multifactorial analyses of morphological traits have proven to be suitable in assessing the variation within a population and can discriminate different population types when morphological variables are considered simultaneously [4]. Various multivariate techniques such as Principal Component Analysis (PCA), cluster analysis, multivariate regression analysis, canonical correlation analysis and others have been applied for multivariate variable data analysis in the field of animal science and other related fields. PCA is designed to transform original variables into new, uncorrelated variables (axes) called principal components which are linear combinations of the original variables [5]. PCA has capacity to reduce the original variables measured into few components/factors to provide information on the most meaningful parameters which will describe a whole set affording data reduction with minimum loss of original information [6]. Independent factor scores derived from this multivariate technique can be used to estimate body weight [7], functional traits [8], as a selection criterion for the improvement of body size [9] and also used in reducing the number of individual variables in the prediction of genomic breeding values [10].

Cluster analysis includes a broad suite of techniques designed to find groups of similar items within a data set [11]. Hierarchical cluster methods produce a hierarchy of clusters from small clusters of very similar items to large clusters that include more dissimilar items. Hierarchical methods are particularly useful in that they are not limited to a pre-determined number of clusters and can display similarity of samples across a wide range of scales. Agglomerative hierarchical methods are particularly common in the natural sciences.

As a result of the paucity of information on the genetic structure and improvement, the present study was conducted to evaluate the magnitude of genetic diversity in helmeted Guinea fowl *N. meleagris* varieties and ascertain the most important characters causing variation. This study was also conducted to evaluate the interdependence of the morphological traits with the aim of reducing the number of characters for genetic and breeding purposes using multifactorial analyses.

2. MATERIAL AND METHODS

2.1 Description of the study areas

Kaduna and Katsina States are located in the North - Central Geo-political zone of Nigeria. Kaduna State is located in Northern Guinea Savannah. The global location of the state is between longitude 30 east of the Greenwich Meridian, latitude 0900 and 1130 north of the equator. The study was undertaken in the months of May and June, 2011. The average rainfall, relative humidity and ambient temperature ranged from 120.60 – 146.20 mm, 43.13 - 95.25 and 22.5 - 35.6°C, respectively, during data collection period (Nigerian Meteorological Agency). Katsina state is situated on 12°15'N, 7°30'E. The climate of the zone is tropical continental, having an annual rainfall of 198.3mm and a seasonal average temperature of more than 28°C.

2.2. Research design

A cross – sectional research design was employed in this study whereby all the information was collected once in the same period. Purposive sampling method was used in sampling and selection of the Local Government Areas (LGAs). The selected LGAs were Markarfi, Kubau, Ikara and Lere in Kaduna State; Charanchi, Kankara and Faskari in Katsina State.

2.3 Data collection

Morphological data were taken from a total of 2005 sexually matured randomly-selected birds from 5 varieties comprising of Ash (669), Black (552), Pearl Ash (395), Pearl Black

Comment [SJN3]: Too long, reduce to one paragraph

Comment [SJN4]: Lacks clarity

Comment [SJN5]: Add a map to show study sites

Comment [SJN6]: Were the birds pure breeds or a mixture of both pure breeds and cross breeds?

(342) and White (47). The sampled population comprised of males (49%) and females (51%). The ranges for body weight (kg) of helmeted guinea fowl across varieties also ranged from 1.30 to 2.10 and 1.30 to 2.00 in males and females, respectively.

The phenotypic characteristics and features of helmeted guinea fowl were collected using a modified version of structured survey form for morphological description [12]. Data on qualitative traits such as breast colour (BRC), shank colour (SHC), neck colour (NKC), eye colour (EYC), head colour (HDC), beak colour (BKC), eye lobe colour (ELC), inclination of wattle to upper jaw (WAI) and feather colour (FEC) and quantitative traits were collected using a standard adaptor [12].

Body weight of individual birds was determined using Mettler Toledo® top loading scale sensitive at 1g. Morphometric traits were determined in cm using a measuring tape and vernier caliper, according to adopted methods [1] as defined below:

- (a) Body length (BOL) was taken as the distance between the posterior end of the nasal cavity and the pygotyle.
- (b) The shank length (SKL) was taken as the distance from the tip of the metatarsus to the hock.
- (c) The shank circumference (SKC) was obtained using a measuring tape around the middle region of the shank.
- (d) Neck length (NKL) was determined as the distance between the atlas and the first thoracic vertebrae.
- (e) Wing span (WGS) was taken as the measurement between the coracoid and humerus to the tip of the phalanges.
- (f) Wattle length (WTL) was taken the longest distance between the attachment to the head and the tip of the wattle.
- (g) The number of tail feathers (NTF) was obtained by counting the mature tail feathers.
- (h) Helmet height (HEH) was taken as the distance from the base to the tip of the helmet using a pair of vernier calipers.
- (i) Helmet thickness (HET) was taken at the thickest portion of the helmet using a pair of vernier calipers.

2.4. Data analysis

The data on quantitative and qualitative traits obtained were subjected to principal component analysis using CANOCO version 4.5 for Windows [13]. The original values of predictor variables were transformed as $Y = \log(\text{original variable values})$ and also were standardized before the analysis so as to eliminate the effect of differences of scales of measurements. Hierarchical cluster method was used to describe qualitative variables that were similar between varieties of helmeted guinea fowl with the aid of dendrogram.

3. RESULTS

3.1. Determination of principal components

~~The percentages of total variation accounted for by the first four principal component axes are presented in Table 1. Principal component (PC) loadings show how the original variables influence the principal components. The cumulative percentage of variance accounted for, was the criterion that was used to select the principal factors. The cumulative percentage was determined as that which accounts for about fifty percent of the total variation. The proportion of variations explained in principal components in all the locations reduced with each loading thus indicating the relative importance of each loading. The first and second principal components explained 32.60 and 20.50% of the total variation~~

Comment [SJN7]: How?

respectively, and only the next two (PC 3 and 4) explained 21.20% of the total variation. The Eigen values of the four principal components in all the location was 1.00 (Table 1).

Table 1: Percentages of total variation accounted for by the first four principal component axes of ordination of varieties of helmeted guinea fowl

Location	Principal Axis	Proportion of total variance accounted for (%)	Cumulative percentage of variance (%)
Kaduna and Katsina States	1	32.60	-
	2	20.50	53.10
	3	15.00	68.10
	4	6.20	74.30

*Eigen value of the four components is 1.00

Comment [SJN8]: Sampled where?

3.2. Description of variations of quantitative and qualitative traits in all the study zones

A biplot of the first two principal components of variables responsible for the variations in helmeted guinea fowl in all study zones is shown in Figure 1. The directions of the arrows show the relative loadings of the variables on the first and second principal components. It can be observed that FEC and BKC have strong positive loadings on PC1 while BOL has negative loadings. SKL, HEH, NKL and WGS also had negative loadings on PC1 but were moderate. PC2 has strong positive loadings from AGE and negative loadings from NKC and ELC. The directions pointed by the variables indicate the squared multiple correlation with the principal components. The length of the vector is proportional to the squared multiple correlation between the fitted values for the variable and the variable itself.

Comment [SJN9]: Move to materials and methods





Figure 1: A biplot of the first two Principal components of helmeted guinea fowl *N. meleagris* in all the study zones

Comment [SJN10]: Which are?

3.3. Classification of the helmeted guinea fowl based on all qualitative morphological variables

The relatedness of the varieties of helmeted guinea fowl based on qualitative morphological variables is presented in Figure 2. The morphological features of Ash variety were more closely related to Pearl Black variety while Pearl Ash variety was more similar to the White variety. However, the Black variety was more distinct from all the other four varieties but there were some morphological features in this variety that may be in the other four varieties (Figure 2).



Key: VAR1=Ash, VAR2= Black, VAR3=Pearl Ash, VAR4= Pearl Black, VAR5=White

Figure 2: Dendrogram of five varieties of helmeted guinea fowl based on qualitative morphological features

Comment [SJN11]: From where?

3.4. Variation in varieties found in study locations

There were strong similarities in the varieties of helmeted guinea fowls present in Ikara and Kankara LGAs during the study (Figure 3). The varieties present in Lere LGA were distantly similar to the varieties found in Ikara and Kankara LGAs. However, the varieties of helmeted guinea fowls found in Kubau, Charanchi, Faskari and Markarfi LGAs were not similar (Figure 3).




Figure 3: Distribution of varieties of helmeted guinea fowls found in the study areas

Comment [SJN12]: Which are?

4. DISCUSSION

The higher proportion of variation explained by the first principal component as well as the reduced percentage of variance explained with each loading agrees with the report of others authors [7] indicating the relative importance of each loading.

Comment [SJN13]: Who reported what?

Positive loadings indicate a variable and a principal component are positively correlated: an increase in one results in an increase in the other. Negative loadings indicate the reverse, a negative correlation. In the entire study areas, traits such as FEC, BKC, BOL, AGE, NKL and ELC were responsible for most of the variations among populations of helmeted guinea fowl. The magnitude of variation for a given character dictates the kind of breeding plan that would be employed for the improvement [14]. Thus genetic improvement in body weight can be achieved faster by selecting for BOL, SKL, HEH, NKL and WGS which have direct and positive relationship with body weight. FEC, BRC and BKC were observed to be negatively correlated with body weight; thus indicating an inverse relationship between the traits and body weight.

FEC or plumage colour has been the basis for classification of the helmeted guinea fowl [15,16,17,18,19]. The presence of such noticeable variations in plumage colour could be as a result of their geographical isolation as well as the influence of artificial and natural selection [20,21].

Comment [SJN14]: Start with your own results

There are two factors to consider in the inheritance of feather colour in guinea fowl are their background colour and the amount of spotting or pearling that a bird has [22]. The Pearl Black variety has a dark grey background colour (GG) which is dominant over other background colours such as bluish grey background colour (II) in the Pearl Ash variety. If these birds are mated, each of the offspring in a single hatch will have a heterozygous GI genotype (one dark grey G - gene and one bluish grey I - gene). During further mating of these heterozygous GI helmeted guinea fowl cocks and hens, one chick in four would be pure Pearl Black (GG), one in four will be pure bluish grey (II) and two of the four chicks would be phenotypically Pearl Black but with heterogeneous dark grey / bluish grey genes(GI). These heterogeneous GI helmeted guinea fowls would be ready for further breeding, hence phenotypically, there would always seem to be a higher population of Pearl Black helmeted guinea fowl than Pearl Ash variety [22] which may be the reasons for dissimilarity in the types of varieties found in the different study locations.

Comment [SJN15]: Discuss your own results, do not speculate

5. CONCLUSION

The findings of this study suggest that the important characters accountable for diversity in the varieties of helmeted guinea fowl in the study zones could be grouped in four Principal Components with the first two components being more important than the other two. Since variation in body weight was associated with body, shank and neck lengths, wing spans and helmet height, therefore these characters could be useful at the genomic level or as markers for improving body weight. The varieties (FEC) were negatively correlated with body weight of the helmeted guinea fowl, thus indicating the importance of variety choice in selection for body weight. The varieties (FEC) of helmeted guinea fowl were not evenly spread across the entire study zones indicating that some varieties may not be found in some locations.

COMPETING INTERESTS

There are no competing interests

REFERENCES

1. Ayorinde KL. Characteristics and genetic improvement of the grey breasted helmeted guinea fowl, *Numida meleagris galeata pallas* in for growth and meat production. PhD Thesis submitted to University of Ibadan, Nigeria; 1987.
2. Smith J. Guinea fowl, Diversification Data Base, Scottish Agricultural College. 2000. Available: <http://www.sac.ac.uk/management/external/diversification/Tableofcontents>.
3. Dieng A, Gue'ye EF, Mahoungou-Mouelle NM and Buldgen A. Effect of diet and poultry species on feed intake and digestibility of nutrients in Senegal. *Int Netw Fam Poult Dev Newsl*. 1998; 8:4–9.
4. Yakubu A, Ibrahim IA. Multivariate analysis of morphostructural characteristics of Nigerian indigenous sheep. *Ital J Anim Sci*. 2011; 10(2):83–86.
5. Shrestha S, Kazama F, Nakamura T. Use of principal component analysis, factor analysis and discriminant analysis to evaluate spatial and temporal variations in water quality of the Mekong River. *J Hydroinform*. 2008; 10(1):43–56.
6. Helena B, Pardo R, Vega M, Barrado E, Fernandez J, Fernandez L. Temporal evolution of ground water composition in an alluvial aquifer (Pisuerga River, Spain) by principal component analysis. *Water Res*. 2000; 34(3):807–816.
7. Yakubu A, Ayoade JA. Application of principal component factor analysis in quantifying size and morphological indices of domestic rabbits. *Int J Morpho*. 2009; 27(4):1013–1017.
8. Karacaoren B, Kadarmideen H. Principal component and clustering analysis of functional traits in Swiss dairy cattle. *Turk J Vet Anim Sci*. 2008; 32:163–171.

- 254 9. Pinto L, Dacker I, De Melo C, Ledur M, Coutinho LL. Principal components analysis
255 applied to performance and carcass traits in chicken. Anim Res. 2006; 55:419–425.
- 256 10. Macciotta N, Gaspa G. Use of principal component and factor analysis to reduce the
257 number of independent variables in the prediction of genomic breeding values. Ital J
258 Anim Sci. 2009; 8(2):105–107.
- 259 11. Holland SM. Cluster Analysis. 2006. [http://strata.uga.edu/software/pdf/-](http://strata.uga.edu/software/pdf/-clusterTutorial.pdf)
260 [clusterTutorial.pdf](http://strata.uga.edu/software/pdf/-clusterTutorial.pdf)
- 261 12. Food and Agriculture Organisation. Animal Genetic Resources Data Banks. 3:
262 Descriptor lists for poultry. FAO Animal Production and Health No.59/3, Rome, Italy.
263 1986.
- 264 13. Braak C, Smilauer P. CANOCO Reference manual and CanoDraw for Windows
265 User's guide: Software for Canonical Community Ordination (version 4.5). In:
266 Microcomputer Power, Ithaca, NY, USA; 2002.
- 267 14. Ibe SN. An Introduction to Genetics and Animal Breeding. Publ. Longman Nigeria
268 Plc.;1998. ISBN 978–139–78–29.
- 269 15. Ghigi A. On the Inheritance of colour in the guinea fowl. Proceedings of the 2nd
270 World's Poultry Congress, May 10-18, 1924. Barcelona; 18-19. 1924.
- 271 16. Ghigi A. The breeding of the guinea fowl in Italy. Proceedings of the 13th World Poultry
272 congress (Kiev), 137. 1966.
- 273 17. Ogundipe SO. The raising of guinea fowl. Extension Guide No. 83. Poultry Series No.
274 10 NAERLS, ABU, Zaria, Nigeria. 1976.
- 275 18. Ogundipe SO. Potential for the development of commercial guinea fowl production in
276 Nigeria. The helmeted guinea fowl varieties (*Numida melagris galeata* Pallas) in
277 Nigeria. Edited by Ayeni J.S.O., Olumu, J.M. and Aire T.A. Pp 66 – 72. 1983.
- 278 19. Ayorinde KL. The spice of life. The 71st Inaugural Lecture of University of Ilorin. Pp 12
279 – 17. 2004.
- 280 20. Duguma R. Phenotypic characterization of some indigenous chicken ecotypes of
281 Ethiopia. Livest Res Rural Dev. 2006; 18(9).
- 282 21. Mogesse H. Phenotypic and Genetic Characterization of indigenous chicken
283 populations in Northwest Ethiopia. PhD Thesis submitted to the Faculty of Natural and
284 Agricultural Sciences, Department of Animal, Wildlife and Grassland Sciences,
285 University of the Free State, Bloemfontein, South Africa. 2007.
- 286 22. Smith J. Colour Genetics of Guinea fowl. 2013. [Online]. Available:
287 <http://www.guineafowlinternational.org/articles/genetics.php>.
288