

A Review of Household Products Induced Secondary Organic Aerosols Formation in Indoor Environment

Adeniran, J.A.¹, Jimoda, L.A.^{1*}, Sonibare, J.A.²

¹Environmental Engineering Research Laboratory, Ladok Akintola University of Technology, Ogbomoso, NIGERIA, ² Environmental Engineering Research Laboratory, Obafemi Awolowo University, Ile – Ife, NIGERIA,

^{1*} - Corresponding Author – Jimoda, L.A. (lukumanjimoda@yahoo.com)

Abstract

This paper reviews the process of formation of secondary organic aerosols (SOAs) in the indoor environment. SOAs have been found to have more alarming health effects on humans and environment. To be able to assess and quantify the risk, it is important to have a comprehensive information on their mechanism of formation. The photochemical reactions of volatile organic compounds (VOCs) released from the use of the consumer products with ozone (O₃), hydroxy group (OH), and oxides of nitrogen NO_x were considered. Environmental factors such as temperature, relative humidity and light were also found to have impact on the physical and chemical composition of the resulting SOAs. Gas – phase partitioning was identified as responsible for the formation of SOAs. Common health effects associated with SOAs and mitigation measures were suggested.

Keywords: Aerosols, formation, indoor, environment, consumer products

1.0 Introduction

A variety of inorganic and organic chemical compounds are present in the particulate phase of aerosols from household consumer products. The organic fraction in aerosols can account for a 20 - 90 % of the finest fraction [1] and, therefore, the knowledge of this fraction is important to prevent human health risks associated with them [2]. Both inorganic and organic aerosols can be directly emitted (primary aerosols) or can be formed in the atmosphere as a consequence of multiple physical and chemical processes - secondary aerosols [3 -4]. Human exposure to aerosols takes place both in the indoor and outdoor environment [5-7]. However, because people spend majority of their time in the indoor environment, exposure to aerosols and the consequent effects on human health have been the topics of research interest in recent years [8-11].

Domestic aerosol products such as insecticides, air fresheners and perfumes are commonly used all over the world. Domestic aerosols are consumer aerosol products that are dispersed under pressure from disposable containers. Aerosol product has three main components: including the active ingredient(s), the propellant, and miscellaneous additives such as solvents, plasticizers, and emulsifiers which are used to improve the product [12]. Active ingredients in the domestic aerosol products include terpenes. Terpenes and propellants in domestic aerosol products are volatile organic compounds (VOCs) [12-13].

The use of air fresheners and perfumes indoors can cause an increase in the airborne concentrations of gaseous and particulate species in the indoor environment. Terpenes are unsaturated volatile organic compounds that occur in nature and are presumed to be safe. However, many terpenes have been found to have serious health effects [14][15] while concentrations of terpenes emitted from air fresheners are far greater than those found in nature [16]. Terpenes like α -pinene, d-limonene, linalool, and citronellol react with transient compounds in the air [17]. Ozone-terpenes reactions produce carbonyls such as formaldehyde and acetaldehyde, organic acids, hydrogen peroxide, secondary organic aerosol, and hydroxyl (OH) radical [18]. Subsequent reactions of OH group with the terpenes can generate additional particulate matter and VOCs [19]. Investigations have shown that the reaction between ozone and terpenes from air fresheners and perfumes could result in the formation of particulate matter in indoor atmosphere [14-23].

Most propellants in spraying products are often viewed to be chemically unreactive. However, European Commission report stated that butane and isobutane(used as propellants) in consumer products may contain 1,3-butadiene at a concentration greater than or equal to 0.1% [24]. [25]established in their study that SOAs may be formed when 1, 3, butadiene photodecompose in the presence of air containing nitric oxide. The aim of the present work is to present a basic overview of how secondary secondary organic aerosols are formed from spraying consumer products and their implications on humans and indoor environment. Some of the existing literatures are reviewed therein to give a clearer picture of the factors and mechanisms involved in SOAs formation.

2.1 Organic Compounds in Indoor Environment

According to [26] [27], organic compounds present in indoor air include very volatile organic compounds (VVOCs), volatile organic compounds (VOCs), semi-volatile organic compounds (SVOCs) and non-volatile organic compounds (NVOCs), or organic compounds associated with particulate matter or particulate organic matter (POM) [26]. Table 1 shows the detailed classification of inorganic pollutants as adapted from [26]. VOCs are carbon-based chemicals which are volatile at normal room temperatures, and whose vapours are readily inhaled [28].

Table 1: Classification of Organic Pollutants

Description	Abbreviation	Boiling Point Range (°C)	Example Compounds
Very Volatile (gaseous) organic Compound	VVOC	< 0 to 50 – 100	Propane , Butane, Methyl chloride
Volatile Organic Compounds	VOC	50 -100 to 240 – 260	Formaldehyde, d – limonene, toluene, ehanol(ethyl alcohol), 2 – propanol (isopropyl alcohol, hexanal)

Semi volatile organic compounds	SVOC	240 – 260 to 380 – 400	Pesticides(DDT, Chlordane, plasticizers(phthalates), fire retardants)
---------------------------------	------	---------------------------	---

Source: WHO [24]

VOCs is an important group of indoor air pollutants whose concentration levels are elevated indoors [24]. Typical boiling points of VOCs are in the range of 50-100 ° C [28]. According to Wolkoff [30], VOCs comprise a range of substances including alkanes, alkenes, aromatic hydrocarbons, chlorinated hydrocarbons, esters and alcohols (derived in part from microbiological sources), aldehydes and ketones (many from precursor terpene compounds and fatty acids). Products used at home can release VOCs into indoor air when used or stored [31] [32] [33]. These VOCs released as a result of solvent and propellants used in consumer aerosol products has been referred to as reactive organic compounds (ROGs) because of their high reactivity with the major oxidizing species (O_3 , $OH^\cdot$, and NO_3) present in the atmosphere [34]. ROGs with double bonds (Alkenes, cycloalkenes, dienes, and terpenes) are more reactive and can thus react with the three commonest oxidizing agents present in the atmosphere. [34]. The reactivity of ROGs towards the major oxidizing species are as shown in Table 2

Table 2: Reactivity of ROGs towards oxidizing species in air (units: $cm^3 \text{ molecule}^{-1} s^{-1}$ at 289K)

	Ozone	OH radical	NO_3 radical
Alkanes, Cycloalkanes	$\leq 10^{-23}$	$0.3 - 8.0 \times 10^{-11}$	$\leq 10^{-17}$
Oxygenated Aliphatics	$\leq 2.2 \times 10^{-21}$	$0.2 - 6.0 \times 10^{-11}$	$\leq 1.4 \times 10^{-16}$
Aromatics	$\leq 6 \times 10^{-11}$	$0.1 - 6.0 \times 10^{-11}$	$\leq 10^{-17}$
Alkenes, cycloalkenes, other olefins	$2 \times 10^{-18} - 1.5 \times 10^{-15}$	$0.8 - 12 \times 10^{-11}$	$6 \times 10^{-17} - 3 \times 10^{-11}$

Source: Grosjean and Seinfeld [33]

EPA's Total Exposure Assessment Methodology (TEAM) studies found levels of about a dozen common organic pollutants to be 2 to 5 times higher inside homes than outside, regardless of whether the homes were located in rural or highly industrial areas [36]. Additional EPA TEAM studies [37] established that while people are using products containing organic chemicals, they can expose themselves and others to very high pollutant levels, and elevated concentrations can persist in the air long after the activity is completed. The level of VOCs present in indoor environments is also affected by the phenomenon of so-called secondary emissions [27].


VOCs emitted from the use of consumer spraying products are converted by reaction in photochemical oxidizing agents to form Secondary Organic Aerosols(SOAs) over time. Secondary emission is any process that releases new airborne pollutants from existing sources, changes the total emitted mass of existing pollutants, or results in chemical reactions between compounds on surfaces and those in the air [22].

2.2 Photo – Chemical Reactions of Volatile Organic Compounds

VOCs in consumer spraying products such as terpenes react with $\text{OH}^\cdot$, NO_x and O_3 radicals in the indoor environment to give stable and condense phase compounds which may include the SOAs. Terpenes/ Reactive radicals reaction which results in the formation of SOAs in the presence/ absence of light are highlighted in the following subsections.

2.2.1 Reaction with Hydroxyl Radicals ($\text{OH}^\cdot$)

Hydroxyl radicals have been known to have a very high reactivity with most indoor atmospheric compounds [38 - 40]. They are present in indoor air at concentrations sufficient enough to react with VOCs [41]. Ozone reacts with unsaturated volatile organic compounds such as d-limonene and α -pinene to give hydroxyl radical as an intermediate product[42].Hydroxyl radicals in turn react with VOCs to form Toxic Air Contaminants such as formaldehyde and acetyldehyde [41] [33]. Terpenes have a very high reactivity with $\text{OH}^\cdot$ radicals that the oxidation initiated by $\text{OH}^\cdot$ produces carbonyls and hydroxycarbonyls, such as glyoxal, methylglyoxal, and glycolaldehyde [43].Taking limonene as an example, the reaction could take the following pattern:



2.2.2 Reaction with Ozone (O_3)

Ozone enters indoor environments with outdoor air [44]and typically is present indoors at levels that are 10 to 50% of concurrent outdoor levels [45]. Typical indoor-outdoor ozone ratios are 0.2 - 0.7 [45], so peak indoor levels in the range of 30-100 ppb are expected when ambient pollution levels are high [41]. The concentration of O_3 in the indoor air will have a direct effect with number concentration of SOA formed. This can be depicted in Figure 1.





Figure 1: Effect of O_3 concentration on particle concentration

Source:Nøjgaard *et al.* [46]


The figure shows the total particle concentration following 60 min of ozonolysis of d-limonene and α -pinene in the absence of NO_2 . The experiments were conducted in 1000 l teldar bags to study the effects of the nitrogen dioxide (NO_2) concentration on particle formation during

ozonolysis of two abundant monoterpenes indoors, α -pinene and d-limonene at 21 ± 2 °C and atmospheric pressure. The nucleation potential of NO_2 was found to be much lower than ozone. Therefore, the role of O_3 in SOA formation was more significant as compared to NO_2 . The result indicated that particle concentrations from ozonolysis of the monoterpenes do not depend on O_3 concentration linearly [46]. Table 3 summarizes the half-life for each of a number of processes that affect the ozone concentration indoors. Findings have also revealed that ozone are released indoor as a result of the use of electrical equipment [11, 47 -50]. In the presence of light, O_3 photodecompose to give $\text{OH}^\cdot$ radical.



M is a neutral third body usually N_2 or O_2 [49]

Terpenes, such as d-limonene and α -pinene, are common VOC constituents of products that are used indoors. Constituents of consumer household spraying products react with ozone, which is considered a good initiator for indoor gas – phase oxidation process [52] [32] [53] [41]. The type of SOAs formed from this process is termed ozonolysis product [54-59].



O_3 reacts considerably faster with VOCs containing unsaturated carbon-carbon bonds. [60] pointed out that ozonolysis products also known as ozonides formed from ozone- terpenes reaction are unstable and may easily decompose. The unstable products were termed Criegee intermediates ($\text{CH}_2\text{OO}^\cdot$) [61] [62]. Criegee intermediates may undergo decomposition such that it will yield hydroxyl radical. [60]. Aschmann *et al.* [58] described the process using hydroperoxide channel. The reactions are as follow:



On reaction with water vapour, it yields α -hydroxyhydroperoxides




Table 3: Half-Life of Indoor Ozone for a Number of Processes

Process	Rate coefficient	Range of Half – Lives (min)
Air Exchange rates	$0.2\text{-}2.0 \text{ (h}^{-1}\text{)}$	21 – 210
Removal by surfaces	$1.4 – 7.6 \text{ (h}^{-1}\text{)}$	6 – 30
Reaction with limonene	$4.9 \times 10^{-6} \text{ (ppb}^{-1}\text{s}^{-1}\text{)}$	60 – 500
Reaction with α -pinene	$2.1 \times 10^{-6} \text{ (ppb}^{-1}\text{s}^{-1}\text{)}$	130 – 1100

Source: EC [61]

2.2.3 Reaction with NO_x

SOAs can either be formed in the presence of NO_x or in their absence [64-67]. SOAs growth is affected by NO_x concentration in indoor air [41] [68]. Few investigations have shown that more SOAs are formed under high NO_x conditions [40 – 43, 67]. However, more experiments have shown that low NO_x condition will favour SOAs formation [68 -73]. The proportionality of low NO_x has also been investigated and buttressed using computational simulation [74]. EC [61] showed that when the concentration of NO₂ and O₃ are high, the reaction could proceed as follow:




And also,


NO₃ reacts with rapidly with NO to give NO₂:




When NO concentration is low,



N₂O₅ could react with moist surface to give HNO₃ as indicated in the following reaction



In the presence of light, the reaction could also follow this pattern:



2.3 Secondary Organic Aerosols (SOA)

The aerosols directly emitted into the atmosphere from the source are named primary aerosols while those formed through a series of reactions in the atmosphere are called secondary aerosols [75-77]. SOA is an organic liquid or solid particle suspended in the air, which is formed through photochemical reactions of VOCs or primary aerosols. Investigations [14 -23] have shown that the reaction between ozone and terpenes are responsible for formation of particulate matter indoor with limonene and α – pinene as most reactive compounds. Clausen *et al.* 2005 [20] investigated indoor ozone/terpenes reaction in a ventilated apartment under normal conditions. The result of their investigation showed that even under normal condition of temperature between 22 – 25°C, relative humidity of between 32 – 40% and average air exchange rate of between 0.7 and 0.8/h, high particle concentration could be formed. Some of the previous studies on the formation of secondary organic aerosols in the indoor environment are as presented in Table 4.

2.4 Factors Affecting Secondary Organic Aerosol Formation

Aside the reactive oxidizing agents, atmospheric conditions are factors that may influence the physical and chemical composition of SOAs [78]. Concentration of the reactive groups and terpenes have been acknowledged to have influence on growth and number concentration of SOAs [22] [55]. Other factors that affect SOAs formation discussed in the subsections below.

2.4.1 Light

Ozonolysis proceeds faster in the presence of illumination than in darkness. Oxidation of terpenes are mostly described to occur in the presence of light [77-81]. However, the investigation of Cao and Jang [66] and Presto *et al.* [69] showed that the presence of UV radiation has a negative

effect on the ozonolysis of terpenes, especially α -pinene. Though, it enhances the photolysis of H_2O_2 to OH radicals [80]. The investigations of Leungsakul, *et al.* [81 - 82] on limonene/ozone reaction in a film chamber in both the presence of light and darkness indicated that SOA are formed in both cases.

Table4: Summary of some of the previous studies on SOA formation in Indoor Environment

Study	Location	Type of Reaction	Sampling method	Microclimatic Condition (Temperature, RH and AER)
Chen and Hopke [55]	Dynamic chamber (2.5 m ³)	linalool ozonolysis	Scanning Mobility Particle Sizer (SMPS) coupled with Condensation Particle Counter (CPC)	(22–23 °C); 0.67 ± 0.01 h ⁻¹ AER;
Matsunaga et al. [85]	A 5900 Litre PTFE environmental chamber	OH-radical initiated reactions of linear alkenes in the presence of NO _x	Thermal desorption particle beam mass spectrometer (TDPBMS)	<1% RH; at 25 °C;
Vivanco et al. [86]	EUPHORE photoreactor	Mixture of 1,3,5-trimethylbenzene, o-xylene, octane and toluene) with an oxidant .	TEOM and SMPS	10-20% RH;
Hatfield and Hartz [87]	5.5 m ³ Teflon smog chamber (Welch Fluorocarbon)	Mixtures containing reactive and non-reactive volatile organic compounds (VOCs)	ATSI SMPS coupled with CPC	23 -29°C; 1 - 4% RH
Parikh et al. [88]	Outdoor aerosol smog chamber	toluene oxidation	SMPS coupled with CPC	15 – 37°C; 8-92%RH;
Ng et al. [70]	Environmental chamber	Photooxidation of terpenes		25°C; < 6% RH
Song et al. [89]	Environmental chamber	Dark ozonolysis of α -pinene		28°C; ,10% RH
Shilling et al. [90]	Environmental chamber	Dark ozonolysis		25°C; 40% RH
Presto and Donahue [91]	Environmental chamber	UV		23°C; <10%
Lee et al. [47]	Carnegie Mellon University smog chamber, a 12m ³ Teflon	Ozonolysis of monoterpenes	aerosol mass spectrometer (AMS) + CPC	70 - 90°C;

	reactor and thermodenuder (TD)			
Lauraguais et al.[92]	8 m ³ evacuable Plexiglas (PolyMethylMethacrylate PPMA) reaction chamber	gas-phase reaction of hydroxyl radical with syringol (2,6-dimethoxyphenol)	(SMPS GRIMM, CPC5.404)	21 - 23°C; 2 - 4% RH.
Updyke et al.[93]	a 17 L glass flow tube reactor	Photo-oxidation of VOCs from biogenic and anthropogenic sources	SMPS	20 - 24°C; <2% (RH), and 750±50 mbar pressure
Stewart et al. [94]	80L collapsible Teflon Static reaction chamber	Gas phase reactions of selected monoterpenes and cyclo-alkenes with ozone and NO ₃ radical		25 - 27°C;
Alfarra et al. [95]	27m ³ transparent Teflon® bag	Photooxidation of anthropogenic and biogenic precursors	Aerodyne Aerosol Mass Spectrometer (AMS), SMPS and CPC.	-
Angove et al.[25]	18.1m ³ CSIRO indoor environmental chamber	photodecomposition of 1,3-butadiene in air containing nitric oxide	Impactor jet connected to CPC and TSI 3030 electrical aerosol size analyser	21°C; < 6% RH
Broske et al. [96]	Aerosol flow tubes	reactions of O ₃ with α -pinene, limonene or catechol and OH radicals with toluene or limonene.	SMPS-system (TSI 3071 electrostatic classifier and TSI 3022A condensation particle counter).	25°C; 50 – 57% RH
Cao and Jang, [68]	2m ³ indoor Teflon film chamber	Toluene photooxidation under various NO _x conditions	SMPS and CPC	22 - 32°C; 15 – 115% RH
Carter et al.[97]	two collapsible 90 m ³ FEP Teflon film reactors	photolysis reactions of NO _x and formaldehyde	SMPS coupled with CPC and tandem differential mobility analyzer (TDMA)	5 - 45°C; unhumidified air
Chen and Chang, [98]	2m ³ indoor Teflon film chamber equipped with UV- Visible lamps	Photooxidation of a mixture of dimethyl sulfide and isoprene	Scanning Mobility Particle Sizer (SMPS) combined with	21 - 24°C; 12 - 80% RH

			a condensation nuclei counter(CNC).	
Cocker III et al. [99]	dual 28 m ³ Teflon reactor.	α -pinene/ozone system	Scanning electrical mobility spectrometers (SEMS), equipped with condensation particle counter	20 – 50°C; <2 -89% RH
Coleman et al. [21]	a Teflon-lined, 198-L rectangular chamber	Ozone Initiated Reactions With Terpenes – Rich Household Products	SMPS coupled with OPC	23.0±0.5°C; 3 h ⁻¹ AER
Fadeyi et al. [123]	236 m ³ field environmental chamber (FEC)	ozone of outdoor origin reacting with limonene of indoor origin.	Fast Mobility Particle Sizer (FMPS)	23°C; 1 – 2 h ⁻¹ AER
Huang et al. [53]	18.26 m ³ stainless-steel environmental chamber	Ozonolysis of pollutants from cleaning products	a scanning mobility particle sizer (SMPS) system with a differential mobility analyzer coupled to a condensation particle counter	ACH of 0.36 h ⁻¹ , RH of 50% and T of 23 °C.
Huang et al. [76]	100 L small scale chamber	Ozone initiated SOA emissions from painted wooden panels		25°C; RH of 50%; ACH of 0 – 0.5 h ⁻¹ ;
Huang et al. [131]	aromatherapy spa	Aromatic and volatile organic compounds were the dominant handling materials.	SMPS coupled with CPC	0.9 – 1.3 h ⁻¹ AER
Irei et al. [100]	8m ³ indoor YorkU smog chamber	Photooxidation of toluene	SMPS	25 °C Two reaction chambers (1 and 2) and RH < 6%
Kleindienst et al. [65]	The reaction Chamber is a 14.5-m ³ parallelepiped, stainless-steel chamber with interior walls fused with a 40µm TFE Teflon coating.	isoprene + OH reaction in the absence of NO _x	SMPS and Volatility Differential Mobility Analysis technique.	50 - 250 °C
Langer et al. [22]	Two stainless steel reaction chambers (1 and 2)	ozone/limonene reactions in indoor air	SMPS coupled with CPC	23±2°C; 33±3% RH

	and 14 m ³)			
Lee et al. [101]	Two suspended 28m ³ Flexible Teflon chambers	The ozonolyses of six monoterpenes (α -pinene, β - pinene, 3-carene, terpinolene, α -terpinene, and myrcene), two sesquiterpenes (α - humulene and β - caryophyllene), and two oxygenated terpenes (methyl chavicol and linalool)	SOA yield was calculated theoretically	20 $\pm$ 1°C; and <10% RH
Lu et al. [117]	The cuboid reactor, with a volume of 2m ³	m-xylene/NO _x photooxidation system	SMPS; SOA yield theoretically calculated according to Pankow (1994)	10 - 60°C; RH < 20%
Na et al. [109]	18m ³ Teflon environmental chamber	Reaction of styrene with ozone in the presence and absence of ammonia and water	Scanning Electrical Mobility Spectrometer (SEMS) with differential mobility analyzer (DMA)	20 $\pm$ 1°C; 0 – 56% RH
Nøjgaard et al. [46,57,58]	1000 L Tedlar bags	ozonolysis of α -pinene and d-limonene	A differential mobility particle sizer (DMPS) and a CPC	21 $\pm$ 2°C; RH < 1%
Qi et al. [102]	Dual 90m ³ Teflon reactors	m-xylene photooxidation	Aerosol Particle Mass Analyzer (APM), Scanning Mobility Particle Sizers (SMPS) and a custom-built Hygroscopic Tandem Differential Mobility Analyzer (HTDMA).	0 - 40°C;
Rickard et al. [103]	27m ³ Teflon bag	1,3,5- trimethylbenzene photooxidat	Chemical Ionisation	20 °C; 48 – 62% RH;

		ion	Reaction Time-of-Flight Mass Spectrometer (CIR-TOF-MS), Condensation Particle Counters (CPCs) and a Scanning Mobility Particle Sizer(SMPS)	
Rossignol et al. [104]	4.2m ³ stainless steel chamber in 200m ³ EuphorePhotoreactor	limonene ozonolysis	Gravimetric analysis using 47mm quartz fibre filter	23 °C; 0 – 80% RH;
Salo et al. [105]	The AIDA chamber consists of an aluminium vessel of 84.5m ³	ozonolysis of tetramethylethylene (TME)	AIDA and SAPHIR	-90 °C – 60°C; 0.01–1000hPa
Sarwar et al. [40]	11m ³ stainless steel chamber.	ozone–limonene reaction	A commercial corona discharge ozone generator (Living Air, BORA-IV) Particle Measuring Systems Inc., LASAIRs—Model 1002; TSI TM P-TRAK TM—Model 8525.	23 °C -25 °C; 39–67% RH
Sarwar et al. [123]	11m ³ stainless steel chamber.	homogeneous reactions between ozone and pinene,	A commercial corona discharge ozone generator (Living Air, BORA-IV) Particle Measuring Systems Inc., LASAIRs—Model 1002;	24 °C -33 °C;

			TSI TM P-TRAK TM—Model 8525.	
Sarwar et al. [124]	11m ³ stainless steel chamber.	homogeneous reactions between O ³ and terpenes	A commercial corona discharge ozone generator (Living Air, BORA-IV) Particle Measuring Systems Inc., LASAIRs—Model 1002; TSI TM P-TRAK TM—Model 8525.	23 °C -25 °C;
Sleimann et al. [125]	an 18-m ³ environmental chamber	Ozone initiated reactions with nicotine and secondhand tobacco smoke	SMPS that consisted of electrostatic classifier coupled to an ultrafine CPC	25 – 150°C; 0 – 50% RH

Shilinget *al.* [85] also showed that SOAs are formed from the dark ozonolysis of α - pinene.

2.4.2 Temperature

Ambient temperature has been found to have clear effect on the formation of secondary organic aerosols in terms of particle volume and chemical composition[86]. It has been established that SOAs formation is temperature dependent. The studies indicated that increase in temperature is directly proportional to secondary organic aerosols formation [106 -110]. Studies have also established that ozonolysis of terpenes can occur at room temperature [85 -87;90 -91].

2.4.3 Relative Humidity (RH)

RH affects the formation, size and deposition of SOAs [92]. The absorption of water by aerosols affects their physicochemical properties such as particle size[93], phase, deposition [94] , atmospheric life time and chemical reactivity [80]. The hygroscopicity of aerosols depends on their chemical compositions [80]. Na *et al.* [110] have established that the presence of water vapour may inhibit the the formation of SOAs. The concentration of SOAs may depend on the water composition and distribution of the final oxidation products [110]. Under high humidity condition, SOAs adsorb water from the atmosphere and thus inhibit the reaction surface[95]. In contrast, investigation of the effect of RH on the formation of SOAs generated from ozonolysis of limonene in an environmental chamber by [96] indicated that the number concentration and growth of SOAs were enhanced as RH increases.

2.4.4 Air Exchange Rate (AER)

Air exchange rate is an important factor that determines the concentration of ozone indoor which has a typical half life of between 7 to 10 minutes [43]. Sarwaret *al.* [43] found out in their experiments that increase in air exchange rate increases the concentration of hydroxyl radicals in the indoor environment with alkene/ ozone reactions dominating. Weschler and Shields [97] studied the effect of air exchange rate on SOAs in indoor air. They observed that at higher AER, the mass and number concentrations of particles were smaller and no excess particle formation was detected. The study and that of Weschler [75] indicated that lower air exchange rates will favour accretion and coagulation of smaller particles and that at higher air exchange rate, the small sized particles are likely to be formed within a short period.

2.4.5 Time

The residence time of air in the indoor environment is a factor that affects the growth of aerosol particle (accretion) and coagulation processes [114]. The more the time, the higher the concentration of airborne SOAs formed. High residence time and low air exchange rates are factors that may account for an increment in SOAs particle mass [115]. SOAs will continue to be formed in the indoor environment due to consumer product usage in as much as there are terpenes and the reactive group are in the indoor atmosphere. Waringet *al.* [116] noted that nucleation of particles will commence immediately terpenes are introduced into the indoor environment and that the time of reaction is almost inconclusive as new particles may be formed. Table 3 highlights the lifetime of reaction of some common terpenes, their reaction rate and OH⁻ formation as adapted from Fick [117].

Table 5: Reaction rate, Lifetime and OH Formation from Ozonolysis of Common Terpenes

Monoterpenes	Reaction rate $10^{18} \times k$ (at 298 K) ($\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	Lifetime	OH radical formation (M)
α -pinene	86.6	4.6 h	0.85
β -pinene	15	1.1 day	0.35
Limonene	200	2h	0.86
Δ^3 -carene	37	11h	1.06
Δ^2 -carene	230	1.8h	No reported value
Myrcene	470	50min	1.15
Terpinolene	1880	12 min	1.03
Camphene	0.9	18 days	< 0.18
Sabinene	86	4.6 h	0.26
β -Phellandrene	47	8.0h	0.14

Source: Fick [117]

Lifetime in Table 5 indicates time for decay of compound (terpenes) (37%) of its initial concentration at an ozone concentration of 28 ppb. Unit M is the molar yield of OH⁻.

2.5 Mechanisms of SOA Formation

Gas – particle partition stage is the second stage that will follow the oxidation of VOCs in the formation of SOAs [118]. Active ingredients in consumer product will react with the reactive

species in the gas phase followed by formation of new particles through coagulation, nucleation or accretion. SOAs are formed due to gas to particle conversion [119].

2.5.1 Gas Phase Reaction

Volatile components of consumer household spraying products (propellants and the active ingredients) enter the gas phase during and after use [41]. The gas phase reaction between terpenes and the reactive species such as O_3 , $OH^\cdot$ radical and NO_x will lead to the formation of lower vapour pressure products [103]. Investigations have shown that the gas phase reaction between terpenes and ozone contributes significantly to the growth of SOAs in the indoor atmosphere [95] [68] [104] [105] [106] [107] as ozone is usually removed in the indoor air as a result of the gas phase reaction [108].

2.5.2 Particle Phase Reaction

Gas/ Particle phase partitioning is a major phenomenon in SOA mass formation and occurs throughout the reaction time [97]. Low volatile oxidation products from gas phase reaction yield secondary organic aerosols either by new particle formation [109] [110], or through condensation and nucleation of the already existing particles [81-82, 111]. The nucleation and growth of aerosol particles can occur anywhere in the atmosphere [112 - 113] and most occurs at the beginning of ozone/ terpene reaction [45] [97]. They are presumed to be responsible for the number concentration of SOAs that will be formed [114]. A major product of the particle phase reaction between ozone and terpenes is formaldehyde [115]. According to European Commission [63], gas to particle conversion has two perspectives including (i) a sink mechanism where reactive carbon is lost from the indoor atmosphere and (ii) the accretion of oxygenated material onto existing particle. These are summarized in Figure 2. Gas-particle partitioning of products is usually calculated using the theory of Pankow [154] which assumes that particulate organic matter (POM) is a single, liquid organic phase [85, 155].

3.1 Impacts of SOAs on Human Health

Epidemiological studies have revealed that increase in exposure to aerosols is directly proportional to morbidity and mortality rates [116] [117] and SOAs are believed to have more alarming effect on human health than the primary organic aerosols [77] [118]. They have been associated with respiratory ailments [119] [120] such as Asthma [121], cardiovascular health effects [122] [123] [106] and psychological stress [124]. They have also been noted for inflicting upper airway and pulmonary ailments on individuals [125]. Headaches, fatigue, eye, nose and throat irritation are also among the list of health impacts SOAs may have on humans [126].




Figure 2 : Route of Formation of Secondary Organic Aerosols
Source: Wang [11]

3.2 Impact of SOAs on Indoor Environment

Secondary organic pollutants may be airborne for a long period of time. Singer *et al.* [30] noticed in a study that ozone consumption and OH radical persisted in the indoor air for about 10 – 12 hours following a brief cleaning exercise. SOAs may also be deposited on walls, and filters of the ventilation system[38]. Weschler[41] established that SOAs may be responsible for soiling of walls. In same experiments, it was noticed that the white board in the ozone/ limonene office turned slightly yellow due to SOAs deposition.

3.3 Mitigation Measures

Exposure to Secondary Organic aerosols from the use of consumer products can be reduced by removing ozone from ventilation air and limiting the use of terpenes emitting products. Proper and constant hygienic measures should be taken to replace the use of fragranced products. The use of room air cleaners has also been proved to be an effective means to improve indoor air quality [127] [153]. Fadeyi *et al.* [132] suggested that high efficiency particulate filters could also be used to remove SOAs and ultrafine particles. The use of household and office electrical equipment that produce ozone should be minimized. Good ventilation and well-designed HVAC (Heating, Ventilation and Cooling) system should be given a top priority when designing a building. Regulation and legislation on manufacturing and declaration of contents in household products should be enforced as some contents are not often disclosed [149 -151]. Bernstein *et al.* [118] opined that people should be informed about the existing air quality standards and rational environmental control.

4.0 Conclusion

The use of household spraying products may result in the increase in concentration of VOCs especially terpenes in the indoor environment. Terpenes react with the reactive radicals such as O_3 , $OH^\cdot$ and NO_x in the presence/ absence of light to form SOAs which are presumed to have more adverse health effects on humans than the terpenes themselves. It is therefore recommended that a sensitational measure should be put in place to educate consumers on the use of household spraying products. Legislations should be made in such a way that ingredients used in manufacturing these products will be fully declared. Efforts should be made to develop products that will have a drastically reduced VOCs.

5.0 Research Gaps and Recommendations for Future Research

Qualitative estimate of the exact quantity of SOAs produced as a result of household products use has not been fully addressed in the literature. The available methods and techniques for quantifying SOAs may be insufficient to account for it. There is need to develop an approximate technique / equipment that will determine the approximate amount of SOAs yield. Uncertainties still arise as regards the effects of some factors affecting SOAs yield. Such uncertainties should be subjects of future research work. Formation of SOAs for consumer products should be considered as a critical area for further research work as little work found in the literature addressed this at present. Determination of undisclosed and unregulated VOC content recently researched into in some advanced countries (Steinemann[148;149] and Dodson *et al.* [151] should also be adopted by researchers and the regulatory agencies in the under-developed nations of the world.

References

1. Kanakidou, M., Seinfeld, J.H., Pandis, S.N., Barnes, I., Dentener, F.J., Facchini, M.C., *et al.* Organic Aerosol and Global Climate Modelling: A Review. *Atmos. Chem. Phys.* 2005; 5: 1053–1123. doi:10.5194/acp-5-1053-2005.
2. Vivanco, M.G. and Santiago, M., 2010. Secondary Organic Aerosol Formation from the Oxidation of a Mixture of Organic Gases in a Chamber, *Air Quality*, Ashok Kumar (Ed.), 2010; ISBN: 978-953-307-131-2, InTech, Available from: <http://www.intechopen.com/books/air-quality/secondary-organic-aerosols-experiments-in-an-outdoor-chamber>. Downloaded on 2012-06-21.
3. Nazaroff, W.W. and Weschler, C.J. Cleaning Products and Air Fresheners: Exposure to Primary and Secondary Air Pollutants. *Atmospheric Environment*. 2004;38 : 2841 – 2865
4. Jiang, L., Xu, Y., Yin, B., and Bai, Z., (2012). Theoretical Study on The Reaction Mechanism of Ozone Addition to the Double Bonds of Keto-Limonene. *Journal of Environmental Sciences*, 24(1) 147–151.
5. Koutrakis, P., Brauer, M., Briggs, S.L.K., Leaderer, B.P. Indoor Exposures to Fine Aerosols and Acid Gases. *Environmental Health Perspectives*. 1991; 95: 23-28.
6. He, C., Morawska, L., and Gilbert, D. L. (2005). Particle Deposition Rates in Residential Houses. *Atmospheric Environment*, 39(21), 3891-3899. Sonibare, J.A. A Critical Review of Natural Gas Flares-Induced Secondary Air Pollutants. *Global NEST Journal*. 2011; Vol 13, No 1: pp 74-89
7. Wallace, L., Emmerich, S.J. and Howard – Reed, C. Effect of Central Fans and Induct filters on deposition rates of ultrafine and fine particles in an occupied townhouse. *Atmospheric Environment*. 2004 38: 405 – 413. doi:10.1016/j.atmosenv.2003.10.003
8. He, C., Morawska, L., and Gilbert, D.L. Particle Deposition Rates in Residential Houses. *Atmospheric Environment* . 2005;39(21): 3891- 3899.
9. Mitsakou, C., Housiadas, C., Eleftheriadis, K., Vratolis, S., Helmis, C., and Asimakopoulos, D. Lung Deposition of Fine and Ultrafine Particles Outdoors and Indoors During A Cooking Event and A No Activity Period. *Indoor Air*. 20; 17: 143–152.
10. Wang, H. Formation Mechanisms of Secondary Organic Aerosols in Relation to Laser Printer Emissions. Doctoral Thesis, International Laboratory for Air Quality and Health, Queensland University of Technology, Australia. 2011
11. United States Environmental Protection Agency. Office of Air and Radiation (OAR). Indoor Environments Division. Characterizing Air Emissions from Indoor Sources. 1995; EPA-600-F-95-005.
12. Sharief, R.A., 2010. Investigations into Domestic Household Aerosols Air Freshener and Body Spray. ILASS – Europe 2010, 23rd Annual Conference on Liquid Atomization and Spray Systems, Brno, Czech Republic.
13. Wainman, T., Zhang, J., Weschler, C.J., & Li, P.J. Ozone and Limonene in Indoor Air: A Source of Submicron Particle Exposure. *Environmental Health Perspectives*. 2000;108(12): 1139 – 1145
14. Liu, X., Mason, M., Krebs, M., Sparks. Full-Scale Chamber Investigation and Simulation of Air Freshener Emissions in the Presence of Ozone. *Environmental Science and Technology*. 2004;38: 2802–2812.
15. Freed, L., and Wilson, D. The Science of Air Fresheners Part 1: From the Product to the Air, NEPHC, Washington. 2009.

16. Destailats, H., Lunden, M.M., Singer, B.C., Coleman, B.K., Hodgson, A.T., Weschler, C.J., *et al.* Indoor Secondary Pollutants from Household Product Emissions in the Presence of Ozone: A Bench-Scale Chamber Study. *Environmental Science & Technology*.2008; 40: 4421 – 4428.
17. Singer, B.C., Destailats, H., Hodgson, A.T., Nazaroff, W.W. Cleaning products and air fresheners: emissions and resulting concentrations of glycol ethers and terpenoids. *Indoor Air*.2006a; 16: 179–191.
18. Clausen, P.A., Larsen, K., Kofoed – Sorenson V., Jensen, K.A., Afshari, A., Gunnarsen, L., *et al.* Indoor Ozone/Terpene Reactions in An Apartment Under Normal Conditions. *Geophysical Research*.2005; 7: 1078 - 1086.
19. Coleman, B.K., Lunden, M.M., Destailats, H., Nazaroff, W.W. Secondary Organic Aerosols From Ozone Initiated Reactions With Terpenes – Rich Household Products. *Atmospheric Environment*.2008; 42: 8234 –8245.
20. Langer, S., Moldanova, J., Arrhenius, K., Ljungstrom, E., and Ekberg, L. Ultrafine Particles Produced by Ozone/Limonene Reactions in Indoor Air Under Low/Closed Ventilation Conditions. *Atmospheric Environment*.2008; 42: 4149 – 4159.
21. Hao, L.Q., Yli-Piri, P., Tiitta, P., Romakkaniemi, S., Vaattovaara, P., Kajos, M.K. *et al.* New Particle Formation From the Oxidation of Direct Emissions of Pine Seedlings. *Atmos. Chem. Phys.* 2009;9, 8121–8137.
22. European Commission. Butane. Commission Directive 2001/59/EC of 6 August 2001. Annex IB. Official Journal of the European Union. 21.08.2001. L225/24. European Commission. 28th ATP. Available from: <http://ecb.jrc.it/documents/Classification-Labelling/ATPS OF DIRECTIVE 67-548-EEC>. retrieved April 21 2012.
23. Angove, D.E., Fookes, C.J.R., Hynes, R.G., Walters, C.K., and Azzi, M. The Characterisation of Secondary Organic Aerosol Formed During the Photodecomposition of 1,3-Butadiene in Air Containing Nitric Oxide. *Atmospheric Environment* .2006;40:4597–4607.
24. World Health Organization, 1989. WHO Indoor Air Quality: Organic Pollutants. Euro Reports and Studies, No 111, World Health Organization, Copenhagen.
25. Zabiegala, B. Organic Compounds in Indoor Environments. *Polish Journal of Environ. Stud.*2006; 15(3): 383-393.
26. Baechler, M.C., Hadley, D.L., Marseille, T.J., Stenner, R.D., Peterson, M.R., Naugle, D., *et al.*1991. Sick Building Syndrome - Sources, Health Effects, Mitigation. *Pollution Technology Review* No. 205, Noyes Data Corporation, New Jersey.
27. Walgraeve, C., Demeestere, K., Dewulf, J., Van Huffel, K., and Van Langenhove, H. Diffusive Sampling of 25 Volatile Organic Compounds in Indoor Air: Uptake Rate Determination and Application in Flemish Homes for the Elderly. *Atmospheric Environmen*.2011;45: 5828 – 5836
28. Wolkoff, P., (1995). Volatile Organic Compounds - Sources, Measurements, Emissions and the Impact on IAQ. *Indoor Air Suppl.*, 3, 9–73.
29. Clobes, A.L., Ananth, G.P., Hood, A.L., Schroeder, J.A., Lee, K.A. Human activities as sources of volatile organic compounds in residential environments: *Ann NY Acad Sci.* 1992; 641:79 – 86.
30. Singer, B.C., Coleman, B.K., Destailats, H., Hodgson, A.T., Lunden, M.M., Weschler, C.J. and Nazaroff, W.W. Indoor Secondary Pollutants From Cleaning Product And Air

- Freshener Use in The Presence of Ozone, *Atmospheric Environment*.2006b; 40: 6696–6710.
31. Heroux, M.E., Gauvin, D., Gilbert, N.L., Guay, M., Dupuis, G., Legris, M., *et al.* Housing Characteristics and Indoor Concentrations of Selected Volatile Organic Compounds (VOCs) in Quebec City, Canada. *Indoor and Built Environment*.2008;17 (2):128–137.
 32. Dusek, U. 2000. Secondary Organic Aerosol – Formation Mechanisms and Source Contributions in Europe. Interim Report - IR-00-066, International Institute for Applied Systems Analysis(IIASA), Austria
 33. Grosjean, D., and Seinfeld, J.H. Parameterization of the Formation Potential of Secondary Organic Aerosols, *Atmos. Environ*1989;. 23: 1733-1747
 34. United States Environmental Protection Agency, 2012a. The Inside Story: A Guide to Indoor Air Quality', U.S. EPA/Office of Air and Radiation Office of Radiation and Indoor Air (6609J) retrieved from <http://www.epa.gov/iaq/pubs/insidestory.html> on April 24, 2012.
 35. ? USEPA 2012B
 36. Kwok, E.S.C., Atkinson, R., and Arey, J. Observation of Hydroxycarbonyls from the OH Radical-Initiated Reaction of Isoprene, *Environ. Sci. Technol.*1995; 29: 2467– 2469. doi:10.1021/es00009a046.
 37. Finlayson-Pitts B.J., Pitts, J.N., 1996 Atmospheric Chemistry.Wiley, New York.
 38. Sarwar, G., Corsi, R., Kimura, Y., Allen, D., and Weschler, C. J. Hydroxyl Radicals in Indoor Environments. *Atmospheric Environment*. 2006;36: 3973–3988.
 39. Nazaroff, W.W., Coleman, B.K., Destailats, H., Hodgson, A.T., Liu, D.L., Lunden, M.M., Singer, B.C. and Weschler, C.J. 2006. Indoor Air Chemistry: Cleaning Agents, Ozone and Toxic Air Contaminants. A Technical Report prepared forCalifornia Air Resources Board and the California Environmental Protection Agency Final Report: Contract No. 01-336.
 40. Britigan, N., Alshawwa, A., Nizkorodov, S.A. Quantification of Ozone Levels in Indoor Environments Generated by Ionization and Ozonolysis Air Purifiers.*J. Air & Waste Manage. Assoc.*2006;56: 601– 10.
 41. Galloway, M.M., Huisman, A.J., Yee, L.D., Chan, A.W.H., Loza, C.L., Seinfeld, J.H. *et al.* Yields of Oxidized Volatile Organic Compounds During the OH Radical Initiated Oxidation of Isoprene, Methyl Vinyl Ketone, and Methacrolein under High-NO_x Conditions. *Atmos. Chem. Phys.*2011; 11: 10779 – 10790.
 42. Schripp, T., Langer, S., and Salthammer, T. 2012. Interaction of Ozone with Wooden Building Products, Treated Wood Samples, and Exotic Wood Species. *Atmospheric Environment*, 54,365 – 54,372. Doi:10.1016/j.atmosenv.2012.02.064.
 43. Weschler, C. J. Ozone in Indoor Environments: Concentration and Chemistry. *Indoor Air*'2000;10: 269-288.
 44. Nøjgaard, J.K., Bilde, M., Stenby, C.,Nielsen, O.J., and Wolkoff, P. The Effect of Nitrogen Dioxide on Particle Formation DuringOzonolysis of Two Abundant Monoterpenes Indoors. *Atmospheric Environment*.2006b;40: 1030– 1042.doi:10.1016/j.atmosenv.2005.11.029.
 45. Lee, S.C., Lam, S., and Fai, H.K. Characterization of VOCs, Ozone, and PM₁₀ Emissions from Office Equipment in an Environmental Chamber. *Building and Environment* .2001; 36: 837– 842.PII: S0360-1323(01)00009-9.

46. Weschler, C.J., and Shields, H.C. Indoor Ozone/Terpene Reactions as a Source of Indoor Particles. *Atmospheric Environment*.1999; 33: 2301-2312.
47. Uhde, E., and Salthammer. Impact of Reaction Products from Building Materials and Furnishings on Indoor Air Quality—A Review of Recent Advances in Indoor Chemistry. *Atmospheric Environment*.2007; 41: 3111 – 3128.
48. Rohr, A.C. 2006. Indoor Secondary Organic Aerosols Formation and Health Effects. Health Effects of Organic Aerosols Workshop, Electric Power Research Institute, Palo Alto, CA
49. Seinfeld, J. H. and Pandis, S. N. *Atmospheric Chemistry and Physics*, John Wiley & Sons Inc., New York, 1326.1998
50. Destailats, H., Lunden, M. M. , Singer, B. C., Coleman, B. K., Hodgson, A. T., Weschler, C. J. and Nazaroff, W. W. (2006). Indoor Secondary Pollutants from Household Product Emissions in the Presence of Ozone: A Bench-Scale Chamber Study. *Environmental Science and Technology*, 40, 4421 – 4428
51. Huang, Y., Lee, S.C., Ho, K.F., Ho, S.S.H., Cao, N., Cheng, Y. *et al.* Effects of ammonia on Ozone Initiated Formation of Indoor Secondary Products with Emissions from Cleaning Product. *Atmospheric Environment*.2012a; 59: 224 – 31.
52. Fan, Z., Liou, P., Weschler, C.J., Fiedler, N., Kipen, H., Zhang, J. Ozone-Initiated Reactions with Mixtures of Volatile Organic Compounds under Simulated Indoor Conditions. *Environ. Sci. Technol.*2003; 37:1811-1121.
53. Chen, X., and Hopke, P.K. A Chamber Study of Secondary Organic Aerosol Formation by Linalool Ozonolysis. *Atmospheric Environment*.2009; 43: 3935–3940.
54. Perraud, V., Brunsa, E.A., Ezella, M.J., Johnson, S.N., Yu, Y., Alexander, M.L., Zelenyuk, A., Imre, D., Chang, W.L., Dabdub, D., Pankow, J.F. and Finlayson-Pitts, B.J. Nonequilibrium Atmospheric Secondary Organic Aerosol Formation and Growth.2012; *PNAS*, Vol. 109, No. 8: 2836 – 2841.
55. Nøjgaard, J.K., Nøjgaard, A.W., Larsen, K., Spørring, S., Wilkins, C.K., Clausen, P.A., and Wolkoff, P. Secondary Limonene Endo: Ozonide : A Major Product from Gas- Phase Ozonolysis of R-(+)- Limonene at Ambient Temperature. *Atmospheric Environment*.2006a; 40: 3460 – 3466.
56. Nøjgaard, J.K, Nøjgaard, A.W., and Wolkoff, P. On line Analysis of Secondary Ozonides from Cyclohexenes and d- limonene Ozonolysis Using Atmospheric Townsend Discharge Ionisation Mass Spectrometry. *Atmospheric Environment*.2007; 41: 8345 – 8354.
57. Vibenholt, A., Nøjgaard, A.W., Clausen, P.A., and Wolkoff, P. Formation and Stability of Secondary Ozonides from Monoterpenes Studied by Mass Spectrometry. *Chemosphere*.2009; 76: 572 – 577.
58. Aschmann, S.M., Martin, P., Tuazon, E.C., Arey, J., and Atkinson, R. Kinetic and product studies of the reactions of selected glycol ethers with OH radicals. *Environmental Science and Technology* .2001;35: 4080-4088.
59. Welz, O., Savee, J.D., Osborn, D.L., Vasu, S.S., Percival, C.J., Shallcross, D.E., and Taatjes, C.A. Direct Kinetic Measurements of Criegee Intermediate (CH_2OO) Formed by Reaction of CH_2I with O_2 . *Science*.2012;335(6065): 204-207.DOI: 10.1126/science.1213229.
60. Taatjes, C.A., Welz, O., Eskola, A.J., Savee, J.D., Osborn, D.L., Lee, E.P.F., Dyke, J.M., Mok, D.W.K., Shallcross, D.E., and Percival, C.J. Direct Measurement of

- Criegee Intermediate (CH_2OO) Reactions with Acetone, Acetaldehyde, and Hexafluoroacetone. *Phys. Chem. Chem. Phys.* 2012;14, 10391-10400. DOI: 10.1039/C2CP40294G.
61. European Commission, Joint Research Centre - Institute for Health & Consumer Protection Physical & Chemical Exposure Unit. Impact of Ozone-initiated Terpene Chemistry on Indoor Air Quality and Human Health. Report No. 26, EUR 23052 EN, Italy 2007.
 62. Cao, G., and Jang, M. Effects of Particle Acidity and UV Light on Secondary Organic Aerosol Formation from Oxidation of Aromatics in the Absence of NO_x . *Atmospheric Environment*. 2007; 41: 7603 – 7613.
 63. Kleindienst, T.E., Lewandowski, M., Offenberg, J.H., Jaoui, M., and Edney, E.O. The Formation of Secondary Organic Aerosol from the Isoprene + OH Reaction in the Absence of NO_x . *Atmos. Chem. Phys.* 2009; 9: 6541– 6558
 64. Kroll, J.H., Ng, N.L., Murphy, S.M., Flagan, R.C., and Seinfeld, J.H. Secondary Organic Aerosol Formation from Photooxidation of Isoprene, *Environ. Sci. Technol.* 2006; 40: 1869–77.
 65. Kwan, A.J. Investigating Sources and Sinks of Organic Aerosol: NO_3 -Initiated Oxidation of Isoprene and Heterogeneous Oxidation of Organic Aerosol. Doctoral Thesis, California Institute of Technology. 2011.
 66. Cao, G., and Jang, M. Secondary Organic Aerosol Formation from Toluene Photooxidation Under Various NO_x Conditions and Particle Acidity. *Atmos. Chem. Phys.* 2008; 8: 14467 – 14495.
 67. Zhang, J., Hartz, K. E. H., Pandis, S. N., and Dohanue, N. M. Secondary Organic Aerosol Formation from Limonene Ozonolysis: Homogeneous and Heterogeneous Influences as a Function of NO_x , *J. Phys. Chem.* 2006; 110: 11053-11063.
 68. Ng, N.L., Kroll, J.H., Chan, A.W.H., Chhabra, P.S., Flagan, R.C. and Seinfeld, J.H. Secondary Organic Aerosol Formation from m-Xylene, Toluene, and Benzene, *Atmos. Chem. Phys.* 2007; Discuss., 7: 4085-4126.
 69. Presto, A. A., Huff Hartz, K. E. and Donahue, N. M. Secondary Organic Aerosol Production from Ozonolysis: 2. Effect of NO_x Concentration, *Environ. Sci. Technol.* 2005; 39: 7046- 7054.
 70. Kiendler-Scharr, A., Andres, S., Bachner, M., Behnke, K., Broch, S., Hofzumahaus, A., *et al.* Isoprene in Poplar Emissions: Effects on New Particle Formation and OH Concentrations. *Atmos. Chem. Phys.* 2012; 12: 1021–30.
 71. Kim, H., Barkey, B., and Paulson, S.E. Real Refractive Indices and Formation Yields of Secondary Organic Aerosol Generated from Photooxidation of Limonene and α -Pinene: The Effect of the HC/ NO_x Ratio. *J. Phys. Chem.* 2012; A, 116 (24): 6059 – 67. doi:10.1021/jp301302z
 72. Liu, Y. Computational Simulation of Secondary Organic Aerosol (SOA) Formation from Toluene Oxidation. A Master's thesis submitted to Rice University, Houston, Texas. 2012
 73. Chang, X. Module for Simulating Composition Effects on Secondary Organic Aerosol Partitioning and its Evaluation in the Southeastern United States. Published PhD thesis, Graduate School of Vanderbilt University, Nashville, Tennessee. 2006.
 74. Huang, Y.T., Chen, C.C., Chen, Y.K., Chiang, C.M., and Lee, C.Y. Environmental Test Chamber for the Elucidation of Ozone – Initiated Secondary Pollutants Emission from Painted Wooden Panels in Buildings. *Building and Environment*. 2012b; 50: 135 – 40.

75. Weschler, C.J. Reactions among Indoor Pollutants. *The Scientific World*. 2001; 1: 443 – 457.
76. Forster, P., Ramaswamy, V., Artaxo, P., Bernsten, T., Betts, R.W., Fahey, D. *et al.* Changes in Atmospheric Constituents and in Radiative Forcing, *Climate Change: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. 2007
77. Styler, S.A., Loiseaux, M.E. and Donaldson, D.J. Substrate Effects in the Photoenhanced Ozonation of Pyrene. *Atmos. Chem. Phys.* 2011; 11: 1243 – 1253.
78. Varutbangkul, V., Brechtel, F. J., Bahreini, R., Ng, N. L., Keywood, M. D., Kroll, J. H. , Flagan, R. C., Seinfeld, J. H., Lee, A., and Goldstein, A. H. Hygroscopicity of Secondary Organic Aerosols Formed by Oxidation of Cycloalkenes, Monoterpenes, Sesquiterpenes, and Related Compounds. *Atmos. Chem. Phys.* 2006; 6: 2367–2388.
79. Weschler, C.J. Chemistry in Indoor Environment: 20 Years of Research. *Indoor Air*. 2011; 21: 205 – 208.
80. Librando, V., and Tringali, G. Atmospheric fate of OH Initiated Oxidation of Terpenes. Reaction Mechanism of α -pinene Degradation and Secondary Organic Aerosol Formation. *Journal of Environmental Management*. 2005; 75: 275 – 282.
81. Leungsakul, S., Jaoui, M., and Kamens, R.M. Kinetic Mechanism for Predicting Secondary Organic Aerosol Formation from the Reaction of D-Limonene with Ozone. *Environmental Science & Technology*. 2005a; 39, (24): 9583–94.
82. Leungsakul, S., Jeffries, H.E, and Kamens, R.M. A Kinetic Mechanism for Predicting Secondary Aerosol Formation from the Reactions of D-limonene in the Presence of Oxides of Nitrogen and Natural Sunlight. *Atmospheric Environment*. 2005b; 39: 7063–7082.
83. Matsunaga, A., Docherty, K.S., Lim, Y.B. and Ziemann, P.J. Composition and yields of secondary organic aerosols formed from OH radical-initiated reactions of linear alkenes in the presence of NO_x: Modeling and measurements. *Atmospheric Environment*. 2009; 43: 1349–1357.
84. Vivanco, M.G., Santiago, M. Martínez-Tarifa, A., Borrás, E., Ródenas, M., García-Diego, C., Sánchez, M. SOA formation in a photoreactor from a mixture of organic gases and HONO for different experimental conditions. *Atmospheric Environment*. 2011; 45: 708-715.
85. Hatfield, M.L. and Hartz, K.E.H. Secondary Organic Aerosol from biogenic volatile organic compounds mixtures. *Atmospheric Environment*. 2011; 45: 2211 – 2219.
86. Parikh HM, Carlton AG, Vizuette W and Kamens RM. Modeling secondary organic aerosols using dynamic partitioning approach incorporating particle aqueous –phase chemistry. *Atmospheric Environment* 2011; 45: 1126 – 1137.
87. Song, C., Zaveri, R.A., Alexander, M.L., Thornton, J. A., Madronich, S., Ortega, J.V., Zelenyuk, A., Yu, X.Y., Laskin, A., and Maughan, D.A. Effect of hydrophobic primary organic aerosols on secondary organic aerosols formation from ozonolysis of α -pinene. *Geophysical Research Letters* 34.
88. Shilling, J. E., Chen, Q., King, S. M., Rosenoern, T., Kroll, J.H., Worsnop, D. R., McKinney, K. A., and Martin, S. T. Particle Mass Yield in Secondary Organic Aerosol Formed by the Dark Ozonolysis of α -Pinene, *Atmos. Chem. Phys.* 2008; 8: 2073–2088.

89. Presto, A.A. and Donahue, N.M. Investigation of alpha-pinene plus ozone secondary organic aerosols formation at low total aerosol mass. *Environmental Science and Technology*. 2006;40: 3536 – 3543.
90. Lauraguais, A., Coeur-Tourneur, C., Cassez, A., and Seydi, A. Rate constant and secondary organic aerosol yields for gas-phase reaction of hydroxyl radicals with syringol (2,6-dimethoxyphenol). *Atmospheric Environment*. 2012;55: 43-48.
91. Updyke, K.M., Nguyen, T.B. and Nizkorodov, S.A. Formation of brown carbon via reactions of ammonia with secondary organic aerosols from biogenic and anthropogenic precursors. *Atmospheric Environment*. 2012; 63: 22 -31.
92. Stewart, D.J., Almarok, S.H., Lockhart, J.P., Mohamed, O.M., Nutt, N.R., Pfrang, C. and Marston, G. The kinetics of the gas-phase reactions of selected monoterpenes and cycloalkenes with ozone and the NO₃ radical. *Atmospheric Environment*; 70 :227 -235.
93. Alfarra MR, Paulsen D, Gysel M, Garforth AA, Dommen J, Prevot ASH, Worsnop DR, Baltensperger U and Coe H. A mass spectrometric study of secondary organic aerosols formed from the photooxidation of anthropogenic and biogenic precursors in a reaction chamber. *Atmospheric Chemistry and Physics* 2006; 6: 5279 – 5293.
94. Broske, R., Kleffmann, J. and Wiesen, P. Heterogeneous conversion of NO₂ on secondary organic aerosol surfaces: A possible source of nitrous acid (HONO) in the atmosphere? *Atmospheric Chemistry and Physics*. 2003;3: 469–474.
95. Carter, W.P.L., Cocker III, D.R., Fitz, D.R., Malkina, I.L., Bumiller, K., Sauer, C.G., Pisano, J.T., Bufalino, C., and Song, C. A new environmental chamber for evaluation of gas-phase chemical mechanisms and secondary aerosol formation. *Atmospheric Environment*. 2005; 39 :7768–7788.
96. Chen, T. and Jang, M. Secondary organic aerosol formation from photooxidation of a mixture of dimethyl sulphide and isoprene. *Atmospheric Environment* 2012; 46: 271 -278.
97. Cocker III, D.R., Clegg, S.L., Flagan, R.C. and Seinfeld, J.H. The effect of water on gas-particle partitioning of secondary organic aerosol. Part I: a-pinene/ozone system. *Atmospheric Environment*, 2001;35: 6049–6072.
98. Irei S, Rudolph J., Huang L., Auld J, and Hastie D. Stable isotope ratio of secondary particulate organic matter formed by photooxidation of toluene in indoor smog chamber. *Atmospheric Environment* 2011; 45: 856 – 862.
99. Lee A., Goldstein AH, Keywood MD, Gao S, Varutbangkul V, Bahreini R, Ng NL, Flagan RC, and Seinfeld JH. Gas-phase products and secondary aerosol yields from the ozonolysis of ten different terpenes, *J. Geophys. Res.*, 2006, 111, D07302.
100. Qi, L., Nakao, S., Tang, P., and Cocker, D. R. Temperature Effect on Physical and Chemical Properties of Secondary Organic Aerosol from M-Xylene Photooxidation. *Atmos. Chem. Phys.* 2010; 10: 3847–3854.
101. Rickard AR, Wyche KP, Metzger A, Monks PS, Ellis AM, Dommen J, Baltensperger U, Jenkin ME, Pilling MJ. Gas phase to anthropogenic secondary organic aerosol: Using the Master Chemical Mechanism to probe detailed observations of 1,3,5- trimethylbenzene photooxidation. *Atmospheric Environment* 2010;44: 5423 -5433.
102. Rossignol S, Chiappini L, Perraudin E, Rio C, Fable S, Valorso R, and Doussin JF. Development of a parallel sampling and analysis method for the elucidation of gas/particle partitioning of oxygenated semi-volatile organics: a limonene ozonolysis study. *Atmos. Meas. Tech.* 2012, 5: 1459–1489.

103. Salo K, Hallquist M, Jonsson AM, Saathoff H, Naumann KH, Spindler C, Tillmann R, Fuchs H, Bohn B, Rubach F, Mentel TH F, Müller L, Reinnig M, Hoffmann T, and Donahue NM. Volatility of secondary organic aerosol during OH radical induced ageing. *Atmos. Chem. Phys.* 2011, 11, 11055–11067.
104. Vutukuru, S., Griffin, R.J., and Dabdub, D., 2006. Simulation and Analysis of Secondary Organic Aerosol Dynamics in the South Coast Air Basin of California. *Journal of Geophysical Research*, Vol. III, D10S12, doi:10.1029/2005JD006139.
105. Lane, T.E., and Pandis, S.N. Predicted Secondary Organic Aerosol Concentrations from the Oxidation of Isoprene in the Eastern United States. *Environ. Sci. Technol.* 2007;41: 3984-3990.
106. Takewawa, H. Secondary Organic Aerosol Formation from Photochemical Reaction of Aromatic Hydrocarbons. Research Report, R&D Review of Toyota CRDL.2003; 38(1): 57- 62.
107. Berndt, T., Boge, O., Stratmann, F. Gas-Phase Ozonolysis of α -Pinene: Gaseous Products and Particle Formation, *Atmos. Environ.*2003; 37: 3933 –3945.
108. Pathak, R. K., Presto, A. A., Lane, T. E., Stanier, C. O., Donahue, N. M., and Pandis, S. N. Ozonolysis of α -pinene: Parameterization of Secondary Organic Aerosol Mass Fraction. *Atmos.Chem. Phys.*2007;7: 3811–3821.
109. Wolkoff, P., and Kjærgaard, S.K. The Dichotomy of Relative Humidity on Indoor Air Quality. *Environment International*.2007;33: 850–857.
110. Na, K., Song, C., and Cocker III, D.R. Formation of Secondary Organic Aerosol from the Reaction of Styrene with Ozone in the Presence and Absence of Ammonia and Water. *Atmospheric Environment*.2006; 40: 1889–1900.
111. Chan, H.K., Eber, S., Daviskas, E., Constable, C., Young, I.. Changes in Lung Deposition of Aerosols due to Hygroscopic Growth: A Fast SPECT Study, *Journal of Aerosol Medicine Deposition Clearance and Effects in the Lung*.2002; 15: 307 – 311.
112. Petrick, L.M., Svidovsky, A., and Dubowski, Y Thirdhand Smoke: Heterogeneous Oxidation of Nicotine and Secondary Aerosol Formation in the Indoor Environment. *Environ. Sci. Technol.*2011;45: 328–333.
113. Yu, K.P., Lin, C.C., Yang, S.C., and Zhao, P. Enhancement Effect of Relative Humidity on the Formation and Regional Respiratory Deposition of Secondary Organic Aerosol. *Journal of Hazardous Materials*.2011;191: 94–102.
114. Weschler, C.J., and Shield, H. Experiments Probing the Influence of Air Exchange Rates on Secondary Organic Aerosols Derived from Indoor Chemistry. *Atmospheric Environment*.2003;37: 5621–5631.
115. EC, European Commission, Joint Research Centre - Institute for Health & Consumer Protection Physical & Chemical Exposure Unit. Impact of Ozone-initiated Terpene Chemistry on Indoor Air Quality and Human Health. Report No. 26, EUR 23052 EN, Italy 2006.
116. Waring, M.S., Wells, R.J., and Siegel, J.A. Secondary Organic Aerosol Formation from Ozone Reactions with Single Terpenoids and Terpenoid Mixtures. *Atmospheric Environment*.2011; 45: 4235 – 4242.
117. Fick, J. Chemical Reactions in Ventilation Systems -Ozonolysis of Monoterpenes. Doctoral Thesis, Umea University, Sweden.2003; ISBN 91-7305-511-5

118. Lu, Z., Hao, J., Takekawa, H., Hu, L., and Li, J. Effect of High Concentration of Inorganic Seed Aerosols on Secondary Organic Aerosols Formation in m-xylene/NO_x Photooxidation System. *Atmospheric Environment*, 2009; 43: 897–904.
119. He, C., Morawska, L., Hitchins, J., and Gilbert, D. Contribution from Indoor Sources to Particle Number and Mass Concentrations in Residential Houses. *Atmospheric Environment*. 2004; 38(21): 3405 – 3415
120. Muller, L., Reinnig, M.C., Warnke, J., and Hoffmann, T. Unambiguous Identification of Esters as Oligomers in Secondary Organic Aerosol Formed from Cyclohexene and Cyclohexene/ α -Pinene Ozonolysis. *Atmos. Chem. Phys.* 2000; 80: 1423–1433. 101
121. Sarwar, G., Corsi, R., Allen, D., Weschler, C.J.,. The significance of secondary organic aerosol formation and growth in buildings: experimental and computational evidence. *Atmospheric Environment* 2003a, 37, 1365–1381.
122. Sarwar, M.G., Olson, D., Corsi, R., Weschler, C.J., Indoor fine particles: the role of terpene emissions from consumer products. *Journal of Air and Waste Management Association*, 2003b.
123. Sleiman M, Destailats H, Smith JD, Liu CL, Ahmed M, Wilson KR and Gundel LA. Secondary organic aerosols from ozone-initiated reactions with nicotine and secondhand tobacco smoke. *Atmospheric Environment* 2010; 44: 4191 – 4198.
124. Rohr, A. C., Weschler, C. J., Koutrakis, P., Spengler, J. D. Generation and Quantification of Ultrafine Particles through Terpene/Ozone Reaction in a Chamber Setting. *Aerosol Science and Technology*. 2003; 37, 65–78.
125. Sleiman, M., Destailats, H., Smith, J.D., Liu, C.L. Ahmed, M., Wilson, K. R., and Gundel, L.A. 2010. Secondary Organic Aerosols from Ozone Initiated Reactions with Nicotine and Secondhand Tobacco Smoke. *Atmospheric Environment*, 44: 4191 – 4198.
126. Weschler, C.J. Ozone's Impact on Public Health: Contributions from Indoor Exposure to Ozone and Products of Ozone – Initiated Chemistry. *Environ. Health Persp.* 2006; 114: 1489 – 1496.
127. Fadeyi, M.O., Weschler, C.J., Tham, K.W. The Impact of Recirculation, Ventilation, and Filters on Secondary Organic Aerosols Generated by Indoor Chemistry. *Atmospheric Environment* 2009; 43: 3538 – 3547.
128. Finlayson-Pitts, B.J., Pitts, J.N. 2000. *Chemistry of the Upper and Lower Atmosphere*, Academic Press, San Diego, 969.
129. Nazaroff, W.W. Indoor Particles Dynamics. *Indoor Air*. 2004; 14 (supplement 7): 175 – 183.
130. Fan, Z., Weschler, C.J., Han, I., Zhang, J. Co- Formation of hydroperoxides and Ultra fine Particles during the Reaction of Ozone with a Complex VOC mixture under Simulated Indoor Conditions. *Atmospheric Environment* .2005; 39: 5171 – 5182.
131. Hovorka, J., and Branis, M. New particle Formation and Condensational Growth in a Large Indoor Space. *Atmospheric Environment*. 2011; 45: 2736 – 49. doi.org/10.1016/j.atmosenv.2011.02.039.
132. Ziemann, C.J. Formation of Alkoxyhydroxyl Aldehydes and Cyclic Peroxylhemiacetals from Reactions of Cyclic Alkenes with O₃ in the Presence of Alcohols. *Journal of Physical Chemistry A*. 2003; 107: 2048 – 2060
133. Huang, H.L., Tsai, T.J., Hsu, N.Y., Lee, C.C., Wu, P.C., and Su, H.J. 2012c. Effects of Essential Oil on the Formation of Formaldehyde and Secondary Organic Aerosols in an Aromatherapy Environment. *Building and Environment*. 2012c; 57: 120-25.

134. Asgharian, B., and Price, O.T. Deposition of Ultrafine (Nano) Particle in the Human Lung. *Inhalation Toxicology*.2007;19 (13): 1045 – 1054.
135. Katsouyanni, K., Touloumi, G., Samoli, E., Gryparis, A., Le Tertre, A., Monopolis, Y., *et al.* Confounding and Effect Modification in Short Term Effects of Ambient Particles on Total Mortality: Results from 29 European Cities within the APHEA2 Project. *Epidemiology*.2001;12 (5): 521 – 531.
136. International Consumer Research and Testing 2005. Emission of chemicals by air fresheners: Tests on 74 consumer products sold in Europe. ICRT, United Kingdom.
137. Bello, A., Quinn M.M., Perry M.J., Milton D.K. Quantitative Assessment of Airborne Exposures Generated during Common Cleaning Tasks: A Pilot Study. *Environ Health*.2010; 9:76. doi:10.1186/1476-069X-9-76
138. Franklin, P., Dingle, P., and Stick, S. Raised exhaled nitric oxide in healthy children is associated with domestic formaldehyde levels. *Am J RespirCrit Care Med* .2000; 161: 1757 – 1759.
139. Quirce, S.,andBarranco, P. Cleaning Agents and Asthma. *J. Investig. Allergol. Clin. Immunol*.2010; 20(7): 542 – 550.
140. Mehta, A.J., Adam, M.,Schaffner, E., Barthélémy, J.C.,Carballo, D.,Gaspoz, J.M.,Rochat, T., Schindler,C., Schwartz, J., Zock, J.P.,Künzli,N.,Probst-Hensch, Probst-Hensch, N., and the SAPALDIA Team .Heart Rate Variability in Association with Frequent Use of Household Sprays and Scented Products in SAPALDIA.*Environ. Health Perspect.*.2012; 120:958–964. <http://dx.doi.org/10.1289/ehp.1104567>.
141. Pope, C. A., and Dockery, D. W. Health Effects of Fine Particulate Air Pollution: Lines that Connect. *Journal of the Air& Waste Management Association*.2006; 56: 709-742
142. Fiedler, N., Laumbach, R., Kelly- McNeil, K., Liroy, P., Fan, Z.H., Zhang, J., Ottenweller, J, *et al.* Health Effects of a Mixture of Indoor Air Volatile Organics, Their Ozone Oxidation Products, and Stress.*Environmental Health Perspective*.,2005;113, 11, 1542 – 1548.
143. Rohr, A., Wilkins, C.K., Clausen, P.A., Hammer, M., Nielsen, G.D., Wolkoff, P., and Spengler, J.D. Upper Airway and Pulmonary Effects of Oxidation Products of (+)- α -pinene , d- limonene, and Isoprene in BALB/c Mice. *Inhal. Toxicol*. 2002; 14: 663 – 684.
144. Anderson, S.E., Wells, J.R., Fedorowicz, A., Butterworth, L.F., Meade, B.J., and Munson, A.E. Evaluation of the Contact and Respiratory Sensitization Potential of Volatile Organic Compounds Generated by Simulated Indoor Air Chemistry.*Toxicological Sciences* .2007;97(2): 355-363.
145. Fisk, W., Faulkner, D.,Palonen, J., Seppanen, O. Performance and Costs of Particle Air filtration Technologies. *Indoor Air* .2002; 12: 223 – 234.
146. Hacker, D.W., and Sparrow, E.M. Use of Air-Cleaning Devices to Create Airborne Particle-Free Spaces Intended to Alleviate Allergic Rhinitis and Asthma During Sleep. *Indoor Air*.2005; 15:420 – 431.
147. Caress, S.M, and Steinemann, A.C. Prevalence of Frangrance Sensitivity in the American Population. *Journal of Environmental Health*.2009; 71(7): 46 – 50.
148. Steinemann, A.C. Fragranced Consumer Products and Undisclosed Ingredients. *Environmental Impact Assessment Review*.2009; 29: 32–38. doi:/10.1016/j.eiar.2008.05.002.
149. Steinemann, A.C., MacGregor, I.C., Gordon S.M., Gallagher, L.G., Davis, A.L., Ribeiro, D.S., and Wallace, L.A. Fragranced Consumer Products: Chemicals Emitted, Ingredients

- Unlisted, Environmental Impact Assessment Review.2011; 31: 328–333.
doi:10.1016/j.eiar.2010.08.002
150. Bernstein, J.A, Alexis, N., Bacchus, H., Bernstein, I.L, Fritz, P., Horner, E., *etal*.The Health Effects of Non-industrial Indoor Air Pollution. J. Allergy Clin. Immunol.2008; 121(3): 585 – 591.
 151. Dodson, R.E., Nishioka, M., Standley, L.J., Perovich, L.J.,Brody, J.G. and Rudel, R.A., 2012. Endocrine Disruptors and Asthma-Associated Chemicals in Consumer Products. Environmental Health Perspectives 120, 7, 935 – 943.
 152. Seinfeld, J.H.,Erdakos,G.B., Asher,W.E., Pankow,J.F. Modeling the formation of secondary organic aerosols(SOA).2.The predicted effects of relative humidity on aerosol formation in α -pinene, β -pinene,sabinene-, Δ^3 -carene-,and cyclohexene –ozone systems. Environmental Science and Technology. 2001; 35:1806 -1817.
 153. Pankow,J.F.An absorption model of the gas/aerosol partitioning involved in the formation of secondary organic aerosol. Atmospheric Environment.1994; 28:189 -193.