

Discovery of pyroclastic surge deposits in Bamenda volcano (Bambili, West-Cameroon, Cameroon volcanic Line): importance in chronostratigraphy of the massif

ABSTRACT

Field studies in Bambili locality (NE of Bamenda city) have recently permitted us to discover above the welded massive lapilli tuff (mlT), the remnant blocks (up to 6.5 x 11 m) of pyroclastic surge deposits characterized by well sorted and distinctly stratified layers with thicknesses ranging between 8 to 35 cm and showing a graded bedding. In addition to ashy matrix, the different layers of the deposits are made up of lithic fragments of various natures, represented by devitrified fiammes, vitrophyres, trachytes, rhyolites, granites and ignimbrites. The mineralogy of these surge deposits is quasi identical to those of the welded mlT, they are made up which consists of alkali feldspar (sanidine), quartz, plagioclase, clinopyroxene, biotite and Fe-Ti oxides. The discovery of these remnant blocks of pyroclastic surge deposits of in the Bambili locality, emitted probably from Mt Oku vent permit to reconsider the chronostratigraphy of the Bamenda volcano. In fact, after the trachytic lava flow emitted from????, the pyroclastic flow deposits originating from???? have then permitted the formation of the welded ignimbrites represented by dark gray and whitish units; pulsating hydrostatic and magma pressures have subsequently produced alternating hydromagmatic pyroclastic surges, which have afterward cover the massif followed by basaltic flows.

Keywords: surge deposits, welded ignimbrites, Bamenda volcano, West-Cameroon

1. INTRODUCTION

The intraplate Cameroon Volcanic Line (CVL) forms one of the major Lineament of the African continent (fig. 1). It can be compared to the East African Rift System, which is currently defined by an almost SW-NE geological lineament (mean value: N30°E). It is made up of a continental part and an oceanic part, making it a unique feature in Africa and even in the world [1]. The continental part includes both plutonic complexes and volcanic massifs. The oceanic part is entirely volcanic and consists of four well-studied islands (Bioko, Principe, São Tomé, and Annobon) and two large seamounts [2].

Recent work [3] reported the existence of ignimbritic deposits only in the continental part of the CVL; these rocks (deposits?) are found mainly in the central area of the CVL, where the ignimbrites of especially within the mountains Bambouto and the Mount Bamenda are located and form the most important deposits with respectively 13.5 and 6.3 km³. With the alteration and poor condition of these rocks outcrops, the estimated volumes would be significantly higher. Outcrops of smaller sizes (<1 km³) were identified in Oku-Oku [4, 5], Nkogam-Nkogam [6] and Nganha-Nganha [7] volcanoes. Studies in these massifs so far have shown that these ignimbrites are only deposits of pyroclastic flows with no internal stratification.

Comment [BCT1]: This cannot be a discovery if already mentioned in a published paper [3] for example. Consider change the title of the work. You can say Characteristics of

Comment [BCT2]: Yellow highlight, I would say: "The layers are matrix-supported and heterolithic, with the accidental lithic fragments consisting of devitrified fiammes, vitrophyres, trachytes, rhyolites, granites and ignimbrites"

Comment [BCT3]: Turquoise highlight: What do you mean by mineralogy of the surge deposit? Is it the mineralogy of the juvenile clasts? If yes, then say it like that, otherwise Please specify.

Comment [BCT4]: Green highlight: if the surges are from Mt Oku, how it affects the chronostratigraphy of Mt Bamenda located far away from each other.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Comment [BCT5]: If you mentioned that it is an intraplate volcanism, you should cite Deruelle et al. 2007 e.g.

Comment [BCT6]: Insufficient, add more references

Comment [BCT7]: Which rocks

Comment [BCT8]: Are these the volume of the volcanoes or the ignimbrite deposits? Please specify.

Comment [BCT9]: Add references that report these volumes.

Comment [BCT10]: What condition?

Formatted: Superscript

Comment [BCT11]: Compare to what?

Formatted: Superscript

Comment [BCT12]: I do not think this author did any volume estimate. Please verify and update

Comment [BCT13]: Reference

33




Fig. 1. Map of Cameroon showing the distribution of the Cameroon Volcanic Line (CVL) magmatism. Locations of seamounts are after Burke [2]. C.A.R.: Central African Republic. Inset: location map of CVL in Africa.

Deposits of pyroclastic density currents, known as a particulate gaseous volcanic flow moving along the ground (Freundt et al., 2000), have been generally categorized according to lithology and sedimentary structure, as ignimbrites, block-and-ash flow deposits and pyroclastic surge deposits [8]. Until now, only welded and non-welded ignimbrites with their massive lapilli tuff (mLT) and massive lithic breccias (mLBr) facies have been described in the Mount Bamenda [3]. These ignimbrites represent the deposits of pyroclastic flows. In this work, we highlight the existence of pyroclastic surge in the Bamenda volcano, at Bambili locality, situated in the NE of Bamenda town. These surge deposits are investigated here in detail in order to understand their mode of deposition and give their importance in the chronostratigraphy of the Bamenda volcano.

2. GEOLOGICAL SETTING

The Mount Bamenda (600 km²) constitutes in volumetric importance the fourth largest volcano of the CVL after Mt Cameroon, Mt Manengouba and Mt Bambouto. They constitute with the Bambouto mountains to the SW and the Oku massif to the NE (Fig. 2), the three central volcanoes belonging to the volcanic centres of the Western Cameroon Highlands. Their basement consists of Pan-African granitoids [9, 10, 11]. The Bamenda volcano, on top of which exists the lake Bambili, lies between longitudes 10°00'E

Comment [BCT14]:

Comment [BCT15]: Then, define what is a pyroclastic surge deposit because you found it in your area and will focus your investigation on it.

Comment [BCT16]: This should come more earlier. You can even use it as the first sentence of your introduction.

Comment [BCT17]: Why Only, is there another type? If yes specify otherwise delete.

Comment [BCT18]: Why this conclusion here.
Do you mean that they were interpreted as pyroclastic flows? In that case please state it well and say who interpreted it like that.

Comment [BCT19]:

Formatted: Highlight

and 10°30'E and latitudes 05°45'N and 06°10'N and culminates at 2621 masl at lake Bambili borders.

The Mount Bamenda is characterized by the presence of two calderas: calderas of Santa-Mbu (6 x 4 km width) and Lefo (4 x 3 km width) (Fig. 2). They are small sizes and open respectively to the west and south-east respectively [12]. Their floors located respectively at elevation of 550 m and 400 m is are mainly composed of trachytic domes, which are also abundant on the external slopes of the massif (volcano?). Many craters caused by explosive volcanism, are occupied by lakes generated by phreatomagmatism like Bambili Lake, Lake Awing and Lake Oku.



Fig. 2. Digital Elevation Model (DEM) of Mt Bamenda and surroundings.

Petrographic and geochemical studies [13] show that the mts-Mt Bamenda lavas consist of basanites, basalts, hawaiites, mugearites, benmoreites, trachytes, rhyolites and ignimbrites [13]. These ignimbrite overlays the granite and gneiss basement rock (constituted by granite and gneiss) and are covered by laterized basalt. The felsic lavas are most abundant than intermediate types (mugearites and benmoreites). The radiometric dating of this active volcano gives ages ranging from the current 0 Ma to 17.4 Ma for the basaltic lavas and from 18.98 Ma to 27.40 Ma for the felsic lavas. As in the case of neighboring Mts Bambouto, the ignimbrites of the Bamenda volcano belong to Miocene formations.

3. FIELD (Facies?) STUDY (characteristics?) AND PETROGRAPHY of the Bambili ignimbrite and pyroclastic surge remnants

The ignimbrites sheets of Bamenda volcano constitute about 7.5% of the rocky outcrops of the volcano representing approximately 45 km² with a volume estimated at 6.3 km³; these deposits lie on the trachytic flows and the granito-gneissic basement and covered in places by columnar jointed basaltic flows.

The ignimbrite of Bambili (Fig. 3 and 4) consists of mIT facies in its basal part represented by a single cooling unit constituted by two flow units (dark gray unit and whitish unit), covered by multiple layers of poorly sorted and matrix-poor BrIm facies (Fig. 5). The Pyroclastic surge deposits of Bambili are characterized by stratified ashy layers insert in BrIm facies. Anisotropy of magnetic susceptibility study of Bamenda volcano showed that

Comment [BCT20]: Small compare to what?

Comment [BCT21]: Which massif? the Oku massif? Or the

Comment [BCT22]: Are there references that justified this phreatomagmatic origin? If yes please mentioned them here.

Comment [BCT23]: Reference is needed.

Comment [BCT24]: I think this is the result section already.
Where is the method section?

Formatted: Font: 11 pt, Bold, Highlight

Formatted: Font: 11 pt, Bold, Highlight

Formatted: Font: 11 pt, Bold, Highlight

Comment [BCT25]: This is not a result of the current study and should be used rather for comparison in the discussion section.

Comment [BCT26]: I am confuse here, what are you studying? the ignimbrite or the pyroclastic surge? Please be clear in what you want to show.

In the introduction you said you will investigate in detail the pyroclastic flow but here you emphasise on the ignimbrite.

Comment [BCT27]: What does this means ?

these deposits were emitted from Oku vent (1.8 x 2.2 km) situated in mount Oku, the neighboring NE volcano [3].



Fig. 3. a) Google map of Bambili town showing the ignimbrite outcrops; b) Landscape with ignimbritic remnants of whitish unit of welded mIT lithofacies.




Fig. 4. Ignimbrite outcrops of Bambili. a) and b) Remnant blocks of pyroclastic surge deposits of Bambili; c) Fine-poor matrix of mIBr; d) Remnant block of whitish unit of welded mIT. mIBr: massive lithic breccia; mIT: massive lithic tuff; sT: stratified tuff/ash

3.1 Base of the The Lower unit of the mIT facies

The base of the lower unit of the mIT facies is a flow unit whose color is dark gray and rubefied in places (Fig. 5 and 6a). This rubefaction is reflected by the presence of reddish areas in the deposit. The rock is dense and massive.

Comment [BCT28]: Not your results, should be used in discussion for comparison

Comment [BCT29]: Ameliorate the quality of your figure

Comment [BCT30]: Where is the pyroclastic surge location.

Comment [BCT31]: Show the pyroclastic surge outcrops

Comment [BCT32]: Which rock?




Fig. 5. A) Geological map of mounts Bamenda and surroundings.**B)** Stratigraphic section of Bambillignimbrites; the numbers 24, 25, 26, 33 represents sampled sites for anisotropy of magnetic susceptibility (AMS) studies (after [3]).

The volcanic clast population includes both apyric and porphyritic rhyolite (whose dimensions do not exceed 0.5 x 0.3 cm and represent 5% of the facies) and many apyric black and white fiammes (30 to 35% of the facies; up to 1.2 x 0.2 cm;) which are wispy, lenticular to much stretched and generally distorted displaying a eutaxitic texture; these fiammes are interpreted to be flattened, juvenile pumice lapilli.

Comment [BCT33]: The scripts on the figure are hard to read. Please ameliorate the quality of all your figures.

Comment [BCT34]: Given the title and the objective of your study, a detailed stratigraphic log of the pyroclastic surge section (3b) of this existing log from [3] is necessary.

Comment [BCT35]: What about the pyroclastic surge section in the upper part of this section?

Comment [BCT36]: How do you inferred these values? A method section is needed




Fig. 6. **A)** Sample of rubefied dark grey unit (DGU) of mIT and **b)** the corresponding microphotograph showing the eutaxitic texture with imbricated fiammes; **c)** Crystal of titanohematite with thin exsolution lamellae of ilmenorutile in the DGU; **d)** Sample of mIT whitish unit (WU) and **e)** the corresponding microphotograph showing xenolith of granite; **f)** Crystals of titanohematite with thin exsolution lamellae of ilmenorutile in the WU; **g)** Devitrified matrix of WU showing relict shards, lithic fragments and multiple alkali feldspar fragments; **h)** Basal DGU showing subhorizontal compaction foliation defined by the fiammes and deformation of fiammes around the margins of lithic clasts; **i)** Devitrification of large fiamme in DGU showing micropoikilitic texture characterized by crystallization of microlites (mainly cristobalite and alkali feldspar) along the boundaries of the glass shards or within glassy masses.

Comment [BCT37]: This picture is copy from [3] and should be mentioned and explained why it is used here, otherwise this will be consider as plagiarism

In thin sections, this eutaxitic texture is well characterized by preferentially oriented or imbricated devitrified fiammes showing that transport direction of pyroclastic current was roughly directed toward azimuth 235 (Fig. 6b). The chemistry of these fiammes shows that there are rhyolitic in composition (SiO_2 : 73.65 – 76.44% and $\text{Na}_2\text{O} + \text{K}_2\text{O}$: 8.5 – 9.75%) (Fig. 7). The matrix is devitrified and comprises various proportions of relict shards (Fig. 6g) and lithic fragments (mainly trachytic) with a diameter inferior to the millimeter. The ignimbrite underwent hot welding compaction, as indicated by the subhorizontal compaction foliation defined by the fiamme, deformation of fiamme around the margins of lithic clasts (Fig. 6h), plastic deformation of cusped matrix shards and the presence of spherulitic and micropoikilitic textures (Fig. 6i) within the fiamme (resulting from high-temperature crystallization of a glassy precursor; [14, 15, 16]. The minerals such as quartz and alkali feldspar (sanidine), are abundant; plagioclase, clinopyroxene, biotite and Fe-Ti oxides (ilmenorutile; Fig. 6c) are less represented.

Comment [BCT38]: Please specified if these are thin sections of a juvenile pyroclasts or if these are from a slab of a bed/layer of the deposit. This information is important to interpret the origin and mode of deposition you claimed in the objective of your study




Fig. 7. TAS classification graph [33] for whole rock pumice and vitrophyres samples collected from the Bambili ignimbrites during this study.

Comment [BCT39]: Table of analytical data is need otherwise remove the graph

3. 2. The second unit of the mIT facies

The second unit of the mIT facies is whitish and has low density (Fig. 5 and 6d). The rock is characterized by the presence of numerous vesicular pumice, with chalk-white color and also rhyolitic in composition (Fig. 7; SiO_2 : 74.36 - 76.11% and $\text{Na}_2\text{O} + \text{K}_2\text{O}$: 8.1 - 9.23%). Pumice shapes are subrounded to ovoid and they constitute about 35 to 40% of the volume of the rock; their dimensions vary from 2.2 x 1.9 cm to 0.3 x 0.2 cm. Trachytic enclaves are rare (less than 5%); enclaves of black vitrophyres represent less than 2% of the rock; xenoliths of ignimbrite (2%) are difficult to identify because of their advanced status of alteration. Xenoliths of granite (1%) (Fig. 6e) are generally well preserved in this unit. The dimensions of these enclaves do not exceed 0.4 x 0.2 cm. The matrix is made up of submillimetric devitrify fiammes, small rock fragments smaller than one millimeter and minerals such as: sanidine, plagioclase, biotite and Fe-Ti oxides (ilmenorutile; Fig. 6f). Subvertical elutriation pipes (0.5 to 0.7 cm in diameter) are present in the surface regions of the outcrop.

Comment [BCT40]: Please specify what you call rock in this study. You are describing pyroclastic deposits. The rock in the deposit are both juvenile and lithics thus take care.

3. 2 Surface The upper unit: the Brlm facies

The surface Brlm facies (Fig. 5 and 8a-8a) is represented by an ignimbrite whose overall color is white with a low consistency (Fig. 4a-d) compared to the previous facies. This lithic-rich ignimbrite generated by the second eruption episode has a sharp contact with precedent mIT unit. Consistency depends on the abundance of enclaves, so areas of high rocky enclaves are more resistant to erosion and that's why the remnant blocks are predominantly made up. Stratification is observable in some of these blocks and is characterized by alternating coarse-grained deposit of more than 35 cm thick and fine particle size deposit not exceeding 8 cm of thickness (Fig. 4a). Pyroclastic surge divisions are ash- to lapilli-sized, grey to whitish in colour with low proportion of accretionary lapilli (Fig. 4b). The lithic breccia is generally poorly sorted and diffusely stratified in some outcrop.

Comment [BCT41]: In Fig. 5 this is the pyroclastic surge unit.

Formatted: Highlight

Comment [BCT42]: This is already an interpretation. On the basis of what you decide here that this is a pyroclastic surge ?




Fig. 8. a) Detail of mlBr facies and **b)** microphotograph of ignimbrite enclave **c)** Enclave of vitrophyre and **d)** the corresponding microphotograph.

Rock elements—fragments constituting this ignimbrite are varied in nature. The most represented are enclaves of vitrophyres (25 to 35%) with black color (Fig. 8a,c); some pockets can contain up to 70% of these rock inclusion. Their dimensions range from 22 x 19 cm to 5 x 2.5 cm. These enclaves are essentially prismatic elements with chilled margins. Under microscope, the matrix of vitrophyre is essentially vitreous with some crystals of quartz and feldspar (Fig. 8d). The geochemistry of these vitrophyres (Fig. 7) show that they have rhyolitic in composition (SiO_2 : 71.03 - 73.23% and $\text{Na}_2\text{O} + \text{K}_2\text{O}$: 7.12 - 7.88%). Enclaves of rhyolite are light gray colored and represent 5% of the rock volume with sizes ranging from 14 x 12 cm to 0.8 x 0.6 cm; some elements reach up to 45 x 35 cm. The basement rock enclaves are rare and usually represented by large quartz crystals (11 x 9 cm) constituting about 5% of the rock. Enclaves of first ignimbritic phase (Fig. 8a,b) included in the second phase were found in this facies, it measures 18 x 16.5 cm. These enclaves of ignimbrites contain the same rock elements than the country rock. Matrix, whitish in color, consists essentially of consolidated volcanic ashes in which are embedded small rocks and minerals fragments.

4. DISCUSSION AND CONCLUSION

The studied deposit succession shows evidence of multiple changes of eruptive regime and emplacement modes which likely occurred during a single eruptive event. The stratified character of the deposits observed at Bambili can be attributed to a spatial and temporal variation of the nature of pyroclastic current, which would have been transformed into pyroclastic surge with a current, became becoming weakly concentrated and rich in ashy and gaseous fraction.

In particular, hydromagmatic deposits may form during late eruptive stages following disruption of the top of the magma chamber (e.g., [17]). The presence of vesicular tuff layers, along with dominance of blocky, poorly vesicular ash particles with hydration cracks, indicates an origin as wet pyroclastic surges of hydromagmatic origin [18].

The Brmdeposits unitare is composed of an alternation of stratified clast-supported layers and laminated ash layers, which can be interpreted as a medial-to-distal deposits.

Formatted: Highlight

Formatted: Highlight

Comment [BCT43]: Can you support this assertion with a reference ?

Comment [BCT44]: Please use phreatomagmatic or specify if you are talking about phreatic or phreatomagmatic

Comment [BCT45]: Yes, but what is the relation to the first or following sentence?

Formatted: Highlight

Comment [BCT46]: Nothing have been said before about these particular characteristics.

Comment [BCT47]: Which part of the deposit are you interpreting here?

UNDER PEER REVIEW

produced by ~~hydromagmatic-phreatomagmatic~~ surges and contemporaneous fallout [19]. On other volcanoes, similar deposits were interpreted as fallout deposits or as pyroclastic surge and co-surge fallout deposits, due to the contemporaneous presence of characteristics normally pertaining to both fallout and surge deposits (e.g. [20, 21, 22, 23]). However, ~~it is difficult to discriminate between the two~~ confirm this depositional mechanisms, especially in the absence of ~~a large number of~~ extending outcrops. Nevertheless, given that ~~a single surge layer is composed by a well-graded coarse lapilli-rich bed covered by a more fine ash bed~~ [23, 24], it is likely that our remnant blocks that ~~present the same characteristic were deposited by a phreatomagmatic surge, reflect the~~. In our case, only remnant blocks reflect the presence of pyroclastic surge deposits; in addition, these deposits are partly covered by more recent basaltic lava flow. Nevertheless, a single couplet constituted by both coarse and fine-grained layers is interpreted as deposited from a single surge [23, 24] and this lithofacies is interpreted as inferred to be a medial-to-distal deposit from hydromagmatic surges and concurrent simultaneous fallouts.

The large crater of Mount Oku in which is located the lake Oku, ~~who-which~~ is suspected to be the center of emission of these ignimbrites [3] will justify the presence of accretionary lapilli in the late surge deposits of this volcano.

In the last phase of the eruption, magma has encountered water and thus generated a large explosion which was suspended the huge amount of ash and lithic fragments. Pulsating hydrostatic and magma pressure produced alternating **hydromagmatic** pyroclastic surges. The vitrophyres abundantly found in these deposits would be the lava cooled rapidly by water, and then sprayed by the explosion. The complex dynamics of water-magma interaction defines the nature of explosive activity, characterized by variable energy outputs and different degrees of magmatic or hydromagmatic fragmentation [25, 26, 27, 28, 29]. Depending on the extent of water-magma interaction, deposits can be formed by fallout layers made of couplets of ash and lapilli, by the alternation of fallout layers and surge deposits, or by the simultaneous concurrent deposition from both processes, leading to the formation of highly complex deposits made of numerous (from few tens to over thousands) of thin tephra beds (e.g. [23, 30, 31, 32]). As a consequence, hydromagmatic deposits show remarkable variability in grain-size from layer-to-layer and within layers. Definitely, we inferred that the eruptions of Bamenda volcano were affected by interaction with lake water, generating phreatomagmatic pyroclastic flows and surges. The new chronological succession of geological formations of Bambili can be considered as follows: trachytic flow at the base, follow by welded ignimbrites represented by dark gray and whitish units and pyroclastic surges deposits covered by basaltic flow.

REFERENCES

1. Déruelle B, Ngounouno I, Demaiffe D. The "Cameroon Hot line" (CHL): a unique example of active alkaline intraplate structure in both oceanic and continental lithospheres. *Compte Rendus Géoscience*. 2007;339:589-600.
2. Burke K. Origin of the Cameroon Line of volcano-capped swells. *Journal of Geology*. 2001;109:349-362.
3. GountiéDedzo M, Nédélec A, Nono A, Njanko T, Font E, Kamgang P, Njonfang E, Launeau P. Magnetic fabrics of the Miocene ignimbrites from West-Cameroon: implications for pyroclastic flow source and sedimentation. *Journal of Volcanology and Geothermal Research*. 2011;203:113-132.

Formatted: Highlight

Comment [BCT48]: Please give evidences that let conclude that your remnant blocks are pyroclastic surges

Formatted: Highlight

Formatted: Font: 10 pt, Highlight

Comment [BCT49]: Did you found the lava flow on top the remnants blocks? If yes, How do you explain the presence of lava flow on top the remnant blocks?

Formatted: Highlight

Formatted: Font: 10 pt, Highlight

Comment [BCT50]: Why nevertheless here? You can say that: "According to [23] and [24], a single surge layer is composed by a well-graded coarse lapilli-rich bed covered by a more fine ash bed.

Formatted: Highlight

Formatted: Font: 10 pt, Highlight

Comment [BCT51]: I suggest that you delete this part for a more better comprehension or rephrase.

Comment [BCT52]: Why and how the emission from Mt Oku will explain the presence of accretionary lapilli?

Comment [BCT53]: I cannot see the transition between the begging of eruption and the last phase. Please make it clear.

Formatted: Font: 10 pt, Highlight

Comment [BCT54]: Which units? State is clearly so that someone can follow you up.

Comment [BCT55]: What is the use of this bibliographic description. Which part of the interpretation are you trying to support with this ?

Comment [BCT56]: Why ?

UNDER PEER REVIEW

- 276 4. Dunlop HM. Strontium isotope geochemistry and potassium-argon studies on
277 volcanic rocks from the Cameroon Line, West Africa. PhD Thesis, University of
278 Edinburgh, Edinburgh; 1983.
- 279 5. Lissom J. Etude pétrologique des laves alcalines du massif d'Oku: un ensemble
280 volcanique de la "Ligne du Cameroun". Thèse Université Pierre et Marie Curie,
281 (Paris VI) ; 1991. French.
- 282 6. Kamgang P. Contribution à l'étude géochimique et pétrologique du massif de
283 Nkogam (pays Bamoun, Ouest-Cameroun). Thèse Doctorat 3^e cycle, Université de
284 Yaoundé; 1986. French.
- 285 7. Nono A, Déruelle B, Demaiffe D, Kambou R. Tchabal Nganha volcano in Adamawa
286 (Cameroon): petrology of a continental alkaline lava series. Journal of Volcanology
287 and Geothermal Research. 1994;60:147-178.
- 288 8. Branney MJ, Kokelaar P. Pyroclastic density currents and the sedimentation of
289 ignimbrites. Geological Society of London Memoirs, 2002;27.
- 290 9. Toteu SF, Van Schmus WR, Penaye J, Michard A. New U–Pb and Sm–Nd data from
291 north Cameroon and its bearing on the pre-Pan African history of central Africa.
292 Precambrian Research. 2001;108:45-73.
- 293 10. Nzolang C, Kagami H, Nzenti P, Hotz F. Geochemistry and preliminary Sr–Nd
294 isotopic data on the Neoproterozoic granitoids from the Bantoum Area, West
295 Cameroon: evidence for a derivation from a Paleoproterozoic to Archean crust. Polar
296 Geoscience. 2003;16:196-226.
- 297 11. Nzolang C. Crustal evolution of the Precambrian basement in west Cameroon:
298 inference from geochemistry, Sr–Nd and experimental investigation of some
299 granitoids and metamorphic rocks. Ph.D. thesis, Graduate School of Science and
300 Technology, Niigata University, Japan; 2005.
- 301 12. Gountié Dedzo M, Njonfang E, Kamgang P, Nono A, Zangmo Tefogoum G, Kagou
302 Dongmo A, Nkouathio DG. Dynamic and evolution of the Mounts Bambouto and
303 Bamenda calderas by study of ignimbritic deposits (West-Cameroon, Cameroon
304 Line). Syllabus Review, Science Series. 2012;3:11-23.
- 305 13. Kamgang P, Njonfang E, Nono A, Gountié Dedzo M, Tchoua FM. Petrogenesis of a
306 silicic magma system: Geochemical evidence from Bamenda Mountains, NW
307 Cameroon, Cameroon Volcanic Line. Journal of African Earth Sciences.
308 2010;58:285-304.
- 309 14. Lofgren G. Experimentally produced devitrification textures in natural rhyolite glass.
310 Geological Society of America Bulletin, 1971a;82:553-560.
- 311 15. Lofgren G. Spherulitic textures in glassy and crystalline rocks. Journal of
312 Geophysical Research. 1971b;76:5635-5648.
- 313 16. Lofgren G. An experimental study of plagioclase crystal morphology: isothermal
314 crystallization. American Journal of Science. 1974;274:243-273.
- 315 17. Sigurdsson H, Carey S, Cornell W, Pescatore T. The eruption of Vesuvius in A.D. 79,
316 National Geographic Research. 1985;1:332-387.
- 317 18. Palladino DM, Simeì S. Eruptive dynamics and caldera collapse during the Onano
318 eruption, Vulturno, Italy. Bulletin of Volcanology. 2005;67:423-440.
- 319 19. Zanon V, Pacheco J, Pimentel A. Growth and evolution of an emergent tuff cone:
320 Considerations from structural geology, geomorphology and facies analysis of São
321 Roque volcano, São Miguel (Azores). Journal of Volcanology and Geothermal
322 Research 2009;180:277-291.
- 323 20. Wohletz KH, Sheridan MF. A model of pyroclastic surge. Geological Society of
324 American Special Paper. 1979;180:177-194.
- 325 21. Fisher RV, Schmincke HU. Pyroclastic rocks. Springer-Verlag, New York. 1984.
- 326 22. Cole PD, Guest JE, Duncan AM, Pacheco JM. Capelinhos 1957–1958, Faial,
327 Azores: deposits formed by an emergent surtseyan eruption. Bulletin of Volcanology.
328 2001;63:204-220.

Formatted: Font: 10 pt, French (France)

Formatted: Font: 10 pt, French (France)

UNDER PEER REVIEW

- 329 23. Dellino P, Isaia R, La Volpe L, Orsi G. Interaction between particles transported by
330 fallout and surge in the deposits of the Agnano–Monte Spina eruption (CampiFlegrei,
331 Southern Italy). *Journal of Volcanology and Geothermal Research*. 2004;133:193-
332 210.
- 333 24. Sohn YK, Chough SK. Depositional processes of the Suwolbong tuff ring, Cheju
334 Island (Korea). *Sedimentology*. 1989;36:837-855.
- 335 25. Wohletz KH, Sheridan MF. Hydrovolcanic eruptions: II. Evolution of basaltic tuff rings
336 and tuff cones. *American Journal of Science*. 1983;283:385-413.
- 337 26. Houghton BF, Hackett WR. Strombolian and phreatomagmatic deposits of Ohakune
338 craters, Ruapehu, New Zealand: a complex interaction between external water and
339 rising basaltic magma. *Journal of Volcanology and Geothermal Research*.
340 1984;21:207-231.
- 341 27. Kokelaar P. Magma–water interactions in subaqueous and emergent basaltic
342 volcanism. *Bulletin of Volcanology*. 1986;48:275-289.
- 343 28. White JDL, Houghton B. Surtseyan and related phreatomagmatic eruptions. In:
344 Sigurdsson, H., Houghton, B.F., McNutt, S.R., Rymer, H., Stix, J. (Eds.),
345 *Encyclopedia of Volcanoes*, Academic Press, San Diego. 2000;495-511.
- 346 29. Mastin LG, Christiansen RL, Thornber C, Lowenstern J, Beeson M. What makes
347 hydromagmatic eruptions violent? Some insights from the Keanakākoī Ash, Kīlauea
348 Volcano, Hawaii. *Journal of Volcanology and Geothermal Research*. 2004;137:15-31.
- 349 30. Chough SK, Sohn YK. Depositional mechanics and sequences 705 of base surges,
350 Songaksan tuff ring, Cheju Island, Korea. *Sedimentology*. 1990;37:1115-1135.
- 351 31. Dellino P, La Volpe L. Structures and grain size distribution in surge deposits as a
352 tool for modelling the dynamics of dilute pyroclastic density currents at La Fossa di
353 Vulcano. *Journal of Volcanology and Geothermal Research*. 2000;96:57-78.
- 354 32. Dellino P, Isaia R, La Volpe L, Orsi G. Statistical analysis of textural data from
355 complex pyroclastic sequences: implications for fragmentation processes of the
356 Agnano–Monte Spina tephra (4.1 ka), CampiFlegrei, southern Italy. *Bulletin of*
357 *Volcanology*. 2001;63:443-461.
- 358 33. Le Bas M, Le Maitre R, Streckeisen A, Zanettin B. A chemical classification of
359 volcanic rocks based on the total alkali–silica diagram. *Journal of Petrology*.
360 1986;27:745-750.
- 361