

Original Research Article

Synthesis of novel imidazole derivatives based on camalexin scaffold and anti-fungal activity against rice blast

ABSTRACT (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)

Aims: Rice blast disease (RBD), which is caused by the pathogenic fungi of *Magnaporthe oryzae*. Plant diseases caused by plant pathogenic fungi is a leading constraint in world's crop production. One of the most devastating diseases of cultivated rice (*Oryza sativa* L) is the rice blast disease (RBD), which is caused by the pathogenic fungi of *Magnaporthe oryzae*. Although many strategies have been applied to control the RBD, chemical anti-fungicide fungal agents is are the most common solution effectively solutions to minimize the severity of RBD. The aim of this study is to explore new anti-fungal agents for RBD control.

Study design: Natural products have been widely used as are rich important source for of biological active compounds. Thus, we used camalexin (3-thiazol-2'-yl-indole), which is a major well known phytoalexin natural product of *Arabidopsis thaliana* with potent anti-fungal activity, as a molecular scaffold to In the present work, we designed a new series of imidazoles based on camalexin as a molecular scaffold and the anti-fungal activity of the synthesized compounds against *Magnaporthe oryzae* were as candidates of new agents for rice blast control determined.

Place and Duration of Study: All the experiments were conducted from October 2014 to March 2015 at Akita Prefectural University, Japan.

Methodology: A series of new imidazole derivatives was designed and synthesized based on the camalexin as a molecular scaffold. The antifungal activity of the synthesized compounds against *Magnaporthe oryzae* was determined by using *in vitro* mycelial growth inhibition tests.

Results: Among 3 newly synthesized camalexin based imidazole compounds, we found that 3-[3-(1*H*-imidazol-1-yl) propyl]-1*H*-Indole (**2c**) exhibits potent inhibitory activity with an IC₅₀ value approximately 7.8±1.2 μM.

Conclusion: We discovered a new lead compound with potent activity for rice blast control.

Keywords: camalexin, antifungal agent, imidazole derivatives, rice blast

1. INTRODUCTION

Plants experience a broad range of environmental stresses including biotic and abiotic stresses are major external stimuli which greatly affect the growth of plants. Unfavorable temperature, drought, chemical pollutions are major abiotic stresses. Among the biotic stresses, pathogen infection is one of the most severe stresses. Rice blast diseases (RBD) caused by pathogenic fungi of *Magnaporthe oryzae* is a leading constraint in world's rice production [1-3]. The high incidence of plant mortality and the lack of effective control methods make it responsible for billions of dollars losses worldwide each year.

To control the RBD, tremendous efforts have been made and various management strategies have been applied [4], including controlled use of nitrogen fertilizers, application of silica and flooding of paddy fields have been applied in use [4]. Currently, chemical fungicide is the most common solution effectively to minimize the severity of RBD. Many systemic

fungicides with different mode of actions have been used for RBD control. Like anti-mitotic fungal compounds [5], melanin inhibitors [6], ergosterol biosynthesis inhibitor (EBI) and other organic compounds were discovered and applied have been widely used for RBD control [7]. With the increasing use of chemical fungicides, the appearance of pathogens that resistant to existing fungicides raised new challenges for diseases control in agricultural industry. Hence, development new anti-microbial agent for RBD control is of significant importance in modern agricultural industry.


Different from vertebrate, the cell order and integrity of fungal membrane are significantly important in fungi. Hence, chemicals that directly or indirectly target fungal membranes functions or their components synthesis are quite effective to cause the lethality of the fungi thereby controlling the fungal disease. Some of these anti-fungal compounds affect the synthesis of specific membrane components (e.g., sterol biosynthesis inhibitors) are among the most effective fungicides in plant disease control.

To meet the demands for plant diseases control, variety strategies have been applied for fungicides development. Natural products are useful sources of novel chemical structures for development of new fungicides [8-11]. Until now, major advances in modern medicine and the pharmaceutical industry were driven by natural products or their derivatives [12]. Many experts believe that using natural products as leads is a feasible way to find the core structure of biological active compounds in drug discovery [13].

Based on these facts, using the core structure of natural products which display anti-microbial activity is a straightforward approach to explore new agents for RBD control. Phytoalexins are heterogeneous low molecular mass secondary metabolites with antimicrobial activity at the infection site in response to pathogen invasion [14, 15].

The biosynthesis of phytoalexins is commonly regulated by a complex mechanism of plant defense signal transduction networks. The plant hormones, jasmonic acid, salicylic acid and ethylene, are important signal mediators that involve in plant defense response and phytoalexins synthesis [16]. These plant hormones activate the expression of defense phytoalexins synthesis genes, causing rapidly accumulation of phytoalexins at the areas of pathogen infection [17]. Indeed, the structural diversity of phytoalexins are in different plant species and most of the phytoalexins are belong to terpenoids [18], glycosteroids and alkaloids [19].

One of the well-known phytoalexin is camalexin (3-thiazol-2'-yl-indole, the structure is shown in figure 1), a primary phytoalexin of Arabidopsis thaliana [20], is induced by a great variety of plant pathogens. The biosynthesis and the regulation of the biosynthesis of camalexin have been studied extensively [21, 22]. It has been demonstrated that camalexin is biosynthesized from tryptophan [23] and its biosynthesis involves the cytochrome P450 enzymes CYP79B2 [24] and CYP71B15 (PAD3) [25]. Accumulation of camalexin upon pathogen infection, as well as its anti-microbial nature supports the role in disease resistance [26].



n=1,2,3

Camalexin

Target compounds

Figure 1. Design of new imidazole derivatives based on camalexin scaffold.

To explore new agents for RBD control, we report ~~herein a chemical approach of the~~ synthesis ~~of~~ new compounds based on the camalexin scaffold. ~~The indole structure of camalexin is proposed to be derived from tryptophan biosynthesis intermediate of indole-3-glycerol phosphate. Hence we~~ design the target compound with an indole skeleton. ~~On the other hand, The the~~ thiazol moiety was replaced by an imidazole group with a spacer structure as shown in figure 1. The biological activity of synthesized compounds against *Magnaporthe oryzae* was evaluated by using in vitro mycelial growth inhibition tests.


2. MATERIALS AND METHODS

2.1 Subheading Subheading

Stock solutions of all the test compound were dissolved in DMSO at a concentration of 100 mM and stocked at -30 °C. Other reagents were of the highest grade and purchased from Wako Pure Chemical Industries, Ltd. (Tokyo, Japan). ¹H-NMR spectra were recorded with a JEOL ECP-400 spectrometer (Tokyo, Japan), chemical shifts being expressed in ppm downfield from TMS as an internal standard. High resolution electrospray ionization Fourier transform ion cyclotron resonance mass spectra (ESI-FTICR) were recorded on an Exactive MS System (Thermo Fisher Scientific, Waltham, MA, USA).

2.2 Chemical synthesis

Preparation of 3-(1H-imidazol-1-ylmethyl)-1H-Indole (2a) was carried out by a method as previously described [11]. a solution of 1H-indole-3-methanol (1a; 294 mg, 2 mmol) and CDI (420 mg, 2.6 mmol, 1.3 equiv) in anhyd MeCN (5 ml) was refluxed for 5 h (the reactions were monitored by TLC and terminated after complete consumption of the hydroxyl compound) the reaction mixture was diluted with cold water (20 ml) and the resulting precipitate was collected, washed with water and dried under vacuum at 40-50 °C. recrystallization were carried out by using anhyd MeCN to give pure 3-(1H-imidazol-1-ylmethyl)-1H-Indole (2a) (Yield 92%), ¹H NMR(CDCl₃), δ : 5.31 (2H, s), 6.96 (1H, s), 7.05 (1H, s), 7.13 (1H, t, J=7.1), 7.18 (1H, d, J=2.3), 7.23(1H, d, J=8.2), 7.43 (2H, q, J₁=8.1, J₂=12.0), 7.59(1H, s), 8.23(1H, s).



101	n=2: 1b	n=2: 2b
102	n=3: 1c	n=3: 2c

103
104
105
106
107

Scheme 1. Synthesis of target compounds: reagents and conditions: (a), CDI; CH₃CN (anhydride), reflux, 5 hour.

108 Compound 2b and 2c were prepared in a similar way by using different alcohols as a starting
109 material as shown below.

110 3-[2-(1H-imidazol-1-yl) ethyl]-1H-Indole (2b). (Yield 90%), ¹H NMR (CDCl₃), δ : 3.28 (2H, t,
111 J=7.1), 4.68 (2H, t, J=7.1), 7.06(1H, t, J=1.3), 7.10 (1H, d, J=2.3), 7.16 (1H, d, J=0.92), 7.18-
112 7.25 (1H, m), 7.39-7.41 (2H, m), 7.64 (1H, d, J=8.0), 8.11 (2H, d, J=6.4). 2c were prepared
113 in a similar way by using corresponding indole derivatives.

114 3-[3-(1H-imidazol-1-yl) propyl]-1H-Indole (2c). (Yield 94%), ¹H NMR (CDCl₃), δ : 2.20-2.27
115 (2H, m), 2.94 (2H, t, J=7.5), 4.48 (2H, t, J=6.5), 7.03 (1H, d, J=2.3), 7.07 (1H, q, J1=0.92,
116 J2=1.6), 7.13 (1H, t, J=0.92), 7.15-7.23 (1H, m), 7.38-7.39 (2H, m), 7.60 (1H, d, J=8.2), 8.05
117 (2H, d, J=23.8).

118 **2.3 Magnaporthe-oryzae strain**

119 Pathogenic fungi of Rice blast (*Magnaporthe oryzae*) isolate designated APU00-093A (race
120 007.0) was obtained by mono-spore isolation from diseased rice panicle on the paddy field
121 of Akita Prefecture Japan in 2000. This isolate was kept on potato dextrose agar at 15oC.

122 **2.4 Antifungal activity assay**

123 Poisoned food technique was performed to investigate antifungal effect of test compounds
124 against *Magnaporthe oryzae*. The Mycelial growth inhibition tests were carried out. Each test
125 compound dissolved and diluted in DMSO was added to potato sucrose agar (PSA) medium
126 (kept at 50oC after autoclaving) to the appropriate concentration. The final concentration of
127 DMSO of each medium was 0.1%. Three mycelial pellet (1mm in diameter) of *Magnaporthe*
128 *oryzae* pre-cultured on potato dextrose agar (PDA) medium were placed on the PSA
129 medium containing the given concentrations of the test compound. The diameter of the
130 mycelial mat of *Magnaporthe oryzae* was measured when the diameter of each
131 corresponding untreated control reached about 20-30mm. Concentration for 50% inhibition
132 (IC₅₀, μM) of mycelial growth was calculated by the linear regression formula obtained from
133 the logarithm of concentration and the inhibition rate at each concentration against untreated
134 control. All experiments were carried out in triplicate and the report data represents average
135 values.

136 **3. RESULTS AND DISCUSSION**

137 **3.1 Chemistry**

138 Method for synthesis target compounds is shown in Scheme 1 in a condition as previously
139 reported [27]. 3-Indolemethanol (1a) and its derivatives (2b, 2c) were used as starting
140 materials. Compound 2a was prepared by reacting compound 1a with 1.3 equivalent of 1, 1'-
141 carbonylbis-1H-imidazole in anhydride acetonitrile under a condition of reflux for 5 hours.
142 The reaction was monitored by using TLC until the starting material 1a is disappeared.

Compound 2b and 2c were prepared in a similar way by using 1b and 1c as starting materials.

3.2 Biology

The chemical structures of the test compounds were listed in Table 1. The concentration of all the test compounds were adjusted to a final concentration of 100 μM in potato sucrose agar (PSA) medium while propiconazole was used as a positive control. The key structure




Figure 2. Antifungal activity of compound 2c against *Magnaporthe oryzae*. (A) Dose dependent effect of 2c on growth of *Magnaporthe oryzae*, filled circle: compound 2c; filled diamond: propiconazole. (B) Effect of compound 2c on growth of *Magnaporthe oryzae*, upper left: 10 μM propiconazole, upper right: control; lower left: 1 mM 2c, lower second from left: 100 μM 2c, lower second from right, 10 μM 2c, lower right, 1 μM . All the experiments were taken three times to establish the repeatability.



were the indole moiety and an imidazole moiety in this synthetic series. Thus, we introduced three kinds of chain structure with different carbon atoms. As shown in table 1, introducing single carbon atom (2a) exhibited antifungal activity with a fungal growth rate at approximately $48.2 \pm 4.2\%$. When two carbon atoms were introduced (2b), we found that the antifungal activity was significantly decreased. The fungal growth rate was found approximately $85.6 \pm 8.2\%$ compared to none chemical treated control. Interestingly, when


three carbon atoms chain was introduced (**2c**), we found that the antifungal activity was enhanced greatly. The fungal growth rate was found approximately 18.2±1.6%.

Next, we used compound **2c** to determine the dose-dependent effect of antifungal activity against *Magnaporthe oryzae* of this synthetic series. As shown in Figure 2, compound **2c** inhibits *Magnaporthe oryzae* growth in a dose dependent manner. The IC₅₀ was found approximately 7.8 ± 1.2 μM. while the IC₅₀ of the positive control of propiconazole was found approximately 3.7 ± 0.2 μM in our assay system.

4. CONCLUSION

In the present work, we synthesized a series of new imidazole derivatives based on the

TABLE I: ANTIFUNGAL ACTIVITY OF TEST COMPOUNDS




Compound No.	n	Fungal growth (%) [*]
2a	1	48.2±4.2
2b	2	85.6±8.2
2c	3	18.2±1.6
Propiconazole		3.3±2.8

^{*}The concentration of all the test compounds and positive control Propiconazole were 100 μM. Fungal growth was determined by measure the mycelial diameter of none chemical treated line as 100%. All the experiment were done three times to establish the repeatability.

camalexin (3-thiazol-2'-yl-indole) scaffold. Camalexin is a primary phytoalexin of *Arabidopsis thaliana* that display antifungal activity against many kinds of pathogenic fungi. Evaluation the antifungal activity of newly synthesized imidazole derivatives against *Magnaporthe oryzae* indicated that the carbon number of the chain structure between indole and imidazole moiety significantly influence the antifungal activity. Among the synthesized compounds, compound **2c** exhibits potent antifungal activity with an IC₅₀ value approximately 7.8 ± 1.2 μM. The antifungal potency was found approximately half of that of propiconazole. We expect further structure-activity relationship studies may provide in slight information about the structural requirements of this synthetic series on antifungal activity against rice blast pathogen *Magnaporthe oryzae*.

TABLE I: ANTIFUNGAL ACTIVITY OF TEST COMPOUNDS



<u>Compound No.</u>	<u>n</u>	<u>Fungal growth (%)*</u>
<u>2a</u>	<u>1</u>	<u>48.2±4.2</u>
<u>2b</u>	<u>2</u>	<u>85.6±8.2</u>
<u>2c</u>	<u>3</u>	<u>18.2±1.6</u>
<u>Propiconazole</u> <u>e</u>		<u>3.3±2.8</u>

**The concentration of all the test compounds and positive control Propiconazole were 100 µM. Fungal growth was determined by measure the mycelial diameter of none chemical treated line as 100%. All the experiment were done three times to establish the repeatability.*

4. CONCLUSION

In the present work, we synthesized a series of new imidazole derivatives based on the camalexin (3-thiazol-2'-yl-indole) scaffold. Camalexin is a primary phytoalexin of Arabidopsis thaliana that display antifungal activity against many kinds of pathogenic fungi. Evaluation the antifungal activity of newly synthesized imidazole derivatives against Magnaporthe oryzae indicated that the carbon number of the chain structure between indole and imidazole moiety significantly influence the antifungal activity. Among the synthesized compounds, compound 2c exhibits potent antifungal activity with an IC50 value approximately 7.8 ± 1.2 µM. The antifungal potency was found approximately half of that of propiconazole. We expect further structure-activity relationship studies may provide in slight information about the structural requirements of this synthetic series on antifungal activity against rice blast pathogen Magnaporthe oryzae.

REFERENCES

- 1 Talbot, NJ. On the trail of a cereal killer: Exploring the biology of Magnaporthe grisea. Annu. Rev. Microbiol. Vol 57, pp.177-202, June 2003.
- 2 Clergeot, PH., Gourgues, M., Cots, J., Laurans, F., Latorse, MP., Pepin, R., Tharreau, D., Notteghem, JL. and Lebrun, MH. PLS1, a gene encoding a tetraspanin-like protein, is required for penetration of rice leaf by the fungal pathogen Magnaporthe grisea. Proc. Natl. Acad. Sci. USA. Vol 98, pp.6963-6968. March, 2001.
- 3 Bechinger, C., Giebel, KF., Schnell, M., Leiderer, P., Deising, HB. and Bastmeyer, M. Optical measurements of invasive forces exerted by appressoria of a plant pathogenic fungus. Science, Vol 285, pp.1896-1899. Sep. 1999.
- 4 Pageau K., Reisdorf-Cren M., Morot-Gaudry JF. and Masclaux-Daubresse C. The nutrient supply of pathogenic fungi, a fertile field for study. Mol. Plant Pathol. Vol 4, pp.203–210. May, 2003.
- 5 Davidse, LC. Benzimidazole Fungicides: Mechanism of Action and Biological Impact, Annu. Rev. Phytopathol. Vol. 24, pp. 43-65. Sept. 1986.
- 6 Knight, SC., Anthony, VM., Brady, AM., Greenland, AJ., Heaney, SP., Murray, DC., Powell, KA., Schulz, MA., Spinks, CA., Worthington, PA. and Youle, D. Rationale and

- 250 perspectives on the development of fungicides. *Annu. Rev. Phytopathol.* Vol. 35, pp.
251 349-372. Sept. 1997.
- 252 7 Baldwin, BC. and Rathmell, W. G. Evolution of concepts for chemical control of plant
253 disease. *Annu. Rev. Phytopathol.* vol. 26, pp. 265-283. Sept. 1988.
- 254 8 Shu YZ. Recent natural products based drug development: a pharmaceutical industry
255 perspective. *J Nat Prod.* Vol. 61(8), pp. 1053-1071. Aug. 1998.
- 256 9 Dias DA., Urban S. and Roessner U., A Historical Overview of Natural Products in Drug
257 Discovery. *Metabolites* vol. 2, pp. 303-336, Apr. 2012.
- 258 10 Lahlou M., The Success of Natural Products in Drug Discovery. *Pharmacology &
259 Pharmacy*, vol. 4, pp. 17-31. Aug. 2013.
- 260 11 Ji HF., Xue-Juan Li, XU. and Zhang HZ. Natural products and drug discovery. Can
261 thousands of years of ancient medical knowledge lead us to new and powerful drug
262 combinations in the fight against cancer and dementia? *EMBO Rep.* vol. 10 (3),
263 pp. 194–200, Mar. 2009.
- 264 12 Kingston DGI. Modern Natural Products Drug Discovery and its Relevance to
265 Biodiversity. *J Nat Prod.* Vol. 74 (3), pp. 496–511. Mar. 2011.
- 266 13 Cragg GM. and Newman DJ. NATURAL PRODUCTS: A CONTINUING SOURCE OF
267 NOVEL DRUG LEADS. *Biochim Biophys Acta.* Vol. 1830(6), pp. 3670–3695. Jun. 2013.
- 268 14 Rocío González-Lamothe, Gabriel Mitchell, Mariza Gattuso, Moussa S. Diarra, François
269 Malouin, and Kamal Bouarab, Plant Antimicrobial Agents and Their Effects on Plant
270 and Human Pathogens. *Int J Mol Sci.* vol. 10(8), pp. 3400–3419, Aug. 2009.
- 271 15 Found Symp. Vol. 223, pp. 175-187, Aug. 1999.
- 272 16 Pieterse, CMJ. Antonio LR., Sjoerd VE. and Saskia CM. Wees. V. Networking by small-
273 molecule hormones in plant immunity. *Nature Chemical Biology* vol. 5, pp. 308-316 Apr.
274 2009.
- 275 17 Jeandet P., Douillet-Breuil AC., Bessis R., Debord S., Sbaghi M., and Adrian M.
276 Phytoalexins from the Vitaceae: biosynthesis, phytoalexin gene expression in
277 transgenic plants, antifungal activity, and metabolism. *J. Agric. Food Chem.* Vol. 50(10),
278 pp. 2731-2741. Oct. 2007.
- 279 18 Vaughan MM., Christensen S., Schmelz EA., Huffaker A., McAuslane HJ., Alborn HT.,
280 Romero M., Allen LH., and Teal PE. Accumulation of terpenoid phytoalexins in maize
281 roots is associated with drought tolerance. *Plant Cell Environ.* 2014 Nov 13. doi:
282 10.1111/pce.12482.
- 283 19 Angelova S., Buchheim M., Frowitter D., Schierhorn A., and Roos W. Overproduction of
284 alkaloid phytoalexins in California poppy cells is associated with the co-expression of
285 biosynthetic and stress-protective enzymes. *Mol Plant.* Vol. 3(5), pp. 927-939. Sep.
286 2010.
- 287 20 Glawischnig E. Camalexin. *Phytochem.* Vol. 68(4), pp. 401-416. Feb. 2007.

- 288 21 Lin YM., Shih SL., Lin WC., Wu JW., Chen YT., Hsieh CY., Guan LC., Lin L., and
289 Cheng CP. Phytoalexin biosynthesis genes are regulated and involved in plant
290 response to *Ralstonia solanacearum* infection. *Plant Sci.* vol. 224, pp. 86-94. Jul. 2014.
- 291 22 Ruszkowska J., and Wróbel JT. Tryptophan-derived sulfur-containing phytoalexins--a
292 general overview. *Adv. Exp. Med. Biol.* Vol. 527, pp. 629-636. Aug. 2003.
- 293 23 Böttcher C., Westphal L., Schmotz C., Prade E., Scheel D., and Glawischnig E. The
294 multifunctional enzyme CYP71B15 (PHYTOALEXIN DEFICIENT3) converts cysteine-
295 indole-3-acetonitrile to camalexin in the indole-3-acetonitrile metabolic network of
296 *Arabidopsis thaliana*. *Plant Cell* Vol. 21(6), pp.1830-1845. Jun. 2009.
- 297 24 Mikkelsen MD., Fuller VL, Hansen BG., Nafisi M., Olsen CE., Nielsen HB., and Halkier
298 BA. Controlled indole-3-acetaldoxime production through ethanol-induced expression of
299 CYP79B2. *Planta* vol. 229(6), pp. 1209-1217. May. 2009.
- 300 25 Nawrath C., and Métraux JP. Salicylic acid induction-deficient mutants of *Arabidopsis*
301 express PR-2 and PR-5 and accumulate high levels of camalexin after pathogen
302 inoculation. *Plant Cell* vol. 11(8), pp. 1393-1404, Aug. 1999.
- 303 26 Ahuja I., Kissen R., and Bones AM. Phytoalexins in defense against pathogens. *Trends*
304 *Plant Sci.* vol. 17(2), pp. 73-90. Feb. 2012.
- 305 27 Njar V.CO. High-yield synthesis of novel imidazoles and triazoles from alcohols and
306 phenols. *Synthesis* vol. 14, pp. 2019-2028. Oct. 2000.

307