

An iterative method for approximating the common solutions of nonexpansive semigroups, a system of variational inequalities , variational inequalities and equilibrium problems

Meng Wen , Changsong Hu*

Department of Mathematics, Hubei Normal University, Huangshi, 435002, P.R.China,

Abstract. In this work, we consider an general iterative method for finding a common element of the set of common fixed points of a one-parameter nonexpansive semigroup problem, a system of variational inequality problems, a variational inequality problem and the set of solutions of a suitable equilibrium problem in a real Hilbert space. Further we establish a strong convergence theorem based on this method. Our results improve and extend corresponding ones announced by many others.

Keywords. Nonexpansive semigroup; Equilibrium problem; Variational inequality problem; System of variational inequality problems; Projection

MSC. 47H09; 47H10

1 Introduction

Throughout the paper unless otherwise stated, let H be a real Hilbert space with norm $\|\cdot\|$ and inner product $\langle \cdot, \cdot \rangle$. Let C be a nonempty, closed and convex subset of H . Let $\{x_n\}$ be any sequence in H , then $x_n \rightarrow x$ (respectively, $x_n \rightharpoonup x$) will denote strong (respectively, weak) convergence of the sequence $\{x_n\}$.

A mapping $T : C \rightarrow C$ is called nonexpansive if

$$\|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in C. \quad (1.1)$$

One parameter family $\Gamma := \{T(t) : 0 \leq t < \infty\}$ is said a (continuous) Lipschitzian semigroup on C of mappings from C into C if the following conditions are satisfied:

- (1) $T(0)x = x$ for all $x \in C$;
- (2) $T(s+t) = T(s)T(t)$ for all $s, t \geq 0$;
- (3) for each $t > 0$, there exists a bounded measurable function $L_t : (0, \infty) \rightarrow [0, \infty)$ such that $\|T(t)x - T(t)y\| \leq L_t\|x - y\|$, $x, y \in C$;
- (4) for each $x \in C$, the mapping $T(\cdot)x$ from $[0, \infty)$ into C is continuous.

A Lipschitzian semigroup Γ is called nonexpansive (or contractive) if $L_t = 1$ for all $t > 0$ and asymptotically nonexpansive if $\limsup_{t \rightarrow \infty} L_t \leq 1$, respectively. Let $F(\Gamma)$ denote the common fixed point set of the semigroup Γ , i.e., $F(\Gamma) := \{x \in C : T(t)x = x, \forall t > 0\}$.

Let $A : C \rightarrow H$ be a nonlinear mapping. Then A is called

- (1) monotone, if

$$\langle Ax - Ay, x - y \rangle \geq 0, \quad \forall x, y \in H;$$

* Corresponding author.

E-mail address: wen5495688@163.com(M.Wen), huchang1004@yahoo.com.cn(C.S.Hu)

(2) α -strongly monotone, if there exists a constant $\alpha > 0$ such that

$$\langle Ax - Ay, x - y \rangle \geq \alpha \|x - y\|^2, \quad \forall x, y \in H;$$

(3) α -inverse strongly monotone, if there exists a constant $\alpha > 0$ such that

$$\langle Ax - Ay, x - y \rangle \geq \alpha \|Ax - Ay\|^2, \quad \forall x, y \in H;$$

(4) k -Lipschitz continuous, if there exists a constant $k > 0$ such that

$$\|Ax - Ay\| \leq k \|x - y\|, \quad \forall x, y \in H.$$

An operator T is strongly positive on H if there is a constant $\bar{\gamma}$ with property

$$\langle Tx, x \rangle \geq \bar{\gamma} \|x\|^2, \quad \forall x \in H.$$

Given a nonlinear mapping $\varphi : C \rightarrow H$. Then the variational inequality problem (in short, VIP) is to find $x \in C$ such that

$$\langle \varphi x, y - x \rangle \geq 0, \quad \forall y \in C. \quad (1.1)$$

The solution of VIP(1.1) is denoted by $VI(C, \varphi)$. It is well known that if φ is strongly monotone and Lipschitz continuous mapping on C then VIP(1.1) has a unique solution. There are several different approaches towards solving this problem in finite dimensional and infinite dimensional spaces see, e.g. [1-9] and the research in this direction is intensively continued.

Next we consider the following system of variational inequality problems (in short, SVIP): find $(x^*, y^*) \in C \times C$ such that

$$\begin{cases} \langle \rho_1 B_1 y^* + x^* - y^*, x - x^* \rangle \geq 0, & \forall x \in C, \\ \langle \rho_2 B_2 x^* + y^* - x^*, x - y^* \rangle \geq 0, & \forall x \in C, \end{cases} \quad (1.2)$$

where $B_i : C \rightarrow C$ is a nonlinear mapping and $\rho_i > 0$ for each $i = 1, 2$. The set of solutions of SVIP(1.2) is denoted by Ψ .

Some special cases of SVIP(1.2):

(i) If $B_1 = B_2 = B$ then SVIP(1.2) is reduced to the system of problems of finding $(x^*, y^*) \in C \times C$ such that

$$\begin{cases} \langle \rho_1 B y^* + x^* - y^*, x - x^* \rangle \geq 0, & \forall x \in C, \\ \langle \rho_2 B x^* + y^* - x^*, x - y^* \rangle \geq 0, & \forall x \in C, \end{cases} \quad (1.3)$$

which is considered and studied by Verma [10].

(ii) If $x^* = y^*$ in problem (1.3) then problem (1.3) is reduced to the following classical variational inequality problem of finding $x^* \in C$ such that

$$\langle Bx^*, x - x^* \rangle \geq 0, \quad \forall x \in C.$$

In order to find the solution of SVIP(1.2), Ceng et al. [11] introduced an approximation method known as relaxed extragradient method. Some related works, we refer to see [12-14, 28-33].

Let F be a bifunction from $C \times C$ to R , where R is the set of real numbers. Moudafi in [15] studies the equilibrium problem

$$\text{to find } x^* \in C \quad \text{such that } F(x^*, y) + \langle Ax^*, y - x^* \rangle \geq 0, \quad \forall y \in C$$

where A is an α -inverse strongly monotone operator. In [16-18], the authors study the mixed problem

$$\text{to find } x^* \in C \quad \text{such that } F(x^*, y) + \varphi(x^*) - \varphi(y) \geq 0, \quad \forall y \in C$$

with φ being an opportune mapping.

Here, we study the equilibrium problem

$$\text{to find } x^* \in C \quad \text{such that } F(x^*, y) + h(x^*, y) \geq 0, \quad \forall y \in C$$

that includes all previous equilibrium problems as particular cases.

In this paper motivated by the work of K.R. Kazmi and S.H. Rizvi [19] and Giuseppe Marino, Luigi Muglia and YongHong Yao [20], we introduce a iterative scheme for finding a common element of the set of common fixed points of a one-parameter nonexpansive semigroup, a system of variational inequality problems, a variational inequality problem and the set of solutions of suitable equilibrium problem in a real Hilbert space. We establish a strong convergence theorem based on this method. Our results extend and improve the corresponding results of K.R. Kazmi and S.H. Rizvi [19] and many others.

2 Preliminaries

Let H be a real Hilbert space. It is well known that

$$\|x - y\|^2 = \|x\|^2 - \|y\|^2 - 2\langle x - y, y \rangle \quad (2.1)$$

and

$$\|\lambda x + (1 - \lambda)y\|^2 = \lambda\|x\|^2 + (1 - \lambda)\|y\|^2 - \lambda(1 - \lambda)\|x - y\|^2 \quad (2.2)$$

for all $x, y \in H$ and $\lambda \in [0, 1]$.

Let C be a nonempty closed convex subset of H . For every point $x \in H$, there exists a unique nearest point in C , denoted by $P_C x$, such that

$$\|x - P_C x\| \leq \|x - y\| \quad \text{for all } y \in C. \quad (2.3)$$

P is called a metric projection of H onto C . It is well known that P_C is a nonexpansive mapping of H onto C and satisfies

$$\langle x - y, P_C x - P_C y \rangle \geq \|P_C x - P_C y\|^2 \quad (2.4)$$

for every $x, y \in H$. Moreover, $P_C x$ is characterized by the following properties: $P_C x \in C$ and

$$\langle x - P_C x, y - P_C x \rangle \leq 0, \quad (2.5)$$

$$\|x - y\|^2 \geq \|x - P_C x\|^2 + \|y - P_C x\|^2 \quad (2.6)$$

for all $x \in H, y \in C$. It is easy to see that

$$u \in VI(C, \varphi) \Leftrightarrow u = P_C(u - \lambda \varphi u) \quad \text{for all } \lambda > 0.$$

A set-valued mapping $T : H \rightarrow 2^H$ is called monotone if for all $x, y \in H, f \in Tx$ and $g \in Ty$ imply $\langle x - y, f - g \rangle \geq 0$. A monotone mapping $T : H \rightarrow 2^H$ is maximal if the graph $G(T)$ of T is not properly contained in the graph of any other monotone mapping. It is known that a monotone mapping T is maximal if and only if, for $(x, f) \in H \times H, \langle x - y, f - g \rangle \geq 0$ for every $(y, g) \in G(T)$ implies $f \in Tx$. Let A be a monotone mapping of C into H and let $N_C v$ be the normal cone to C at $v \in C$, i.e.,

$$N_C v = \{\omega \in H : \langle v - u, \omega \rangle \geq 0, \forall u \in C\}$$

and define

$$Tv = \begin{cases} Av + N_C v, & v \in C, \\ \emptyset, & v \notin C. \end{cases}$$

Then T is maximal monotone and $0 \in Tv$ if and only if $v \in VI(C, A)$ (see, for example [27]).

The following lemmas will be useful for proving the convergence results of this paper.

Lemma 2.1. ([11]). For any $(x^*, y^*) \in C \times C$, (x^*, y^*) is a solution of SVIP(1.2) if and only if x^* is a fixed point of the mapping $Q : C \rightarrow C$ defined by

$$Q(x) = P_C[P_C(x - \mu_2 B_2 x) - \mu_1 B_1 P_C(x - \mu_2 B_2 x)], \quad \forall x \in C, \quad (2.7)$$

where $y^* = P_C(x^* - \mu_2 B_2 x^*)$, $\mu_i \in (0, 2\beta_i)$ and $B_i : C \rightarrow H$ is a nonlinear mapping for each $i = 1, 2$.

Lemma 2.2. ([21]). Let C be a convex closed subset of a Hilbert space H .

Let $F : C \times C \rightarrow R$ be a bi-function such that

(f1) $F(x, x) = 0$ for all $x \in C$;

(f2) F is monotone and upper hemicontinuous in the first variable;

(f3) F is lower semicontinuous and convex in the second variable.

Let $h : C \times C \rightarrow R$ be a bi-function such that

(h1) $h(x, x) = 0$ for all $x \in C$;

(h2) h is monotone and weakly upper semicontinuous in the first variable;

(h3) h is convex in the second variable

Moreover, let us suppose that

(H) for fixed $r > 0$ and $x \in C$, there exists a bounded $D \subset C$ and $a \in D$ such that for all $z \in C \setminus D$, $-F(a, z) + h(z, a) + \frac{1}{r} \langle a - z, z - x \rangle < 0$.

For $r > 0$ and $x \in H$, let $T_r : H \rightarrow 2^C$ be a mapping defined by

$$T_r x = \{z \in C, F(z, y) + h(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \forall y \in C\} \quad (2.8)$$

called resolvent of F and h .

Then

(1) $T_r x \neq \emptyset$;

(2) $T_r x$ is a singleton;

(3) T_r is firmly nonexpansive;

(4) $EP(F, h) = \text{Fix}(T_r)$ and it is closed and convex, where $EP(F, h) = \{z \in C, F(z, y) + h(z, y) \geq 0, \forall y \in C\}$.

Lemma 2.3. ([21]). Let us suppose that (f1)-(f3), (h1)-(h3) and (H) hold. Let $x, y \in H$, $r_1, r_2 > 0$. Then

$$\|T_{r_2} y - T_{r_1} x\| \leq \|y - x\| + \left| \frac{r_2 - r_1}{r_2} \right| \|T_{r_2} y - y\|.$$

Remark 2.4. In the sequel, given a sequence $\{z_n\}$, we will denote with $\omega_w\{z_n\}$ the set of cluster points of $\{z_n\}$ with respect to the weak topology, i.e.

$$\omega_w\{z_n\} = \{q \in H : \text{there exists } n_k \rightarrow \infty \text{ for which } z_{n_k} \rightharpoonup q\}.$$

Lemma 2.5. (see [20, Lemma 2.5]). Suppose that the hypotheses of Lemma 2.2 are satisfied. Let $\{r_n\}$ a sequence in $(0, +\infty)$ with $\liminf_{n \rightarrow \infty} r_n > 0$. Suppose that $\{x_n\}$ is a bounded sequence. Then the following statements are equivalent and true:

(a) if $\|x_n - T_{r_n} x_n\| \rightarrow 0$, as $n \rightarrow \infty$, the weak cluster points of $\{x_n\}$ satisfies the problem

$$F(x, y) + h(x, y) \geq 0 \quad \forall y \in C$$

i.e. $\omega_w\{x_n\} \subseteq EP(F, h)$.

(b) the demiclosedness principle holds in the sense that, if $x_n \rightharpoonup x$. and $\|x_n - T_{r_n} x_n\| \rightarrow 0$, as $n \rightarrow \infty$, then $(I - T_{r_k})x^* = 0$, for all $k \in \mathbb{N}$.

Lemma 2.6. (*Shimizu and Takahashi, [22]*). Let D be a nonempty, bounded, closed and convex subset of a real Hilbert space H and let $\mathfrak{S} := \{T(u) : 0 \leq u < \infty\}$ a nonexpansive semigroup on D , then for any $h \geq 0$,

$$\limsup_{t \rightarrow \infty, x \in D} \|T(h)(\frac{1}{t} \int_0^t T(u)x du) - (\frac{1}{t} \int_0^t T(u)x du)\| = 0.$$

Lemma 2.7. (*[23]*). Let H be a Hilbert space, C a closed convex subset of H , and $T : C \rightarrow C$ a nonexpansive mapping with $F(T) \neq \emptyset$; if $\{x_n\}$ is a sequence in C weakly converging to x and if $\{(I-T)x_n\}$ converges strongly to y , then $(I-T)x = y$.

Lemma 2.8. (*[Xu, 24]*). Let $\{a_n\}$ be a sequence of non-negative real numbers satisfying

$$a_{n+1} \leq (1 - \delta_n)a_n + b_n + \sigma_n, \quad n \in \mathbb{N},$$

where $\{\delta_n\}_{n=1}^\infty \subset (0, 1)$ and $\{b_n\}_{n=1}^\infty, \{\sigma_n\}_{n=1}^\infty$ is a sequence in $\mathbb{R}$ such that : (i) $\lim_{n \rightarrow \infty} \delta_n = 0$ and $\sum_{n=0}^\infty \delta_n = \infty$, (ii) $\limsup_{n \rightarrow \infty} \frac{b_n}{\delta_n} \leq 0$, (iii) $\sigma_n \geq 0, \sum_{n=0}^\infty \sigma_n < \infty$. Then $\lim_{n \rightarrow \infty} a_n = 0$.

Lemma 2.9. (*[25]*). Assume that T is a strongly positive linear bounded operator on a Hilbert space H with coefficient $\bar{\gamma} > 0$ and $0 < \rho \leq \|T\|^{-1}$. Then $\|I - \rho T\| \leq 1 - \rho \bar{\gamma}$.

Lemma 2.10. Let H be a real Hilbert space. Then the following inequalities hold:

$$(1) \|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y \rangle;$$

$$(2) \|x + y\|^2 \geq \|x\|^2 + 2\langle y, x \rangle,$$

for all $x, y \in H$.

3 Main result

Theorem 3.1. Let C be a nonempty, closed and convex subset of a real Hilbert space H . For each $i = 1, 2$, let $A, B_i : C \rightarrow H$ be α, β_i -inverse strongly monotone mappings, respectively. Let F, h be a bi-function from $C \times C$ to $\mathbb{R}$ satisfying the hypotheses of Lemma 2.2. , and $\mathfrak{S} := \{T(u) : 0 \leq u < \infty\}$ be a one-parameter nonexpansive semigroup on H such that $\Omega := F(\mathfrak{S}) \cap EP(F, h) \cap VI(C, A) \cap \Psi \neq \emptyset$. Let $\phi : H \rightarrow H$ be a nonexpansive mapping and T be a strongly bounded linear operator on H with coefficient $\bar{\gamma}$ such that $0 < \gamma < \theta \bar{\gamma}$ and $0 < \theta \leq \|T\|^{-1}$. Let $\{r_n\} \subset (0, \infty)$ be a real sequence such that $\liminf_{n \rightarrow \infty} r_n > 0$. Suppose $\{x_n\}, \{u_n\}, \{y_n\}$ and $\{z_n\}$ be sequences generated by

$$\begin{cases} x_1 = x \in H, \\ F(u_n, y) + h(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in C, \\ z_n = P_C(u_n - \lambda_n A u_n), \\ y_n = \delta_n z_n + (1 - \delta_n) P_C[P_C(z_n - \mu_2 B_2 z_n) - \mu_1 B_1 P_C(z_n - \mu_2 B_2 z_n)], \\ x_{n+1} = \beta_n x_n + (1 - \beta_n) (\frac{1}{t_n} \int_0^{t_n} T(u) [\alpha_n \gamma \phi(x_n) + (I - \alpha_n \theta T) y_n] du), \end{cases} \quad (3.1)$$

where $\mu_i \in (0, 2\beta_i)$, for each $i = 1, 2$, $\lambda_n \in (0, 2\alpha)$, and $\{\beta_n\}, \{\alpha_n\}, \{\delta_n\}$ are sequences in $[0, 1]$. Assume that the following conditions are satisfied:

$$(i) \lim_{n \rightarrow \infty} \alpha_n = 0, \sum_{n=1}^\infty \alpha_n = \infty, \sum_{n=1}^\infty |\alpha_{n+1} - \alpha_n| < \infty;$$

$$(ii) \lim_{n \rightarrow \infty} \beta_n = 0, \sum_{n=1}^\infty |\beta_{n+1} - \beta_n| < \infty;$$

$$(iii) \lim_{n \rightarrow \infty} \delta_n = 0, \sum_{n=1}^\infty |\delta_{n+1} - \delta_n| < \infty;$$

$$(iv) \sum_{n=1}^\infty |r_{n+1} - r_n| < \infty;$$

$$(v) \lim_{n \rightarrow \infty} \frac{|t_n - t_{n-1}|}{t_n} \frac{1}{\alpha_n(1 - \beta_n)} = 0;$$

$$(vi) 0 < \liminf_{n \rightarrow \infty} \lambda_n \leq \limsup_{n \rightarrow \infty} \lambda_n < 2\alpha, \sum_{n=1}^\infty |\lambda_{n+1} - \lambda_n| < \infty.$$

Then $\{x_n\}$ converges strongly to w , where $w := P_\Omega(\gamma \phi + (I - \theta T))w$.

Proof. First, we show that $I - \lambda_n A$ is nonexpansive. For any $x, y \in C$,

$$\begin{aligned} \|(I - \lambda_n A)x - (I - \lambda_n A)y\|^2 &= \|x - y - \lambda_n(Ax - Ay)\|^2 \\ &= \|x - y\|^2 - 2\lambda_n \langle Ax - Ay, x - y \rangle + \lambda_n^2 \|Ax - Ay\|^2 \\ &\leq \|x - y\|^2 - \lambda_n(2\alpha - \lambda_n) \|Ax - Ay\|^2 \leq \|x - y\|^2. \end{aligned} \quad (3.2)$$

Similarly we can show that the mappings $(I - \mu_i B_i)$ are nonexpansive for each $i = 1, 2$.

Next, we show that $\{x_n\}$ is bounded.

Observe that $\{u_n\}$ can be re-written as $u_n = T_{r_n}(x_n)$, $n \geq 1$. Let $x^* \in \Omega$, then

$$\|u_n - x^*\| = \|T_{r_n}(x_n) - x^*\| \leq \|x_n - x^*\|. \quad (3.3)$$

Since $x^* \in \Omega$, we have

$$x^* = P_C[P_C(x^* - \mu_2 B_2 x^*) - \mu_1 B_1 P_C(x^* - \mu_2 B_2 x^*)].$$

Putting

$$y^* = P_C(x^* - \mu_2 B_2 x^*),$$

we see that

$$x^* = P_C(y^* - \mu_1 B_1 y^*). \quad (3.4)$$

Since the mapping $A : C \rightarrow H$ is α -inverse strongly monotone, we have

$$\begin{aligned} \|z_n - x^*\|^2 &= \|P_C(u_n - \lambda_n A u_n) - P_C(x^* - \lambda_n A x^*)\|^2 \leq \|(u_n - \lambda_n A u_n) - (x^* - \lambda_n A x^*)\|^2 \\ &\leq \|(u_n - x^*) - \lambda_n(A u_n - A x^*)\|^2 \leq \|u_n - x^*\|^2 - \lambda_n(2\alpha - \lambda_n) \|A u_n - A x^*\|^2 \\ &\leq \|u_n - x^*\|^2 \leq \|x_n - x^*\|^2. \end{aligned} \quad (3.5)$$

Setting $t'_n = P_C[P_C(z_n - \mu_2 B_2 z_n) - \mu_1 B_1 P_C(z_n - \mu_2 B_2 z_n)]$ and $v_n = P_C(z_n - \mu_2 B_2 z_n)$. It follows that

$$\begin{aligned} \|v_n - y^*\|^2 &= \|P_C(z_n - \mu_2 B_2 z_n) - P_C(x^* - \mu_2 B_2 x^*)\|^2 \leq \|(z_n - \mu_2 B_2 z_n) - (x^* - \mu_2 B_2 x^*)\|^2 \\ &\leq \|z_n - x^*\|^2 - \mu_2(2\beta_2 - \mu_2) \|B_2 z_n - B_2 x^*\|^2 \leq \|z_n - x^*\|^2 \leq \|x_n - x^*\|^2. \end{aligned} \quad (3.6)$$

Further, we have

$$\begin{aligned} \|t'_n - x^*\|^2 &= \|P_C(v_n - \mu_1 B_1 v_n) - P_C(y^* - \mu_1 B_1 y^*)\|^2 \\ &\leq \|v_n - y^* - \mu_1(B_1 v_n - B_1 y^*)\|^2 \\ &\leq \|v_n - y^*\|^2 - \mu_1(2\beta_1 - \mu_1) \|B_1 v_n - B_1 y^*\|^2 \\ &\leq \|z_n - x^*\|^2 - \mu_2(2\beta_2 - \mu_2) \|B_2 z_n - B_2 x^*\|^2 - \mu_1(2\beta_1 - \mu_1) \|B_1 v_n - B_1 y^*\|^2 \\ &\leq \|z_n - x^*\|^2 \leq \|x_n - x^*\|^2. \end{aligned} \quad (3.7)$$

So

$$\begin{aligned} \|y_n - x^*\| &= \|\delta_n z_n + (1 - \delta_n) P_C[P_C(z_n - \mu_2 B_2 z_n) - \mu_1 B_1 P_C(z_n - \mu_2 B_2 z_n)] - x^*\| \\ &\leq \delta_n \|z_n - x^*\| + (1 - \delta_n) \|t'_n - x^*\| \\ &\leq \delta_n \|x_n - x^*\| + (1 - \delta_n) \|x_n - x^*\| \leq \|x_n - x^*\|. \end{aligned} \quad (3.9)$$

Now, put $w_n = \alpha_n \gamma \phi(x_n) + (I - \alpha_n \theta T)y_n$, $n \geq 1$. So,

$$\begin{aligned} \|w_n - x^*\| &= \|\alpha_n \gamma \phi(x_n) + (I - \alpha_n \theta T)y_n - x^*\| \\ &= \|\alpha_n(\gamma \phi(x_n) - \theta T x^*) + (I - \alpha_n \theta T)(y_n - x^*)\| \\ &\leq (1 - \alpha_n \theta \bar{\gamma}) \|y_n - x^*\| + \alpha_n (\|\gamma \phi(x_n) - \gamma \phi(x^*)\| + \|\gamma \phi(x^*) - \theta T x^*\|) \\ &\leq (1 - \alpha_n \theta \bar{\gamma}) \|x_n - x^*\| + \alpha_n \gamma \|x_n - x^*\| + \alpha_n \|\gamma \phi(x^*) - \theta T x^*\| \\ &= [1 - \alpha_n(\theta \bar{\gamma} - \gamma)] \|x_n - x^*\| + \alpha_n(\theta \bar{\gamma} - \gamma) \frac{\|\gamma \phi(x^*) - \theta T x^*\|}{\theta \bar{\gamma} - \gamma}. \end{aligned} \quad (3.10)$$

From (3.1), we obtain

$$\begin{aligned}
\|x_{n+1} - x^*\| &= \|\beta_n(x_n - x^*) + (1 - \beta_n)\left(\frac{1}{t_n} \int_0^{t_n} [T(u)w_n - T(u)x^*]du\right)\| \\
&\leq \beta_n\|x_n - x^*\| + (1 - \beta_n)\|w_n - x^*\| \\
&\leq \beta_n\|x_n - x^*\| + (1 - \beta_n)([1 - \alpha_n(\theta\bar{\gamma} - \gamma)]\|x_n - x^*\| + \alpha_n(\theta\bar{\gamma} - \gamma)\frac{\|\gamma\phi(x^*) - \theta Tx^*\|}{\theta\bar{\gamma} - \gamma}) \\
&= (1 - \alpha_n(\theta\bar{\gamma} - \gamma)(1 - \beta_n))\|x_n - x^*\| + \alpha_n(\theta\bar{\gamma} - \gamma)(1 - \beta_n)\frac{\|\gamma\phi(x^*) - \theta Tx^*\|}{\theta\bar{\gamma} - \gamma} \\
&\leq \max\{\|x_n - x^*\|, \frac{1}{\theta\bar{\gamma} - \gamma}\|\gamma\phi(x^*) - \theta Tx^*\|\} \\
&\vdots \\
&\leq \max\{\|x_1 - x^*\|, \frac{1}{\theta\bar{\gamma} - \gamma}\|\gamma\phi(x^*) - \theta Tx^*\|\}
\end{aligned}$$

So, $\{x_n\}$ is bounded. Hence, $\{u_n\}, \{z_n\}, \{y_n\}, \{\theta Ty_n\}, \{w_n\}$ and $\{\frac{1}{t_n} \int_0^{t_n} T(u)w_n\}$ are bounded.

From (3.10), we have

$$\begin{aligned}
\|w_n - x^*\| &\leq [1 - \alpha_n(\theta\bar{\gamma} - \gamma)]\|x_n - x^*\| + \alpha_n(\theta\bar{\gamma} - \gamma)\frac{\|\gamma\phi(x^*) - \theta Tx^*\|}{\theta\bar{\gamma} - \gamma} \\
&\leq \max\{\|x_1 - x^*\|, \frac{1}{\theta\bar{\gamma} - \gamma}\|\gamma\phi(x^*) - \theta Tx^*\|\} + \frac{1}{\theta\bar{\gamma} - \gamma}\|\gamma\phi(x^*) - \theta Tx^*\| \\
&\leq \|x_1 - x^*\| + \frac{2}{\theta\bar{\gamma} - \gamma}\|\gamma\phi(x^*) - \theta Tx^*\|.
\end{aligned}$$

Put $D = \{\omega \in H : \|\omega - x^*\| \leq \|x_1 - x^*\| + \frac{2}{\theta\bar{\gamma} - \gamma}\|\gamma\phi(x^*) - \theta Tx^*\|\}$. Then D is a nonempty, bounded, closed and convex subset of H . Since $T(u)$ is nonexpansive for any $u \in [0, \infty)$, D is $T(u)$ -invariant for each $u \in [0, \infty)$ and contains $\{w_n\}$. Without loss of generality, we may assume that $\mathfrak{S} := T(u) : 0 \leq u < \infty$ is a nonexpansive semigroup on D . By Lemma 2.6, we get

$$\lim_{n \rightarrow \infty} \left\| \left(\frac{1}{t_n} \int_0^{t_n} T(u)w_n du \right) - T(h) \left(\frac{1}{t_n} \int_0^{t_n} T(u)w_n du \right) \right\| = 0 \quad (3.11)$$

for every $h \in [0, \infty)$. Furthermore, observe that

$$\begin{aligned}
\|x_{n+1} - T(h)x_{n+1}\| &\leq \|x_{n+1} - \frac{1}{t_n} \int_0^{t_n} T(u)w_n du\| + \left\| \left(\frac{1}{t_n} \int_0^{t_n} T(u)w_n du \right) - T(h) \left(\frac{1}{t_n} \int_0^{t_n} T(u)w_n du \right) \right\| \\
&\quad + \|T(h) \left(\frac{1}{t_n} \int_0^{t_n} T(u)w_n du \right) - T(h)x_{n+1}\| \\
&\leq 2\|x_{n+1} - \frac{1}{t_n} \int_0^{t_n} T(u)w_n du\| + \left\| \left(\frac{1}{t_n} \int_0^{t_n} T(u)w_n du \right) - T(h) \left(\frac{1}{t_n} \int_0^{t_n} T(u)w_n du \right) \right\| \\
&= 2\beta_n\|x_n - \frac{1}{t_n} \int_0^{t_n} T(u)w_n du\| + \left\| \left(\frac{1}{t_n} \int_0^{t_n} T(u)w_n du \right) - T(h) \left(\frac{1}{t_n} \int_0^{t_n} T(u)w_n du \right) \right\|.
\end{aligned}$$

From $\lim_{n \rightarrow \infty} \beta_n = 0$ and (3.11), we get $\lim_{n \rightarrow \infty} \|x_{n+1} - T(h)x_{n+1}\| = 0$ and hence

$$\lim_{n \rightarrow \infty} \|x_n - T(h)x_n\| = 0. \quad (3.12)$$

We next show that $\lim_{n \rightarrow \infty} \|x_n - x_{n+1}\| = 0$.

From the nonexpansivity of the mapping $(I - \lambda_n A)$, we have

$$\begin{aligned}
\|z_n - z_{n-1}\| &= \|P_C(u_n - \lambda_n A u_n) - P_C(u_{n-1} - \lambda_{n-1} A u_{n-1})\| \\
&\leq \|(u_n - \lambda_n A u_n) - (u_{n-1} - \lambda_{n-1} A u_{n-1})\| \\
&= \|(u_n - u_{n-1}) - \lambda_n(A u_n - A u_{n-1}) + (\lambda_n - \lambda_{n-1})A u_{n-1}\| \\
&\leq \|(u_n - u_{n-1}) - \lambda_n(A u_n - A u_{n-1})\| + |\lambda_n - \lambda_{n-1}| \|A u_{n-1}\| \\
&\leq \|u_n - u_{n-1}\| + |\lambda_n - \lambda_{n-1}| \|A u_{n-1}\|.
\end{aligned} \tag{3.13}$$

We next estimate

$$\begin{aligned}
\|t'_n - t'_{n-1}\|^2 &= \|P_C(v_n - \mu_1 B_1 v_n) - P_C(v_{n-1} - \mu_1 B_1 v_{n-1})\|^2 \\
&\leq \|(v_n - \mu_1 B_1 v_n) - (v_{n-1} - \mu_1 B_1 v_{n-1})\|^2 \\
&\leq \|v_n - v_{n-1}\|^2 - \mu_1(2\beta_1 - \mu_1) \|B_1 v_n - B_1 v_{n-1}\|^2 \\
&\leq \|v_n - v_{n-1}\|^2 \\
&\leq \|z_n - z_{n-1}\|^2 - \mu_2(2\beta_2 - \mu_2) \|B_2 z_n - B_2 z_{n-1}\|^2 \\
&\leq \|z_n - z_{n-1}\|^2.
\end{aligned} \tag{3.14}$$

From (3.13) and (3.14), we have

$$\|t'_n - t'_{n-1}\| \leq \|u_n - u_{n-1}\| + |\lambda_n - \lambda_{n-1}| \|A u_{n-1}\|. \tag{3.15}$$

We observe that

$$\begin{aligned}
\|y_n - y_{n-1}\| &= \|\delta_n z_n + (1 - \delta_n)t'_n - \delta_{n-1} z_{n-1} - (1 - \delta_{n-1})t'_{n-1}\| \\
&= \|\delta_n(z_n - z_{n-1}) + (\delta_n - \delta_{n-1})z_{n-1} + (1 - \delta_n)(t'_n - t'_{n-1}) + [(1 - \delta_n) - (1 - \delta_{n-1})]t'_{n-1}\| \\
&\leq \delta_n \|z_n - z_{n-1}\| + |\delta_n - \delta_{n-1}| \|z_{n-1}\| + (1 - \delta_n) \|t'_n - t'_{n-1}\| + |\delta_n - \delta_{n-1}| \|t'_{n-1}\| \\
&\leq \delta_n \|z_n - z_{n-1}\| + |\delta_n - \delta_{n-1}| \|z_{n-1}\| + (1 - \delta_n) \|z_n - z_{n-1}\| + |\delta_n - \delta_{n-1}| \|t'_{n-1}\| \\
&= \|z_n - z_{n-1}\| + |\delta_n - \delta_{n-1}| (\|z_{n-1}\| + \|t'_{n-1}\|) \\
&\leq \|z_n - z_{n-1}\| + |\delta_n - \delta_{n-1}| M \\
&\leq \|u_n - u_{n-1}\| + |\lambda_n - \lambda_{n-1}| \|A u_{n-1}\| + |\delta_n - \delta_{n-1}| M,
\end{aligned}$$

where $M = \sup_{n \geq 1} \{\|z_n\| + \|t'_n\|\}$.

On the other hand, $u_n = T_{r_n}(x_n)$ and $u_{n-1} = T_{r_{n-1}}(x_{n-1})$, by Lemma 2.3, we have

$$\|u_n - u_{n-1}\| \leq \|x_n - x_{n-1}\| + L \left|1 - \frac{r_{n-1}}{r_n}\right|, \tag{3.16}$$

where $L = \sup_n \|u_n - x_n\|$. Hence,

$$\|t'_n - t'_{n-1}\| \leq \|x_n - x_{n-1}\| + L \left|1 - \frac{r_{n-1}}{r_n}\right| + |\lambda_n - \lambda_{n-1}| \|A u_{n-1}\|, \tag{3.17}$$

and

$$\|y_n - y_{n-1}\| \leq \|x_n - x_{n-1}\| + L \left|1 - \frac{r_{n-1}}{r_n}\right| + |\lambda_n - \lambda_{n-1}| \|A u_{n-1}\| + |\delta_n - \delta_{n-1}| M. \tag{3.18}$$

From (3.18), we have

$$\begin{aligned}
\|w_n - w_{n-1}\| &= \|\alpha_n \gamma \phi(x_n) + (I - \alpha_n \theta T)y_n - \alpha_{n-1} \gamma \phi(x_{n-1}) - (I - \alpha_{n-1} \theta T)y_{n-1}\| \\
&= \|\alpha_n \gamma (\phi(x_n) - \phi(x_{n-1})) + (\alpha_n - \alpha_{n-1}) \gamma \phi(x_{n-1}) + (I - \alpha_n \theta T)(y_n - y_{n-1}) \\
&\quad + [(I - \alpha_n \theta T) - (I - \alpha_{n-1} \theta T)]y_{n-1}\| \\
&\leq \alpha_n \gamma \|x_n - x_{n-1}\| + |\alpha_n - \alpha_{n-1}| \|\gamma \phi(x_{n-1})\| + (1 - \alpha_n \theta \bar{\gamma}) \|y_n - y_{n-1}\| \\
&\quad + |\alpha_n - \alpha_{n-1}| \|\theta T y_{n-1}\| \\
&\leq \alpha_n \gamma \|x_n - x_{n-1}\| + |\alpha_n - \alpha_{n-1}| (\|\gamma \phi(x_{n-1})\| + \|\theta T y_{n-1}\|) \\
&\quad + (1 - \alpha_n \theta \bar{\gamma}) [\|x_n - x_{n-1}\| + L|1 - \frac{r_{n-1}}{r_n}| + |\lambda_n - \lambda_{n-1}| \|A u_{n-1}\| + |\delta_n - \delta_{n-1}| M] \\
&\leq [1 - \alpha_n (\theta \bar{\gamma} - \gamma)] \|x_n - x_{n-1}\| + |\alpha_n - \alpha_{n-1}| (\|\gamma \phi(x_{n-1})\| + \|\theta T y_{n-1}\|) \\
&\quad + L|1 - \frac{r_{n-1}}{r_n}| + |\lambda_n - \lambda_{n-1}| \|A u_{n-1}\| + |\delta_n - \delta_{n-1}| M.
\end{aligned} \tag{3.19}$$

Let $\theta_n := \frac{1}{t_n} \int_0^{t_n} T(u) w_n du$, $n \geq 1$. Then, we have

$$\|\theta_n - \theta_{n-1}\| = \left\| \frac{1}{t_n} \int_0^{t_n} [T(u) w_n - T(u) w_{n-1}] du + \left(\frac{1}{t_n} - \frac{1}{t_{n-1}} \right) \int_0^{t_{n-1}} T(u) w_{n-1} du + \frac{1}{t_n} \int_{t_{n-1}}^{t_n} T(u) w_{n-1} du \right\|$$

if $x^* \in \Omega$, we can write

$$\begin{aligned}
\|\theta_n - \theta_{n-1}\| &= \left\| \frac{1}{t_n} \int_0^{t_n} [T(u) w_n - T(u) w_{n-1}] du + \left(\frac{1}{t_n} - \frac{1}{t_{n-1}} \right) \int_0^{t_{n-1}} [T(u) w_{n-1} - T(u) x^*] du \right. \\
&\quad \left. + \frac{1}{t_n} \int_{t_{n-1}}^{t_n} [T(u) w_{n-1} - T(u) x^*] du \right\|
\end{aligned}$$

Thus,

$$\|\theta_n - \theta_{n-1}\| \leq \|w_n - w_{n-1}\| + \left(\frac{2|t_n - t_{n-1}|}{t_n} \right) \|w_{n-1} - x^*\|. \tag{3.20}$$

Substituting (3.19) into (3.20), we obtain

$$\begin{aligned}
\|\theta_n - \theta_{n-1}\| &\leq [1 - \alpha_n (\theta \bar{\gamma} - \gamma)] \|x_n - x_{n-1}\| + |\alpha_n - \alpha_{n-1}| (\|\gamma \phi(x_{n-1})\| + \|\theta T y_{n-1}\|) \\
&\quad + L|1 - \frac{r_{n-1}}{r_n}| + |\lambda_n - \lambda_{n-1}| \|A u_{n-1}\| + |\delta_n - \delta_{n-1}| M + \left(\frac{2|t_n - t_{n-1}|}{t_n} \right) \|w_{n-1} - x^*\|,
\end{aligned} \tag{3.21}$$

From (3.1), we have $x_{n+1} = \beta_n x_n + (1 - \beta_n) \theta_n$ and this implies that

$$\begin{aligned}
\|x_{n+1} - x_n\| &= \|\beta_n x_n + (1 - \beta_n) \theta_n - \beta_{n-1} x_{n-1} - (1 - \beta_{n-1}) \theta_{n-1}\| \\
&\leq \beta_n \|x_n - x_{n-1}\| + (1 - \beta_n) \|\theta_n - \theta_{n-1}\| + |\beta_n - \beta_{n-1}| (\|x_{n-1}\| + \|\theta_{n-1}\|).
\end{aligned} \tag{3.22}$$

Using (3.21) in (3.22), we obtain

$$\begin{aligned}
\|x_{n+1} - x_n\| &\leq \beta_n \|x_n - x_{n-1}\| + (1 - \beta_n) [1 - \alpha_n (\theta \bar{\gamma} - \gamma)] \|x_n - x_{n-1}\| + |\alpha_n - \alpha_{n-1}| (\|\gamma \phi(x_{n-1})\| + \|\theta T y_{n-1}\|) \\
&\quad + L|1 - \frac{r_{n-1}}{r_n}| + |\lambda_n - \lambda_{n-1}| \|A u_{n-1}\| + |\delta_n - \delta_{n-1}| M + \left(\frac{2|t_n - t_{n-1}|}{t_n} \right) \|w_{n-1} - x^*\| \\
&\quad + |\beta_n - \beta_{n-1}| (\|x_{n-1}\| + \|\theta_{n-1}\|) \\
&\leq [1 - (\theta \bar{\gamma} - \gamma) \alpha_n (1 - \beta_n)] \|x_n - x_{n-1}\| + |\alpha_n - \alpha_{n-1}| (\|\gamma \phi(x_{n-1})\| + \|\theta T y_{n-1}\|) \\
&\quad + L|1 - \frac{r_{n-1}}{r_n}| + |\lambda_n - \lambda_{n-1}| \|A u_{n-1}\| + |\delta_n - \delta_{n-1}| M + \left(\frac{2|t_n - t_{n-1}|}{t_n} \right) \|w_{n-1} - x^*\| \\
&\quad + |\beta_n - \beta_{n-1}| (\|x_{n-1}\| + \|\theta_{n-1}\|).
\end{aligned}$$

If we call $D' := \max\{\sup_{n \geq 1}(\|\gamma\phi(x_{n-1})\| + \|\theta T y_{n-1}\|), \sup_{n \geq 1}(\|x_{n-1}\| + \|\theta_{n-1}\|), \sup_{n \geq 1}\|w_{n-1} - x^*\|, L, M, \sup_{n \geq 1}\|Au_{n-1}\|\}$ and $b > 0$ a minorant for $\{r_n\}$, we have

$$\begin{aligned} \|x_{n+1} - x_n\| &\leq [1 - (\theta\bar{\gamma} - \gamma)\alpha_n(1 - \beta_n)]\|x_n - x_{n-1}\| + D'[\alpha_n - \alpha_{n-1}] + \frac{|r_n - r_{n-1}|}{b} \\ &\quad + |\lambda_n - \lambda_{n-1}| + |\delta_n - \delta_{n-1}| + \left(\frac{2|t_n - t_{n-1}|}{t_n}\right) + |\beta_n - \beta_{n-1}|. \end{aligned}$$

From Lemma 2.8, taking

$$\varepsilon_n = (\theta\bar{\gamma} - \gamma)\alpha_n(1 - \beta_n), \quad b_n = \frac{2D'|t_n - t_{n-1}|}{t_n},$$

$$\sigma_n = D'[\alpha_n - \alpha_{n-1}] + \frac{|r_n - r_{n-1}|}{b} + |\lambda_n - \lambda_{n-1}| + |\delta_n - \delta_{n-1}| + |\beta_n - \beta_{n-1}|$$

by using conditions (i), (ii), (iii), (iv), (v) and (vi) it follows that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \quad (3.23)$$

We show that $\lim_{n \rightarrow \infty} \|u_n - y_n\| = 0$, $\lim_{n \rightarrow \infty} \|w_n - y_n\| = 0$ and $\lim_{n \rightarrow \infty} \|x_n - u_n\| = \|x_n - T_{r_n} x_n\| = 0$. Now, by the convexity of $\|\cdot\|^2$, we have

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &= \|\beta_n(x_n - x^*) + (1 - \beta_n)\left(\frac{1}{t_n} \int_0^{t_n} [T(u)w_n - T(u)x^*]du\right)\|^2 \\ &\leq \beta_n\|x_n - x^*\|^2 + (1 - \beta_n)\|w_n - x^*\|^2 \\ &= \beta_n\|x_n - x^*\|^2 + (1 - \beta_n)\|\alpha_n\gamma\phi(x_n) + (I - \alpha_n\theta T)y_n - x^*\|^2 \\ &= \beta_n\|x_n - x^*\|^2 + (1 - \beta_n)\|\alpha_n(\gamma\phi(x_n) - \theta T x^*) + (I - \alpha_n\theta T)(y_n - x^*)\|^2 \\ &\leq \beta_n\|x_n - x^*\|^2 + (1 - \beta_n)[(1 - \alpha_n\theta\bar{\gamma})^2\|y_n - x^*\|^2 + 2\alpha_n\langle\gamma\phi(x_n) - \theta T x^*, w_n - x^*\rangle] \\ &\leq \beta_n\|x_n - x^*\|^2 + (1 - \beta_n)[\|y_n - x^*\|^2 + 2\alpha_n\|\gamma\phi(x_n) - \theta T x^*\|\|w_n - x^*\|] \\ &\leq \beta_n\|x_n - x^*\|^2 + (1 - \beta_n)\|y_n - x^*\|^2 + 2\alpha_n M_1 \end{aligned} \quad (3.24)$$

where M_1 is an appropriate constant such that $M_1 := \sup_{n \geq 1}\{\|\gamma\phi(x_n) - \theta T x^*\|\|w_n - x^*\|\}$. From (3.7), we obtain

$$\begin{aligned} \|y_n - x^*\|^2 &\leq \delta_n\|z_n - x^*\|^2 + (1 - \delta_n)\|t'_n - x^*\|^2 \\ &\leq \delta_n\|x_n - x^*\|^2 + (1 - \delta_n)\|t'_n - x^*\|^2 \\ &\leq \delta_n\|x_n - x^*\|^2 + (1 - \delta_n)[\|z_n - x^*\|^2 - \mu_2(2\beta_2 - \mu_2)\|B_2 z_n - B_2 x^*\|^2 \\ &\quad - \mu_1(2\beta_1 - \mu_1)\|B_1 v_n - B_1 y^*\|^2] \\ &\leq \delta_n\|x_n - x^*\|^2 + (1 - \delta_n)[\|u_n - x^*\|^2 - \lambda_n(2\alpha - \lambda_n)\|Au_n - Ax^*\|^2 \\ &\quad - \mu_2(2\beta_2 - \mu_2)\|B_2 z_n - B_2 x^*\|^2 - \mu_1(2\beta_1 - \mu_1)\|B_1 v_n - B_1 y^*\|^2] \\ &\leq \delta_n\|x_n - x^*\|^2 + (1 - \delta_n)[\|x_n - x^*\|^2 - \lambda_n(2\alpha - \lambda_n)\|Au_n - Ax^*\|^2 \\ &\quad - \mu_2(2\beta_2 - \mu_2)\|B_2 z_n - B_2 x^*\|^2 - \mu_1(2\beta_1 - \mu_1)\|B_1 v_n - B_1 y^*\|^2] \\ &\leq \|x_n - x^*\|^2 - (1 - \delta_n)[\lambda_n(2\alpha - \lambda_n)\|Au_n - Ax^*\|^2 \\ &\quad + \mu_2(2\beta_2 - \mu_2)\|B_2 z_n - B_2 x^*\|^2 + \mu_1(2\beta_1 - \mu_1)\|B_1 v_n - B_1 y^*\|^2]. \end{aligned} \quad (3.25)$$

Substituting (3.25) into (3.24), we obtain

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &\leq \beta_n\|x_n - x^*\|^2 + (1 - \beta_n)\{\|x_n - x^*\|^2 - (1 - \delta_n)[\lambda_n(2\alpha - \lambda_n)\|Au_n - Ax^*\|^2 \\ &\quad + \mu_2(2\beta_2 - \mu_2)\|B_2 z_n - B_2 x^*\|^2 + \mu_1(2\beta_1 - \mu_1)\|B_1 v_n - B_1 y^*\|^2]\} + 2\alpha_n M_1 \\ &\leq \|x_n - x^*\|^2 - (1 - \beta_n)(1 - \delta_n)[\lambda_n(2\alpha - \lambda_n)\|Au_n - Ax^*\|^2 \\ &\quad + \mu_2(2\beta_2 - \mu_2)\|B_2 z_n - B_2 x^*\|^2 + \mu_1(2\beta_1 - \mu_1)\|B_1 v_n - B_1 y^*\|^2] + 2\alpha_n M_1. \end{aligned}$$

It follows that

$$(1 - \beta_n)(1 - \delta_n)[\lambda_n(2\alpha - \lambda_n)\|Au_n - Ax^*\|^2 + \mu_2(2\beta_2 - \mu_2)\|B_2z_n - B_2x^*\|^2 + \mu_1(2\beta_1 - \mu_1)\|B_1v_n - B_1y^*\|^2] \\ \leq \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2 + \alpha_n M_1 \leq \|x_n - x_{n+1}\|(\|x_n - x_{n+1}\| + \|x_{n+1} - x^*\|) + 2\alpha_n M_1. \quad (3.26)$$

Since $0 < \liminf_{n \rightarrow \infty} \lambda_n \leq \limsup_{n \rightarrow \infty} \lambda_n < 2\alpha$, $\|x_n - x_{n+1}\| \rightarrow 0$, $\alpha_n \rightarrow 0$, $\beta_n \rightarrow 0$ and $\delta_n \rightarrow 0$, we obtain $\lim_{n \rightarrow \infty} \|Au_n - Ax^*\| = 0$, $\lim_{n \rightarrow \infty} \|B_1v_n - B_1y^*\| = 0$, $\lim_{n \rightarrow \infty} \|B_2z_n - B_2x^*\| = 0$. By (2.1) and (2.4), we also have

$$\begin{aligned} \|z_n - x^*\|^2 &= \|P_C(u_n - \lambda_n Au_n) - P_C(x^* - \lambda_n Ax^*)\|^2 \\ &\leq \langle u_n - \lambda_n Au_n - (x^* - \lambda_n Ax^*), z_n - x^* \rangle \\ &\leq \frac{1}{2}[\|u_n - \lambda_n Au_n - (x^* - \lambda_n Ax^*)\|^2 + \|z_n - x^*\|^2 - \|u_n \\ &\quad - x^* - \lambda_n(Au_n - Ax^*) - (z_n - x^*)\|^2] \\ &\leq \frac{1}{2}[\|u_n - x^*\|^2 + \|z_n - x^*\|^2 - \|u_n - z_n - \lambda_n(Au_n - Ax^*)\|^2] \\ &\leq \frac{1}{2}[\|u_n - x^*\|^2 + \|z_n - x^*\|^2 - \|u_n - z_n\|^2 + 2\lambda_n \langle u_n - z_n, Au_n - Ax^* \rangle - \lambda_n^2 \|Au_n - Ax^*\|^2] \\ &\leq \frac{1}{2}[\|x_n - x^*\|^2 + \|z_n - x^*\|^2 - \|u_n - z_n\|^2 + 2\lambda_n \|u_n - z_n\| \|Au_n - Ax^*\|]. \end{aligned}$$

Hence

$$\|z_n - x^*\|^2 \leq \|x_n - x^*\|^2 - \|u_n - z_n\|^2 + 2\lambda_n \|u_n - z_n\| \|Au_n - Ax^*\|. \quad (3.27)$$

Next, we estimate

$$\begin{aligned} \|v_n - y^*\|^2 &= \|P_C(z_n - \mu_2 B_2 z_n) - P_C(x^* - \mu_2 B_2 x^*)\|^2 \\ &\leq \langle (z_n - \mu_2 B_2 z_n) - (x^* - \mu_2 B_2 x^*), v_n - y^* \rangle \\ &\leq \frac{1}{2}[\|z_n - x^* - \mu_2(B_2 z_n - B_2 x^*)\|^2 + \|v_n - y^*\|^2 \\ &\quad - \|z_n - x^* - \mu_2(B_2 z_n - B_2 x^*) - (v_n - y^*)\|^2] \\ &\leq \frac{1}{2}[\|z_n - x^*\|^2 + \|v_n - y^*\|^2 - \|z_n - v_n - \mu_2(B_2 z_n - B_2 x^*) - (x^* - y^*)\|^2] \\ &\leq \frac{1}{2}[\|z_n - x^*\|^2 + \|v_n - y^*\|^2 - \|z_n - v_n - (x^* - y^*)\|^2 \\ &\quad + 2\mu_2 \langle z_n - v_n - (x^* - y^*), B_2 z_n - B_2 x^* \rangle - \mu_2^2 \|B_2 z_n - B_2 x^*\|^2] \\ &\leq \frac{1}{2}[\|z_n - x^*\|^2 + \|v_n - y^*\|^2 - \|z_n - v_n - (x^* - y^*)\|^2 \\ &\quad + 2\mu_2 \|z_n - v_n - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\|]. \end{aligned}$$

Hence

$$\begin{aligned} \|v_n - y^*\|^2 &\leq \|z_n - x^*\|^2 - \|z_n - v_n - (x^* - y^*)\|^2 + 2\mu_2 \|z_n - v_n - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| \\ &\leq \|x_n - x^*\|^2 - \|u_n - z_n\|^2 + 2\lambda_n \|u_n - z_n\| \|Au_n - Ax^*\| \\ &\quad - \|z_n - v_n - (x^* - y^*)\|^2 + 2\mu_2 \|z_n - v_n - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\|. \end{aligned} \quad (3.28)$$

Similarly, we also estimate

$$\begin{aligned}
\|t'_n - x^*\|^2 &= \|P_C(v_n - \mu_1 B_1 v_n) - P_C(y^* - \mu_1 B_1 y^*)\|^2 \\
&\leq \langle (v_n - \mu_1 B_1 v_n) - (y^* - \mu_1 B_1 y^*), t'_n - x^* \rangle \\
&\leq \frac{1}{2} [\|v_n - y^* - \mu_1(B_1 z_n - B_1 y^*)\|^2 + \|t'_n - x^*\|^2 \\
&\quad - \|v_n - y^* - \mu_1(B_1 v_n - B_1 y^*) - (t'_n - x^*)\|^2] \\
&\leq \frac{1}{2} [\|v_n - y^*\|^2 + \|t'_n - x^*\|^2 - \|v_n - t'_n - \mu_1(B_1 v_n - B_1 y^*) + (x^* - y^*)\|^2] \\
&\leq \frac{1}{2} [\|v_n - y^*\|^2 + \|t'_n - x^*\|^2 - \|v_n - t'_n + (x^* - y^*)\|^2] \\
&\quad + 2\mu_1 \langle v_n - t'_n + (x^* - y^*), B_1 v_n - B_1 y^* \rangle - \mu_1^2 \|B_1 v_n - B_1 y^*\|^2 \\
&\leq \frac{1}{2} [\|v_n - y^*\|^2 + \|t'_n - x^*\|^2 - \|v_n - t'_n + (x^* - y^*)\|^2] \\
&\quad + 2\mu_1 \|v_n - t'_n + (x^* - y^*)\| \|B_1 v_n - B_1 y^*\|.
\end{aligned}$$

Hence

$$\begin{aligned}
\|t'_n - x^*\|^2 &\leq \|v_n - y^*\|^2 - \|v_n - t'_n + (x^* - y^*)\|^2 + 2\mu_1 \|v_n - t'_n + (x^* - y^*)\| \|B_1 v_n - B_1 y^*\| \\
&\leq \|x_n - x^*\|^2 - \|u_n - z_n\|^2 + 2\lambda_n \|u_n - z_n\| \|Au_n - Ax^*\| \\
&\quad - \|z_n - v_n - (x^* - y^*)\|^2 + 2\mu_2 \|z_n - v_n - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| \\
&\quad - \|v_n - t'_n + (x^* - y^*)\|^2 + 2\mu_1 \|v_n - t'_n + (x^* - y^*)\| \|B_1 v_n - B_1 y^*\|. \tag{3.29}
\end{aligned}$$

Substituting (3.29) into (3.24), we have

$$\begin{aligned}
\|x_{n+1} - x^*\|^2 &\leq \beta_n \|x_n - x^*\|^2 + (1 - \beta_n) \|y_n - x^*\|^2 + 2\alpha_n M_1 \\
&\leq \beta_n \|x_n - x^*\|^2 + (1 - \beta_n) [\delta_n \|z_n - x^*\|^2 + (1 - \delta_n) \|t'_n - x^*\|^2] + 2\alpha_n M_1 \\
&\leq \beta_n \|x_n - x^*\|^2 + (1 - \beta_n) [\delta_n \|z_n - x^*\|^2 + (1 - \delta_n) [\|x_n - x^*\|^2 - \|u_n - z_n\|^2 \\
&\quad + 2\lambda_n \|u_n - z_n\| \|Au_n - Ax^*\| - \|z_n - v_n - (x^* - y^*)\|^2 \\
&\quad + 2\mu_2 \|z_n - v_n - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| - \|v_n - t'_n + (x^* - y^*)\|^2 \\
&\quad + 2\mu_1 \|v_n - t'_n + (x^* - y^*)\| \|B_1 v_n - B_1 y^*\|] + 2\alpha_n M_1 \\
&\leq \beta_n \|x_n - x^*\|^2 + (1 - \beta_n) [\delta_n \|x_n - x^*\|^2 + (1 - \delta_n) [\|x_n - x^*\|^2 - \|u_n - z_n\|^2 \\
&\quad + 2\lambda_n \|u_n - z_n\| \|Au_n - Ax^*\| - \|z_n - v_n - (x^* - y^*)\|^2 \\
&\quad + 2\mu_2 \|z_n - v_n - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| - \|v_n - t'_n + (x^* - y^*)\|^2 \\
&\quad + 2\mu_1 \|v_n - t'_n + (x^* - y^*)\| \|B_1 v_n - B_1 y^*\|] + 2\alpha_n M_1 \\
&\leq \|x_n - x^*\|^2 + (1 - \beta_n)(1 - \delta_n) [2\lambda_n \|u_n - z_n\| \|Au_n - Ax^*\| \\
&\quad + 2\mu_2 \|z_n - v_n - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| + 2\mu_1 \|v_n - t'_n + (x^* - y^*)\| \|B_1 v_n - B_1 y^*\| \\
&\quad - (1 - \beta_n)(1 - \delta_n) [\|u_n - z_n\|^2 + \|z_n - v_n - (x^* - y^*)\|^2 + \|v_n - t'_n + (x^* - y^*)\|^2] + 2\alpha_n M_1.
\end{aligned}$$

Which yields

$$\begin{aligned}
&(1 - \beta_n)(1 - \delta_n) [\|u_n - z_n\|^2 + \|z_n - v_n - (x^* - y^*)\|^2 + \|v_n - t'_n + (x^* - y^*)\|^2] \\
&\leq \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2 + (1 - \beta_n)(1 - \delta_n) [2\lambda_n \|u_n - z_n\| \|Au_n - Ax^*\| \\
&\quad + 2\mu_2 \|z_n - v_n - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| + 2\mu_1 \|v_n - t'_n + (x^* - y^*)\| \|B_1 v_n - B_1 y^*\|] + 2\alpha_n M_1 \\
&= \|x_n - x_{n+1}\| (\|x_n - x^*\| + \|x_{n+1} - x^*\|) + (1 - \beta_n)(1 - \delta_n) [2\lambda_n \|u_n - z_n\| \|Au_n - Ax^*\| \\
&\quad + 2\mu_2 \|z_n - v_n - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| + 2\mu_1 \|v_n - t'_n + (x^* - y^*)\| \|B_1 v_n - B_1 y^*\|] + 2\alpha_n M_1. \tag{3.30}
\end{aligned}$$

Since $0 < \lambda_n < 2\alpha$, $\|x_n - x_{n+1}\| \rightarrow 0$, $\alpha_n \rightarrow 0$, $\beta_n \rightarrow 0$, $\delta_n \rightarrow 0$ and $\lim_{n \rightarrow \infty} \|Au_n - Ax^*\| = 0$, $\lim_{n \rightarrow \infty} \|B_1v_n - B_1y^*\| = 0$, $\lim_{n \rightarrow \infty} \|B_2z_n - B_2x^*\| = 0$. it follows from the boundedness of $\{x_n\}$, $\{u_n\}$, $\{z_n\}$, $\{t'_n\}$ and $\{v_n\}$ that

$$\lim_{n \rightarrow \infty} \|u_n - z_n\| = 0, \quad \lim_{n \rightarrow \infty} \|z_n - v_n - (x^* - y^*)\| = 0, \quad \lim_{n \rightarrow \infty} \|v_n - t'_n + (x^* - y^*)\| = 0. \quad (3.31)$$

Consequently, it follows that

$$\|z_n - t'_n\| \leq \|z_n - v_n - (x^* - y^*)\| + \|v_n - t'_n + (x^* - y^*)\| \rightarrow 0 \text{ (as } n \rightarrow \infty). \quad (3.32)$$

Also $\lim_{n \rightarrow \infty} \|u_n - t'_n\| = 0$ and $\lim_{n \rightarrow \infty} \|y_n - t'_n\| = \delta_n \|z_n - t'_n\| = 0$ as $n \rightarrow \infty$

$$\|u_n - y_n\| \leq \|u_n - t'_n\| + \|t'_n - y_n\| \rightarrow 0 \text{ (as } n \rightarrow \infty) \quad (3.33)$$

and

$$\|w_n - y_n\| = \alpha_n \|\gamma\phi(x_n) - \theta Ty_n\| \rightarrow 0. \quad (3.34)$$

Next, we show that $\|x_n - u_n\| = \|x_n - T_{r_n}x_n\| \rightarrow 0$. By the firm nonexpansivity of T_{r_n} , a standard calculation (see [26]) shows that if $p \in EP(F, h)$

$$\|u_n - p\|^2 \leq \|x_n - p\|^2 - \|x_n - u_n\|^2.$$

So, let $x^* \in \Omega$; then by (3.24)

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &\leq \beta_n \|x_n - x^*\|^2 + (1 - \beta_n) \|y_n - x^*\|^2 + 2\alpha_n M_1 \\ &\leq \beta_n \|x_n - x^*\|^2 + (1 - \beta_n) \|u_n - x^*\|^2 + 2\alpha_n M_1 \\ &\leq \beta_n \|x_n - x^*\|^2 + (1 - \beta_n) [\|x_n - x^*\|^2 - \|x_n - u_n\|^2] + 2\alpha_n M_1 \\ &= \|x_n - x^*\|^2 - (1 - \beta_n) \|x_n - u_n\|^2 + 2\alpha_n M_1, \end{aligned}$$

which implies that

$$\begin{aligned} (1 - \beta_n) \|x_n - u_n\|^2 &\leq \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2 + 2\alpha_n M_1 \\ &\leq \|x_{n+1} - x_n\| (\|x_n - x^*\| + \|x_{n+1} - x^*\|) + 2\alpha_n M_1. \end{aligned}$$

Since $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\lim_{n \rightarrow \infty} \beta_n = 0$ and (3.23), one gets

$$\lim_{n \rightarrow \infty} \|x_n - u_n\| = 0. \quad (3.35)$$

So

$$\|y_n - x_n\| \leq \|y_n - u_n\| + \|u_n - x_n\| \rightarrow 0. \quad (3.36)$$

Furthermore, from (3.33), (3.34) and (3.35), we have for every $h \in [0, \infty)$ that

$$\begin{aligned} \|T(h)w_n - T(h)x_n\| &\leq \|w_n - x_n\| \\ &\leq \|x_n - u_n\| + \|u_n - y_n\| + \|w_n - y_n\| \rightarrow 0. \end{aligned} \quad (3.37)$$

So, we obtain from (3.12) and (3.37) that

$$\|T(h)w_n - x_n\| \leq \|T(h)w_n - T(h)x_n\| + \|T(h)x_n - x_n\| \rightarrow 0.$$

Hence, we have for every $h \in [0, \infty)$ that

$$\|T(h)w_n - w_n\| \leq \|T(h)w_n - x_n\| + \|x_n - u_n\| + \|u_n - y_n\| + \|y_n - w_n\| \rightarrow 0. \quad (3.38)$$

Next, putting $w = P_\Omega(\gamma\phi + (I - \theta T))w$, we show that $\limsup_{n \rightarrow \infty} \langle (\gamma\phi - \theta T)w, w_n - w \rangle \leq 0$. Observing that $P_\Omega(\gamma\phi + (I - \theta T))$ is a contraction. Indeed, for any $x, y \in H$, by Lemma 2.7 we have

$$\begin{aligned} \|P_\Omega(\gamma\phi + (I - \theta T))x - P_\Omega(\gamma\phi + (I - \theta T))y\| &\leq \|(\gamma\phi + (I - \theta T))x - (\gamma\phi + (I - \theta T))y\| \\ &\leq \|\gamma\phi(x) - \gamma\phi(y)\| + \|(I - \theta T)x - (I - \theta T)y\| \\ &\leq \gamma\|x - y\| + (1 - \theta\bar{\gamma})\|x - y\| \\ &\leq (1 - (\theta\bar{\gamma} - \gamma))\|x - y\|. \end{aligned}$$

Banachs Contraction Mapping Principle guarantees that $P_\Omega(\gamma\phi + (I - \theta T))w$ has a unique fixed point. Say $w \in H$. That is, $w = P_\Omega(\gamma\phi + (I - \theta T))w$. To do this, we choose a subsequence $\{w_{n_j}\}$ of $\{w_n\}$ such that

$$\limsup_{n \rightarrow \infty} \langle (\gamma\phi - \theta T)w, w_n - w \rangle = \lim_{j \rightarrow \infty} \langle (\gamma\phi - \theta T)w, w_{n_j} - w \rangle. \quad (3.39)$$

Since $\{w_{n_j}\}$ is bounded, without loss of generality, we can assume that $w_{n_j} \rightharpoonup q$. From (3.34), we obtain that $y_{n_j} \rightharpoonup q$. It follows from (3.33) and (3.35) that $u_{n_j} \rightharpoonup q$ and $x_{n_j} \rightharpoonup q$. Since $\{u_{n_j}\} \subset C$ and C is closed and convex, we obtain $q \in C$. Next we prove that $q \in \Omega = F(\mathfrak{S}) \cap EP(F, h) \cap VI(C, A) \cap \Psi$. (a) First, we prove that $q \in \Psi$. Take any $x, y \in C$. Using (2.7), we estimate

$$\begin{aligned} \|Q(x) - Q(y)\|^2 &= \|P_C[P_C(x - \mu_2 B_2 x) - \mu_1 B_1 P_C(x - \mu_2 B_2 x)] \\ &\quad - P_C[P_C(y - \mu_2 B_2 y) - \mu_1 B_1 P_C(y - \mu_2 B_2 y)]\|^2 \\ &\leq \|[P_C(x - \mu_2 B_2 x) - P_C(y - \mu_2 B_2 y)] - \mu_1[B_1 P_C(x - \mu_2 B_2 x) - B_1 P_C(y - \mu_2 B_2 y)]\|^2 \\ &\leq \|P_C(x - \mu_2 B_2 x) - P_C(y - \mu_2 B_2 y)\|^2 \\ &\quad - \mu_1(2\beta_1 - \mu_1)\|B_1 P_C(x - \mu_2 B_2 x) - B_1 P_C(y - \mu_2 B_2 y)\|^2 \\ &\leq \|(x - \mu_2 B_2 x) - (y - \mu_2 B_2 y)\|^2 \leq \|x - y\|^2 - \mu_2(2\beta_2 - \mu_2)\|B_2 x - B_2 y\|^2 \leq \|x - y\|^2. \end{aligned}$$

This implies that $Q : C \rightarrow C$ is nonexpansive.

Now, we have

$$\begin{aligned} \|y_n - Q(y_n)\| &\leq \delta_n \|z_n - Q(y_n)\| + (1 - \delta_n) \|P_C[P_C(z_n - \mu_2 B_2 z_n) - \mu_1 B_1 P_C(z_n - \mu_2 B_2 z_n)] - Q(y_n)\| \\ &\leq \delta_n \|z_n - Q(y_n)\| + (1 - \delta_n) \|Q(z_n) - Q(y_n)\| \\ &\leq \delta_n \|z_n - Q(y_n)\| + (1 - \delta_n) \|z_n - y_n\|. \end{aligned} \quad (3.40)$$

Since $\delta_n \rightarrow 0$ and $\|z_n - y_n\| \rightarrow 0$ as $n \rightarrow \infty$, (3.40) implies $\liminf_{n \rightarrow \infty} \|y_n - Q(y_n)\| = 0$ and hence by Lemma 2.7, it follows that $q \in Q(q)$. Further, it follows from Lemma 2.1 that $q \in \Psi$.

(b) Next, we prove that $q \in VI(C, A)$. Indeed, let

$$Tx = \begin{cases} Ax + N_C x, & x \in C, \\ \emptyset, & x \notin C. \end{cases}$$

For any given $(x, u) \in G(T)$, $u - Ax \in N_C x$. Since $z_n \in C$, by the definition of N_C , we have

$$\langle x - z_n, u - Ax \rangle \geq 0. \quad (3.41)$$

On the other hand, since $z_n = P_C(u_n - \lambda_n A u_n)$, we have $\langle x - z_n, z_n - (u_n - \lambda_n A u_n) \rangle \geq 0$, and so

$$\langle x - z_n, \frac{z_n - u_n}{\lambda_n} + A u_n \rangle \geq 0.$$

By (3.41) and the monotonicity of A , we have

$$\begin{aligned}
\langle x - z_{n_j}, u \rangle &\geq \langle x - z_{n_j}, Ax \rangle \\
&\geq \langle x - z_{n_j}, Ax \rangle - \langle x - z_{n_j}, \frac{z_{n_j} - u_{n_j}}{\lambda_{n_j}} + Au_{n_j} \rangle \\
&= \langle x - z_{n_j}, Ax - Az_{n_j} \rangle + \langle x - z_{n_j}, Az_{n_j} - Au_{n_j} \rangle - \langle x - z_{n_j}, \frac{z_{n_j} - u_{n_j}}{\lambda_{n_j}} \rangle \\
&\geq \langle x - z_{n_j}, Az_{n_j} - Au_{n_j} \rangle - \langle x - z_{n_j}, \frac{z_{n_j} - u_{n_j}}{\lambda_{n_j}} \rangle.
\end{aligned} \tag{3.42}$$

Since $z_{n_j} \rightarrow q$ and $\|z_{n_j} - u_{n_j}\| \rightarrow 0$, we have

$$\lim_{j \rightarrow \infty} \langle x - z_{n_j}, u \rangle = \langle x - q, u \rangle \geq 0.$$

Again since T is maximal monotone, we have $q \in T^{-1}(0)$ and hence $q \in VI(C, A)$.

(c) Next, we prove that $q \in EP(F, h)$.

By $\lim_{n \rightarrow \infty} \|x_n - u_n\| = \|x_n - T_{r_n}x_n\|$ and Lemma 2.5, we note that $q \in EP(F, h)$.

(d) We show that $q \in F(\mathfrak{S})$. Assume the contrary that $q \neq T(h)q$ for some $h \in [0, \infty)$. Then by Opial's condition, we obtain from (3.38) that

$$\begin{aligned}
\liminf_{n \rightarrow \infty} \|z_{n_j} - q\| &< \liminf_{j \rightarrow \infty} \|z_{n_j} - T(h)q\| \\
&\leq \liminf_{j \rightarrow \infty} (\|z_{n_j} - T(h)z_{n_j}\| + \|T(h)z_{n_j} - T(h)q\|) \\
&\leq \liminf_{j \rightarrow \infty} \|z_{n_j} - q\|.
\end{aligned}$$

This is a contradiction. Hence, $q \in F(\mathfrak{S})$. Thus $q \in \Omega = F(\mathfrak{S}) \cap EP(F, h) \cap VI(C, A) \cap \Psi$. By (3.39) and property of metric projection, we obtain

$$\limsup_{n \rightarrow \infty} \langle (\gamma\phi - \theta T)w, w_n - w \rangle = \lim_{j \rightarrow \infty} \langle (\gamma\phi - \theta T)w, w_{n_j} - w \rangle = \langle (\gamma\phi - \theta T)w, q - w \rangle \leq 0.$$

Finally, we prove that $x_n \rightarrow w$, since $w_n - w = \alpha_n(\gamma\phi(x_n) - \theta Tw) + (I - \alpha_n\theta T)(y_n - w)$

$$\begin{aligned}
\|w_n - w\|^2 &= \|\alpha_n(\gamma\phi(x_n) - \theta Tw) + (I - \alpha_n\theta T)(y_n - w)\|^2 \\
&\leq (1 - \alpha_n\theta\bar{\gamma})^2 \|y_n - w\|^2 + 2\alpha_n \langle \gamma\phi(x_n) - \theta Tw, w_n - w \rangle \\
&\leq (1 - \alpha_n\theta\bar{\gamma})^2 \|x_n - w\|^2 + 2\alpha_n \langle \gamma\phi(x_n) - \gamma\phi(w), w_n - w \rangle + 2\alpha_n \langle \gamma\phi(w) - \theta Tw, w_n - w \rangle \\
&\leq (1 - \alpha_n\theta\bar{\gamma})^2 \|x_n - w\|^2 + \alpha_n\gamma(\|x_n - w\|^2 + \|w_n - w\|^2) + 2\alpha_n \langle \gamma\phi(w) - \theta Tw, w_n - w \rangle.
\end{aligned}$$

This implies that

$$\begin{aligned}
\|w_n - w\|^2 &\leq \frac{(1 - \alpha_n\theta\bar{\gamma})^2 + \alpha_n\gamma}{1 - \alpha_n\gamma} \|x_n - w\|^2 + \frac{2\alpha_n}{1 - \alpha_n\gamma} \langle \gamma\phi(w) - \theta Tw, w_n - w \rangle \\
&= (1 - \frac{2\alpha_n(\theta\bar{\gamma} - \gamma)}{1 - \alpha_n\gamma}) \|x_n - w\|^2 + \frac{\alpha_n^2(\theta\bar{\gamma})^2}{1 - \alpha_n\gamma} \|x_n - w\|^2 \\
&\quad + \frac{2\alpha_n}{1 - \alpha_n\gamma} \langle \gamma\phi(w) - \theta Tw, w_n - w \rangle.
\end{aligned} \tag{3.43}$$

$$\begin{aligned}
\|x_{n+1} - w\|^2 &\leq \beta_n \|x_n - w\|^2 + (1 - \beta_n) \left\| \left(\frac{1}{t_n} \int_0^{t_n} [T(u)w_n - T(u)w] du \right) \right\|^2 \\
&\leq \beta_n \|x_n - w\|^2 + (1 - \beta_n) \|w_n - w\|^2 \\
&\leq \beta_n \|x_n - w\|^2 + (1 - \beta_n) \left[\left(1 - \frac{2\alpha_n(\theta\bar{\gamma} - \gamma)}{1 - \alpha_n\gamma} \right) \|x_n - w\|^2 + \frac{\alpha_n^2(\theta\bar{\gamma})^2}{1 - \alpha_n\gamma} \|x_n - w\|^2 \right. \\
&\quad \left. + \frac{2\alpha_n}{1 - \alpha_n\gamma} \langle \gamma\phi(w) - \theta Tw, w_n - w \rangle \right] \\
&= [1 - (1 - \beta_n) \frac{2\alpha_n(\theta\bar{\gamma} - \gamma)}{1 - \alpha_n\gamma}] \|x_n - w\|^2 + \frac{\alpha_n(1 - \beta_n)}{1 - \alpha_n\gamma} [\alpha_n(\theta\bar{\gamma})^2 \|x_n - w\|^2 \\
&\quad + 2\langle \gamma\phi(w) - \theta Tw, w_n - w \rangle] \\
&= (1 - \delta_n) \|x_n - w\|^2 + b_n.
\end{aligned}$$

where $\delta_n = (1 - \beta_n) \frac{2\alpha_n(\theta\bar{\gamma} - \gamma)}{1 - \alpha_n\gamma}$ and $b_n = \frac{\alpha_n(1 - \beta_n)}{1 - \alpha_n\gamma} [\alpha_n(\theta\bar{\gamma})^2 \|x_n - w\|^2 + 2\langle \gamma\phi(w) - \theta Tw, w_n - w \rangle]$. It is easily verified that $\sum_{n=1}^{\infty} \delta_n = \infty$ and $\limsup_{n \rightarrow \infty} \frac{b_n}{\delta_n} \leq 0$. then by Lemma 2.8, we obtain $\lim_{n \rightarrow \infty} \|x_n - w\| = 0$. $\square$

Corollary 3.2. *Let C be a nonempty, closed and convex subset of a real Hilbert space H . For each $i = 1, 2$, let $A, B_i : C \rightarrow H$ be α, β_i -inverse strongly monotone mappings, respectively. Let F, h be a bi-function from $C \times C$ to R satisfying the hypotheses of Lemma 2.2, and $\phi, Y : C \rightarrow C$ are nonexpansive mappings such that $\Omega := F(Y) \cap EP(F, h) \cap VI(C, A) \cap \Psi \neq \emptyset$. Let T be a strongly bounded linear operator on H with coefficient $\bar{\gamma}$ such that $0 < \gamma < \theta\bar{\gamma}$ and $0 < \theta \leq \|T\|^{-1}$. Let $\{r_n\} \subset (0, \infty)$ be a real sequence such that $\liminf_{n \rightarrow \infty} r_n > 0$. Suppose Let $\{x_n\}$, $\{u_n\}$, $\{y_n\}$ and $\{z_n\}$ be sequences generated by*

$$\begin{cases} x_1 = x \in H, \\ F(u_n, y) + h(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in C \\ z_n = P_C(u_n - \lambda_n A u_n), \\ y_n = \delta_n z_n + (1 - \delta_n) P_C[P_C(z_n - \mu_2 B_2 z_n) - \mu_1 B_1 P_C(z_n - \mu_2 B_2 z_n)], \\ x_{n+1} = \beta_n x_n + (1 - \beta_n) \left(\frac{1}{n+1} \sum_{j=0}^n Y^j [\alpha_n \gamma \phi(x_n) + (I - \alpha_n \theta T) y_n] \right), \end{cases} \quad (3.1)$$

where $\mu_i \in (0, 2\beta_i)$, for each $i = 1, 2$, $\lambda_n \in (0, 2\alpha)$, and $\{\beta_n\}$, $\{\alpha_n\}$, $\{\delta_n\}$ are sequences in $[0, 1]$. Assume that the following conditions are satisfied:

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\sum_{n=1}^{\infty} \alpha_n = \infty$, $\sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty$;
- (ii) $\lim_{n \rightarrow \infty} \beta_n = 0$, $\sum_{n=1}^{\infty} |\beta_{n+1} - \beta_n| < \infty$;
- (iii) $\lim_{n \rightarrow \infty} \delta_n = 0$, $\sum_{n=1}^{\infty} |\delta_{n+1} - \delta_n| < \infty$;
- (iv) $\sum_{n=1}^{\infty} |r_{n+1} - r_n| < \infty$;
- (v) $0 < \liminf_{n \rightarrow \infty} \lambda_n \leq \limsup_{n \rightarrow \infty} \lambda_n < 2\alpha$, $\sum_{n=1}^{\infty} |\lambda_{n+1} - \lambda_n| < \infty$.

Then $\{x_n\}$ converges strongly to w , where $w := P_{\Omega}(\gamma\phi + (I - \theta T))w$.

Corollary 3.3. *Let C be a nonempty, closed and convex subset of a real Hilbert space H and $A : C \rightarrow H$ be α -inverse strongly monotone mappings. Let F, h be a bi-function from $C \times C$ to R satisfying the hypotheses of Lemma 2.2, and $\mathfrak{S} = \{T(u) : 0 \leq u < \infty\}$ be a one-parameter nonexpansive semigroup on H such that $\Omega := F(\mathfrak{S}) \cap EP(F, h) \cap VI(C, A) \neq \emptyset$. Let $\phi : H \rightarrow H$ be a nonexpansive mapping and T be a strongly bounded linear operator on H with coefficient $\bar{\gamma}$ such that $0 < \gamma < \theta\bar{\gamma}$ and $0 < \theta \leq \|T\|^{-1}$. Let $\{r_n\} \subset (0, \infty)$ be a real sequence such that $\liminf_{n \rightarrow \infty} r_n > 0$. Suppose Let $\{x_n\}$, $\{u_n\}$ and $\{z_n\}$ be sequences generated by*

$$\begin{cases} x_1 = x \in H, \\ F(u_n, y) + h(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in C \\ z_n = P_C(u_n - \lambda_n A u_n), \\ x_{n+1} = \beta_n x_n + (1 - \beta_n) \left(\frac{1}{t_n} \int_0^{t_n} T(u) [\alpha_n \gamma \phi(x_n) + (I - \alpha_n \theta T) z_n] du \right), \end{cases} \quad (3.1)$$

where $\lambda_n \in (0, 2\alpha)$, and $\{\beta_n\}$, $\{\alpha_n\}$ are sequences in $[0, 1]$. Assume that the following conditions are satisfied:

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\sum_{n=1}^{\infty} \alpha_n = \infty$, $\sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty$;
- (ii) $\lim_{n \rightarrow \infty} \beta_n = 0$, $\sum_{n=1}^{\infty} |\beta_{n+1} - \beta_n| < \infty$;
- (iii) $\sum_{n=1}^{\infty} |r_{n+1} - r_n| < \infty$;
- (iv) $\lim_{n \rightarrow \infty} \frac{|t_n - t_{n-1}|}{t_n} \frac{1}{\alpha_n(1-\beta_n)} = 0$;
- (v) $0 < \liminf_{n \rightarrow \infty} \lambda_n \leq \limsup_{n \rightarrow \infty} \lambda_n < 2\alpha$, $\sum_{n=1}^{\infty} |\lambda_{n+1} - \lambda_n| < \infty$.

Then $\{x_n\}$ converges strongly to w , where $w := P_{\Omega}(\gamma\phi + (I - \theta T))w$.

References

- [1] L.C. Ceng, J.C. Yao, An extragradient like approximation method for variational inequality problems and fixed point problems, Appl. Math. Comput. 190 (2007) 205-215.
- [2] L.C. Ceng, J.C. Yao, On the convergence analysis of inexact hybrid extragradient proximal point algorithms for maximal monotone operators, J. Comput. Appl. Math. 217 (2008) 326-338.
- [3] L.C. Ceng, J.C. Yao, Approximate proximal algorithms for generalized variational inequalities with pseudomonotone multifunctions, J. Comput. Appl. Math. 213 (2008) 423-438.
- [4] R. Glowinski, Numerical Methods for Nonlinear Variational Problems, Springer-Verlag, New York, 1984.
- [5] C. Jaiboon, P. Kumam, H.W. Humphries, Weak convergence theorem by extragradient method for variational inequality, equilibrium problems and fixed point problems, Bull. Malaysian Math. Sci. Soc. 2 (32) (2009) 173-185.
- [6] F. Liu, M.Z. Nasheed, Regularization of nonlinear ill-posed variational inequalities and convergence rates, Set Valued Anal. 6 (1998) 313-344.
- [7] G. Stampacchia, Formes bilineaires coercitives sur les ensembles convexes, C.R. Acad. Sci. Paris 258 (1964) 4413-4416.
- [8] R. Wangkeeree, R. Wangkeeree, A general iterative methods for variational inequality problems and mixed equilibrium problems and fixed point problems of strictly pseudocontractive mappings in Hilbert spaces, Fixed Point Theory Appl. (2007) 32 (Article ID 519065).
- [9] S.S. Zeng, N.C. Wong, J.C. Yao, Convergence of hybrid steepest-descent methods for generalized variational inequalities, Acta Math. Sin. English Ser. 22 (1) (2006) 1-12.
- [10] R.U. Verma, On a new system of nonlinear variational inequalities and associated iterative algorithms, J. Math. Sci. Res. Hot-Line 3 (8) (1999) 65-68.
- [11] L.C. Ceng, C.Y. Wang, J.C. Yao, Strong convergence theorems by a relaxed extragradient method for a general system of variational inequalities, Math. Methods Oper. Res. 67 (2008) 375-390.
- [12] W. Kumam, P. Kumam, Hybrid iterative scheme by a relaxed extragradient method for solutions of mixed equilibrium problems and general system of variational inequalities with application to optimization, Nonlinear Anal. Hybrid Syst. 3 (2009) 640-656.
- [13] R. Wangkeeree, U. Kamraks, An iterative approximation for solving general system of variational inequality problems, mixed equilibrium problems, Nonlinear Anal. Hybrid Syst. 3 (2009) 615-630.

- [14] Y. Yao, Y.C. Liou, S.M. Kang, Y. Yu, Algorithms with strong convergence for a system of nonlinear variational inequalities in Banach space, *Nonlinear Anal.* 74 (2011) 6024-6034.
- [15] A. Moudafi, Weak convergence theorems for nonexpansive mappings and equilibrium problems, *J. Nonlinear Convex Anal.* 9 (1) (2008) 37-43.
- [16] J.W. Peng, Iterative algorithms for mixed equilibrium problems, strictly pseudo-contractions and monotone mappings, *J. Optim. Theory Appl.* 144 (1) (2010) 107-119.
- [17] L.C. Ceng, J.C. Yao, A hybrid iterative scheme for mixed equilibrium problems and fixed point problems, *J. Comput. Appl. Math.* 214 (1) (2008) 186-201 (English summary).
- [18] J.W. Peng, J.C. Yao, A new hybrid-extragradient method for generalized mixed equilibrium problems, fixed point problems and variational inequality problems, *Taiwanese J. Math.* 12 (6) (2008) 1401-1432.
- [19] K.R. Kazmi, S.H. Rizvi, A hybrid extragradient method for approximating the common solutions of a variational inequality, a system of variational inequalities, a mixed equilibrium problem and a fixed point problem, *Appl. Math. Comput.* 218 (2012) 5439-5452.
- [20] G. Marino, L. g. Muglia and Y. H. Yao, Viscosity methods for common solutions of equilibrium and variational inequality problems via multi-step iterative algorithms and common fixed points, *Nonlinear Anal.*(2011),doi:10.1016/j.na.2011.09.019.
- [21] F. Cianciaruso, G. Marino, L. Muglia, Y. Yao, A hybrid projection algorithm for finding solutions of mixed equilibrium problem and variational inequality problem, *Fixed Point Theory Appl.* (2010) Art. ID 383740, 19 pp.
- [22] T. Shimizu, W. Takahashi, Strong convergence of common fixed points of families of nonexpansive mappings, *J. Math. Anal. Appl.* 211 (1997) 71-83.
- [23] K. Geobel, W.A. Kirk, Topics in Metric Fixed Point Theory, in: *Cambridge Stud. Adv. Math.*, vol. 28, Cambridge Univ. Press, 1990, pp. 473-504.
- [24] H.K. Xu, Iterative algorithm for nonlinear operators, *J. Lond. Math. Soc.* 66 (2) (2002) 1-17.
- [25] G. Marino, H.K. Xu, A general iterative method for nonexpansive mapping in Hilbert spaces, *J. Math. Anal. Appl.* 318 (2006) 43-52.
- [26] V. Colao, G. Marino, H.K. Xu, An iterative method for finding common solutions of equilibrium and fixed point problems, *J. Math. Anal. Appl.* 344 (1) (2008) 340-352 (English summary).
- [27] H. Iiduka, W. Takahashi, Strong convergence theorems for nonexpansive mappings and inverse-strongly monotone mappings, *Nonlinear Anal.* 61(2005) 341-350.
- [28] P. Katchang and P. Kumam, A composite implicit iterative process with a viscosity method for lipschitzian semigroup in a smooth Banach space, *Bulletin of the Iranian Mathematical Society*, Vol. 37 No. 1 (2011), 143-159.
- [29] W. Kumam, P. Junlouchai, and P. Kumam, Generalized Systems of Variational Inequalities and Projection Methods for Inverse-Strongly Monotone Mappings, *Discrete Dynamics in Nature and Society*, Volume 2011, Article ID 976505, 24 pages.
- [30] P. Katchang, C.a Jaiboon, and P. Kumam, Convergence theorems for two viscosity iterative algorithms for solving equilibrium problems and fixed point problems, *International Journal of Pure and Applied Mathematics*, Volume 72 No. 2 2011, 217-235.

- [31] Phayap Katchang and Poom Kumam, "Hybrid-extragradient type methods for a generalized equilibrium problem and variational inequality problems of nonexpansive semigroups" *Fixed Point Theory*, 13 (2012), 107-120.
- [32] P. Sunthrayuth and P. Kumam, "An iterative method for solving a system of mixed equilibrium problems, system of quasivariational inclusions and fixed point problems of nonexpansive semigroups with application to optimization problems," *Abstract and Applied Analysis*, Volume 2012, Article ID 979870, 30 pages.
- [33] P. Katchang and P. Kumam "A system of mixed equilibrium problems, a general system of variational inequality problems for relaxed cocoercive and fixed point problems for nonexpansive semigroup and strictly pseudo-contractive mappings," *Journal of Applied Mathematics*, Volume 2012, Article ID 414831, 36 pages.