

Generalized Growth and Approximation of Pseudoanalytic Functions on the Disk

Devendra Kumar¹

Department of Mathematics
M.M.H. College, Model Town, Ghaziabad-201001, U.P., India
d_kumar001@rediffmail.com

Vandna Jain

Applied Science Department (Mathematics)
Bhai Gurdas Institute of Engineering and Technology, Sangrur.
vandnajain.mittal@gmail.com

Balbir Singh

Department of Mathematics
Aryabhatta College, Barnala, (Punjab).
drbalbirdhami@gmail.com

Abstract. McCoy [15] considered the approximation of pseudoanalytic functions (PAF) on the disk. Pseudoanalytic functions are constructed as complex combination of real - valued analytic solutions to the Stokes-Beltrami System. These solutions include the generalized biaxisymmetric potentials. McCoy obtained some coefficient and Bernstein type growth theorems on the disk. The aim of this paper is to generalize the results of McCoy [15]. Moreover, we study the generalized order and generalized type of PAF in terms of Fourier coefficients occurring in its local expansion and optimal approximation errors in Bernstein sense on the disk. Our approach and method are different from those of McCoy [15].

2010 Mathematics Subject classification: 30E10, 41A20.

Keywords and phrases. Pseudoanalytic functions, generalized growth, approximation errors, Fourier coefficients and Stokes - Beltrami System.

1 Introduction

In Bernstein work [5,8,16,18] there is an implicit characterization of the solutions of a system of partial differential equations since the real and imaginary parts of the real analytic function solve the Cauchy-Riemann system. Thus, there is a notion of the harmonic conjugate and its role in the approximation process. However, this concept is not found in the function theory of the generalized biaxisymmetric potentials (GBASP) equation and is necessary for a complete function theoretic generalization of Bernstein type. GBASP that are harmonic at the origin may be expanded, in analogy with the Taylor's series for analytic functions of a single complex variable, in a convergent series of homogeneous harmonic polynomials on an open set.

Pseudoanalytic functions are constructed as complete combinations of real - valued function pairs that are analytic solutions of Stokes - Beltrami System (SBS); a generalization of the Cauchy - Riemann equations that is linked to the GBASP equation by eliminating one of the dependent variables from the system. Pseudoanalytic functions provide sufficient basis for the transformation of Bernstein's ideas through

¹This work was done in the memory of Prof. H.S.Kasana, Senior Associate, ICTP, Trieste, Italy

transform and special function methods. The real part of pseudoanalytic function i.e., eliminating the harmonic conjugate gives the theory of GBASP. The GBASP equation frequently found in the summability theory of [3] Jacobi series as

$$r \frac{\partial}{\partial r} \left\{ r^{\alpha+\beta+1} \rho^{(\alpha,\beta)}(\theta) r \frac{\partial u}{\partial r} \right\} + \frac{\partial}{\partial \theta} \left\{ r^{\alpha+\beta+1} \rho^{(\alpha,\beta)}(\theta) \frac{\partial u}{\partial \theta} \right\} = 0$$

$$\rho^{(\alpha,\beta)}(\theta) = (\sin \theta/2)^{2\alpha+1} (\cos \theta/2)^{2\beta+1}, \alpha \geq \beta > -\frac{1}{2} \quad (1.1)$$

where (r, θ) are the plane polar coordinates. The domain of the potential u is a simply connected region with smooth boundary in the upper half plane $C^+ = C \cap \text{Re}(z) \geq 0$. The existence of a harmonic conjugate ν of u is implied in the sense of the generalized Stokes - Beltrami System (SBS);

$$\begin{aligned} r \frac{\partial \nu}{\partial r} &= -r^{\alpha+\beta+1} \rho^{(\alpha,\beta)}(\theta) \frac{\partial u}{\partial \theta} \\ \frac{\partial \nu}{\partial \theta} &= r^{\alpha+\beta+1} \rho^{(\alpha,\beta)}(\theta) r \frac{\partial u}{\partial r} \end{aligned}$$

a system that reduces to the Cauchy-Riemann equations of analytic function theory in the limit $\alpha = \beta = -1/2$. Following along the lines of analytic function theory, a pseudoanalytic function [2, 7, 20] (PAF) is defined as the complex combination

$$F(re^{i\theta}) = u(r, \theta) + i\nu(r, \theta)$$

of a real-valued analytic function pair formed from the potential u and the principal branch of its harmonic conjugate $\nu = \nu(r, \theta)$. In particular, a PAF whose domain Ω^+ intersects the x -axis in a single component may be analytically continued across the axis by reflexion to Ω^- as an even function on $\Omega : \Omega^+ \cup \Omega^-$ provided that $\frac{\partial u(r, 0+\pi k)}{\partial \theta} = \frac{\partial \nu(r, 0+\pi k)}{\partial \theta} = 0, 1[7]$.

The Bernstein theorem identify a real analytic function on the closed unit disk as the restriction of an analytic function defined on an open disk of radius $R > 1$ by computing R from the sequence of minimal errors generated from optimal polynomials approximates. Here we use this concept for pseudoanalytic functions (PAF). The disk D_R of maximum radius on which a PAF F exists, is designated by $F \in P(D_R)$. If F is an entire PAF, it has no singularities in the finite C^+ -plane and writes $F \in P(C)$.

The maximum modulus of a pseudoanalytic function $F \in P(D_R)$ is defined on the closure of a disk D_R as

$$M_r(F) = \max\{|F(re^{i\theta})| : re^{i\theta} \in \overline{D_R}\}, r < R.$$

Let p and q be two positive, strictly increasing to infinity differentiable functions $]0 + \infty[$ to $]0, +\infty[$ such that for every $c > 0$ satisfies:

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{p(cx)}{p(x)} &= 1, \\ \lim_{x \rightarrow \infty} \frac{q(1+xw(x))}{q(x)} &= 1, \lim_{x \rightarrow \infty} w(x) = 0, \\ \lim_{x \rightarrow \infty} \frac{d(q^{-1}(cp(x)))}{d(\log x)} &\leq b. \end{aligned}$$

$$p\left(\frac{x}{q^{-1}(cp(x))}\right) = (1 + o(x))p(x), \text{ for } x \rightarrow \infty,$$

where $d(u)$ means differential of u .

The (p, q) -order and (p, q) -type of an entire PAF $F \in P(C)$ (or generalized order and generalized type) are defined respectively by:

$$\rho(p, q) = \limsup_{r \rightarrow \infty} \frac{p(\log M(r, f))}{q(\log r)}$$

and

$$\sigma(p, q) = \limsup_{r \rightarrow \infty} \frac{p(M(r, f))}{[q(r)]^{\rho(p, q)}}$$

where

$$M(r, f) = \sup_{z \in \overline{D}_R} |f(z)|.$$

Now applying the concept introduced by G.R. MacLane [11] to the measures of order and type for an analytic function on a disk $|z| < R$ by normalizing these definitions relative to the boundary under the transformation $r \rightarrow R/(R - r)$. In this direction, a pseudoanalytic function $F \in P(D_R)$ with radial limits is said to be of generalized regular growth $(\rho_0(p, q), \sigma_0(p, q))$ if it satisfies

$$\begin{aligned} \rho_0(p, q) &= \limsup_{r \rightarrow R} \frac{p(\log M_r(F))}{q(R/(R - r))}, \\ \sigma_0(p, q) &= \limsup_{r \rightarrow R} \frac{p(\log M_r(F))}{[q(R/(R - r))]^{\rho_0(p, q)}}, 1 < r < R, \end{aligned}$$

in which case $\rho_0(p, q)$ is referred to as the (p, q) - order of F provided that $0 < \rho_0(p, q) < \infty$ and $\sigma_0(p, q)$ the (p, q) -type. Normalized to the closed unit disk $\overline{D}_1$ the $M_r(F)$ is a special case of the norm, $\|\cdot\|$,

$$\|F\| = \begin{cases} \|F\|_\delta = \left[\int_{\overline{D}_1} |F|^\delta r dr d\phi \right]^{1/\delta}, & 1 \leq \delta < \infty \\ \|F\|_\infty = M_1(F), & \delta = \infty. \end{cases}$$

The approximating pseudoanalytic polynomials of (fixed) degree n are taken from the sets

$$\pi_n = \left\{ P : P(re^{i\theta}) = \sum_{k=0}^n c_k w_k F_k(r, e^{i\theta}), c_k \text{ real} \right\}, n = 0, 1, 2, \dots$$

The optimal approximates minimize the error $\|F - P\|$ for $P \in \pi_n$ in Bernstein sense as

$$E_n(F) = \inf \{ \|F - P\| : P \in \pi_n \}, n = 0, 1, 2, \dots \quad (1.2)$$

P.A.McCoy [12-14] studied the function - theoretic methods based on the integral operator approach with Bernstein's ideas in the case of weakening the norms measuring the best approximates, and through less stringent analyticity requirements such

as replacing the restriction of an entire function on an open disk simply with analyticity to GBASP. Kumar [10] obtained the relationship between the pseudoanalytic functions and Bergmann Gilbert type integral operators for GBASP and polynomial approximation. Bergman [4] and Gilbert [7] generalize the operation of taking the real part. They obtained bounded linear operators which transform analytic functions to solution u , where integral operators are developed to provide the transformation from analytic functions to solutions of corresponding elliptic equation. Bers [6] and Vekua [19] have also extended function theory so that solutions u of elliptic equations can be obtained as $u = \text{Re}(F)$, where f is a pseudoanalytic function sharing many of the properties associated with classical analytic functions of single complex variable.

McCoy [15] considered the approximation of pseudoanalytic functions on the disk and obtained some coefficient and Bernstein type growth theorems. Also Kapoor and Nautiyal [9] characterized the order and type of GBASP's (not necessarily entire) in terms of rates of decay of approximation errors on both sup norm and L^p -norm, $1 \leq p < \infty$. It has been noticed that all these authors have not studied the generalized growth of pseudoanalytic functions on the disk. In this paper we study the generalized order and generalized type of PAF in terms of Fourier coefficients occurring in its local expansion and optimal approximation errors in Bernstein sense on the disk. Our results and methods in the present paper are different from all those of authors mentioned above.

2 Generalized Order and Generalized Type with Fourier Coefficients of Pseudoanalytic Functions

Let the pseudoanalytic function $F \in P(D_R)$. In a neighborhood of the origin, a PAF has the local expansion

$$F(re^{i\theta}) = \sum_{n=0}^{\infty} a_n w_n F_n(re^{i\theta}), re^{i\theta} \in D_R,$$

$F_n(re^{i\theta}) = u_n(r, \theta) + i\nu_n(r, \theta)$, $n = 0, 1, 2, \dots$, and a_n real-valued. Also we have the estimate $|w_n F_n(re^{i\theta})| \sim n^{2\alpha+1} r^{n+\alpha+\beta+1}$ for large n . Put

$$\limsup_{n \rightarrow \infty} \frac{p(n)}{q \left[\frac{n}{\log(n^{2\alpha+1} |a_n| R^n)} \right]} = \mu(p, q). \quad (2.1)$$

Lemma 2.1. For every $r > 1$ and $\mu > 0$, the maximum of the function

$$x \rightarrow w(x, r) = x \log(r/R) + \frac{x}{q^{-1}(p(x)/\mu)}$$

is reached for $x = x_r$ solution of the equation

$$x = p^{-1} \left\{ \mu q \left[\frac{1 - d \log(q^{-1}(p(x)/\mu))/d(\log x)}{\log(R/r)} \right] \right\}.$$

Proof. Let $G(x, \mu) = q^{-1}(p(x)/\mu)$, then

$$w(x, r) = x \log(r/R) + \frac{x}{G(x, \mu)}.$$

The maximum of the function $x \rightarrow w(x, r)$ is reached for $x = x_r$, solution of the equation of $\frac{dw(x, r)}{dx} = 0$.

We have

$$\frac{dw(x, r)}{dx} = 0 \Leftrightarrow \log(r/R) + \frac{G(x, \mu) - x \frac{dG(x, \mu)}{dx}}{(G(x, \mu))^2} = 0$$

or

$$G(x, \mu) = \frac{1 - \frac{x}{G(x, \mu)} \frac{dG(x, \mu)}{dx}}{\log(R/r)}.$$

Since

$$\frac{dG(x, \mu)}{dx} = \frac{dG(x, \mu)}{d(\log x)} \cdot \frac{d(\log x)}{dx} = \frac{1}{x} \frac{dG(x, \mu)}{d(\log x)},$$

we get

$$G(x, \mu) = \frac{1 - \frac{1}{G(x, \mu)} \frac{dG(x, \mu)}{d(\log x)}}{\log(R/r)} = \frac{1 - \frac{d \log G(x, \mu)}{d(\log x)}}{\log(R/r)},$$

we obtain

$$x = x_r = p^{-1} \left\{ \mu q \left[\frac{1 - d \log(q^{-1}(p(x)/\mu))/d(\log x)}{\log(R/r)} \right] \right\}. \quad (2.2)$$

Lemma 2.2. Let $F(re^{i\theta}) = \sum_{n=0}^{\infty} a_n w_n F_n(re^{i\theta})$, $F \in P(D_R)$. For every $1 < r < R$, we put

$$\begin{aligned} \overline{M}(r, F) &= \sup_{n \in N} \{ \|a_n w_n\| \cdot r^n, r > 0 \}, \\ \bar{\rho}_0(p, q) &= \limsup_{r \rightarrow R} \frac{p(\log \overline{M}(r, F))}{q(R/(R-r))}, \end{aligned}$$

then

$$\bar{\rho}_0(p, q) \leq \mu(p, q) \quad \text{and} \quad \rho_0(p, q) \leq \bar{\rho}_0(p, q).$$

Proof. From (2.1) we have for r sufficiently close to R and $\bar{\mu} = \mu + \varepsilon$,

$$\log(n^{2\alpha+1} |a_n| R^n) \leq \frac{n}{q^{-1} \left(\frac{p(n)}{\bar{\mu}} \right)}$$

or

$$\log(n^{2\alpha+1} |a_n| R^n) \leq n \log(r/R) + \frac{n}{q^{-1} \left(\frac{p(n)}{\bar{\mu}} \right)}.$$

Using the result

$$\lim_{x \rightarrow \infty} \left| \frac{d(q^{-1}(cp(x)))}{d(\log x)} \right| \leq b, \log(1+t) = (1+o(t)).t, t \rightarrow 0,$$

we get from (2.2)

$$x_r = (1+o(1))p^{-1}(\mu q(R/(R-r))) \quad \text{as } r \rightarrow R.$$

It gives

$$\log(n^{2\alpha+1}|a_n|r^n) \leq C_0 p^{-1}(\mu q(R/(R-r))),$$

or

$$\log \overline{M}(r, F) \leq C_0 p^{-1}(\mu q(R/(R-r))),$$

using the property of p , we obtain

$$\frac{p(\log \overline{M}(r, F))}{q(R/(R-r))} \leq \bar{\mu}.$$

Proceeding the limit supremum as $r \rightarrow R$, we get

$$\bar{\rho}_0(p, q) \leq \mu. \quad (2.3)$$

Now consider

$$\begin{aligned} f(z) &= \sum_{n=0}^{\infty} a_n z^n \\ M(r, f) &\leq \sum_{n=0}^{\infty} |a_n| r^n \leq \sum_{n=0}^{\infty} |a_n| w_n r^n, \end{aligned}$$

substituting $r = \sqrt{r.R} \cdot \sqrt{(r/R)}$ in above we have

$$M(r, f) \leq \sum_{n=0}^{\infty} |a_n| w_n \left(\sqrt{r.R}\right)^n \left(\sqrt{(r/R)}\right)^n, (r/R) < 1,$$

or

$$M(r, f) \leq \sum_{n=0}^{\infty} \sup \left(|a_n| w_n \left(\sqrt{r.R}\right)^n \right) \cdot \left(\sqrt{(r/R)}\right)^n$$

or

$$\begin{aligned} M(r, f) &\leq \overline{M}(r', F) \sum_{n=0}^{\infty} \left(\sqrt{(r/R)}\right)^n \\ &\leq \overline{M}(r', F) \left(\frac{1}{1 - \sqrt{(r/R)}} \right), r' = \sqrt{(r.R)} \end{aligned}$$

It gives

$$\log M(r, f) \leq \log \overline{M}(r', F) - \log \left(1 - \sqrt{(r/R)} \right)$$

or

$$\frac{p(\log M(r, f))}{q(R/(R-r))} \leq \frac{p(\log \overline{M}(r', F) - \log(1 - \sqrt{(r/R)}))}{q(R/(R - \sqrt{(r.R)}))} \cdot \frac{q(R/(R - \sqrt{(r.R)}))}{q(R/(R-r))}.$$

Passing to limit we get

$$\rho_0(p, q) \leq \bar{\rho}_0(p, q). \quad (2.4)$$

From (2.3) and (2.4) we get $\rho_0(p, q) \leq \mu(p, q)$.

Theorem 2.1. Let $F(re^{i\theta}) = \sum_{n=0}^{\infty} a_n w_n F_n(re^{i\theta})$, $F \in P(D_R)$, $re^{i\theta} \in D_R$ such that

$$\limsup_{n \rightarrow \infty} \frac{p(n)}{q \left[\frac{n}{\log(|a_n| n^{2\alpha+1} R^n)} \right]} = \mu(p, q) < \infty.$$

Then F is the restriction of pseudoanalytic function in $P(D_R)$ ($R > 1$) and its (p, q) -order $\rho(p, q) = \mu(p, q)$.

Proof. To study the expansion of a PAF in terms of a complete set of particular solutions of the SBS. Let us consider the normalized Jacobi polynomials $R_n(\cos \theta) = P_n^{(\alpha, \beta)}(\cos \theta) / P_n^{(\alpha, \beta)}(1)$, $n = 0, 1, 2, \dots$, [1, 17] which form an orthogonal set

$$\begin{aligned} \int_0^\pi R_n(\cos \phi) R_m(\cos \phi) \rho^{(\alpha, \beta)}(\phi) d\phi &= w_n^{-1} \delta_{nm} \\ w_n &= \frac{(2n + \alpha + \beta + 1) \Gamma(n + \alpha + \beta + 1) \Gamma(n + \alpha + 1)}{\Gamma(n + \beta + 1) \Gamma(n + 1) \Gamma(\alpha + 1) \Gamma(\beta + 1)} \sim O(n^{2\alpha}) \end{aligned}$$

relative to the measure $\rho^{(\alpha, \beta)}(\phi) d\phi$ on $[0, \pi]$. The Abel - Poisson and conjugate Abel-Poisson means [3, 17] of these polynomials

$$\begin{aligned} u_n(r, \theta) &= A_r(R_n(\cos \theta)) = r^n R_n(\cos \theta) \\ \nu_n(r, \theta) &= \tilde{A}_r(R_n(\cos \theta)) \\ &= nr^{n+\alpha+\beta+1} \rho^{(\alpha, \beta)}(\theta) R_{n-1}^{(\alpha+1, \beta+1)}(\cos \theta) (\cos \theta / 2) (\sin \theta / 2) / (\alpha + 1). \end{aligned} \quad (2.5)$$

$(R_{-1}^{(\alpha+1, \beta+1)}(\cos \theta) \equiv 1)$ produce a set of particular solutions [3] that is complete [17] relative to uniform convergence on compacta of a sufficiently small region. In view of the local expansion of PAF and the estimate

$$|w_n F_n(re^{i\theta})| \sim n^{2\alpha} (1 + nr^{\alpha+\beta+1}) r^n$$

with the bound $|R_n(\cos \theta)| \leq 1$, a standard argument shows that the Fourier coefficients support the natural domain since a necessary and sufficient condition for $F \in P(D_R)$ is that

$$\limsup_{n \rightarrow \infty} |a_n|^{1/n} = R^{-1}.$$

So, for every $1 < r < R$ the series $\sum_{n=0}^{\infty} a_n w_n F_n(re^{i\theta})$ is convergent in D_R when $\sum_{n=0}^{\infty} a_n w_n F_n(re^{i\theta})$ is analytic in D_R .

Now we shall prove that $\mu(p, q)$ is the (p, q) -order of F .

By Lemma 2.2, to complete the proof of the theorem it suffices to show that $\rho(p, q) \geq \mu(p, q)$.

By definition of $\rho(p, q)$, we have for every $\varepsilon > 0$ there exists $1 < r_\varepsilon < R$ such that for every $r_\varepsilon < r < R$

$$\log M_r(F) \leq p^{-1}[(\rho(p, q) + \varepsilon) \cdot q(R/(R - r))]. \quad (2.6)$$

Let $\overline{D}_{r_0}$ be the closure of a disk D_{r_0} in the natural domain of D_R of a PAF. Such a PAF has the representation [3]

$$F(re^{i\theta}) = \{A_r(\cdot) + i\tilde{A}_r(\cdot)\}f(r_0e^{i\theta}), re^{i\theta} \in D_{r_0} \quad (2.7)$$

where the Abel means, as in (2.5), are

$$\begin{aligned} A_r(f(r_0e^{i\theta})) &= \sum_{n=0}^{\infty} a_n w_n u_n(r, \theta), \\ \tilde{A}_r(f(r_0e^{i\theta})) &= \sum_{n=0}^{\infty} a_n w_n \nu_n(r, \theta). \end{aligned}$$

At the boundary (2.7) reduces to

$$F(r_0e^{i\theta}) = f(r_0e^{i\theta}) + ir_0^{(\alpha+\beta+1)}\rho^{(\alpha,\beta)}(\theta)\tilde{f}(r_0e^{i\theta})$$

where the function f and its conjugate $\tilde{f}$ are real-valued. The Fourier coefficients are reclaimed by orthogonality as

$$a_n r^n = \int_0^\pi f(re^{i\phi}) R_n(\cos \phi) \rho^{(\alpha,\beta)}(\phi) d\phi, r < r_0.$$

Integrating this identity, and applying Holder's inequality we get

$$\begin{aligned} |a_n/(n+2)| &= \left| \int_0^1 a_n r^n r dr \right| \leq \int_0^1 \int_0^\pi |f(re^{i\phi})| \rho^{(\alpha,\beta)}(\phi) r dr d\phi \\ &\leq \int_0^1 \int_0^\pi |F(re^{i\phi})| \rho^{(\alpha,\beta)}(\phi) r dr d\phi \\ &\leq \|F\|_\delta \cdot A, A = \|\rho^{(\alpha,\beta)}\|_{\delta'}, \frac{1}{\delta} + \frac{1}{\delta'} = 1, \end{aligned}$$

or

$$|a_n| \leq (n+2)A\|F\|_\delta. \quad (2.8)$$

Now using (2.5) for $1 < r$ sufficiently close to R in above inequality we obtain

$$\log(|a_n|n^{2\alpha+1}R^n) \leq -n \log(r/R) + \log((n+2)n^{2\alpha+1}A) + p^{-1}[(\rho(p, q) + \varepsilon)q(R/(R-r))].$$

The minimum value of right hand side satisfying above inequality is attained at

$$r = R \left\{ 1 - \frac{1}{q^{-1} \left(\frac{1}{(\rho+\varepsilon)} p \left(\frac{n}{q^{-1}(p(n)/(\rho+\varepsilon))} \right) \right)} \right\}, \rho = \rho(p, q),$$

applying the properties of the function p and q

$$p \left(\frac{x}{q^{-1}(cp(x))} \right) \simeq (1 + o(x))p(x) \text{ for } c > 0, x \rightarrow \infty,$$

and the properties of logarithm, we get

$$\log(|a_n|n^{2\alpha+1}R^n) \leq c_1 \frac{n}{q^{-1}(p(n)/(\rho+\varepsilon))}$$

where c_1 is a constant. Hence

$$q \left(\frac{c_1 n}{\log(|a_n|n^{2\alpha+1}R^n)} \right) \geq p(n)/(\rho+\varepsilon).$$

Now proceeding to limit supremum as $n \rightarrow \infty$ we get

$$\mu(p, q) \leq \rho(p, q).$$

Hence the proof is complete.

Remark 2.1. For $p(x) = q(x) = \log x$, Theorem 2.1 gives part(i) of [15, Thm. 2].

Let $F = \sum_{n=0}^{\infty} a_n w_n F_n(re^{i\theta})$ be pseudoanalytic function of (p, q) -order $\rho = \rho(p, q)$, and write

$$T(p, q) = \limsup_{n \rightarrow \infty} \frac{p(n)}{\left\{ q \left(\frac{n}{\log(|a_n| n^{2\alpha+1} R^n)} \right) \right\}^{\rho(p, q)}}.$$

Now we prove

Lemma 2.3. Let $F = \sum_{n=0}^{\infty} a_n w_n F_n(re^{i\theta})$. For every $1 < r < R$,

$$\bar{\sigma}_1(p, q) = \limsup_{r \rightarrow R} \frac{p(\log \bar{M}(r, F))}{(q(R/(R-r)))^{\rho(p, q)}}$$

then

$$\sigma(p, q) \leq \bar{\sigma}_1(p, q).$$

Proof. Using the same reasoning as in the proof of Lemma 2.2. we get

$$\frac{p(\log M(r, f))}{[q(R/(R-r))]^{\rho(p, q)}} \leq \frac{p(\log(\bar{M}(\sqrt{rR}, F) - \log(1 - \sqrt{(r/R)})))}{[q(R/(R - \sqrt{rR}))]^{\rho(p, q)}} \cdot \frac{[q(R/(R - \sqrt{rR}))]^{\rho(p, q)}}{[q(R/(R-r))]^{\rho(p, q)}}.$$

Proceeding to the limit supremum as $r \rightarrow R$, we get

$$\sigma(p, q) \leq \bar{\sigma}_1(p, q).$$

Theorem 2.2. Let $F = \sum_{n=0}^{\infty} a_n w_n F_n(re^{i\theta})$ be a pseudoanalytic function on the closed unit disk. If F is of finite generalized (p, q) -order $\rho(p, q)$, and

$$T(p, q) = \limsup_{n \rightarrow \infty} \frac{p(n)}{\left[q \left(\frac{n}{\log(|a_n| n^{2\alpha+1} R^n)} \right) \right]^{\rho(p, q)}} < \infty.$$

Then F is the restriction of a pseudoanalytic function in $P(D_R)(R > 1)$ and its (p, q) -type $\sigma(p, q) = T(p, q)$.

Proof. The function F is the restriction of a pseudoanalytic function in $P(D_R)$ by using the arguments in Theorem 2.1. Now first we show that $\sigma(p, q) \leq T(p, q)$.

If $T \equiv T(p, q) < \infty$, by definition of T , for every $\varepsilon > 0$, there exists $n \geq n_\varepsilon$

$$p(n) \leq (T + \varepsilon) \left[q \left(\frac{n}{\log(|a_n| n^{2\alpha+1} R^n)} \right) \right]^\rho$$

or

$$\log(|a_n| n^{2\alpha+1} R^n) \leq \frac{n}{q^{-1} \left(\left(\frac{p(n)}{T} \right)^{1/\rho} \right)}, \bar{T} = T + \varepsilon, \quad (2.9)$$

for every $n \geq n_\varepsilon$.

Since

$$\log(|a_n|n^{2\alpha+1}r^n) \leq n \log(r/R) + \log(|a_n|n^{2\alpha+1}R^n).$$

By the relation (2.9), we get

$$\log(|a_n|n^{2\alpha+1}r^n) \leq n \log(r/R) + \frac{n}{q^{-1} \left(\left(\frac{p(n)}{T} \right)^{1/\rho} \right)}. \quad (2.10)$$

For every $1 < r < R$ and r sufficiently close to R , we put

$$\varphi(x, r) = x \log(r/R) + \frac{x}{q^{-1} \left(\left(\frac{p(x)}{T} \right)^{1/\rho} \right)}.$$

If we put $S = S(x, \bar{T}, 1/\rho) = q^{-1} \left(\left(\frac{p(x)}{T} \right)^{1/\rho} \right)$ then

$$\varphi(x, r) = x \log(r/R) + \frac{x}{S},$$

and the maximum of the function $x \rightarrow \varphi(x, r)$ is reached for $x = x_r$ solution of the equation

$$\frac{d\varphi(x, r)}{dx} = \frac{\partial \varphi(x, r)}{\partial x} = \log(r/R) + \frac{d}{dx} \left(\frac{x}{S} \right) = 0.$$

or

$$\frac{d\varphi(x, r)}{dx} = 0 \Leftrightarrow \log(r/R) + \frac{S - x \frac{dS}{dx}}{S^2} = 0$$

or

$$S = \frac{1 - \frac{x}{S} \frac{dS}{dx}}{\log(R/r)}.$$

Since

$$\frac{dS}{dx} = \frac{dS}{d \log x} \cdot \frac{d \log x}{dx} = \frac{1}{x} \frac{dS}{d \log x},$$

we put

$$S = \frac{1 - \frac{1}{S} \frac{dS}{d \log x}}{\log(R/r)} = \frac{1 - \frac{d \log S}{d \log x}}{\log(R/r)},$$

or

$$q^{-1} \left(\left(\frac{p(x)}{T} \right)^{1/\rho} \right) = \frac{1 - \frac{d \log q^{-1} \left(\left(\frac{p(x)}{T} \right)^{1/\rho} \right)}{d \log x}}{\log(R/r)}.$$

It gives

$$x = x_r = p^{-1} \left\{ \left[\bar{T} q \left(\frac{1 - d \log \left(q^{-1} \left(\frac{p(x)}{\bar{T}} \right)^{1/\rho} \right) / d(\log x)}{\log(R/r)} \right) \right]^\rho \right\}.$$

We have

$$\log(r/R) = \log \left(\frac{r-R}{R} + 1 \right) \sim \frac{r-R}{R} \left(\text{as } \frac{r-R}{R} \rightarrow 0 \right),$$

and

$$\left| \frac{d \left[\log \left(q^{-1} \left(\frac{p(x)}{\bar{T}} \right)^{1/\rho} \right) \right]}{d(\log x)} \right| \leq b,$$

where b is a positive constant. Now we have

$$x_r = (1 + o(1)) \rho p^{-1} \left(\bar{T} q(R/(R-r))^\rho \right).$$

Using the relation (2.10), we get

$$\log(|a_n| n^{2\alpha+1} r^n) \leq \sup \varphi(x, r)$$

or

$$\log(|a_n| n^{2\alpha+1} r^n) \leq \frac{(1 + o(1)) p^{-1} \left(\bar{T} (q(R/(R-r)))^\rho \right)}{R/(R-r)}.$$

Since $\frac{R}{R-r} > 1$ and $\frac{\rho-1}{\rho} < 1$, then

$$\log(|a_n| n^{2\alpha+1} r^n) \leq C p^{-1} (\bar{T} (q(R/(R-r)))^\rho).$$

Then

$$\log \bar{M}(r, F) \leq C p^{-1} (\bar{T} (q(R/(R-r)))^\rho).$$

Thus

$$\frac{p(\log \bar{M}(r, F))}{(q(R/(R-r)))^\rho} \leq \bar{T}.$$

Proceeding to limit supremum for $r \rightarrow R$, we get

$$\bar{\sigma}_1(p, q) = \limsup_{r \rightarrow R} \frac{p(\log \bar{M}(r, F))}{(q(R/(R-r)))^\rho} \leq T.$$

Now using Lemma 2.3, we obtain

$$\sigma(p, q) \leq T(p, q).$$

The result obviously holds for $T = \infty$. Now we shall show that $\sigma(p, q) \geq T(p, q)$.

Suppose that $\sigma(p, q) < \infty$. By definition of $\sigma(p, q)$, we have for every $\varepsilon > 0$, there exist $r_\varepsilon \in]1, R[$, such that for every $r > r_\varepsilon$ ($R > r > r_\varepsilon > 1$)

$$\log M(r, F) \leq p^{-1} [\bar{\sigma}(q(R/(R-r)))^\rho].$$

Now using (2.8) for $1 < r$ sufficiently close to R with above inequality we get

$$\log(|a_n|n^{2\alpha+1}R^n) \leq -n \log(r/R) + \log((n+2)n^{2\alpha+1}A) + p^{-1} [\bar{\sigma}(q(R/(R-r)))^\rho]$$

or

$$\frac{\log(|a_n|n^{2\alpha+1}R^n)}{n} \leq \eta(r, n)$$

where

$$\eta(r, n) = -\log(r/R) + \frac{1}{n} \log((n+2)n^{2\alpha+1}A) + \frac{1}{n} p^{-1} [\bar{\sigma}(q(R/(R-r)))^\rho].$$

Since

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log((n+2)n^{2\alpha+1}A) + \frac{1}{n} p^{-1} [\bar{\sigma}(q(R/(R-r)))^\rho] = 0,$$

we get, for r sufficiently close to R

$$\lim_{n \rightarrow \infty} \eta(r, n) = -\log(r/R) = \log(R/r).$$

So we have

$$\eta(r, n) = (1 + o(1)) \log(R/r), n \rightarrow \infty,$$

then

$$\frac{1}{n} \log(|a_n|n^{2\alpha+1}R^n) \leq (1 + o(1)) \log(R/r). \quad (2.11)$$

Choose

$$r = R \frac{q^{-1} \left(\frac{p(n)}{\bar{\sigma}} \right)^{1/\rho}}{1 + q^{-1} \left(\frac{p(n)}{\bar{\sigma}} \right)^{1/\rho}}.$$

Using the relation (2.11) and the properties of the function $t \rightarrow \log(t)$, we get for r sufficiently close to R

$$\frac{\log(|a_n|n^{2\alpha+1}R^n)}{n} \leq (1 + o(1))((R/r) - 1).$$

$$\left[\log(R/r) = \log \left(\frac{R-r+r}{r} \right) = \log \left(1 + \frac{(R-r)}{r} \right) \sim \frac{(R-r)}{r}. (r \rightarrow R) \right].$$

Now we have

$$\frac{R-r}{r} = \frac{1}{q^{-1} \left(\frac{p(n)}{\bar{\sigma}} \right)^{1/\rho}}.$$

Then for r sufficiently close to R and large n , we get

$$\frac{\log(|a_n|n^{2\alpha+1}R^n)}{n} \leq \frac{1}{q^{-1}\left(\frac{p(n)}{\bar{\sigma}}\right)^{1/\rho}}$$

or

$$q^{-1}\left(\frac{p(n)}{\bar{\sigma}}\right)^{1/\rho} \leq \frac{n}{\log(|a_n|n^{2\alpha+1}R^n)}$$

or

$$\frac{p(n)}{\bar{\sigma}} \leq \left\{q\left(\frac{n}{\log(|a_n|n^{2\alpha+1}R^n)}\right)\right\}^\rho$$

or

$$\frac{p(n)}{\left\{q\left(\frac{n}{\log(|a_n|n^{2\alpha+1}R^n)}\right)\right\}^\rho} \leq \bar{\sigma} = \sigma + \varepsilon.$$

Proceeding to limit supremum as $n \rightarrow \infty$, we get

$$\sigma(p, q) \geq T(p, q).$$

The result is obviously holds for $\sigma(p, q) = \infty$. Hence the proof is complete.

Remark 2.2. For $p(x) = q(x) = x$, Theorem 2.2 gives part (iii) of [15, Thm. 2].

3 Generalized (p, q) –Growth and Approximation of Pseudoanalytic Functions

In this section we obtain the generalized order and generalized type in terms of approximation errors defined by (1.2).

First we prove

Lemma 3.1. Let the pseudoanalytic function PAF $F \in P(D_R)$, has the series expansion

$$F(re^{i\theta}) = \sum_{n=0}^{\infty} a_n w_n F_n(re^{i\theta}), re^{i\theta} \in D_R,$$

then

$$\limsup_{n \rightarrow \infty} \frac{p(n)}{\left\{q\left(\frac{n}{\log(|a_n|n^{2\alpha+1}R^n)}\right)\right\}} = \limsup_{n \rightarrow \infty} \frac{p(n)}{\left\{q\left(\frac{n}{\log(E_n(F)n^{2\alpha+1}R^n)}\right)\right\}}.$$

and

$$\limsup_{n \rightarrow \infty} \frac{p(n)}{\left\{ q \left(\frac{n}{\log(|a_n|n^{2\alpha+1}R^n)} \right) \right\}^{\rho(p,q)}} = \limsup_{n \rightarrow \infty} \frac{p(n)}{\left\{ q \left(\frac{n}{\log(E_n(F)n^{2\alpha+1}R^n)} \right) \right\}^{\rho(p,q)}}.$$

Proof. Using the identity [15, eq. 12]

$$\limsup_{n \rightarrow \infty} |a_n|^{1/n} = \limsup_{n \rightarrow \infty} [E_n(F)]^{1/n},$$

with the generalized (p, q) –order and generalized (p, q) –type calculation in Theorem 2.1 and Theorem 2.2, the proof is immediate.

Finally, we have the following theorem.

Theorem 3.1. Let F be pseudoanalytic function on $P(D_R)(R > 1)$. Then

(i) The generalized (p, q) –order $\rho(p, q)$ of F is

$$\rho(p, q) = \limsup_{n \rightarrow \infty} \frac{p(n)}{q \left(\frac{n}{\log(E_n(F)n^{2\alpha+1}R^n)} \right)}.$$

(ii) The generalized (p, q) –type of F is $T(p, q)$, if and only if,

$$T(p, q) = \limsup_{n \rightarrow \infty} \frac{p(n)}{\left\{ q \left(\frac{n}{\log(E_n(F)n^{2\alpha+1}R^n)} \right) \right\}^{\rho(p,q)}}$$

when $0 < \rho(p, q) < \infty$.

Remark 3.1. For $p(x) = q(x) = \log x$ is the classical case for order i.e., part (i) of [15, Thm. 3] and $p(x) = q(x) = x$ is the classical case for type i.e., part (ii) of [15, Thm. 3].

References

- [1] R. Askey, *Orthogonal polynomials and special functions*, Regional Conf. Series in Applied Math. 21 SIAM, Philadelphia, 1975.
- [2] K.W. Bauer and St. Ruscheweyh, *Differential Operators for Partial Differential Equations and Function Theoretic Methods*, Lecture Notes in Math. No. 791, Springer, New York, 1980.
- [3] H. Bavinck, *Jacobi Series and Approximation*, Mathematical Center Tracts No. 39, Mathematisch Centrum, Amsterdam, 1972.
- [4] S. Bergman, *Integral Operators in the Theory of Linear Partial Differential Equations*, (Berlin Springer-Verlag), 1969.
- [5] S.N. Bernstein, *Lecons sur les Proprieties Extremales et la Meilleure Approximation des Fonctions Analytiques d'une Variable Reelle*, Gauthier-Villars, Paris, 1926.

- [6] L. Bers, *Theory of Pseudoanalytic Functions*, Lecture Notes (New York: New York University), 1953.
- [7] R.P. Gilbert, *Function Theoretic Methods in Partial Differential Equations*, (New York: Academic Press), 1969.
- [8] A. Giroux, *Approximation of entire functions over bounded domains*, J. Approx. Theory, 28(1980), 45-53.
- [9] G.P. Kapoor and A. Nautiyal, *Growth and approximation of generalized bi-axially symmetric potentials*, Indian J. Pure Appl. Math., 19(5)(1988), 464-476.
- [10] D. Kumar, *Ultraspherical expansions of generalized bi-axially symmetric potentials and pseudoanalytic functions*, Complex Variables and Elliptic Equations, 53(1) (2008), 53-64.
- [11] G.R. MacLane, *Asymptotic Values of Holomorphic Functions*, Rice University Studies, Houston, 1963.
- [12] P.A. McCoy, *Polynomial approximation of generalized bi-axially symmetric potentials*, J. Approx. Theory, 225(1979), 153-168.
- [13] P.A. McCoy, *Best L^p -approximation of generalized bi-axi symmetric potentials*, Proc. Amer. Math. Soc., 779(1980), 435-44.
- [14] P.A. McCoy, *Approximation of generalized bi-axially symmetric potentials on certain domain*, J. Math. Anal. Appl., 776(1981), 463-469.
- [15] P.A. McCoy, *Approximation of pseudoanalytic functions on the unit disk*, Complex Variables and Elliptic Equations, 6(2)(1986), 123-133.
- [16] A.R. Reddy, *Best polynomial approximation of certain entire functions*, J. Approx. Theory, 5(1972), 97-112.
- [17] G. Szego, *Orthogonal Polynomials*, Amer. Math. Soc., Colloquium Publ. Vol. 23, 3rd(ed) Providence, 1967.
- [18] R.S. Varga, *On an extension of a result of S.N. Bernstein*, J. Approx. Theory, 1(1968), 176-179.
- [19] I. Vekua, *Generalized Analytic Functions*, (Reading, M.A. Addison - Wesley), 1962.
- [20] A. Weinstein, *Transonic flow and generalized axially symmetric potential theory*, Proc. NOL Aeroballistic Res. Symp. Naval Ord Laboratory, White Oak, MD, 1949.