

**A CERTAIN SUBCLASS OF MULTIVALENT ANALYTIC FUNCTIONS
DEFEND BY FRACTIONAL DIFFERINTEGRAL OPERATOR**

ABSTRACT. In this paper we introduce a new subclass of multivalent analytic functions defined by fractional differintegral operator. Such results as subordination and superordination properties, convolution properties, inequality properties and other interesting properties of this subclass are proved.

2000 *Mathematics Subject Classification.* 30C45

Key words and phrases. Analytic function, multivalent function, fractional differintegral operator, convex univalent function, Hadamard product, subordination, Superordination.

1. INTRODUCTION

Let $H(U)$ be the class of functions analytic in $U = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$ and $H[a, k]$ be the subclass of $H(U)$ consisting of functions of the form $f(z) = a + a_k z^k + a_{k+1} z^{k+1} + \dots$, with $H_0 \equiv H[0, 1]$ and $H \equiv H[1, 1]$.

Let $A_p(k)$ denote the class of functions of the form

$$f(z) = z^p + \sum_{n=k}^{\infty} a_{n+p} z^{n+p} \quad (p, k \in \mathbb{N} = \{1, 2, 3, \dots\}; z \in U), \quad (1.1)$$

which are analytic in the open unit disk U , and let $A_p(1) = A_p$ and $A_1(1) = A$.

A function $f(z) \in A_p(k)$ is said to be in the class $S_{p,k}^*(\rho)$ of multivalent (p -valent) starlike of order ρ ($0 \leq \rho < p$), if it satisfies the following inequality:

$$\Re \left\{ \frac{z f'(z)}{f(z)} \right\} > \rho \quad (0 \leq \rho < p, z \in U). \quad (1.2)$$

Let f and F be members of $H(U)$, the function $f(z)$ is said to be subordinate to $F(z)$, or $F(z)$ is said to be superordinate to $f(z)$, if there exists a function $w(z)$ analytic in U with $w(0) = 0$ and $|w(z)| < 1$ ($z \in U$), such that $f(z) = F(w(z))$. In such a case we write $f(z) \prec F(z)$. In particular, if F is univalent, then $f(z) \prec F(z)$ if and only if $f(0) = F(0)$ and $f(U) \subset F(U)$ (see [16, 24]).

For two functions $f(z)$ given by (1.1) and

$$g(z) = z^p + \sum_{n=k}^{\infty} b_{n+p} z^{n+p}, \quad (1.3)$$

The hadmard product (or convolution) of f and g is defined by

$$(f * g)(z) = z^p + \sum_{n=k}^{\infty} a_{n+p} b_{n+p} z^{n+p} = (g * f)(z). \quad (1.4)$$

We recall the definitions of the fractional derivative and integral operators introduced and studied by Saigo (cf. [16] and [18], see also [19]).

Definition 1.1 Let $\alpha > 0$ and $\beta, \gamma \in \mathbb{R}$, then the generalized fractional integral operator $I_{0,z}^{\alpha,\beta,\gamma}$ of order α of a function $f(x)$ is defined by

$$I_{0,z}^{\alpha,\beta,\gamma} f(z) = \frac{z^{-\alpha-\beta}}{\Gamma(\alpha)} \int_0^z (z-t)^{\alpha-1} {}_2F_1\left(\alpha+\beta, -\gamma; \alpha; 1-\frac{t}{z}\right) f(t) dt, \quad (1.5)$$

where the function $f(z)$ is analytic in a simply -connected region of the z -plane containing the origin and where the multiplicity of $(z-t)^{\alpha-1}$ is removed by requiring $\log(z-t)$ to be real when $(z-t) > 0$ provided further that

$$f(z) = O(|z|^\varepsilon), z \rightarrow 0 \text{ for } \varepsilon > \max(0, \beta - \gamma) - 1. \quad (1.6)$$

Definition 1.2 Let $0 \leq \alpha < 1$ and $\beta, \gamma \in \mathbb{R}$, then the generalized fractional derivative operator $J_{0,z}^{\alpha,\beta,\gamma}$ of order α of a function $f(x)$ is defined by

$$\begin{aligned} J_{0,z}^{\alpha,\beta,\gamma} f(z) &= \frac{1}{\Gamma(1-\alpha)} \frac{d}{dz} \left[z^{\alpha-\beta} \int_0^z (z-t)^{-\alpha} {}_2F_1\left(\beta-\alpha, 1-\gamma; 1-\alpha; 1-\frac{t}{z}\right) f(t) dt \right], \\ &= \frac{d^n}{dz^n} J_{0,z}^{\alpha,\beta,\gamma} f(z) \quad (n \leq \alpha < n+1; n \in \mathbb{N}). \end{aligned} \quad (1.7)$$

where the function $f(z)$ be analytic in a simply -connected region of the z -plane containing the origin with the order as given in (1.6) and the multiplicity of $(z-t)^\alpha$ is removed by requiring $\log(z-t)$ to be real when $(z-t) > 0$.

Not that

$$I_{0,z}^{\alpha,-\alpha,\gamma} f(z) = D_z^{-\alpha} f(z), \quad (\alpha > 0) \quad (1.8)$$

$$J_{0,z}^{\alpha,\alpha,\gamma} f(z) = D_z^\alpha f(z), \quad (0 \leq \alpha < 1) \quad (1.9)$$

where $D_z^{-\alpha} f(z)$ and $D_z^\alpha f(z)$ are respectively the known Riemann- Liouville fractional integral and derivative operator (cf. [13] and [14], see also [21]).

Definition 1.3 For real number $\alpha (-\infty < \alpha < 1)$ and $\beta (-\infty < \beta < 1)$ and a positive real number γ , the fractional operator $U_{0,z}^{\alpha,\beta,\gamma} : A_p \rightarrow A_p$ is defined in terms of $J_{0,z}^{\alpha,\beta,\gamma}$ and $I_{0,z}^{\alpha,\beta,\gamma}$ by (see [12] and [5])

$$U_{0,z}^{(\alpha,\beta,\gamma)} f(z) = z^p + \sum_{n=1}^{\infty} \frac{(1+p)_n (1+p+\gamma-\beta)_n}{(1+p-\beta)_n (1+p+\gamma-\alpha)_n} a_{n+p} z^{n+p}, \quad (1.10)$$

which for $f(z) \neq 0$ may be written as

$$U_{0,z}^{\alpha,\beta,\gamma} f(z) = \begin{cases} \frac{\Gamma(1+p-\beta)\Gamma(1+p+\gamma-\alpha)}{\Gamma(1+p)\Gamma(1+p+\gamma-\beta)} z^\beta J_{0,z}^{\alpha,\beta,\gamma} f(z); & 0 \leq \alpha \leq 1 \\ \frac{\Gamma(1+p-\beta)\Gamma(1+p+\gamma-\alpha)}{\Gamma(1+p)\Gamma(1+p+\gamma-\beta)} z^\beta I_{0,z}^{-\alpha,\beta,\gamma} f(z); & -\infty \leq \alpha \leq 0 \end{cases} \quad (1.11)$$

Where $J_{0,z}^{\alpha,\beta,\gamma} f(z)$ and $I_{0,z}^{-\alpha,\beta,\gamma} f(z)$ are, respectively the fractional derivative of f of order α if $0 \leq \alpha < 1$ and the fractional integral of f of order $-\alpha$ if $-\infty \leq \alpha < 0$.

It is easily verified (see Choi [4]) from (1.10) that

$$(p - \beta)U_{0,z}^{\alpha+1,\beta+1,\gamma+1} f(z) + \beta U_{0,z}^{\alpha+1,\beta+1,\gamma+1} f(z) = z \left(U_{0,z}^{\alpha,\beta,\gamma} f(z) \right)' \quad (1.12)$$

Note that

$$U_{0,z}^{\alpha,\beta,\gamma} f(z) = \Omega_z^{(\alpha,p)} f(z) \quad (-\infty < \alpha < 1), \quad (1.13)$$

The fractional differintegral operator $\Omega_z^{(\alpha,p)} f(z)$ and $(-\infty < \alpha < p+1)$ is studied Patel and Mishra [15], and the fractional differential operator $\Omega_z^{(\alpha,p)}$ with $0 < \alpha < 1$ was investigated by Srivastava and Aouf [22]. We, further observe that $\Omega_z^{(\alpha,1)} = \Omega_z^\alpha$ is the operator introduced and studied by Owa and Srivastava [14].

It is interesting to observe that

$$U_{0,z}^{0,0,\gamma} f(z) = f(z) \quad (1.14)$$

$$U_{0,z}^{1,1,\gamma} f(z) = \frac{z}{p} f'(z) \quad (1.15)$$

By making use of the differintegral operator $\Omega_z^{(\alpha,p)}$ and the above mentioned principle of subordination between analytic functions, we introduce and investigate the following subclass of the class of $A_p(k)$ of p -valent analytic functions.

Definition 1.4 A function $f(z) \in A_p(k)$ is said to be in the class $S_{p,k}^{\lambda,\mu}(\alpha; A, B)$ if it satisfies the following subordination condition:

$$(1 - \lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec \frac{1 + Az}{1 + Bz}, \quad (1.16)$$

$$(-\infty < \alpha < p; -1 \leq B < A \leq 1; A \neq B; A \in \mathbb{R}; p, k \in \mathbb{N}; \lambda \in \mathbb{R} \text{ and } \operatorname{Re}(\mu) > 0).$$

It may be noted that for suitable choice of μ, A, B, p, λ and α the class $S_{p,k}^{\lambda,\mu}(\alpha; A, B)$ extends several classes of analytic and p -valent functions studied by several authors such as Aouf and Seoudy [3], D.-G. Yang [24], Z.-Z. Zhou and S. Owa [25] and J.-L. Liu [6].

To prove our results, we need the following definitions and lemmas.

Definition 1.5 ([10]). Denote by $\mathcal{Q}$ the set of all functions $f(z)$ that are analytic and injective on $\bar{U} \setminus E(f)$ where

$$E(f) = \{ \zeta \in \partial U : \lim_{z \rightarrow \zeta} f(z) = \infty \},$$

and are such that $f'(\zeta) \neq 0$ for $\zeta \in \partial U \setminus E(f)$.

Lemma 1.1 ([11]). Let the function $h(z)$ be analytic and convex (univalent) in U with $h(0) = 1$. Suppose also that the function $g(z)$ given by

$$g(z) = 1 + c_k z^k + c_{k+1} z^{k+1} + \dots \quad (1.17)$$

is analytic in U . If

$$g(z) + \frac{zg'(z)}{\gamma} \prec h(z) \quad (\Re(\gamma) > 0; \gamma \neq 0; z \in U), \quad (1.18)$$

then

$$g(z) \prec q(z) = \frac{\gamma}{k} z^{\frac{\gamma}{k}} \int_1^{\frac{\gamma}{k}} h(t) t^{\frac{\gamma}{k}} dt \prec h(t),$$

and $q(z)$ is the best dominant of (1.18).

Lemma 1.2 ([20]). Let $q(z)$ be a convex univalent function in U and let $\alpha \in \mathbb{C}$, $\eta \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}$ with

$$\Re \left\{ 1 + \frac{z q''(z)}{q'(z)} \right\} > \max \left\{ 0, -\Re \left(\frac{\sigma}{\eta} \right) \right\}.$$

If the function $g(z)$ is analytic in U and

$$\sigma g(z) + \eta z g'(z) \prec \sigma q'(z) + \eta z q'(z),$$

then $g(z) \prec q(z)$ and $q(z)$ is the best dominant.

Lemma 1.3 ([11]). Let $q(z)$ be convex univalent function in U and let $k \in \mathbb{C}$. Further assume $\Re(k) > 0$. If $g(z) \in H[0, 1] \cap \mathcal{Q}$, and $g(z) + kzg'(z)$ is univalent in U , then

$$q(z) + kzq'(z) \prec g(z) + kzg'(z),$$

implies $g(z) \prec q(z)$ and $q(z)$ is the best subordinate.

Lemma 1.4 ([21]). Let the function F be analytic and convex in U . If $f, g \in A$ and $f, g \prec F$, then $\lambda f + (1-\lambda)g \prec F$ ($0 \leq \lambda \leq 1$).

Lemma 1.5 ([17]). Let $f(z) = 1 + \sum_{k=1}^{\infty} a_k z^k$, be analytic in U and $g(z) = 1 + \sum_{k=1}^{\infty} b_k z^k$ be analytic and convex in U . If $f(z) \prec g(z)$, then

$$|a_k| < |b_k| \quad (k \in \mathbb{N}).$$

Lemma 1.6 ([8]). Let $0 \neq \delta \in \mathbb{C}$, $\frac{\rho}{\delta} > 0$, $0 \leq \rho < 1$, $g(z) \in H[1, k]$ and

$$g(z) \prec +Lz \quad \left(L = \frac{\nu M}{k\delta + \nu} \right),$$

where

$$M = M_k(\delta, \nu, \rho) = \frac{(1-\rho)|\delta| \left(1 + \frac{k\delta}{\nu} \right)}{|1-\delta + \rho\delta| + \sqrt{1 + \left(1 + \frac{k\delta}{\nu} \right)^2}}.$$

If $h(z) \in H[1, k]$ satisfies the following subordination condition;

$$g(z)[1-\delta + \delta(1-\rho)h(z) + \rho] \prec 1 + Mz,$$

then

$$\Re(h(z)) > 0 \quad (z \in U).$$

2. MAIN RESULT

Theorem 2.1 Let $f(z) \in S_{p,k}^{\lambda,\mu}(\alpha; A, B)$ with $\Re(\lambda) > 0$. Then

$$\left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec q(z) = \frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1+Az u}{1+Bzu} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \prec \frac{1+Az}{1+Bz}, \quad (2.1)$$

and $q(z)$ is the best dominant.

Proof. Define the function $g(z)$ by

$$g(z) = \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \quad (z \in U). \quad (2.2)$$

Then $g(z)$ is of the form (1.17) and analytic in U . Differentiating (2.1) with respect to z and using (1.12), we get

$$(1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu = g(z) + \frac{\lambda z g'(z)}{(p-\beta)\mu} \prec \frac{1+Az}{1+Bz}. \quad (2.3)$$

Applying Lemma 1.1 to (2.3) with $\gamma = \frac{(p-\beta)\mu}{\lambda}$, we get

$$\begin{aligned} \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec q(z) &= \frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1+At}{1+Bt} t^{\frac{(p-\beta)\mu}{\lambda k}-1} dt \\ &= \frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1+Az u}{1+Bzu} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \prec \frac{1+Az}{1+Bz}, \end{aligned} \quad (2.4)$$

and $q(z)$ is the best dominant.

Theorem 2.2 Let $q(z)$ be univalent function in U and let $\lambda \in \mathbb{C}^*$. Suppose also that $q(z)$ satisfies the following inequality:

$$\Re \left\{ 1 + \frac{z q''(z)}{q'(z)} \right\} > \max \left\{ 0, -\Re \left(\frac{(p-\beta)\mu}{\lambda} \right) \right\}. \quad (2.5)$$

If $f \in A_p$ satisfies the following subordination:

$$(1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec q(z) + \frac{\lambda z q'(z)}{(p-\beta)\mu}, \quad (2.6)$$

then

$$\left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec q(z),$$

and $q(z)$ is the best dominant.

Proof. Let the function $g(z)$ be defined by (2.2). we know that (2.3) holds true.

Combining (2.3) and (2.6), we find that

$$g(z) + \frac{\lambda z g'(z)}{(p-\beta)\mu} \prec q(z) + \frac{\lambda z q'(z)}{(p-\beta)\mu}. \quad (2.7)$$

By using Lemma 1.2 and (2.7), we easily get the assertion of Theorem 2.2.

Taking $q(z) = \frac{1+Az}{1+Bz}$ in Theorem 2.2, we get the following result.

Corollary 2.1 Let $\lambda \in \mathbb{C}^*$ and $-1 \leq B < A \leq 1$. Suppose also that

$$\Re \left\{ \frac{1-Bz}{1+Bz} \right\} > \max \left\{ 0, -\Re \left(\frac{(p-\beta)\mu}{\lambda} \right) \right\}.$$

If $f \in A_p$ satisfies the following subordination:

$$(1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec \frac{1+Az}{1+Bz} + \frac{\lambda}{(p-\beta)\mu} \frac{(A-B)z}{(1+Bz)^2},$$

then

$$\left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec \frac{1+Az}{1+Bz},$$

and $\frac{1+Az}{1+Bz}$ is the best dominant.

Theorem 2.3 Let $q(z)$ be convex univalent function in U and let $\lambda \in \mathbb{C}$ with $\Re(\lambda) > 0$. Also let

$$\left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \in H[q(0), 1] \cap \mathcal{Q},$$

and

$$(1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu$$

be univalent in U . If

$$q(z) + \frac{\lambda z q'(z)}{(p-\beta)\mu} \prec (1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu,$$

then

$$q(z) \prec \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu,$$

and $q(z)$ is the best subordinate.

Proof. Let the function $g(z)$ be defined by (2.2). Then

$$\begin{aligned} q(z) + \frac{\lambda z q'(z)}{(p-\alpha)\mu} &\prec (1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \\ &= g(z) + \frac{\lambda z g'(z)}{(p-\beta)\mu}. \end{aligned}$$

By using Lemma 1.3 we easily get the assertion of theorem 2.3.

Taking $q(z) = \frac{1+Az}{1+Bz}$ in Theorem 2.3, we get the following result.

Corollary 2.2 Let $q(z)$ be convex univalent function in U and $-1 \leq B < A \leq 1$, $\lambda \in \mathbb{C}$ with $\Re(\lambda) > 0$. Also let

$$0 \neq \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \in H[q(0), 1] \cap \mathcal{Q},$$

and

$$(1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu$$

be univalent in U . If

$$\frac{1+Az}{1+Bz} + \frac{\lambda}{(p-\beta)\mu} \frac{(A-B)z}{(1+Bz)^2} \prec (1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu,$$

then

$$\frac{1+Az}{1+Bz} \prec \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu,$$

and $\frac{1+Az}{1+Bz}$ is the best subordinate.

Combining the above results of subordination and superordination, we easily get the following "sandwich-type result".

Corollary 2.3 Let $q_1(z)$ be convex function in U and let $q_2(z)$ be univalent function in U , let $\lambda \in \mathbb{C}$ with $\Re(\lambda) > 0$. let $q_2(z)$ satisfy (2.5). If

$$0 \neq \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \in H[q(0), 1] \cap \mathcal{Q},$$

and

$$(1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu$$

is univalent in U , and also

$$\begin{aligned} q_1(z) + \frac{\lambda z q_1'(z)}{(p-\beta)\mu} &\prec (1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \\ &\prec q_2(z) + \frac{\lambda z q_2'(z)}{(p-\beta)\mu}, \end{aligned}$$

then

$$q_1(z) \prec \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec q_2(z),$$

and $q_1(z)$ and $q_2(z)$ are respectively, the best subordinate and dominant.

Theorem 2.4 If $\lambda, \mu > 0$ and $f(z) \in S_{p,k}^{0,\mu}(\alpha; 1-2\rho, -1)$ ($0 \leq \rho < 1$), then $f(z) \in S_{p,k}^{\lambda,\mu}(\alpha; 1-2\rho, -1)$ for $|z| < R$, where

$$R = \left(\sqrt{\left(\frac{\lambda k}{(p-\alpha)\mu} \right)^2 + 1} - \frac{\lambda k}{(p-\alpha)\mu} \right)^{\frac{1}{k}}. \quad (2.8)$$

The bound R is the best possible.

Proof. We begin by writing

$$\left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{z^p} \right)^\mu = \rho + (1-\rho)g(z) \quad (z \in U; 0 \leq \rho < 1). \quad (2.9)$$

Then, clearly, the function $g(z)$ is of the form (1.17), is analytic and has a positive real part in U . Differentiating (2.9) with respect to z and using the identity (1.12), we get

$$\begin{aligned} \frac{1}{1-\rho} \left\{ (1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu - \rho \right\} \\ = g(z) + \frac{\lambda z g'(z)}{(p-\beta)\mu}. \end{aligned} \quad (2.10)$$

By making use of the following well-known estimate (see [9]):

$$\frac{|z g'(z)|}{\Re(g(z))} \leq \frac{2kr^k}{1-r^{2k}} \quad (|z| < r < 1)$$

In (2.10), we obtain that

$$\begin{aligned} \Re \left\{ \frac{1}{1-\rho} \left\{ (1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu - \rho \right\} \right\} \\ \geq \Re \{ g(z) \} \left\{ 1 - \frac{2kr^k \lambda}{(p-\beta)\mu(1-r^{2k})} \right\}. \end{aligned} \quad (2.11)$$

It is seen that the right-hand side of (2.11) is positive, provided that $r < R$, where R is given by (2.8).

In order to show that the bound R is the best possible, we consider the function

$f(z) \in A_p(k)$ defined by

$$\left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu = \rho + (1-\rho) \frac{1+z^k}{1-z^k} \quad (z \in U; 0 \leq \rho < 1).$$

Noting that

$$\begin{aligned} \frac{1}{1-\rho} \left\{ (1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu - \rho \right\} \\ = \frac{1+z^k}{1-z^k} + \frac{2k \lambda z^k}{(p-\beta)\mu(1+z^k)^2} = 0. \end{aligned} \quad (2.12)$$

for $|z| < R$, we conclude that the bound is the best possible. Theorem 2.4 is thus proved.

Theorem 2.5 Let $f(z) \in S_{p,k}^{\lambda,\mu}(\alpha; A, B)$ with $\Re(\lambda) > 0$. Then

$$f(z) = \left(z^p \left(\frac{1+Aw(z)}{1+Bw(z)} \right)^{\frac{1}{\mu}} \right) * \left(z^p + \sum_{n=1}^{\infty} \frac{(1+p-\beta)_n (1+p+\gamma-\alpha)_n}{(1+p)_n (1+p+\gamma-\beta)_n} z^{n+p} \right), \quad (2.13)$$

where $w(z)$ is an analytic function with $w(0) = 0$ and $|w(z)| < 1$ ($z \in U$).

Proof. Let $f(z) \in S_{p,k}^{\lambda,\mu}(\alpha; A, B)$ with $\Re(\lambda) > 0$. It follows from (2.1) that

$$\left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^{\mu} = \frac{1+Aw(z)}{1+Bw(z)}, \quad (2.14)$$

where $w(z)$ is an analytic function with $w(0) = 0$ and $|w(z)| < 1$ ($z \in U$). By virtue of (2.14), we easily find that

$$U_{0,z}^{(\alpha,\beta,\gamma)} f(z) = z^p \left(\frac{1+Aw(z)}{1+Bw(z)} \right)^{\frac{1}{\mu}}. \quad (2.15)$$

Combining (1.9) and (2.15), we have

$$\left(z^p + \sum_{n=1}^{\infty} \frac{(1+p)_n (1+p+\gamma-\beta)_n}{(1+p-\beta)_n (1+p+\gamma-\alpha)_n} z^{n+p} \right) * f(z) = \left(z^p \left(\frac{1+Aw(z)}{1+Bw(z)} \right)^{\frac{1}{\mu}} \right) \quad (2.16)$$

The assertion (2.13) of Theorem 2.5 can now easily be derived from (2.16).

Theorem 2.6 Let $f(z) \in S_{p,k}^{\lambda,\mu}(\alpha; A, B)$ with $\Re(\lambda) > 0$. Then

$$\begin{aligned} \frac{1}{z^p} \left[\left(1+Be^{i\theta} \right)^{\frac{1}{\mu}} \left(z^p + \sum_{n=1}^{\infty} \frac{(1+p)_n (1+p+\gamma-\beta)_n}{(1+p-\beta)_n (1+p+\gamma-\alpha)_n} z^{n+p} \right) * f(z) \right. \\ \left. - z^p \left(1+Ae^{i\theta} \right)^{\frac{1}{\mu}} \right] \neq 0 \quad (z \in U; 0 < \theta < 2\pi). \end{aligned} \quad (2.17)$$

Proof. Let $f(z) \in S_{p,k}^{\lambda,\mu}(\alpha; A, B)$ with $\Re(\lambda) > 0$. We know that (2.1) holds true, which implies that

$$\left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^{\mu} \neq \frac{1+Ae^{i\theta}}{1+Be^{i\theta}} \quad (z \in U; 0 < \theta < 2\pi). \quad (2.18)$$

It is easy to see that the condition (2.18) can be written as follows:

$$\frac{1}{z^p} \left[U_{0,z}^{(\alpha,\beta,\gamma)} f(z) \left(1+Be^{i\theta} \right)^{\frac{1}{\mu}} - z^p \left(1+Ae^{i\theta} \right)^{\frac{1}{\mu}} \right] \neq 0 \quad (z \in U; 0 < \theta < 2\pi). \quad (2.19)$$

Combining (1.10) and (2.19), we easily get the convolution property (2.17) asserted by Theorem 2.6.

Theorem 2.7 Let $\lambda_2 \geq \lambda_1 \geq 0$ and $-1 \leq B_1 \leq B_2 < A_2 \leq A_1 \leq 1$. then

$$S_{p,k}^{\lambda_2,\mu}(\alpha; A_2, B_2) \subset S_{p,k}^{\lambda_1,\mu}(\alpha; A_1, B_1). \quad (2.20)$$

Proof. Let $f(z) \in S_{p,k}^{\lambda_2,\mu}(\alpha; A_2, B_2)$. Then

$$(1-\lambda_2) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda_2 \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec \frac{1+A_2 z}{1+B_2 z}.$$

Since $-1 \leq B_1 \leq B_2 < A_2 \leq A_1 \leq 1$, we easily find that

$$(1-\lambda_2) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda_2 \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec \frac{1+A_2 z}{1+B_2 z} \prec \frac{1+A_1 z}{1+B_1 z}, \quad (2.21)$$

that is $f(z) \in S_{p,k}^{\lambda_2, \mu}(\alpha; A_1, B_1)$. Thus the assertion (2.20) holds for $\lambda_2 = \lambda_1 \geq 0$. If

$\lambda_2 \geq \lambda_1 \geq 0$, by Theorem 2.1 and (2.21), we know that $f(z) \in S_{p,k}^{\lambda_0, \mu}(\alpha; A_1, B_1)$, that is

$$\left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec \frac{1+A_1 z}{1+B_1 z}, \quad (2.22)$$

At the same time, we have

$$\begin{aligned} & (1-\lambda_1) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda_1 \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \\ &= \left(1 - \frac{\lambda_1}{\lambda_2} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \frac{\lambda_1}{\lambda_2} \left[(1-\lambda_2) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda_2 \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \right]. \end{aligned} \quad (2.23)$$

Moreover, $0 \leq \frac{\lambda_1}{\lambda_2} < 1$, and the function $\frac{1+A_1 z}{1+B_1 z}$ ($-1 \leq B_1 < A_1 \leq 1; z \in U$) is analytic and

convex in U . Combining (2.21)-(2.23) using Lemma 1.4, we find that

$$(1-\lambda_1) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda_1 \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec \frac{1+A_1 z}{1+B_1 z},$$

that is $f(z) \in S_{p,k}^{\lambda_1, \mu}(\alpha; A_1, B_1)$, which implies that the assertion (2.20) of Theorem 2.7 holds.

Theorem 2.8 Let $f(z) \in S_{p,k}^{\lambda, \mu}(\alpha; A, B)$ with $\Re(\lambda) > 0$ and $-1 \leq B_1 < A_1 \leq 1$. Then

$$\begin{aligned} & \frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1-Au}{1-Bu} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \\ & \prec \Re \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu < \frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1+Au}{1+Bu} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du. \end{aligned} \quad (2.24)$$

The extremal function of (2.24) is defined by

$$U_{0,z}^{(\alpha,\beta,\gamma)} f(z) = z^p \left(\frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1+Az u}{1+Bz u} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \right)^{\frac{1}{\mu}}. \quad (2.25)$$

Proof. Let $f(z) \in S_{p,k}^{\lambda, \mu}(\alpha; A, B)$ with $\Re(\lambda) > 0$. From Theorem 2.1 we know that (2.1) holds true, which implies that

$$\Re \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu < \sup_{z \in U} \Re \left\{ \frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1+Az u}{1+Bz u} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \right\}$$

$$\begin{aligned}
&\leq \frac{(p-\beta)\mu}{\lambda k} \int_0^1 \sup_{z \in U} \Re \left(\frac{1+Az u}{1+Bz u} \right) u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \\
&\leq \frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1+Au}{1+Bu} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du,
\end{aligned} \tag{2.26}$$

and

$$\begin{aligned}
\Re \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right) &> \inf_{z \in U} \Re \left\{ \frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1+Az u}{1+Bz u} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \right\} \\
&\geq \frac{(p-\beta)\mu}{\lambda k} \int_0^1 \inf_{z \in U} \Re \left(\frac{1+Az u}{1+Bz u} \right) u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \\
&\geq \frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1+Au}{1+Bu} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du,
\end{aligned} \tag{2.27}$$

Combining (2.26) and (2.27), we get (2.24). By noting that the function $U_{0,z}^{(\alpha,\beta,\gamma)} f(z)$ defined by (2.25) belongs to the class $S_{p,k}^{\lambda,\mu}(\alpha; A, B)$, we obtain that equality (2.24) is sharp. The proof of Theorem 2.8 is evidently completed.

In view of Theorem 2.8, we easily derive the following distortion theorems for the class $S_{p,k}^{\lambda,\mu}(\alpha; A, B)$.

Corollary 2.4 Let $f(z) \in S_{p,k}^{\lambda,\mu}(\alpha; A, B)$ with $\Re(\lambda) > 0$ and $-1 \leq B_1 < A_1 \leq 1$. Then for $|z| = r < 1$, we have

$$\begin{aligned}
&r^p \left(\frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1-Aur}{1-Bur} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \right)^{\frac{1}{\mu}} \\
&< |U_{0,z}^{(\alpha,\beta,\gamma)} f(z)| < r^p \left(\frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1+Aur}{1+Bur} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \right)^{\frac{1}{\mu}}.
\end{aligned} \tag{2.28}$$

The extremal function of (2.28) is defined by (2.25).

By noting that

$$\left(\Re(v) \right)^{\frac{1}{2}} \leq \Re \left(v^{\frac{1}{2}} \right) \leq \left| v^{\frac{1}{2}} \right| \quad (v \in \square; \Re(v) \geq 0).$$

From Theorem 2.8, we easily get the following results.

Corollary 2.5 Let $f(z) \in S_{p,k}^{\lambda,\mu}(\alpha; A, B)$ with $\Re(\lambda) > 0$ and $-1 \leq B_1 < A_1 \leq 1$. Then

$$\begin{aligned}
&\left(\frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1-Au}{1-Bu} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \right)^{\frac{1}{2}} \\
&< \Re \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^{\frac{\mu}{2}} < \left(\frac{(p-\beta)\mu}{\lambda k} \int_0^1 \frac{1+Au}{1+Bu} u^{\frac{(p-\beta)\mu}{\lambda k}-1} du \right)^{\frac{1}{2}}.
\end{aligned}$$

Theorem 2.9 Let $f(z)$ defined by (1.1) be in the class $S_{p,k}^{\lambda,\mu}(\alpha; A, B)$, Then

$$|a_{n+p}| \leq \frac{(1+p-\beta)_k (1+p+\gamma-\alpha)_k}{(1+p)_k (1+p+\gamma-\beta)_k} \left| \frac{A-B}{\lambda k + \mu(p-\beta)} \right| \quad (2.29)$$

The inequality (2.29) is sharp, with the extremal function defined by (2.25).

Proof. Combining (1.1) and (1.16), we obtain

$$\begin{aligned} (1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \\ = 1 + \left[\lambda k + \mu(p-\beta) \right] \frac{(1+p+\gamma-\beta)_k (1+p)_k}{(1+p-\beta)_k (1+p+\gamma-\alpha)_k} a_{p+k} z^k + \dots \\ \prec \frac{1+Az}{1+Bz} = 1 + (A-B)z + \dots \end{aligned} \quad (2.30)$$

An application of Lemma 1.5 to (2.30) yields

$$\frac{(1+p+\gamma-\beta)_k (1+p)_k}{(1+p-\beta)_k (1+p+\gamma-\alpha)_k} \left| \left[\lambda k + \mu(p-\beta) \right] a_{n+p} \right| \leq |A-B|. \quad (2.31)$$

Thus, from (2.31), we easily arrive at (2.29) asserted by Theorem 2.9.

Theorem 2.10 Let $0 \neq \lambda \in \mathbb{C}$, $\mu \in \mathbb{R}$, $\alpha > 1$, $\frac{\mu}{\lambda} > 0$ and $0 \leq \rho < 1$. If $f(z) \in S_{p,k}^{\lambda,\mu}(\alpha; A, 0)$ with

$$A = \frac{(1-\rho) \left| \lambda \right| \left(1 + \frac{k\lambda}{(p-\beta)\mu} \right)}{|1-\lambda+\rho\lambda| + \sqrt{1 + \left(1 + \frac{k\lambda}{(p-\beta)\mu} \right)^2}},$$

then

$$\Omega_z^{(\alpha,p)} f(z) \in S_{p,k}^* (p\rho - (p-\beta)(1-\rho)).$$

Proof. Suppose that $f(z) \in S_{p,k}^{\lambda,\mu}(\alpha; A, 0)$. By (1.16), we have

$$(1-\lambda) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu + \lambda \left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} \right) \left(\frac{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)}{z^p} \right)^\mu \prec 1 + Az. \quad (2.32)$$

Let the function $g(z)$ be defined by (2.2). We then find from (2.1) and (2.32) that

$$\begin{aligned} g(z) &\prec \frac{(p-\beta)\mu}{\lambda k} z^{\frac{(p-\beta)\mu}{\lambda k}} \int_0^z (1+At) t^{\frac{(p-\beta)\mu}{\lambda k}-1} dt \\ &= 1 + \frac{(p-\beta)\mu}{\lambda k + (p-\beta)\mu} z. \end{aligned}$$

We now suppose that

$$\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)} f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)} f(z)} = (1-\rho)h(z) + \rho \quad (\alpha > 1; 0 \leq \rho < 1; z \in U). \quad (2.33)$$

Then $h \in H[1, k]$. It follows from (2.32) and (2.33) that

$$g(z) \prec \left\{ (1-\lambda) + \lambda[(1-\rho)h(z) + \rho] \right\} \prec 1 + Az \quad (z \in U). \quad (2.34)$$

An application of Lemma 1.6 to (2.34) yields

$$\Re(h(z)) > 0 \quad (z \in U). \quad (2.35)$$

Combining (2.33) and (2.35), we find that

$$\Re\left(\frac{U_{0,z}^{(\alpha+1,\beta+1,\gamma+1)}f(z)}{U_{0,z}^{(\alpha,\beta,\gamma)}f(z)}\right) = (1-\rho)\Re(h(z)) + \rho > \rho \quad (\alpha > 1; 0 \leq \rho < 1; z \in U). \quad (2.36)$$

The assertion of Theorem 2.10 can now easily be derived from (1.12) and (2.36).

References

- [1] R. Aghalary, R. M. Ali, S. B. Joshi and V. Ravichandran, Inequalities for analytic functions defined by certain liner operator, *Internat. J. Math. Sci.* 4 (2005), 267-274.
- [2] R. M. Ali, V. Ravichandran, and N. Seenivasagan, Differential subordination and superordination of analytic functions defined by the multiplier transformation, *Math. Inequal. Appl.* 12 (2009), 123-139.
- [3] M. K. Aouf and T. M. Seoudy, Some properties of a certain subclass of multivalent analytic functions involving the Liu - Owa operator, *Comput. Math. Appl.* 60 (2010) 1525-1535.
- [4] J. M. Choi, On differential subordinations of multivalent functions involving a certain fractional derivative operator. *Int. J. Math. Sci.* doi:10.1155/2010/952036.
- [5] S. M. Khainar and M. More, A subclass of uniformly convex functions associated with certain fractional calculus operator, *IAENG International Journal of Applied Math.* 39(3)(2009. IJAM-39-07.
- [6] J.-L. Liu, On subordination for certain subclass of analytic functions, *Internat. J. Math. Math. Sci.* 20 (1997) 225-228.
- [7] M.-S. Liu, On certain subclass of analytic functions, *J. South China Normal Univ.* 4 (2002) 15-20 (in Chinese).
- [8] M.-S. Liu, On the starlikeness for certain subclass of analytic functions and a problem of Ruscheweyh, *Adv. Math. (China)* 34 (2005) 416-424.
- [9] T. H. Macgregor, The radius of univalence of certain analytic functions, *Proc. Amer. Math. Soc.* 14 (1963) 514-520.
- [10] S. S. Miller, P. T. Mocanu, Differential Subordinations: Theory and Applications, in: Monographs and Textbooks in Pure and Applied Mathematics, vol. 225, *Marcel Dekker, New York, Basel*, 2000
- [11] S. S. Miller, P. T. Mocanu, Subordinants of differential superordinations, *Complex Var. Theory Appl.* 48 (10) (2003) 815-826.
- [12] G. Murugusundaramoorthy, T. Rosy and M. Darus, A subclass of uniformly convex functions associated with certain fractional operators, *Appl. Math.* 6(3)(2005) Article 86, [online: <http://jipam.vu.edu.au>].
- [13] S. Owa, On the distortion theorems I, *Kyungpook Math. J.* 18 (1978) 53-59.
- [14] S. Owa and H. M. Srivastava, Univalent and starlike generalized hypergeometric function, *Canad. J. Math.* 39 (1987) 1057-1077.
- [15] J. Patel and A. K. Mishra, On certain subclasses of multivalent functions associated with an extended fractional differintegral operator, *J. Math. Anal. Appl.* 332 (2007) 109-122.
- [16] D.A. Patil and N. K. Thakare, On convex hulls and extreme points of p-valent starlike and convex classes with applications, *Bull. Math. Soc. R. S. Roumania*, 27(75) (1983), 145-160.
- [17] W. Rogosinski, On the coefficients of subordinate functions, *Proc. Lond. Math. Soc., Ser. 2* 48 (1943) 48-82.
- [18] M. Saigo, A certain boundary value problem for the Euler-Darboux equation. *Math.*

- Apon.* 25(1979),377-385.
- [19] M. Saigo, A remark on integral operators involving the Gauss hypergeometric functions. *Math. Rep. Kyushu Univ.* 11(1978),135-143.
 - [20] T. N. Shanmugam, V. Ravichandran, S. Sivasubramanian, Differential sandwich theorems for subclasses of analytic functions, *Aust. J. Math. Anal. Appl.* 3 (2006) 1-11. Art. 8.
 - [21] H. M. Srivastava and S. Owa(Eds.), Univalent Functions, Fractional calculus, and Their Applications, Halsted Press(Ellis Horwood Limited, Chichester), JohnWiley and Sons, New York, 1989.
 - [22] H. M. Srivastava and M. K. Aouf, A certain fractional derivative operator and its applications to a new class of analytic and multivalent functions with negative coefficients I, *J. Math. Anal. Appl.* 171 (1992) 1-13; II, *J. Math. Anal. Appl.* 192 (1995) 637-688.
 - [23] H. M. Srivastava and A. K. Mishra, A fractional differintegral operator and its applications to a nested class of multivalent functions with negative coefficients, *Adv. Stud. Contemp. Math.* 7(2003) 203-214.
 - [24] D.-G. Yang, Properties of a class of analytic functions, *Math. Japonica* 41(1995)371-381.
 - [25] Z.-Z. Zhou and S. Owa, Convolution properties of a class of bounded analytic functions, *Bull. Austral. Math. Soc.* 45(1992) 9-23.