

On the Existence of Positive Solutions for Singular Quasilinear Elliptic Equations with Gradient Terms¹

Jing Zhang Zuodong Yang

Institute of Mathematics, School of Mathematical Sciences,
Nanjing Normal University, Jiangsu Nanjing 210023, China

Abstract

In this paper, we investigate the positive solutions of quasilinear elliptic equations of the form

$$\begin{cases} -\Delta_p u = a(\delta(x))g(u) + f(x, u) + \lambda|\nabla u|^{p-1}, & \text{in } B_R \\ u > 0, & \text{in } B_R \\ u = 0, & \text{on } \partial B_R \end{cases} \quad (1.1)$$

where $B_R(0) \subset \mathbf{R}^N$, $N \geq 2$ is an open ball centered at origin of $\mathbf{R}^N$, g is an unbounded decreasing function, $a(\delta(x))$ is positive and continuous, $\delta(x) = \text{dist}(x, \partial B_R)$, $p \geq 2$, $\lambda < 0$. We emphasis the effect of all these terms in the study of existence and nonexistence of positive solutions.

Keywords: Gradient terms; Quasilinear elliptic equation; Singular; Existence and Nonexistence

Mathematics Subject Classification 35J15, 35J75.

1 Introduction

In this paper, we are concerned with quasilinear elliptic problems in the from

$$\begin{cases} -\Delta_p u = a(\delta(x))g(u) + f(x, u) + \lambda|\nabla u|^{p-1}, & \text{in } B_R \\ u > 0, & \text{in } B_R \\ u = 0, & \text{on } \partial B_R \end{cases} \quad (1.1)$$

where $B_R \subset \mathbf{R}^N$, $N \geq 2$, $\Delta_p u = \text{div}(|\nabla u|^{p-2}\nabla u)$, where B_R is a smooth and bounded domains an open ball centered at origin of $\mathbf{R}^N$, $a((\delta(x)))$ is positive and continuous, $p \geq 2$, $\lambda < 0$. For the convenience, we denote $h(x, u, \nabla u) = a(\delta(x))g(u) + f(x, u) + \lambda|\nabla u|^{p-1}$, and $h(x, u, \nabla u)$ is nonincreasing respect to u .

We assume that $g \in C^1(0, \infty)$ is a positive decreasing function and

¹Project Supported by the National Natural Science Foundation of China(Grant No.11171092). Project Supported by the Natural Science Foundation of the Jiangsu Higher Education Institutions of China (Grant No.08KJB110005)

$$(g1) \lim_{t \rightarrow 0+} g(t) = \infty$$

The function $f : \bar{B}_R \times [0, \infty) \rightarrow [0, \infty)$ is *Hölder* continuous, nondecreasing with respect to the second variable and f fulfills the hypotheses:

$$(f1) \text{ the mapping } (0, \infty) \ni t \mapsto \frac{f(x, t)}{t^{p-1}} \text{ is nonincreasing for all } x \in \bar{B}_R;$$

$$(f2) \lim_{t \rightarrow 0+} \frac{f(x, t)}{t^{p-1}} = \infty \text{ and } \lim_{t \rightarrow \infty} \frac{f(x, t)}{t^{p-1}} = 0, \text{ uniformly for } x \in \bar{B}_R.$$

Such singular boundary value problems arise in the context of chemical heterogeneous catalysts and chemical catalyst, in the theory of heat conduction in electrically conducting materials, singular minimal surfaces, as well as in the study of non-Newtonian fluid theory([21]), non-Newtonian filtration([22]) and the turbulent flow of a gas in porous medium. In the non-Newtonian fluid theory, the quantity p is characteristic of the medium. Media with $p > 2$ are called dilatant fluids and those with $p < 2$ are called pseudoplastics. If $p = 2$, then they are Newtonian fluids.

The main features of the paper are the presence of the convection term $|\nabla u|^{p-1}$ combined with the singular weight $a : (0, \infty) \rightarrow (0, \infty)$ which is assumed to be nonincreasing and *Hölder* continuous.

Many papers have been devoted to the case $a \equiv 1$ or $\lambda = 0$ (see [1-7, 9, 10, 14-17, 19, 20] and the references therein). In [1], the author considered the following problem

$$\operatorname{div}(|\nabla u|^{p-2} \nabla u) + q(x)u^{-\gamma} = 0, \quad x \in \mathbf{R}^N, \quad (1.2)$$

has a positive entire solution if $1 < p < N$, $0 \leq \gamma < p - 1$, and $q(x) \in C(\mathbf{R}_+)$ satisfy some suitable conditions.

In [2], the author studied the existence of the positive solutions of the equation

$$\begin{cases} -\Delta_p u = \lambda f(x, u), & \text{in } \Omega \\ u > 0, & \text{in } \Omega \\ u = 0, & \text{on } \partial\Omega \end{cases} \quad (1.3)$$

where $\Omega \subset \mathbf{R}^N$ is a $C^{1,\omega}$ bounded domain, for some $0 < \omega < 1$, $f : \Omega \times (0, \infty) \rightarrow [0, \infty)$ is a suitable function and allowed to be singular, $\lambda > 0$.

In [10], the existence and uniqueness of positive solutions of the following quasilinear singular equations

$$\begin{cases} -\Delta_p u = \rho(x)f(u), & \text{in } \mathbf{R}^N \\ u > 0, & \text{in } \mathbf{R}^N \\ \lim_{|x| \rightarrow \infty} u = 0. \end{cases} \quad (1.4)$$

and

$$\begin{cases} -\Delta_p u = \rho(x)f(u), & \text{in } \Omega \\ u > 0, & \text{in } \Omega \\ u = 0, & \text{on } \partial\Omega \end{cases} \quad (1.5)$$

are considered.

When $p = 2$, there is a vast literature on stable solutions to the equation (1.1), we refer to [11, 18, 23-27] and references therein. In particular, Marius Ghergu and Taliaferro S.D. [11, 23] dealing with singular weights in connection with singular nonlinearities. The present paper is an part extension of [11, 23] to the p -Laplacian equation. Our main ideas come from the paper [11, 23].

Theorem 1. Assume $(f1)$, $(f2)$, and for some $R > 0$, then the problem (1.1) has at least one solution for all $\lambda < 0$.

2 Preliminary

Definition 2.1. A function $\underline{u} \in C^{1+\alpha}(\Omega) \cap C(\overline{\Omega})$ is called a subsolution of problem (2.3) if

$$-\operatorname{div}(|\nabla \underline{u}|^{p-2} \nabla \underline{u}) \leq h(x, \underline{u}, \nabla \underline{u}), \quad \underline{u} > 0, x \in \Omega, \underline{u} = 0, x \in \partial\Omega.$$

A function $\overline{u} \in C^{1+\alpha}(\Omega) \cap C(\overline{\Omega})$ is called a supersolution of problem (2.3) if

$$-\operatorname{div}(|\nabla \overline{u}|^{p-2} \nabla \overline{u}) \geq h(x, \overline{u}, \nabla \overline{u}), \quad \overline{u} > 0, x \in \Omega, \overline{u} = 0, x \in \partial\Omega.$$

Lemma 2.1. ([8, Theorem 9.5.])

$$Q(u, \phi) = \int_{\Omega} (A(x, u, \nabla u)) \cdot \nabla \phi - B(x, u, \nabla u) \phi dx.$$

for all non-negative $\phi \in C_0^1(\Omega)$. Let $u, v \in C^1(\overline{\Omega})$ satisfy $Qu \geq 0$ in Ω , $Qv \leq 0$ in Ω and $u \leq v$ on $\partial\Omega$, where the functions A, B are continuously differentiable with respect to the z, p variables in $\overline{\Omega} \times \mathbf{R} \times \mathbf{R}^N$, the operator Q is elliptic in Ω , and the function B is non-increasing in z for fixed $(x, p) \in \Omega \times \mathbf{R}^N$. Then, if either

- (i) the vector function A is independent of z ; or
- (ii) the function B is independent of p .

It follows that $u \leq v$ in Ω .

Lemma 2.2. ([12, Lemma 2.2]) Let $h(x, u, \xi)$ satisfy the following two basic conditions:

(A) $h(x, u, \xi)$ is locally Holder continuous function in $\Omega \times \mathbf{R}^+ \times \mathbf{R}^N$ and continuously differentiable with respect to the variables u and ξ ;

(B) For every bounded domain $\Omega_1 \subset \subset \Omega$, for any $M > 0$, $\exists \rho(\Omega_1, M) > 0$, such that

$$|h(x, u, \xi)| < \rho(\Omega, M)(1 + |\xi|^p), \quad x \in \Omega_1, 0 \leq u \leq M, \xi \in \mathbf{R}^N.$$

If problem (1.1) has a supersolution $\overline{u} \in C^1(\Omega)$ and a subsolution $\underline{u} \in C^1(\Omega)$ such that $\underline{u} \leq \overline{u}$ in Ω , the problem (1.1) has at least one solution $u(x) \in C^1(\Omega)$ with $\underline{u}(x) \leq u(x) \leq \overline{u}(x)$.

3 Proof of Theorem 1.

Let us note first that in our setting problem (1.1) reads

$$\begin{cases} -\Delta_p u = a(R - |x|)g(u) + f(x, u) + \lambda|\nabla u|^{p-1}, & |x| < R \\ u > 0, & |x| < R \\ u = 0, & |x| = R. \end{cases} \quad (3.1)$$

In order to provide a supersolution of (3.1), we consider the problem

$$\begin{cases} -\Delta_p u = a(R - |x|)g(u) + 1 + \lambda|\nabla u|^{p-1}, & |x| < R \\ u > 0, & |x| < R \\ u = 0, & |x| = R. \end{cases} \quad (3.2)$$

Lemma 3.1. Assume (g1), problem (3.2) has at least one solution.

Proof. We are looking for radially decreasingly symmetric solution u of (3.2), that is $u = u(r)$, $0 \leq r = |x| \leq R$. In this case, problem (3.2) becomes

$$\begin{cases} -[(|u'|^{p-2}u')' + \frac{N-1}{r}|u'|^{p-2}u'] = a(R - |x|)g(u(r)) + 1 + \lambda|u'|^{p-1}, & |x| < R \\ u(r) > 0, & |x| < R \\ u(r) = 0, & |x| = R. \end{cases} \quad (3.3)$$

Since $u(r)$ is decreasing, that is $u'(r) \leq 0$, then (3.3) gives

$$-[(|u'|^{p-2}u')' + \frac{N-1}{r}|u'|^{p-2}u'] = a(R - |x|)g(u(r)) + 1, \quad 0 \leq r < R.$$

We obtain

$$-(e^{-\lambda r} r^{N-1} |u'|^{p-2} u')' = e^{-\lambda r} r^{N-1} \psi(r, u(r)), \quad 0 \leq r < R, \quad (3.4)$$

where

$$\psi(r, t) = a(R - |x|)g(t) + 1, \quad (r, t) \in [0, R) \times (0, \infty).$$

From (3.4) we obtain

$$u(r) = u(0) - \int_0^r [e^{\lambda t} t^{1-N} \int_0^t e^{-\lambda s} s^{N-1} \psi(s, u(s)) ds]^{\frac{1}{p-1}} dt, \quad 0 \leq r < R. \quad (3.5)$$

On the other hand, due to [6] and to the symmetry of the domain, there exists a solution $\omega = \omega(r) \in C^{1+\alpha}(B_R(0)) \cap C(\bar{B}_R(0))$ of the problem

$$\begin{cases} -\Delta_p \omega = a(R - |x|)g(\omega) + 1, & |x| < R \\ \omega > 0, & |x| < R \\ \omega = 0, & |x| = R \end{cases} \quad (3.6)$$

As above we get

$$\omega(r) = \omega(0) - \int_0^r [t^{1-N} \int_0^t s^{N-1} \psi(s, \omega(s)) ds]^{\frac{1}{p-1}} dt, \quad 0 \leq r < R. \quad (3.7)$$

We claim that there exists a solution $v \in C^{1+\alpha}(B_R(0)) \cap C(\bar{B}_R(0))$ of (3.5) such that $v > 0$ in $[0, R]$.

Let $A = \omega(0)$ and define the sequence $(v_k)_{k \geq 0}$ by

$$\begin{cases} v_k(r) = A - \int_0^r [e^{\lambda t} t^{1-N} \int_0^t e^{-\lambda s} s^{N-1} \psi(s, v_{k-1}(s)) ds]^{\frac{1}{p-1}} dt, & 0 \leq r < R \\ v_0(r) = \omega. \end{cases} \quad (3.8)$$

Since

$$\begin{aligned} v_1(r) &= A - \int_0^r [e^{\lambda t} t^{1-N} \int_0^t e^{-\lambda s} s^{N-1} \psi(s, v_0(s)) ds]^{\frac{1}{p-1}} dt \\ &\geq A - \int_0^r [e^{\lambda t} t^{1-N} e^{-\lambda t} \int_0^t s^{N-1} \psi(s, v_0(s)) ds]^{\frac{1}{p-1}} dt \\ &= v_0(r) \end{aligned}$$

we have $\omega = v_0(r) \leq v_1(r)$, then

$$\begin{aligned} v_2(r) &= A - \int_0^r [e^{\lambda t} t^{1-N} \int_0^t e^{-\lambda s} s^{N-1} \psi(s, v_1(s)) ds]^{\frac{1}{p-1}} dt \\ &\geq A - \int_0^r [e^{\lambda t} t^{1-N} \int_0^t e^{-\lambda s} s^{N-1} \psi(s, v_0(s)) ds]^{\frac{1}{p-1}} dt \\ &= v_1(r) \end{aligned}$$

As the above iteration, we reduce $v_k(r) \geq v_{k-1}(r)$ for all $k \geq 2$. Hence

$$\omega = v_0 \leq v_1 \leq v_2 \leq \dots \leq v_k \leq \dots \leq A, \quad \text{in } B_R(0).$$

Thus, there exists $\bar{v}(r) := \lim_{k \rightarrow \infty} v_k(r)$, for all $0 \leq r < R$ and $v > 0$ in $[0, R]$. We now can pass to the limit in (3.8) in order to get that v is a solution of (3.5). By classical regularity results we also obtain $\bar{v} \in C^{1+\alpha}([0, R]) \cap C([0, R])$. This proves the claim.

Clearly, $\underline{v} = \bar{v} - \bar{v}(R)$ is a subsolution of (3.2). On the other hand, we have obtained a supersolution $\bar{v}$ of (3.2) such that $\bar{v} > \underline{v}$ in $B_R(0)$. So, by the standard sub- and sup-solution method, the problem (3.2) has at least one solution and the proof of Lemma 3.1 is completed.

Proof of Theorem 1. In order to provide a subsolution of (3.1), we consider the following problem

$$\begin{cases} -\Delta_p u = a(R - |x|)g(u) + \lambda |\nabla u|^{p-1}, & \text{in } |x| < R \\ u > 0, & \text{in } |x| < R \\ u = 0, & \text{on } |x| = R \end{cases} \quad (3.9)$$

It is easy to see that the solution of (3.9) is the subsolution of (3.1), next we are looking for the solution of (3.9).

In [9], it is easy to see that there exists $\bar{u}_1 \in C^{1+\alpha}(B_R) \cap C(\bar{B}_R)$ such that

$$\begin{cases} -\Delta_p u = a(R - |x|)g(u), & \text{in } |x| < R \\ u > 0, & \text{in } |x| < R \\ u = 0, & \text{on } |x| = R \end{cases} \quad (3.10)$$

It is obvious that $\bar{u}_1$ is a subsolution of (3.9) for all $\lambda < 0$.

We can obtain easily that there exists $\underline{u}_1$ such that

$$\begin{cases} -\Delta_p u = \lambda |\nabla u|^{p-1}, & \text{in } B_R(0) \\ u > 0, & \text{in } B_R(0) \\ u = 0, & \text{on } \partial B_R(0). \end{cases} \quad (3.11)$$

It is obvious that $\underline{u}_1 > 0$ is a subsolution of (3.9) for all $\lambda < 0$. By comparison principle, we easily obtain

$$\bar{u}_1 > \underline{u}_1.$$

Then, by the standard sub- and sup-solution method, the problem (3.9) has a solution $\underline{u}$ and $\bar{u}_1 > \underline{u} > \underline{u}_1$. It follows that $\underline{u}$ is a subsolution of (3.1).

Let u be a solution of (3.2). For $M > 1$ we have

$$\begin{aligned} -\Delta_p (Mu) &= M^{p-1}a(R - |x|)g(u) + M^{p-1} + \lambda |\nabla (Mu)|^{p-1} \\ &\geq a(R - |x|)g(Mu) + M^{p-1} + \lambda |\nabla (Mu)|^{p-1}. \end{aligned}$$

Since (f_1) , we can choose $M > 1$ such that

$$M^{p-1} \geq f(x, M|u|_\infty), \text{ in } B_R(0).$$

Then $\bar{u}_\lambda := Mu$ satisfies

$$-\Delta_p (\bar{u}_\lambda) \geq a(R - |x|)g(\bar{u}_\lambda) + f(x, \bar{u}_\lambda) + \lambda |\nabla (\bar{u}_\lambda)|^{p-1}, \text{ in } B_R(0).$$

It follows that $\bar{u}_\lambda$ is a supersolution of (3.1). Since Lemma 2.1, we know $\underline{u} \leq \bar{u}_\lambda$ in $B_R(0)$. So, (3.1) has at least one solution.

The proof of Theorem 1 is completed.

References

- [1] Zuodong Yang, Existence of positive entire solutions for singular and non-singular quasi-linear elliptic equation, J.Comput.Appl.Math. 197(2006) 355-364.

- [2] Ahmed Mohammed, Positive solutions of the p -Laplacian equation with singular nonlinearity, *J.Math. Anal. Appl.* 352(2009)234-245.
- [3] Rhouma, N.B., Mosbah, M., on the existence of positive solutions for semilinear elliptic equations with indefinite nonlinearities, *J. Differ. Equ.* 215(2005) 37-51.
- [5] Shi, J., Yao, M., On a singular nonlinear semilinear elliptic problem, *Proc. R. Soc. Edinb. Sect.A, Math.* 128(2008) 1389-1401.
- [4] Yong-Hoon Lee, Inbo Sim, Existence results of sign-changing solutions for singular one -dimensional p -Laplacian problems, *Nonlinear Anal.* 68(2008)1195-1209.
- [6] Siegfried Carl, Kanishka Perera, Generalized solutions of singular p -Laplacian problems in $\mathbf{R}^N$, *Nonlinear Studies* 18(2011)113-124.
- [7] Ying Shen, Jihui Zhang, Existence of positive entire solutions of a p -Laplacian problem with a gradient term, *Differential Equations Applications* 2(2011) 225-233.
- [8] David Gilbar and Neil S Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin Heidelberg New York 1997.
- [9] Junli Yuan, Zuodong Yang, Elliptic and asymptotic behavior of radially symmetric ground states of quasi-linear singular elliptic equations, *Applied Mathematics and computation* 216(2010)213-220.
- [10] Josévaldo Goncalves, Carlos Alberto P.santos, Positive solutions for a class of quasilinear singular equations, *Electronic J. of Differential Equations* 2004(2004)1-15.
- [11] Marius Ghergu, Singular semilinear elliptic equations with subquadratic gradient terms, M.Reissig, M.Ruzhansky(eds.), *Progress in Partial Differential Equations, Springer Proceedings in Math. Statistics* 44(2013) 75-91.
- [12] Zuodong Yang, On the existence of multiple positive entire solutions for a class of quasilinear elliptic equations, *International Journal of Mathematics and Mathematical Sciences* 2006(2006), Article ID 34538, 1-19.
- [13] Ravi P. Agarwal, Haishen Lü, Donal O'Regan, Existence theorems for the one -dimensional p -Laplacian equation with sign changing nonlinearities, *Applied Mathematics and computation* 143(2003)15-18.
- [14] Daqing Jiang, Xiaojie Xu, Multiple positive solution to a class of singular boundary value problems for the one -dimensional p -Laplacian, *Computers and Math. with Applications* 47(2004)667-681.
- [15] Wenshu Zhou, Existence of positive solution for a singular p -Laplacian Dirichlet problem, *Electronic Journal Of Differential Equations* 2008(102)(2008) 1-6.
- [16] Kanishka Perera, Elves A.B. Silva, Existence and multiplicity of positive solution for singular quasilinear problems, *J. Math. Anal. Appl.* 323(2006) 1238-1252.
- [17] J.V.A. Goncalves, M.C. Rezende, C.A. Santos, Positive solution for a mixed and singular quasilinear problems, *Nonlinear Anal.* 74(2011) 132-140.
- [18] A. Porretta, S.Segura de León, Nonlinear elliptic equations having a gradient term with natural growth, *J. Math. Pures Appl.* 85(2006) 465-492.
- [19] Marcelo Montenegro, Existence and nonexistence of solutions for quasilinear elliptic equations, *Journal Mathematical Analysis and Applications* 245(2000) 303-316.
- [20] Kanishka Perera, Zhitao Zhang, Multiple positive solutions of singular p -Laplacian problems by variational methods, *Journal Mathematical Analysis and Applications* 3(2005) 377-382.

- [21] M.A. Herrero and J. Vazquez, On the propagation properties of a nonlinear degenerate parabolic equation, *Communication in Partial Differential Equations* 7(12) (1982), 1381-1402.
- [22] J. R. Esteban, J. L. Vazquez, On the equation of turbulent filtration in one-dimensional porous media, *Nonlinear Anal.* 10(1982), 1303-1325.
- [23] Taliaferro, S.D., A nonlinear singular boundary value problem, *Nonlinear Anal. TAM* 3(1979) 897-904.
- [24] Ghergu, M., Radulescu, V., Sublinear singular elliptic problems with two parameters, *J. Differ. Equ.* 195(2003) 520-536.
- [25] Shi, J., Yao, M., Positive solutions for elliptic equations with singular nonlinearity, *Electro. J. Differ. Equ.* 4(2008) 1-11.