

On Some New Nonlinear Retarded Integral Inequalities With Iterated Integrals And Their Applications In Integro-Differential Equations

A. Abdeldaim¹ and A. A. El-Deeb²

¹*Department of Mathematics and Computer Science, Faculty of Science, Port Said University, Port Said, Egypt.*

²*Department of Mathematics, Faculty of Science, Al-Azhar University, Nasr City (11884), Cairo, Egypt.*

Abstract

In this paper, some new generalized retarded nonlinear integral inequalities of Gronwall-Bellman type are discussed. The upper bounds estimation of the embedded unknown functions are discussed by integral and differential techniques. Our results generalize those of H. El-Owaidy et al. [6] to some inequalities with both retard and nonlinear integral. Some applications are also presented in order to illustrate the usefulness of some of our results.

keywords: Retarded integral inequalities, Estimation, Iterated integrals, Analysis technique, Integro-Differential equations.

1 Introduction

Integral inequalities that give explicit bounds on unknown functions provide a very useful and important device in the study of many qualitative as well as quantitative properties of solutions of nonlinear differential equations and integral equations. In 1919, Gronwall [7] introduced the famous Gronwall inequality (Theorem 1.1) while investigating the dependence of systems of differential equations with respect to a parameter. Integral inequalities of Gronwall type are important tools in the study of existence, uniqueness, boundedness, stability, invariant manifolds and other qualitative properties of solutions of differential equations. Since then, a lot of contributions have been achieved by many researchers. The original Gronwall inequality has been extended to the more general cases including the generalized linear and nonlinear Gronwall type inequalities and mixed Gronwall-Bellman inequalities (see [1,3-6,8,10-13,19]).

Theorem 1.1 (*Gronwall*[7]). Let $u(t)$ be a continuous function defined on the interval $D = [\alpha, \alpha + h]$ and

$$0 \leq u(t) \leq \int_{\alpha}^t [bu(s) + a]ds, \forall t \in D,$$

where α, a, b and h are nonnegative constants. Then

$$0 \leq u(t) \leq ahe^{bh}, \forall t \in D.$$

Recently, H. El-Owaidy et al. [6] studied the following nonlinear integral inequalities of Gronwall type

Theorem 1.2(H. El-Owaidy et al. [6]). Let $u(t)$ be a real-valued positive continuous function and $f(t), g(t)$ are real-valued non-negative continuous functions defined on $I = [0, \infty)$ and satisfy the inequality

$$u^p(t) \leq u_0 + \int_0^t f(s) \left[u^p(s) + \int_0^s g(\lambda) u(\lambda) d\lambda \right] ds, \forall t \in I,$$

where u_0 and p are positive constant. Then

$$u(t) \leq u_0^{\frac{1}{p}} \left[1 + \int_0^t f(s) \exp \left(\int_0^s [f(\lambda) + g(\lambda)] d\lambda \right) ds \right]^{\frac{1}{p}}, \quad \forall t \in I.$$

Theorem 1.3(H. El-Owaidy et al. [6]). Let $u(t)$ be a real-valued positive continuous function and $f(t), g(t)$ are real-valued nonnegative continuous functions defined on $I = [0, \infty)$ and satisfy the inequality

$$u^p(t) \leq u_0 + \int_0^t f(s) \left[u^q(s) + \int_0^s g(\lambda) u(\lambda) d\lambda \right] ds, \forall t \in I,$$

where u_0, p and q are constants such that $u_0 \geq 0$ and $p > q \geq 0$. Then

$$u(t) \leq \left[u_0 + \int_0^t f(s) K(s) \exp \left(\int_0^s g(\lambda) d\lambda \right) ds \right]^{\frac{1}{p}}, \quad \forall t \in I,$$

where

$$K(t) = \left[u_0^{\frac{q[p-q]}{p}} + \left[\frac{q[p-q]}{p} \right] \int_0^t f(s) \exp \left(-[p-q] \int_0^s g(\lambda) d\lambda \right) ds \right]^{\left[\frac{1}{p-q} \right]}, \forall t \in I.$$

Theorem 1.4 (H. El-Owaidy et al. [6]). Let $u(t)$ be a real-valued positive continuous function and $f(t), g(t)$ are nonnegative real-valued continuous functions defined on $I = [0, \infty)$, and satisfy the inequality

$$u(t) \leq u_0 + \int_0^t f(s) u(s) \left[u^q(s) + \int_0^s g(\lambda) u(\lambda) d\lambda \right]^p ds, \forall t \in I,$$

where u_0, p and q , are positive constants such that $p + q > 1$, . Then

$$u(t) \leq u_0 \exp \left(\int_0^t f(s) K_1(s) ds \right), \quad t \in I,$$

where

$$K_1(t) = \frac{u_0^{pq} \exp \left(p \int_0^t g(s) ds \right)}{\left[1 - q[p+q-1] u_0^{q[p+q-1]} \int_0^t f(s) \exp([p+q-1] \int_0^s g(\lambda) d\lambda) ds \right]^{\left[\frac{p}{p+q-1} \right]}},$$

such that, $q[p+q-1]u_0^{q[p+q-1]} \int_0^t f(s) \exp([p+q-1] \int_0^s g(\lambda)d\lambda)ds < 1, \forall t \in I$.

Theorem 1.5(H. El-Owaidy et al. [6]). Let $u(t)$ be a real-valued positive continuous function and $f(t), g(t)$ are nonnegative real-valued continuous functions defined on $I = [0, \infty)$, and $n(t)$ be a positive monotonic nondecreasing continuous function on $I = [0, \infty)$ and satisfy the inequality

$$u(t) \leq n(t) + \int_0^t f(s) \left[u(s) + \int_0^s g(\lambda)u(\lambda)d\lambda \right]^p ds, \forall t \in I,$$

where p be a constant such that $p \in (0, 1)$. Then

$$u(t) \leq n(t) \left[1 + \int_0^t f(s) K_2(s) n^{[p-1]}(s) \right], \quad \forall t \in I,$$

where

$$\begin{aligned} K_2(t) = & \exp \left(p(1-p) \int_0^t g(s)ds \right) \left[1 + (1-p) \int_0^t f(s) n^{[p-1]}(s) \right. \\ & \times \exp \left(-(1-p) \int_0^s g(\lambda)d\lambda \right) ds \left. \right]^{\left[\frac{p}{1-p} \right]}, \quad \forall t \in I. \end{aligned}$$

However, many real life problems that have in the past sometimes been modeled by initial value problems for differential equations, actually involve a significant memory effect that can be represented in a more refined model using a differential equation incorporating retarded or delayed arguments. In this situations, we need to discuss some retarded nonlinear integral inequalities (see [2,9,14-18]). So, the given bound on such inequalities in [6] are not directly applicable in the study of certain retarded nonlinear differential and integral equations. It is desirable to establish new inequalities of the above type, which can be used more effectively in the study of certain classes of retarded nonlinear differential, integral and integro-differential equations.

In this paper, we extend certain results that where proved in [6] to obtain new generalizations of formerly famous Gronwall type inequalities where non retarded case t in [6] is changed into retarded case $\alpha(t)$, and give upper bound estimation of the unknown function by integral and differential techniques in order to study retarded differential, integral and integro-differential equations.

2 Main results

In this section, several new retarded integral inequalities of Gronwall-Bellman type are introduced.

Throughout this article, $\mathbb{R}$ denoted the set of real numbers; $I = [0, \infty)$, $\mathbb{R}_+ = (0, \infty)$, $\mathbb{R}_1 = [1, \infty)$ and $'$ denotes the derivative. $\mathcal{C}(I, I)$ denotes the set of all nonnegative real-valued continuous functions from I into I and $\mathcal{C}^1(I, I)$ denotes the set of all nonnegative real-valued continuously differentiable functions from I into I .

Theorem 2.1. Let $u(t), g(t), f(t) \in \mathcal{C}(I, I), \alpha(t) \in \mathcal{C}^1(I, I)$ be nondecreasing with $\alpha(t) \leq t$ on I and satisfy the inequality

$$u^p(t) \leq u_0 + \int_0^{\alpha(t)} f(s) \left[u^p(s) + \int_0^s g(\lambda) u(\lambda) d\lambda \right] ds, \quad \forall t \in I, \quad (2.1)$$

where u_0 and p are positive constants. Then

$$u(t) \leq u_0^{\frac{1}{p}} \left[1 + \int_0^{\alpha(t)} f(s) \exp \left(\int_0^s [f(\lambda) + g(\lambda)] d\lambda \right) ds \right]^{\frac{1}{p}}, \quad \forall t \in I. \quad (2.2)$$

Proof. Let $J^p(t)$ equal the right hand side in (2.1), we have $J(0) = u_0^{\frac{1}{p}}$ and

$$u(t) \leq J(t), u(\alpha(t)) \leq J(\alpha(t)) \leq J(t), \quad \forall t \in I. \quad (2.3)$$

Differentiating $J^p(t)$ with respect to t , and using (2.3) leads to

$$pJ^{[p-1]}(t) \frac{dJ(t)}{dt} \leq \alpha'(t) f(\alpha(t)) Y(t), \quad \forall t \in I, \quad (2.4)$$

where $Y(t) = J^p(t) + \int_0^{\alpha(t)} g(s) J(s) ds$, thus we have $Y(0) = J^p(0) = u_0$ and

$$J(t) \leq Y(t), \quad \forall t \in I. \quad (2.5)$$

Differentiating $Y(t)$ with respect to t , and using (2.4), (2.5) leads to

$$\frac{dY(t)}{dt} \leq \alpha'(t) [f(t) + g(t)] Y(t), \quad \forall t \in I.$$

By taking $t = s$ in the above inequality and integrating from 0 to t , we get

$$Y(t) \leq u_0 \exp \left(\int_0^{\alpha(t)} [f(s) + g(s)] ds \right), \quad \forall t \in I, \quad (2.6)$$

from (2.6) and (2.4), we obtain

$$J^{[p-1]}(t) \frac{dJ(t)}{dt} \leq \frac{1}{p} u_0 \alpha'(t) f(\alpha(t)) \exp \left(\int_0^{\alpha(t)} [f(s) + g(s)] ds \right), \quad \forall t \in I.$$

The above inequality implies an estimation of $J(t)$ as follows

$$J(t) \leq u_0^{\frac{1}{p}} \left[1 + \int_0^{\alpha(t)} f(s) \exp \left(\int_0^s [f(\lambda) + g(\lambda)] d\lambda \right) ds \right]^{\frac{1}{p}}, \quad \forall t \in I. \quad (2.7)$$

Using (2.7) in (2.3), we get the required inequality in (2.2). The proof is completed. $\square$

Remark 2.1. If $\alpha(t) = t$, then Theorem 2.1 reduces to Theorem 1.2.

Theorem 2.2. Let $u(t), g(t), f(t) \in \mathcal{C}(I, I), \alpha(t) \in \mathcal{C}^1(I, I)$ be nondecreasing with $\alpha(t) \leq t, \alpha(0) = 0$ on I and satisfy the inequality

$$u^p(t) \leq u_0 + \int_0^{\alpha(t)} f(s) \left[u^q(s) + \int_0^s g(\lambda) u(\lambda) d\lambda \right] ds, \quad \forall t \in I, \quad (2.8)$$

where $u_0 > 0, p > q \geq 1$ are constants. Then

$$u(t) \leq \left[u_0 + \int_0^{\alpha(t)} f(s) k(\alpha^{-1}(s)) ds \right]^{\frac{1}{p}}, \quad \forall t \in I, \quad (2.9)$$

where

$$\begin{aligned} k(t) = & \exp \left(\int_0^{\alpha(t)} g(s) ds \right) \left[u_0^{\frac{q[p-q]}{p}} \right. \\ & \left. + \left[\frac{q[p-q]}{p} \right] \int_0^{\alpha(t)} f(s) \exp \left(-(p-q) \int_0^s g(\lambda) d\lambda \right) ds \right]^{\left[\frac{1}{p-q} \right]}. \end{aligned} \quad (2.10)$$

Proof. Let $J_1^p(t)$ equal the right hand side in (2.8), we have $J_1(0) = u_0^{\frac{1}{p}}$ and

$$u(t) \leq J_1(t), u(\alpha(t)) \leq J_1(\alpha(t)) \leq J_1(t), \quad \forall t \in I. \quad (2.11)$$

Differentiating $J_1^p(t)$, with respect to t , using (2.11) we get

$$p J_1^{[p-1]}(t) \frac{dJ_1(t)}{dt} \leq \alpha'(t) f(\alpha(t)) Y_1(t), \quad \forall t \in I, \quad (2.12)$$

where $Y_1(t) = J_1^q(t) + \int_0^{\alpha(t)} g(s) J_1(s) ds$, thus we have $Y_1(0) = J_1^q(0) = u_0^{\frac{q}{p}}$, and $J_1^q(t) \leq Y_1(t)$, but $q \geq 1$, thus

$$J_1(t) \leq Y_1(t), \quad \forall t \in I. \quad (2.13)$$

Differentiating $Y_1(t)$ with respect to t , and using (2.12), (2.13) leads to

$$\frac{dY_1(t)}{dt} \leq \frac{q}{p} \alpha'(t) f(\alpha(t)) Y_1^{[q-p+1]}(t) + \alpha'(t) g(\alpha(t)) Y_1(t), \quad \forall t \in I,$$

but $Y_1(t) > 0$, thus we have

$$Y_1^{[p-q-1]}(t) \frac{dY_1(t)}{dt} - \alpha'(t) g(\alpha(t)) Y_1^{[p-q]}(t) \leq \frac{q}{p} \alpha'(t) f(\alpha(t)), \quad \forall t \in I. \quad (2.14)$$

Now, if we put

$$Z(t) = Y_1^{[p-q]}(t), \quad \forall t \in I, \quad (2.15)$$

then we have $Z(0) = Y_1^{[p-q]}(0) = u_0^{\frac{q[p-q]}{p}}$ and $Y_1^{[p-q-1]}(t) \frac{dY_1(t)}{dt} = \left[\frac{1}{p-q} \right] \frac{dZ}{dt}$, thus from (2.14) we obtain

$$\frac{dZ(t)}{dt} - [p-q] \alpha'(t) g(\alpha(t)) Z(t) \leq \left[\frac{q[p-q]}{p} \right] \alpha'(t) f(\alpha(t)), \quad \forall t \in I. \quad (2.16)$$

The above inequality implies the following estimations of $Z(t)$, such that

$$Z(t) \leq k^{[p-q]}(t), \quad \forall t \in I,$$

where $k(t)$ is as defined in (2.10). From (2.8) and the above inequality we have

$$Y_1(t) \leq k(t), \quad \forall t \in I,$$

thus from (2.12) and the above inequality we have

$$pJ_1^{[p-1]}(t) \frac{dJ_1(t)}{dt} \leq \alpha'(t)f(\alpha(t))k(t), \quad \forall t \in I.$$

By taking $t = s$ in the above inequality and integrating from 0 to t , gives

$$J_1(t) \leq \left[u_0 + \int_0^{\alpha(t)} f(s)k(\alpha^{-1}(s)) \right]^{\frac{1}{p}}, \quad \forall t \in I. \quad (2.17)$$

Using (2.17) in (2.11) leads to the required inequality in (2.9). The proof is completed. $\square$

Remark 2.2. If $\alpha(t) = t$, then Theorem 2.2 reduces to Theorem 1.3.

Theorem 2.3. Let $u(t), g(t), f(t) \in \mathcal{C}(I, I), \alpha(t) \in \mathcal{C}^1(I, I)$ be nondecreasing with $\alpha(t) \leq t, \alpha(0) = 0$, on I and satisfy the inequality

$$u(t) \leq u_0 + \int_0^{\alpha(t)} f(s)u(s) \left[u^q(s) + \int_0^s g(\lambda)u(\lambda)d\lambda \right]^p ds, \quad \forall t \in I, \quad (2.18)$$

where u_0, p and q are positive constants such that $p + q > 1$. Then

$$u(t) \leq u_0 \exp \left(\int_0^{\alpha(t)} f(s)k_1(\alpha^{-1}(s))ds \right), \quad \forall t \in I, \quad (2.19)$$

where

$$k_1(t) = \frac{u_0^{pq} \exp \left(p \int_0^{\alpha(t)} g(s)ds \right)}{\left[1 - q[p+q-1]u_0^{q[p+q-1]} \int_0^{\alpha(t)} f(s) \exp([p+q-1] \int_0^s g(\lambda)d\lambda)ds \right]^{\left[\frac{p}{p+q-1} \right]}}, \quad (2.20)$$

such that, $q[p+q-1]u_0^{q[p+q-1]} \int_0^{\alpha(t)} f(s) \exp([p+q-1] \int_0^s g(\lambda)d\lambda)ds < 1, \forall t \in I$.

Proof. Let $J_2(t)$ equal the right hand side of (2.18), which is a positive and nondecreasing function on I with $J_2(0) = u_0$. Then (2.18) reduces to

$$u(t) \leq J_2(t), u(\alpha(t)) \leq J_2(\alpha(t)) \leq J_2(t), \quad \forall t \in I. \quad (2.21)$$

Differentiating $J_2^p(t)$, with respect to t , using (2.21) we have

$$\frac{dJ_2(t)}{dt} \leq \alpha'(t)f(\alpha(t))J_2(t)Y_2^p(t), \quad \forall t \in I, \quad (2.22)$$

where $Y_2(t) = J_2^q(t) + \int_0^{\alpha(t)} g(s)J_2(s)ds$, hence $Y_2(0) = J_2^q(0) = u_0^q$, and

$$J_2(t) \leq Y_2(t) \quad \forall t \in I. \quad (2.23)$$

Differentiating $Y_2(t)$ with respect to t , and using (2.22), (2.23), leads to

$$\frac{dY_2(t)}{dt} \leq q\alpha'(t)f(\alpha(t))Y_2^{[p+q]}(t) + \alpha'(t)g(\alpha(t))Y_2(t), \quad \forall t \in I,$$

but $Y_2(t) > 0$, then we have

$$Y_2^{-[p+q]}(t) \frac{dY_2(t)}{dt} - \alpha'(t)g(\alpha(t))Y_2^{[1-(p+q)]}(t) \leq q\alpha'(t)f(\alpha(t)), \quad \forall t \in I. \quad (2.24)$$

If we let

$$Y_2^{[1-(p+q)]}(t) = Z_1(t), \quad \forall t \in I, \quad (2.25)$$

we have $Z_1(0) = Y_2^{-[p+q-1]}(0) = u_0^{-q[p+q-1]}$, and $Y_2^{-[p+q]}(t) \frac{dY_2(t)}{dt} = \frac{-1}{[p+q-1]} \frac{dZ_1}{dt}$, then we can write the inequality (2.24) as follows

$$\frac{dZ_1}{dt} + (p+q-1)\alpha'(t)g(\alpha(t))Z_1(t) \geq -q(p+q-1)\alpha'(t)f(\alpha(t)), \quad \forall t \in I. \quad (2.26)$$

The above inequality implies the following estimation of $Z_1(t)$ such that

$$Z_1(t) \geq [k_1(t)]^{\frac{-(p+q-1)}{p}}, \quad \forall t \in I,$$

where $k_1(t)$ as defined in (2.20). From (2.25) and the above inequality, we have

$$Y_2^p(t) \leq k_1(t), \quad \forall t \in I,$$

thus from the above inequality in (2.22), we have

$$\frac{dJ_2(t)}{dt} \leq \alpha'(t)f(\alpha(t))k_1(t), \quad \forall t \in I.$$

By taking $t = s$ in the above inequality and integrating from 0 to t , produces

$$J_2(t) \leq u_0 \exp\left(\int_0^{\alpha(t)} f(s)k_1(\alpha^{-1}(s))ds\right), \quad t \in I. \quad (2.27)$$

Using (2.27) in (2.21), we get the required inequality in (2.19). The proof is completed. $\square$

Remark 2.3. If $q = 1$, then Theorem 2.3 reduces to Theorem 2.3 in [15].

Remark 2.4. If $\alpha(t) = t$, then Theorem 2.3 reduces to Theorem 1.4.

Remark 2.5. If $q = 1$ and $\alpha(t) = t$, then Theorem 2.3 reduces to Theorem 3.2 in [1].

Theorem 2.4. Let $u(t), g(t), f(t) \in \mathcal{C}(I, I), \alpha(t) \in \mathcal{C}^1(I, I)$ be nondecreasing with $\alpha(t) \leq t, \alpha(0) = 0$, on $I, n(t) \in \mathcal{C}(I, \mathbb{R}_+)$ be nondecreasing and satisfy the inequality

$$u(t) \leq n(t) + \int_0^{\alpha(t)} f(s) \left[u(s) + \int_0^s g(\lambda) u(\lambda) d\lambda \right]^p ds, \quad \forall t \in I, \quad (2.28)$$

where p be a constant such that $p \in (0, 1)$. Then

$$u(t) \leq n(t) \left[1 + \int_0^{\alpha(t)} f(s) n^{[p-1]}(s) k_2(\alpha^{-1}(s)) \right], \quad \forall t \in I, \quad (2.29)$$

where

$$\begin{aligned} k_2(t) = & \exp \left(p(1-p) \int_0^{\alpha(t)} g(s) ds \right) \left[1 + (1-p) \int_0^{\alpha(t)} f(s) n^{[p-1]}(s) \right. \\ & \times \exp \left(-(1-p) \int_0^s g(\lambda) d\lambda \right) ds \left. \right]^{\left[\frac{p}{1-p} \right]}, \quad \forall t \in I. \end{aligned} \quad (2.30)$$

Proof. Since $n(t)$ is a positive monotonic nondecreasing function, we observe from the inequality (2.28)

$$\left[\frac{u(t)}{n(t)} \right] \leq 1 + \int_0^{\alpha(t)} f(s) n^{[p-1]}(s) \left[\left(\frac{u(s)}{n(s)} \right) + \int_0^{\alpha(s)} g(\lambda) \left(\frac{u(\lambda)}{n(\lambda)} \right) d\lambda \right]^p ds, \quad \forall t \in I.$$

Let

$$m(t) = \frac{u(t)}{n(t)}, \quad m(0) \leq 1, \quad \forall t \in I. \quad (2.31)$$

Hence

$$m(t) \leq 1 + \int_0^{\alpha(t)} f(s) n^{[p-1]}(s) \left[m(s) + \int_0^s g(\lambda) m(\lambda) d\lambda \right]^p ds, \quad \forall t \in I. \quad (2.32)$$

Let $J_3(t)$ equal the right hand side of (2.32), which is a positive and nondecreasing function on I with $J_3(0) = 1$. Then (2.32) reduces to

$$m(t) \leq J_3(t), m(\alpha(t)) \leq J_3(\alpha(t)) \leq J_3(t) \quad \forall t \in I. \quad (2.33)$$

Differentiating $J_3(t)$ with respect to t , using (2.33), we get

$$\frac{dJ_3(t)}{dt} \leq \alpha'(t) f(\alpha(t)) n^{[p-1]}(\alpha(t)) Y_3^p(t), \quad \forall t \in I, \quad (2.34)$$

where $Y_3(t) = J_3(t) + \int_0^{\alpha(t)} g(s) J_3(s) ds$, $Y_3(0) = J_3(0) = 1$, and

$$J_3(t) \leq Y_3(t), \quad \forall t \in I. \quad (2.35)$$

Differentiating $Y_3(t)$ with respect to t and using (2.34), (2.35) we get

$$\begin{aligned}\frac{dY_3(t)}{dt} &= \frac{dJ_3(t)}{dt} + \alpha'(t)g(\alpha(t))J_3(t) \\ &\leq \alpha'(t)f(\alpha(t))n^{[p-1]}(\alpha(t))Y_3^p(t) + \alpha'(t)g(\alpha(t))Y_3(t), \forall t \in I.\end{aligned}$$

The above inequality implies the following estimation of $Y_3(t)$, such that

$$Y_3^p(t) \leq k_2(t), \quad \forall t \in I,$$

where $k_2(t)$ is as given in (2.30). From the above inequality in (2.34), we have

$$\frac{dJ_3(t)}{dt} \leq \alpha'(t)f(\alpha(t))n^{[p-1]}(t)k_2(t), \quad \forall t \in I,$$

the above inequality implies the following estimation of $J_3(t)$ such that

$$J_3(t) \leq 1 + \int_0^{\alpha(t)} f(s)k_2(\alpha^{-1}(s))n^{[p-1]}ds, \quad \forall t \in I.$$

Using the above inequality in (2.33), we get

$$m(t) \leq 1 + \int_0^{\alpha(t)} f(s)k_2(\alpha^{-1}(t))n^{[p-1]}ds, \quad \forall t \in I. \quad (2.36)$$

We get the desired bound in (2.29) from (2.31) and (2.36). The proof is completed. $\square$

Remark 2.6. If $\alpha(t) = t$, then Theorem 2.4 reduces to Theorem 1.5.

Theorem 2.5. Let $u(t), g(t), f(t) \in \mathcal{C}(I, I)$ be nondecreasing, we assume that $\psi_1(t), \psi_2(t)$ are nondecreasing and continuous functions defined on I with $\psi_i(t) > 0, \forall t > 0, i = 1, 2$, and $\alpha(t) \in \mathcal{C}^1(I, I)$ is a nondecreasing function with $\alpha(t) \leq t, \alpha(0) = 0$ and satisfy the inequality

$$u(t) \leq u_0 + \int_0^{\alpha(t)} f(s)\psi_1(u(s)) \left[u^q(s) + \int_0^s g(\lambda)\psi_2(u(\lambda))d\lambda \right]^p ds, \forall t \in I, \quad (2.37)$$

where $u_0 > 0, p > 0$ and $q \geq 1$, are constants such that $p + q > 1$. Then

$$u(t) \leq \Psi_1^{-1} \left[\Psi_2^{-1} \left(\Psi_2 \left(\Psi_1(u_0^q) + \int_0^{\alpha(t)} g(s)ds \right) + \int_0^{\alpha(t)} f(s)ds \right) \right], \quad (2.38)$$

for all $t \in [0, T_1]$, where

$$\Psi_1(r) = \int_1^r \frac{dt}{\psi_2(t)}, r > 0, \quad (2.39)$$

$$\Psi_2(r) = \int_1^r \frac{\psi_2(\Psi_1^{-1}(s))ds}{\psi_1(\Psi_1^{-1}(s))(\Psi_1^{-1}(s))^{p+q-1}}, r > 0, \quad (2.40)$$

and T_1 is the largest number such that

$$\begin{aligned} \Psi_2 \left(\Psi_1(u_0^q) + \int_0^{\alpha(t)} g(s) ds \right) + \int_0^{\alpha(t)} f(s) ds &\leq \frac{\psi_2(\Psi_1^{-1}(s)) ds}{\psi_1(\Psi_1^{-1}(s))(\Psi_1^{-1}(s))^{p+q-1}}, \\ \Psi_2^{-1} \left(\Psi_2 \left(\Psi_1(u_0^q) + \int_0^{\alpha(t)} g(s) ds \right) + \int_0^{\alpha(t)} f(s) ds \right) &\leq \frac{dt}{\psi_2(t)}. \end{aligned}$$

Proof. Let $J_4(t)$ equal the right hand side in (2.37), which is a positive and nondecreasing function on I , we have

$$u(t) \leq J_4(t), u(\alpha(t)) \leq J_4(\alpha(t)) \leq J_4(t), J_4(0) = u_0, \forall t \in I. \quad (2.41)$$

Differentiating $J_4(t)$ with respect to t and using (2.41), we get

$$\frac{dJ_4(t)}{dt} \leq \alpha'(t) f(\alpha(t)) \psi_1(J_4(t)) Y_4^p(t), \quad \forall t \in I, \quad (2.42)$$

where $Y_4(t) = J_4^q(t) + \int_0^{\alpha(t)} g(s) \psi_2(J_4(s)) ds$, hence $Y_4(0) = J_4(0) = u_0^q$, and $J_4^q(t) \leq Y_4(t)$ but $q \geq 1$, thus

$$J_4(t) \leq Y_4(t) \quad \forall t \in I. \quad (2.43)$$

Differentiating $Y_4(t)$ with respect t and using (2.42), (2.43), leads to

$$\frac{dY_4(t)}{dt} \leq q\alpha'(t) f(\alpha(t)) \psi_1(Y_4(t)) Y_4^{[p+q-1]}(t) + \alpha'(t) g(\alpha(t)) \psi_2(Y_4(t)), \quad (2.44)$$

for all $t \in I$, since $\psi_2(Y_4(t)) > 0, \forall t > 0$, from (2.44) we get

$$\frac{dY_4(t)}{\psi_2(Y_4(t))} \leq q\alpha'(t) f(\alpha(t)) \frac{\psi_1(Y_4(t)) Y_4^{[p+q-1]}(t)}{\psi_2(Y_4(t))} dt + \alpha'(t) g(\alpha(t)) dt, \forall t \in I.$$

By taking $t = s$ in the last inequality and integrating both sides from 0 to t , and using (2.39), we have

$$\begin{aligned} \Psi_2(Y_4(t)) &\leq \Psi_2(u_0^q) + \int_0^t q\alpha'(s) f(\alpha(s)) \frac{\psi_1(Y_4(s)) Y_4^{[p+q-1]}(s)}{\psi_2(Y_4(s))} ds \\ &\quad + \int_0^t \alpha'(s) g(\alpha(s)) ds, \quad \forall t \in I, \end{aligned}$$

where Ψ_1 is defined by (2.39) from the above inequality we have

$$\begin{aligned} \Psi_2(Y_4(t)) &\leq \Psi_2(u_0^q) + \int_0^T \alpha'(s) g(\alpha(s)) ds \\ &\quad + \int_0^t q\alpha'(s) f(\alpha(s)) \frac{\psi_1(Y_4(s)) Y_4^{[p+q-1]}(s)}{\psi_2(Y_4(s))} ds, \end{aligned}$$

for all $t < T$, where $T \in [0, T_1]$ is chosen arbitrary. Let $Y_5(t)$ denote the function on the right-hand side of the above inequality, which is a positive and nondecreasing function on I with $Y_5(t) = \Psi_2(u_0^q) + \int_0^T \alpha'(s)g(\alpha(s))ds$ and

$$Y_4(t) \leq \Psi_1^{-1}(Y_5(t)), \forall t < T. \quad (2.45)$$

Differentiating $Y_5(t)$ with respect to t , and using (2.45) we obtain

$$\begin{aligned} \frac{dY_5(t)}{dt} &\leq q\alpha'(t)f(\alpha(t)) \frac{\psi_1(Y_3(t))Y_4^{[p+q-1]}(t)}{\psi_2(Y_4(t))} \\ &\leq q\alpha'(t)f(\alpha(t)) \frac{\psi_1(\psi_2(Y_5(t)))(\psi_2(Y_5(t)))^{[p+q-1]}(t)}{\psi_2(\psi_2(Y_5(t)))}, \forall t < T. \end{aligned}$$

From the above inequality, by the definition of Ψ_2 in (2.40) and let $t = T$ we have

$$\Psi_2(Y_5(t)) \leq \Psi_2\left(\Psi_1(u_0^q) + \int_0^{\alpha(T)} g(s)ds\right) + \int_0^{\alpha(T)} qf(s)ds, \forall t < T. \quad (2.46)$$

Since $0 < T < T_1$ is chosen, from (2.41), (2.43), (2.45) and (2.46), we get the required inequality in (2.38). The proof is completed. $\square$

Remark 2.7. If $q = 1$, then Theorem 2.5 reduces to Theorem 4 in [15].

Remark 2.8. If $\psi_1(t) = \psi_2(t) = t$, then Theorem 2.5 reduces to Theorem 2.3.

3 Application

In this section we present some applications of the inequalities given in our Theorem 2.1 and Theorem 2.5 in order to illustrate the usefulness of our results.

Now we give an example for the application of our Theorem 2.1 to the following retarded integro-differential equation

Example 3.1 Consider the retarded integro-differential equation :

$$\begin{cases} pu^{p-1}(t) \frac{du(t)}{dt} = M(t, u(\alpha(t)), \int_0^t H(s, u(\alpha(s)))ds), \forall t \in I, \\ u(0) = u_0, \end{cases} \quad (3.1)$$

where $M \in \mathcal{C}(I \times \mathbb{R}^3, \mathbb{R})$, $H \in \mathcal{C}(I \times \mathbb{R}, \mathbb{R})$, $|u_0| > 0$ is a constant, satisfy the following conditions

$$|H(t, u(t))| \leq g(t)|u(t)|, \quad (3.2)$$

$$|M(t, u(\alpha(t)), \int_0^t H(s, u(\alpha(s)))ds)| \leq f(t) \left(|u(t)|^p + \int_0^t |K(s, u(\alpha(s)))|ds \right), \quad (3.3)$$

$$u_0^{\frac{1}{p}} \left[1 + \int_0^{\alpha(t)} \frac{f(\alpha^{-1}(s))}{\alpha'(\alpha^{-1}(s))} \exp \left(\int_0^s \left[\frac{f(\alpha^{-1}(\lambda)) + g(\alpha^{-1}(\lambda))}{\alpha'(\alpha^{-1}(\lambda))} \right] d\lambda \right) ds \right]^{\frac{1}{p}} < \infty, \quad (3.4)$$

where $u(t), f(t), g(t), \alpha(t), p$ as defined in Theorem 2.1. Integrate both sides of the equation (3.1) from 0 to t , we have

$$u^p(t) = u_0 + \int_0^t M(s, u(\alpha(s)), \int_0^s H(\lambda, u(\alpha(\lambda)))d\lambda)ds, \forall t \in I, \quad (3.5)$$

using the conditions (3.2) and (3.3), from (3.5) we get

$$\begin{aligned} |u(t)|^p &\leq |u_0| + \int_0^t f(s) \left[|u(s)|^p + \int_0^s g(\lambda) |u(\lambda)| d\lambda \right] ds, \\ &\leq |u_0| + \int_0^{\alpha(t)} \frac{f(\alpha^{-1}(s))}{\alpha'(\alpha^{-1}(s))} \left[|u(s)|^p + \int_0^s \frac{g(\alpha^{-1}(\lambda))}{\alpha'(\alpha^{-1}(\lambda))} |u(\lambda)| d\lambda \right] ds \end{aligned}$$

Now, a suitable application of the inequality given in Theorem 2.1 to the above inequality yields

$$|u(t)| \leq u_0^{\frac{1}{p}} \left[1 + \int_0^{\alpha(t)} \frac{f(\alpha^{-1}(s))}{\alpha'(\alpha^{-1}(s))} \exp \left(\int_0^s \left[\frac{[f(\alpha^{-1}(\lambda)) + g(\alpha^{-1}(\lambda))]}{\alpha'(\alpha^{-1}(\lambda))} \right] d\lambda \right) ds \right]^{\frac{1}{p}}, \quad (3.6)$$

for all $t \in I$. Thus, from the hypotheses (3.4) and the estimation in (3.6), implies the boundedness of the solution $u(t)$ of (3.1).

Now we give an example for the application of our Theorem 2.5 to the following retarded integro-differential equation

$$\frac{du(t)}{dt} = H(t, u(\alpha(t)), \int_0^t K(s, u(\alpha(s)))ds), \forall t \in I, \quad (3.7)$$

with initial condition $u(0) = u_0$, we assume that $K \in \mathcal{C}(\mathbb{R} \times \mathbb{R}, \mathbb{R})$, $H \in \mathcal{C}(\mathbb{R}^*, \mathbb{R})$, $|u_0| > 0$ is a constant, satisfy the following conditions

$$|K(t, u(t))| \leq g(t)\psi_2(|u(t)|), \quad (3.8)$$

$$|H(t, u(\alpha), \int_0^t K(s, u(\alpha))ds)| \leq f(t)\psi_1(|u(\alpha)|) \left(|u(t)|^q + \int_0^t |K(s, u(\alpha))|ds \right)^p, \quad (3.9)$$

where $f(t), g(t) \in \mathcal{C}(I, I)$.

Example 3.2 Consider the nonlinear retarded integro-differential equation (3.7) and suppose that K, H satisfy the conditions (3.8) and (3.9), and $\psi_1(t), \psi_2(t)$ are nondecreasing and continuous functions defined on I with $\psi_i(t) > 0, \forall t > 0, i = 1, 2$, and $\alpha(t) \in \mathcal{C}^1(I, I)$ is a nondecreasing function with $\alpha(t) \leq t, \alpha(0) = 0$ then all the solutions of the equation (3.7) exist on I and satisfy the following estimation

$$|u(t)| \leq \Psi_1^{-1} \left[\Psi_2^{-1} \left(\Psi_2 \left(\Psi_1(|u_0^q|) + \int_0^{\alpha(t)} \frac{g(\alpha^{-1}(s))}{\alpha'(\alpha^{-1}(s))} ds \right) + \int_0^{\alpha(t)} \frac{f(\alpha^{-1}(s))}{\alpha'(\alpha^{-1}(s))} ds \right) \right], \quad (3.10)$$

for all $t < T_2$, where

$$\Psi_1(r) = \int_1^r \frac{dt}{\psi_2(t)}, r > 0, \quad (3.11)$$

$$\Psi_2(r) = \int_1^r \frac{\Psi_1(\Psi_1^{-1}(s))ds}{\psi_1(\Psi_1^{-1}(s))(\Psi_1^{-1}(s))^{p+q-1}}, r > 0, \quad (3.12)$$

and T_2 is the largest number such that

$$\begin{aligned} & \Psi_2 \left(\Psi_1(|u_0|^q) + \int_0^{\alpha(t)} \frac{g(\alpha^{-1}(s))}{\alpha'(\alpha^{-1}(s))} ds \right) + \int_0^{\alpha(t)} \frac{f(\alpha^{-1}(s))}{\alpha'(\alpha^{-1}(s))} ds \\ & \leq \frac{\Psi_1(\Psi_1^{-1}(s))ds}{\psi_1(\Psi_1^{-1}(s))(\Psi_1^{-1}(s))^{p+q-1}}, \\ & \Psi_2^{-1} \left(\Psi_2 \left(\Psi_1(|u_0|^q) + \int_0^{\alpha(t)} \frac{g(\alpha^{-1}(s))}{\alpha'(\alpha^{-1}(s))} ds \right) + \int_0^{\alpha(t)} \frac{f(\alpha^{-1}(s))}{\alpha'(\alpha^{-1}(s))} ds \right) \leq \frac{dt}{\psi_2(t)}. \end{aligned}$$

Proof. Integrating both side of the Eq. (3.7) from 0 to t , we have

$$u(t) = u_0 + \int_0^t H(s, u(\alpha(s)), \int_0^s K(\lambda, u(\alpha(\lambda)))d\lambda)ds, \forall t \in I, \quad (3.13)$$

using the conditions (3.8) and (3.9), from (3.13) we get

$$|u(t)| \leq |u_0| + \int_0^t f(s)\varphi(|u(\alpha(s))|) \left[|u^q(s)| + \int_0^s g(\lambda)\varphi(|u(\lambda)|)d\lambda \right]^p ds, \quad (3.14)$$

the inequality (3.14) can be written in the form

$$\begin{aligned} |u(t)| & \leq |u_0| + \int_0^{\alpha(t)} \frac{f(\alpha^{-1}(s))}{\alpha'(\alpha^{-1}(s))} \varphi(|u(\alpha(s))|) \\ & \times \left[|u^q(s)| + \int_0^s \frac{g(\alpha^{-1}(\lambda))}{\alpha'(\alpha^{-1}(\lambda))} \varphi(|u(\lambda)|)d\lambda \right]^p ds, \forall t \in I. \end{aligned} \quad (3.15)$$

Now, a suitable application of the inequality given in Theorem 2.5 to (3.15), we get the required inequality in (3.10). The proof is completed. $\square$

Acknowledgments

The authors are very grateful to the editor and the referees for their helpful comments and valuable suggestions.

References

- [1] Abdeldaim A., Yakout M.: On some new integral inequalities of Gronwall-Bellman-Pachpatte type. Appl Math Comput. 217, (2011), 7887-7899. doi:10.1016/j.amc.2011.02.093.

- [2] Agarwal R.P., Deng S., Zhang W.: Generalization of a retarded Gronwall-like inequality and its applications. *Appl Math Comput.* 165,(2005), 599-612. doi:10.1016/j.amc.2004.04.067.
- [3] Bellman R.: The stability of solutions of linear differential equations. *Duke Math J.* 10, (1943), 643-647. doi:10.1215/S0012-7094-43-01059-2.
- [4] Bihari I.: A generalization of a lemma of Bellman and its application to uniqueness problem of differential equation. *Acta Math Acad Sci Hung.* 7,(1956), 81-94. doi:10.1007/BF02022967.
- [5] Cheung W.S.: Some new nonlinear inequalities and applications to boundary value problems. *Nonlinear Anal.* 64,(2006), 2112-2128. doi:10.1016/j.na.2005.08.009.
- [6] El-Owaidy H., Ragab A. A., Abuelela W., Eldeeb A. A.: On some new nonlinear integral inequalities of Gronwall-Bellman type, *Kyungpook Mathematical Journal*, (to appear).
- [7] Gronwall T.H.: Note on the derivatives with respect to a parameter of the solutions of a system of differential equations. *Ann Math.* 20,(1919), 292-296. doi:10.2307/1967124.
- [8] Kim Y.H.: On some new integral inequalities for functions in one and two variables. *Acta Math Sinica.* 21,(2005), 423-434. doi:10.1007/s10114-004-0463-7.
- [9] Lipovan O.: A retarded Gronwall-like inequality and its applications. *J Math Anal Appl.* 252, (2000), 389-401. doi:10.1006/jmaa.2000.7085.
- [10] Louartassi Y., El Mazoudi E., Elalami N.: A new generalization of lemma Gronwall-Bellman. *Applied Mathematical Sciences* 6(13), (2012), 621-628.
- [11] Pachpatte B.G.: *Inequalities for Differential and Integral Equations.* Academic Press London (1998).
- [12] Peng Y., Xiang X., Wei W.: Nonlinear impulsive integro-differential equations of mixed type with time-varying generating operators and optimal controls. *Dynamic Systems and Applications*, 16(2007), 481-496.
- [13] Peng Y., Ying Kang, Mingxiu Yan, Rongfeng Huang, and Liu Yang: Gronwal-type integral inequalities with impulses on time scales. *Advances in Difference Equations*, (2011), 1-9.
- [14] Wang W.S.: A generalized retarded Gronwall-like inequality in two variables and applications to BVP. *Appl Math Comput.* 191,(2007), 144-154. doi:10.1016/j.amc.2007.02.099.
- [15] Wang W.S., Shen C.: On a generalized retarded integral inequality with two variables. *J Inequal Appl* 2008, 9(2008). Article ID 518646.

- [16] Wang W.S., Li Z., Li Y., Huang, Y: Nonlinear retarded integral inequalities with two variables and applications. J Inequal Appl 2010, 8(2010). Article ID 240790.
- [17] Wang W.S., Luo R.C., Li Z.: A new nonlinear retarded integral inequality and its application. J Inequal Appl 2010, 8(2010). Article ID 462163.
- [18] Wang W.S., Shen: Some new generalized retarded nonlinear integral inequalities with iterated integrals and their application. J Inequal Appl 2012, 236(2012). doi:10.1186/1029-242X-2012-236.
- [19] Wei W., Xiang X., Peng Y.: Nonlinear impulsive integro-differential equation of mixed type and optimal controls. Optimization, 55(1-2)(2006), 141-156.