

**Traveling wave solutions of the simplified MCH equation via
 $\exp(-\Phi(\xi))$ -expansion method**

1Md. Nur Alam, 1,*Harun-Or-Roshid 2M. Ali Akbar and 1Rafiqul Islam

**1Department of Mathematics, Pabna University of Science and Technology,
Bangladesh**

2Department of Applied Mathematics, University of Rajshahi, Bangladesh

***Email: harunoroshidmd@gmail.com**

harunoroshidmd@gmail.com

Abstract

The $\exp(-\Phi(\xi))$ -expansion method is an efficient method for obtaining exact traveling wave solutions of nonlinear evolution equations. In this paper, the $\exp(-\Phi(\xi))$ -expansion method is applied to construct exact traveling wave solutions of the simplified MCH equation. The traveling wave solutions are expressed in terms of the hyperbolic functions, the trigonometric functions and the rational functions. It is shown that the method is straightforward and effective mathematical tool for solving nonlinear evolution equations in mathematical physics and engineering.

Keywords: The $\exp(-\Phi(\xi))$ -expansion method; the simplified MCH equation; traveling wave solutions; solitary wave solutions.

PACS: 05.45.Yv, 02.30.Jr, 02.30.Ik.

Introduction

Many complex real world problems in nature are due to nonlinear phenomena. Nonlinear processes are one of the biggest challenges and not easy to control because the nonlinear characteristic of the system abruptly changes due to some small changes of valid parameters including time. Thus the issue becomes more complicated and hence needs ultimate solution. Therefore, the studies of exact solutions of nonlinear evolution equations (NLEEs) play a

crucial role to understand the internal mechanism of nonlinear phenomena. Advance nonlinear techniques are significant to solve inherent nonlinear problems, particularly those involving differential equations, dynamical systems and related areas. In recent years, both the mathematicians and physicists have made significant improvement in finding the exact solutions of NLEEs. They establish many effective and powerful methods to handle the NLEEs. For example, the Jacobi elliptic function expansion method [1, 2], the Backlund transformation method [3], the F-expansion method[4, 5], the Darboux transformation method [6], the inverse scattering transform [7], the Adomian decomposition method [8, 9], the complex hyperbolic function method [10, 11], the homogeneous balance method [12-14], the (G'/G) -expansion method [15-25], the modified simple equation method [26, 27], the auxiliary equation method[28, 29], the exp-functions method[30], the $\exp(-\phi(\eta))$ -expansion method [31] and so on.

The objective of this article is to apply the $\exp(-\phi(\xi))$ -expansion method to construct the exact solutions for nonlinear evolution equations in mathematical physics via the simplified MCH equation.

Our aim in this paper is to present an application of the $\exp(-\phi(\xi))$ -expansion method to the simplified MCH equation to be solved by this method for the first time.

The rest of the paper is organized as follows: In Section 2, we give the description of the $\exp(-\phi(\xi))$ -expansion method. In Section 3, we apply this method to the simplified MCH equation and graphical representations of the solutions. Conclusions are given in the last section.

2. Description of the $\exp(-\Phi(\xi))$ -expansion method

Let us consider a general nonlinear PDE in the form

$$F(u, u_t, u_x, u_{xx}, u_{tt}, u_{tx}, \dots), \quad (1)$$

where $u = u(x, t)$ is an unknown function, F is a polynomial in $u(x, t)$ and its derivatives in which highest order derivatives and nonlinear terms are involved and the subscripts stand for the partial derivatives. In the following, we give the main steps of this method:

Step 1: We combine the real variables x and t by a complex variable ξ

$$u(x, t) = u(\xi), \quad \xi = x \pm V t, \quad (2)$$

where V is the speed of the traveling wave. The traveling wave transformation (2) converts Eq. (1) into an ordinary differential equation (ODE) for $u = u(\xi)$:

$$\Re(u, u', u'', u''', \dots), \quad (3)$$

where $\Re$ is a polynomial of u and its derivatives and the superscripts indicate the ordinary derivatives with respect to ξ .

Step 2. Suppose the traveling wave solution of Eq. (3) can be expressed as follows:

$$u(\xi) = \sum_{i=0}^N \alpha_i (\exp(-\Phi(\xi)))^i, \quad (4)$$

where α_i ($0 \leq i \leq N$) are constants to be determined, such that $\alpha_N \neq 0$ and $\Phi = \Phi(\xi)$ satisfies the following ordinary differential equation:

$$\Phi'(\xi) = \exp(-\Phi(\xi)) + \mu \exp(\Phi(\xi)) + \lambda, \quad (5)$$

Eq. (5) gives the following solutions:

Family 1: When $\mu \neq 0$, $\lambda^2 - 4\mu > 0$,

$$\Phi(\eta) = \ln\left(\frac{-\sqrt{(\lambda^2 - 4\mu)} \tanh\left(\frac{\sqrt{(\lambda^2 - 4\mu)}}{2}(\eta + E)\right) - \lambda}{2\mu}\right) \quad (6)$$

Family 2: When $\mu \neq 0$, $\lambda^2 - 4\mu < 0$,

$$\Phi(\eta) = \ln\left(\frac{\sqrt{(4\mu - \lambda^2)} \tan\left(\frac{\sqrt{(4\mu - \lambda^2)}}{2}(\eta + E)\right) - \lambda}{2\mu}\right) \quad (7)$$

72 **Family 3:** When $\mu = 0$, $\lambda \neq 0$, and $\lambda^2 - 4\mu > 0$,

$$73 \quad \Phi(\xi) = -\ln\left(\frac{\lambda}{\exp(\lambda(\xi + E)) - 1}\right) \quad (8)$$

74 **Family 4:** When $\mu \neq 0$, $\lambda \neq 0$, and $\lambda^2 - 4\mu = 0$,

$$75 \quad \Phi(\xi) = \ln\left(-\frac{2(\lambda(\xi + E) + 2)}{\lambda^2(\xi + E)}\right) \quad (9)$$

76 **Family 5:** When $\mu = 0$, $\lambda = 0$, and $\lambda^2 - 4\mu = 0$,

$$77 \quad \Phi(\xi) = \ln(\xi + E) \quad (10)$$

78 $\alpha_N, \dots, V, \lambda, \mu$ are constants to be determined latter, $\alpha_N \neq 0$, the positive integer N can be
 79 determined by considering the homogeneous balance between the highest order derivatives
 80 and the nonlinear terms appearing in Eq. (3).

81 **Step 3:** We substitute Eq. (4) into Eq. (3) and then we account the function $\exp(-\Phi(\xi))$. As a
 82 result of this substitution, we get a polynomial of $\exp(-\Phi(\xi))$. We equate all the coefficients
 83 of same power of $\exp(-\Phi(\xi))$ to zero. This procedure yields a system of algebraic equations
 84 whichever can be solved to find $\alpha_N, \dots, V, \lambda, \mu$. Substituting the values of $\alpha_N, \dots, V, \lambda, \mu$
 85 into Eq. (4) along with general solutions of Eq. (5) completes the determination of the
 86 solution of Eq. (1).

87 **3. The simplified MCH equation**

88 Now we will bring to bear the $\exp(-\phi(\eta))$ -expansion method to find exact solutions and then
 89 the solitary wave solutions of the simplified MCH equation in the form,

$$90 \quad u_t + 2ku_x - u_{xxt} + \beta u^2 u_x = 0. \quad \text{where } k \in \Re, \beta > 0. \quad (11)$$

91 Details of CH and MCH equations can be found in references [32-36].

92 Now, we use the traveling wave transformation Eq. (2) into Eq. (11), which yields

$$93 \quad -V u' + 2ku' + Vu''' + \beta u^2 u' = 0. \quad (12)$$

94 where the superscripts stand for the derivatives with respect to ξ .

95 Integrating Eq. (12) once with respect to ξ yields:

$$96 \quad (2k - V)u + Vu'' + \frac{\beta}{3}u^3 + P = 0. \quad (13)$$

97 where P is an integral constant that could be determined later.

98 Taking the homogeneous balance between u^3 and u'' in Eq. (13), we obtain $N = 1$. Therefore,
99 the solution of Eq. (13) is of the form

$$100 \quad u(\eta) = \alpha_0 + \alpha_1(\exp(-\Phi(\xi))), \quad (14)$$

101 where α_0, α_1 are constants to be determined such that $\alpha_N \neq 0$, while λ, μ are arbitrary
102 constants.

103 Substituting Eq. (14) into Eq. (13) and then equating the coefficients of $\exp(-\Phi(\xi))$ to zero,
104 we get

$$105 \quad 2V\alpha_1 + \frac{1}{3}\beta\alpha_1^3 = 0, \quad (15)$$

$$106 \quad \beta\alpha_0\alpha_1^2 + 3V\alpha_1\lambda = 0, \quad (16)$$

$$107 \quad -V\alpha_1 + V\alpha_1\lambda^2 + 2V\alpha_1\mu + 2k\alpha_1 + \beta\alpha_0^2\alpha_1 = 0, \quad (17)$$

$$108 \quad 2k\alpha_0 + \frac{1}{3}\beta\alpha_0^3 - V\alpha_0 + P + V\alpha_1\mu\lambda = 0, \quad (18)$$

109

110 Solving the Eq. (15)-Eq. (18) yields

$$111 \quad P = 0, \quad V = \frac{4k}{2 + \lambda^2 - 4\mu}, \quad \alpha_0 = \pm \sqrt{\frac{6k}{(-2\beta - \beta\lambda^2 + 4\beta\mu)}}, \quad \alpha_1 = \pm 2\sqrt{\frac{6k}{(-2\beta - \beta\lambda^2 + 4\beta\mu)}}.$$

112 where λ, μ, β, k are arbitrary constants.

113 Now substituting the values of V, α_o, α_1 into Eq. (14) yields

$$114 \quad u(\xi) = \pm \sqrt{\frac{6k}{(-2\beta - \beta\lambda^2 + 4\beta\mu)}} (1 + 2(\exp(-\Phi(\xi))), \quad (19)$$

115 where $\xi = x - (\frac{4k}{2 + \lambda^2 - 4\mu})t$.

116 Now substituting Eq. (6) - Eq. (10) into Eq. (19) respectively, we get the following five
117 traveling wave solutions of the simplified MCH equation.

118 When $\mu \neq 0, \lambda^2 - 4\mu > 0$,

$$119 \quad u_1(\xi) = \pm \sqrt{\frac{6k}{(-2\beta - \beta\lambda^2 + 4\beta\mu)}} \left(1 - \frac{4\mu}{\sqrt{\lambda^2 - 4\mu} \tanh(\frac{\sqrt{\lambda^2 - 4\mu}}{2}(\xi + E)) + \lambda}\right).$$

120 where $\xi = x - (\frac{4k}{2 + \lambda^2 - 4\mu})t$ and E is an arbitrary constant.

121 When $\mu \neq 0, \lambda^2 - 4\mu < 0$,

$$122 \quad u_2(\xi) = \pm \sqrt{\frac{6k}{(-2\beta - \beta\lambda^2 + 4\beta\mu)}} \left(1 + \frac{4\mu}{\sqrt{4\mu - \lambda^2} \tan(\frac{\sqrt{4\mu - \lambda^2}}{2}(\xi + E)) - \lambda}\right).$$

123 where $\xi = x - (\frac{4k}{2 + \lambda^2 - 4\mu})t$ and E is an arbitrary constant.

124 When $\mu = 0, \lambda \neq 0$, and $\lambda^2 - 4\mu > 0$,

$$125 \quad u_3(\xi) = \pm \sqrt{\frac{6k}{(-2\beta - \beta\lambda^2)}} \left(1 + \frac{2\lambda}{\exp(\lambda(\xi + E)) - 1}\right).$$

126 where $\xi = x - (\frac{4k}{2 + \lambda^2})t$ and E is an arbitrary constant.

127 When $\mu \neq 0, \lambda \neq 0$, and $\lambda^2 - 4\mu = 0$,

128
$$u_4(\xi) = \pm \sqrt{\frac{3k}{-\beta}} \left(1 - \frac{\lambda^2(\xi + E)}{(\lambda(\xi + E) + 2)} \right).$$

129 where $\xi = x - 2kt$ and E is an arbitrary constant.

130 When $\mu = 0$, $\lambda = 0$, and $\lambda^2 - 4\mu = 0$,

131
$$u_5(\xi) = \pm \sqrt{\frac{3k}{-\beta}} \left(1 + \frac{2}{(\xi + E)} \right).$$

132 where $\xi = x - 2kt$ and E is an arbitrary constant.

133 Graphical representation of the solutions

134 The graphical illustrations of the solutions are given below in the figures with the aid of

135 Maple.





Fig. 1: Kink wave solution $u_1(\xi)$ when $\beta=1$, $k=2$, $\mu=1$, $\lambda=3$, $E=1$ and $-10 \leq x, t \leq 10$.
 Fig. 2: Periodic solution $u_2(\xi)$ when $\beta=1$, $k=2$, $\mu=3$, $\lambda=1$, $E=1$ and $-1 \leq x, t \leq 1$.




Fig. 3: Singular soliton solution $u_3(\xi)$ when $\beta=1$, $k=2$, $\mu=0$, $\lambda=2$, $E=1$ and $-10 \leq x, t \leq 10$.




Fig. 4: Singular Kink wave solution $u_4(\eta)$ when $\beta=1$, $k=2$, $\mu=1$, $\lambda=2$, $E=1$ and $-10 \leq x, t \leq 10$.




Fig. 5: Singular Kink wave solution $u_5(\eta)$ when $\beta=1$, $k=2$, $\mu=0$, $\lambda=0$, $E=1$ and $-10 \leq x, t \leq 10$.

Conclusion: In this paper, the $\exp(-\Phi(\xi))$ -expansion method is applied successfully for solving the simplified MCH equation. The procedure is simple, direct and constructive without the help of a computer algebra system. The results show that this method is efficient in finding the exact traveling wave solutions of nonlinear differential equations.

143 **References**

- 144 1. D. Liu, Jacobi elliptic function solutions for two variant Boussinesq equations, Chaos
145 solitons Fractals, 24 (2005) 1373-85.
- 146 2. Y. Chen and Q. Wang, Extended Jacobi elliptic function rational expansion method and
147 abundant families of Jacobi elliptic functions solutions to (1+1)-dimensional dispersive
148 long wave equation, Chaos solitons Fractals, 24 (2005) 745-57.
- 149 3. M.R. Miura, Backlund transformation, Springer, Berlin, 1978.
- 150 4. M.L.Wang and Y.B. Zhou, The periodic wave solutions for the Klein-Gordon-
151 Schrodinger equations, Phys.Lett.A 318 (2003) 84-92.
- 152 5. M.L.Wang and X.Z. Li, Extended F-expansion method and periodic wave solutions for
153 the generalized Zakharov equations, Phys.Lett.A 343 (2005) 48-54.
- 154 6. V.B. Matveev and M.A. Salle, Darboux transformation and solitons, Springer, Berlin,
155 1991.
- 156 7. M.J. Ablowitz and P.A. Clarkson, Soliton, nonlinear evolution equations and inverse
157 scattering (Cambridge University Press, New York, 1991).
- 158 8. G. Adomian, Solving frontier problems of physics: The decomposition method, Kluwer
159 Academic Publishers, Boston, 1994.
- 160 9. A.M. Wazwaz, Partial Differential equations: Method and Applications, Taylor and
161 Francis, 2002.
- 162 10. E.M.E. Zayed, A.M. Abourabia, K.A. Gepreel, M.M. Horbaty, On the rational solitary
163 wave solutions for the nonlinear Hirota-Satsuma coupled KdV system, Appl. Anal. 85
164 (2006) 751-768.
- 165 11. K.W. Chow, A class of exact periodic solutions of nonlinear envelope equation, J. Math.
166 Phys. 36 (1995) 4125-4137.

- 167 12. M. Wang, Solitary wave solutions for variant Boussinesq equations, *Phy. Lett. A*, 199
168 (1995) 169-172.
- 169 13. E.M.E. Zayed, H.A. Zedan and K.A. Gepreel, On the solitary wave solutions for
170 nonlinear Hirota-Sasuma coupled KDV equations, *Chaos, Solitons and Fractals*, 22(2004)
171 285-303.
- 172 14. M.L. Wang, Exact solutions for a compound KdV-Burgers equation, *Phys. Lett.A*
173 213(1996) 279-287.
- 174 15. M.L. Wang, X.Z. Li, J. Zhang, The (G'/G) -expansion method and traveling wave
175 solutions of nonlinear evolution equations in mathematical physics. *Phys. Lett. A* 372
176 (2008) 417-423.
- 177 16. H. Naher and F.A. Abdullah, The Modified Benjamin-Bona-Mahony Equation via the
178 Extended Generalized Riccati Equation Mapping Method, *Applied Mathematical*
179 *Sciences*, Vol. 6, 2012, no. 111, 5495 – 5512.
- 180 17. H. Naher and F.A. Abdullah, The Improved (G'/G) -Expansion Method for the (2+1)-
181 Dimensional Modified Zakharov-Kuznetsov Equation , *Journal of Applied Mathematics*
182 Vol. 2012; 2012 Article ID 438928, 20 pages.
- 183 18. H. Naher and F.A. Abdullah, New approach of (G'/G) -expansion method and new
184 approach of generalized (G'/G) -expansion method for nonlinear evolution equation, *AIP*
185 *Advances* 3, 032116 (2013); doi: 10.1063/1.4794947.
- 186 19. M.N. Alam, M.A. Akbar and S.T. Mohyud-Din, A novel (G'/G) -expansion method and
187 its application to the Boussinesq equation, *Chin. Phys. B*, Vol. 23(2), 2014, 020203-
188 020210, DOI: 10.1088/1674-1056/23/2/020203.

- 189 20. M.N. Alam and M.A. Akbar, Exact traveling wave solutions of the KP-BBM equation by
190 using the new generalized (G'/G) -expansion method, SpringerPlus, 2(1), 2013, 617.
191 DOI: 10.1186/2193-1801-2-617.
- 192 21. M.N. Alam and M.A. Akbar, Traveling wave solutions of nonlinear evolution equations
193 via the new generalized (G'/G) -expansion method, Universal Journal of Computational
194 Mathematics, 1(4), 2013, 129-136 DOI: 10.13189/ujcmj.2013.010403.
- 195 22. M.N. Alam, M.A. Akbar and K. Khan, Some new exact traveling wave solutions to the
196 $(2+1)$ -dimensional breaking soliton equation, World Applied Sciences Journal, 25(3),
197 2013, 500-523.
- 198 23. A. Bekir, Application of the (G'/G) -expansion method for nonlinear evolution equations,
199 Phys. Lett. A 372 (2008) 3400-3406.
- 200 24. H. Kheiri and G. Ebadi, Application of the (G'/G) -expansion method for the Burgers,
201 Fisher and Burgers-Fisher equations, Acta Universitatis Apulensis, ISSN: 1582-5329 no.
202 24/2010 pp. 35-44.
- 203 25. M.A. Akbar, N.H.M. Ali and E.M.E. Zayed, A generalized and improved (G'/G) -
204 expansion method for nonlinear evolution equations, Math. Prob. Engr., Vol. 2012 (2012)
205 22 pages. doi: 10.1155/2012/459879.
- 206 26. A.J.M. Jawad, M.D. Petkovic and A. Biswas, Modified simple equation method for
207 nonlinear evolution equations, Appl. Math. Comput., 217 (2010) 869-877.
- 208 27. K. Khan, M.A. Akbar and M.N. Alam, Traveling wave solutions of the nonlinear
209 Drinfel'd-Sokolov-Wilson equation and modified Benjamin-Bona-Mahony equations, J.
210 Egyptian Math. Soc., 21, 2013, 233-240, <http://dx.doi.org/10.1016/j.joems.2013.04.010>.
- 211 28. Sirendaoreji, J. Sun, Auxiliary equation method for solving nonlinear partial differential
212 equations, Phys. Lett. A 309 (2003) 387-396.

- 213 29. Sirendaoreji, Auxiliary equation method and new solutions of Klein-Gordon
214 equations, Chaos Solitons Fractals 31 (2007) 943-950.
- 215 30. J.H. He and X.H. Wu, Exp-function method for nonlinear wave equations, Chaos,
216 Solitons Fract. 30 (2006) 700-708.
- 217 31. R. Islam, M.N. Alam, A.K.M.K.S. Hossain, H.O. Roshid and M.A. Akbar, Traveling
218 wave solutions of nonlinear evolution equations via $\exp(-\Phi(\eta))$ -expansion method,
219 Global J. Scient. Frontier Resear., Vol. 13 (11) , 2014, 63-71.
- 220 32. X. Liu, L. Tian and Y. Wu, Application of the (G'/G) -expansion method to two
221 nonlinear evolution equations, Appl. Math. Computa., 217 (2010) 1376-1384.
- 222 33. A.M. Wazwaz, New compact and noncompact solutions for variants of a modified
223 Camassa-Holm equation, Appl. Math. Computa., 163 (3) 2005: 1165-1179.
- 224 34. R. Camassa and D. Holm, An integrable shallow water equation with peaked solitons,
225 Phys. Rev. Lett., 71 (11) 1993: 1661-1664.
- 226 35. L. Tian and X. Song, New peaked solitary wave solutions of the generalized Camassa-
227 Holm equation, Chaos Solit. Fract., 19 (2004): 621-637.
- 228 36. J.P. Boyd, Peakons and cashoidal waves: traveling wave solutions of the Camassa-Holm
229 equation, Appl. Math. Computa. 81 (2-3) 1997: 173-187.
- 230
- 231