

Existence of coupled Quasi-solutions of Nonlinear Integro-Differential Equations of Volterra Type in Banach Spaces *

Yan Liang

(Zhengzhou Infant Normal College, Zhengzhou 450000, China)

Abstract We study an initial value problem for a class of integro-differential equations of Volterra type in a real Banach space. Using method of upper and lower solutions and Mönch and Von Harten theorem, we obtain an existence theorem of coupled quasi-solutions, which is an extension of those established by Y. Chen and W. Zhuang in [1].

Keywords Banach space; measure of noncompactness; lower and upper solutions; normal cone; quasi-solution.

2000 Mathematics Subject Classification 45D05;34A12.

1 Introduction and Preliminaries

Let E be a real Banach space with norm $\|\cdot\|$, and let E^* denote the dual of E . Let $K \subset E$ be a cone. By means of K a partial order $\leq$ is defined as $v \leq u$ iff $u - v \in K$. We let $K^* = \{\varphi \in E^* : \varphi(u) \geq 0 \text{ for all } u \in K\}$.

A cone K is said to be normal if there exists a real number $\delta > 0$ such that $0 \leq v \leq u$ implies $\|v\| \leq \delta\|u\|$, where δ is independent of u, v . We shall always assume in this paper that K is a normal cone.

Let β denote the measure of noncompactness of Hausdorff respectively. If F is a subspace of E and $M \subset F$ is bounded then we define

$$\beta_F(M) = \inf\{\varepsilon > 0 : M \subset \bigcup_{i=1}^{n(\varepsilon)} S(z_i^\varepsilon, \varepsilon) \text{ for some } z_i^\varepsilon \in F\}.$$

We have

* E-mail: 306572265@QQ.com

$$\beta(B) \leq \beta_F(B) \leq 2 \cdot \beta(B) \text{ for } B \subset F \text{ bounded.}$$

For these and further properties we refer to Deimling [2] and Sadovskii [7].

For any $v, w \in C[I, E]$ such that $v(t) \leq w(t)$ on I , where $I = [0, T], T > 0$, we define the conical segment

$$[v, w] = \{u \in C[I, E] : v \leq u \leq w\}.$$

From the definition of $[v, w]$ and the normality of the cone K , we know that $[v, w]$ is a bounded closed convex subset of $C[I, E]$.

In this paper, we consider the following initial value problem for nonlinear integro-differential equation (IVP) in a real Banach space, namely

$$u'(t) = H(t, u(t), (Su)(t)), \quad u(0) = u_0 \quad (1.1)$$

where $(Su)(t) = \int_0^t s(t, \tau)u(\tau)d\tau$, $s \in C[I \times I, R^+]$ and $s(t, s) \leq s_0$ on $I \times I$, $s_0 > 0$, $H \in C[I \times E \times E, E]$. We obtain an existence theorem of coupled quasi-solutions via the method of upper and lower solutions and Mönch and Von Harten theorem. The results of this paper are extensions of those established in [1].

In the proof of our main results the following lemmas are necessary. See [3],[4],[5] for details.

Lemma 1.1^[4] (**Mönch and Von Harten theorem**) Let E be a Banach space and β the Hausdorff measure of noncompactness on E . Let $\{x_n\}_{n \geq 1}$ be a sequence of continuously differentiable functions from $J = [a, b]$ to E such that there is some $\mu \in L^1(a, b)$ with $\|x_n(t)\| \leq \mu(t)$ and $\|x'_n(t)\| \leq \mu(t)$ on J . Let $\psi(t) = \beta(\{x_n(t)\}_{n \geq 1})$. Then $\psi(t)$ is absolutely continuous on J and

$$\psi'(t) \leq 2\beta(\{x'_n(t)\}_{n \geq 1}) \quad a.e. \text{ on } J.$$

Lemma 1.2^[4] Let E be a separable Banach space and β the Hausdorff measure of noncompactness on E . Let $\{x_n\}_{n \geq 1}$ be a sequence of continuous functions from $J = [a, b]$ to E such that there is some $\mu \in L^1(a, b)$ with $\|x_n(t)\| \leq \mu(t)$ on J . Let $\psi(t) = \beta(\{x_n(t)\}_{n \geq 1})$. Then $\psi(t)$ is integrable on J and

$$\beta(\{\int_a^b x_n(s)ds\}_{n \geq 1}) \leq \int_a^b \psi(s)ds.$$

Lemma 1.3^[5] Let $y(t) \in C[I, R]$, $y(0) \leq 0$, and satisfy

$$y'(t) \leq -My(t) - N \int_0^t s(t, \tau)y(\tau)d\tau$$

where $M > 0, N \geq 0, s \in C[I \times I, R^+]$, $s(t, \tau) \leq s_0$ for $(t, \tau) \in I \times I$. Suppose further that $Ns_0T(\exp(MT) - 1) \leq M$. Then $y(t) \leq 0$ for all $t \in I$.

To define appropriate classes of upper and lower solutions of (1.1) , we shall suppose that H admits a decomposition of the form

$$H(t, u, Su) = H_0(t, u, Su) + H_1(t, u, Su) + H_2(t, u, Su),$$

where $H_0, H_1, H_2 \in C[I \times E \times E, E]$.

Definition 1.1 Let $v_0, w_0 \in C^1[I, E]$. Then v_0, w_0 are said to be coupled lower and upper quasi-solutions of (1.1) if

$$\begin{cases} v_0' - H_0(t, v_0, Sv_0) - H_1(t, v_0, Sv_0) - H_2(t, w_0, Sw_0) \leq 0, & v_0(0) \leq u_0, \\ w_0' - H_0(t, w_0, Sw_0) - H_1(t, w_0, Sw_0) - H_2(t, v_0, Sv_0) \geq 0, & w_0(0) \geq u_0. \end{cases} \quad (1.2)$$

If in (1.2), equalities hold, then v_0, w_0 are said to be coupled quasi-solutions of (1.1). Clearly one can define, based on definition 1.1, coupled maximal and minimal quasi-solutions of (1.1). We also need a stronger form of coupled upper and lower quasi-solutions of (1.1).

Definition 1.2 Let $v_0, w_0 \in C^1[I, E]$ be such that $v_0(t) \leq w_0(t)$ on I . Then v_0, w_0 are said to be strongly coupled lower and upper quasi-solutions of (1.1) if there exist constants $M > 0, N \geq 0$ such that

$$\begin{cases} v_0' \leq H_0(t, \sigma, S\sigma) + H_1(t, v_0, Sv_0) + H_2(t, w_0, Sw_0) - M(v_0 - \sigma) - N(Sv_0 - S\sigma) \\ w_0' \geq H_0(t, \sigma, S\sigma) + H_1(t, w_0, Sw_0) + H_2(t, v_0, Sv_0) - M(w_0 - \sigma) - N(Sw_0 - S\sigma) \end{cases} \quad (1.3)$$

for all $\sigma \in [v_0, w_0]$.

We list for convenience the following assumptions and suppose that $v_0, w_0 \in C^1[I, E]$ such that $v_0(t) \leq w_0(t)$ on I and $Ns_0T(\exp(MT) - 1) \leq M$.

(A₁) For any bounded set $B \subset [v_0, w_0]$,

$$\beta(\{H_0(t, x, Sx) + H_1(t, x, Sx) : x \in B\}) \leq L\beta(B(t)),$$

$$\beta(\{H_2(t, x, Sx) : x \in B\}) \leq L\beta(B(t)),$$

where $L > 0, B(t) = \{x(t) : x \in B\}$.

(A₁') For any $u, v \in [v_0, w_0]$,

$$\|(H_0 + H_1)(t, u, Su) - (H_0 + H_1)(t, v, Sv)\| \leq L\|u(t) - v(t)\|,$$

$$\|H_2(t, u, Su) - H_2(t, v, Sv)\| \leq L\|u(t) - v(t)\|,$$

where $L > 0$.

(A₂) $H_1(t, x, Sx)$ is nondecreasing in x and $H_2(t, x, Sx)$ is nonincreasing in x relatively to the normal cone K .

Note Clearly, (A₁') implies (A₁).

2 Main results

Our main aim in this paper is to prove the following theorem.

Theorem 2.1 Assume that the cone K is normal and assumptions $(A_1'), (A_2)$ and (1.3) are satisfied. Then there exists a unique solution $u(t)$ of (1.1) on I such that $u \in [v_0, w_0]$, provided $v_0(0) \leq u_0 \leq w_0(0)$.

The proof of the above theorem will be completed by a series lemmas.

First of all, we consider the following linear initial value problem (LIVP):

$$u' = \overline{H}(t, u, Su), \quad u(0) = u_0 \quad (2.1)$$

where $\overline{H}(t, u, Su) = -Mu - N(Su) + H_0(t, \eta_1, S\eta_1) + H_1(t, \eta_1, S\eta_1) + H_2(t, \eta_2, S\eta_2) + M\eta_1 + N(S\eta_1)$, and $\eta_1, \eta_2 \in [v_0, w_0]$. Then we have

Lemma 2.1 For any $\eta_1, \eta_2 \in [v_0, w_0]$, there exists a unique solution $u(t)$ on I of (2.1).

The proof of this lemma is similar to a corresponding result given in [5] with minor modifications.

For any $\eta_1, \eta_2 \in [v_0, w_0]$, we define the mapping A by $A[\eta_1, \eta_2] = u$, where $u = u(t)$ is the unique solution of (2.1) on I corresponding to η_1, η_2 . Then we have the following

Lemma 2.2 Suppose that assumptions (A_2) and (1.3) hold. Then A maps $[v_0, w_0] \times [v_0, w_0]$ into $[v_0, w_0]$.

Proof. Let $\eta_1, \eta_2 \in [v_0, w_0]$ and let $u = A[\eta_1, \eta_2]$. For any $\phi \in K^*$, set $p(t) = \phi(v_0(t) - u(t))$ and note that $p(0) = \phi(v_0(0) - u_0) \leq 0$. Then for all $\sigma \in [v_0, w_0]$, we have

$$\begin{aligned} p'(t) &= \phi(v_0'(t) - u'(t)) \\ &\leq \phi(H_0(t, \sigma, S\sigma) + H_1(t, v_0, Sv_0) + H_2(t, w_0, Sw_0) - M(v_0 - \sigma) - N(Sv_0 - S\sigma) \\ &\quad - H_0(t, \eta_1, S\eta_1) - H_1(t, \eta_1, S\eta_1) - H_2(t, \eta_2, S\eta_2) + M(u - \eta_1) + N(Su - S\eta_1)). \end{aligned}$$

Choosing $\sigma = \eta_1$, then we get

$$\begin{aligned} p'(t) &\leq \phi((H_1(t, v_0, Sv_0) - H_1(t, \eta_1, S\eta_1)) + (H_2(t, w_0, Sw_0) - H_2(t, \eta_2, S\eta_2))) \\ &\quad - \phi(M(v_0 - u) - N(Sv_0 - Su)) \\ &\leq -Mp(t) - N \int_0^t s(t, \tau)p(\tau)d\tau, \end{aligned}$$

which implies $p(t) \leq 0$ by Lemma 1.3. This proves $v_0(t) \leq u(t)$ on I since $\phi \in K^*$ is arbitrary.

A similar argument yields that $u(t) \leq w_0(t)$ on I . Since $\eta_1, \eta_2 \in [v_0, w_0]$ are arbitrary, the proof is complete. $\square$

In view of Lemma 2.2, we can define the sequences $\{v_n\}, \{w_n\}$ as follows:

$$v_{n+1} = A[v_n, w_n], w_{n+1} = A[w_n, v_n] \text{ and } v_n, w_n \in [v_0, w_0], n = 0, 1, 2, \dots$$

We now prove the following lemma.

Lemma 2.3 Suppose that assumptions $(A_1), (A_2)$ and (1.3) hold. Then the sequences $\{v_n\}, \{w_n\}$ are uniformly bounded, equicontinuous and relatively compact on I .

Proof. Since the cone K is normal and $v_n, w_n \in [v_0, w_0]$ for all $t \in I$, it follows that $\{v_n\}, \{w_n\}$ are uniformly bounded on I .

By assumption (A_1) , $H_0 + H_1$ and H_2 map a bounded set into a bounded set. Noting that

$$\begin{aligned} v_n'(t) = & -Mv_n - N(Sv_n) + H_0(t, v_{n-1}, Sv_{n-1}) + H_1(t, v_{n-1}, Sv_{n-1}) \\ & + H_2(t, w_{n-1}, Sw_{n-1}) + Mv_{n-1} + N(Sv_{n-1}), \end{aligned} \quad (2.2)$$

so there exists a constant $M_0 > 0$ such that

$$\|v_n'(t)\| \leq M_0, \quad t \in I, \quad n = 1, 2, \dots$$

Therefore, by the mean value theorem (see theorem 1.3.2 in [6]), we have

$$\|v_n(t_1) - v_n(t_2)\| \leq M_0|t_1 - t_2|, \quad t_1, t_2 \in I.$$

This implies that $\{v_n(t)\}$ is equicontinuous on I . A similar argument shows that $\{w_n(t)\}$ is equicontinuous on I .

Let $\varphi(t) = \beta(\{v_n(t) : n \geq 0\})$, $\psi(t) = \beta(\{w_n(t) : n \geq 0\})$. Clearly $\{v_n(t)\}, \{w_n(t)\}$ satisfy the conditions of Lemma 1.2. Therefore

$$\varphi'(t) \leq 2\beta(\{v_n'(t) : n \geq 0\}), \quad \psi'(t) \leq 2\beta(\{w_n'(t) : n \geq 0\}) \quad \text{a.e. on } I. \quad (2.3)$$

Let $E_1 = \overline{\text{span}}\{v_n(t), w_n(t) : n \geq 0, t \in I \cap Q\}$, where Q is the set of rational numbers. By assumption (A_1) , we have

$$\beta(\{H_0(t, v_{n-1}, Sv_{n-1}) + H_1(t, v_{n-1}, Sv_{n-1}) : n \geq 1\}) \leq L\beta(\{v_{n-1}(t) : n \geq 1\}) = L\varphi(t), \quad (2.4)$$

$$\beta(\{H_2(t, w_{n-1}, Sw_{n-1}) : n \geq 1\}) \leq L\beta(\{w_{n-1}(t) : n \geq 1\}) = L\psi(t). \quad (2.5)$$

By the properties of Hausdorff's measure of noncompactness and Lemma 1.2, we obtain

$$\begin{aligned} \beta(\{(Sv_{n-1})(t) : n \geq 1\}) & \leq \beta_{E_1}(\{(Sv_{n-1}) : n \geq 1\}) = \beta_{E_1}(\{ \int_0^t s(t, \tau) v_{n-1}(\tau) d\tau : n \geq 1 \}) \\ & \leq \int_0^t \beta_{E_1}(\{s(t, \tau) v_{n-1}(\tau) : n \geq 1\}) d\tau = \int_0^t s(t, \tau) \beta_{E_1}(\{v_{n-1}(\tau) : n \geq 1\}) d\tau \\ & \leq 2 \int_0^t s(t, \tau) \beta(\{v_{n-1}(\tau) : n \geq 1\}) d\tau \leq 2 \int_0^t s_0 \varphi(\tau) d\tau. \end{aligned} \quad (2.6)$$

By (2.2)-(2.6), we have

$$\begin{aligned} \varphi'(t) & \leq 2(2M\varphi(t) + 2N \cdot 2s_0 \int_0^t \varphi(\tau) d\tau + L\varphi(t) + L\psi(t)) \\ & = 2(2M + L)\varphi(t) + 2L\psi(t) + 8Ns_0 \int_0^t \varphi(\tau) d\tau \quad \text{a.e. on } I. \end{aligned}$$

A similar argument yields that

$$\psi'(t) \leq 2(2M + L)\psi(t) + 2L\varphi(t) + 8Ns_0 \int_0^t \psi(\tau) d\tau \quad \text{a.e. on } I.$$

Set $m(t) = \varphi(t) + \psi(t)$, then

$$m'(t) \leq 2(2M + L)m(t) + 2Lm(t) + 8Ns_0 \int_0^t m(\tau) d\tau, \quad \text{a.e. on } I,$$

that is,

$$m'(t) \leq 4(M + L)m(t) + 8Ns_0 \int_0^t m(\tau) d\tau \quad \text{a.e. on } I.$$

Noting that $m(0) = 0$, we have

$$\begin{aligned} m(t) &\leq 4(M + L) \int_0^t m(\tau) d\tau + 8Ns_0 \int_0^t \int_0^\tau m(\xi) d\xi d\tau \\ &\leq 4(M + L) \int_0^t m(\tau) d\tau + 8Ns_0 \int_0^t \int_0^t m(\xi) d\xi d\tau \\ &\leq 4(M + L) \int_0^t m(\tau) d\tau + 8Ns_0 T \int_0^t m(\xi) d\xi \\ &= 4(M + L + 2Ns_0 T) \int_0^t m(\tau) d\tau. \end{aligned}$$

Consequently, $m(t) \leq m(0) \exp(4(M + L + 2Ns_0 T)) = 0$, we thus obtain $\varphi(t) = \psi(t) = 0, t \in I$.

Hence, by Ascoli-Arzelà theorem, the sequences $\{v_n\}, \{w_n\}$ are relatively compact in $C[I, E]$.

The proof of Lemma 2.3 is complete. $\square$

Lemma 2.4 Suppose that assumptions (A_1') , (A_2) and (1.3) hold. Then we have $p(t) \equiv 0$ on I for either

$$p(t) = \overline{\lim}_{n \rightarrow \infty} \|v_n(t) - v_{n-1}(t)\|$$

or

$$p(t) = \overline{\lim}_{n \rightarrow \infty} \|w_n(t) - w_{n-1}(t)\|.$$

Proof. Let $m_1(t) = \overline{\lim}_{n \rightarrow \infty} \|v_n(t) - v_{n-1}(t)\|$, $m_2(t) = \overline{\lim}_{n \rightarrow \infty} \|w_n(t) - w_{n-1}(t)\|$, $m(t) = m_1(t) + m_2(t)$. In the following we will prove that $m(t) \equiv 0$ on I .

By the proof of Lemma 2.3, there exists a constant $M_0 > 0$ such that

$$\|v_n'(t)\| \leq M_0, \quad \|w_n'(t)\| \leq M_0, \quad \forall t \in I.$$

Therefore, for all $t_1, t_2 \in I$, we have

$$\begin{aligned} &\|v_n(t_1) - v_{n-1}(t_1)\| - \|v_n(t_2) - v_{n-1}(t_2)\| \\ &\leq \|v_n(t_1) - v_n(t_2)\| + \|v_{n-1}(t_1) - v_{n-1}(t_2)\| \\ &\leq 2M_0|t_1 - t_2|, \end{aligned}$$

thus

$$\begin{aligned} \|v_n(t_1) - v_{n-1}(t_1)\| &\leq \|v_n(t_2) - v_{n-1}(t_2)\| + 2M_0|t_1 - t_2|, \\ \|v_n(t_2) - v_{n-1}(t_2)\| &\leq \|v_n(t_1) - v_{n-1}(t_1)\| + 2M_0|t_1 - t_2|. \end{aligned}$$

Taking the limit, we obtain

$$m_1(t_1) \leq m_1(t_2) + 2M_0|t_1 - t_2|, \quad m_1(t_2) \leq m_1(t_1) + 2M_0|t_1 - t_2|$$

and so

$$|m_1(t_1) - m_1(t_2)| \leq 2M_0|t_1 - t_2|,$$

which proves that $m_1(t)$ is continuous on I . A similar argument yields that $m_2(t)$ is also continuous on I .

Now (A_1') yields

$$\begin{aligned} \|v_{n+1}(t) - v_n(t)\| &\leq \int_0^t [\|H_0(\tau, v_n(\tau), (Sv_n)(\tau)) + H_1(\tau, v_n(\tau), (Sv_n)(\tau)) \\ &\quad - H_0(\tau, v_{n-1}(\tau), (Sv_{n-1})(\tau)) - H_1(\tau, v_{n-1}(\tau), (Sv_{n-1})(\tau))\| \\ &\quad + \|H_2(\tau, w_n(\tau), (Sw_n)(\tau)) - H_2(\tau, w_{n-1}(\tau), (Sw_{n-1})(\tau))\| + M\|v_{n+1} - v_n\| \\ &\quad + N\|Sv_{n+1} - Sv_n\| + M\|v_n - v_{n-1}\| + N\|Sv_n - Sv_{n-1}\|] d\tau \\ &\leq \int_0^t [(L + M)\|v_n - v_{n-1}\| + L\|w_n - w_{n-1}\| + M\|v_{n+1} - v_n\| \\ &\quad + N\|Sv_{n+1} - Sv_n\| + N\|Sv_n - Sv_{n-1}\|] d\tau \\ &\leq \int_0^t [(L + M)\|v_n - v_{n-1}\| + L\|w_n - w_{n-1}\| \\ &\quad + (M + Ns_0T)\|v_{n+1} - v_n\| + Ns_0T\|v_n - v_{n-1}\|] d\tau \\ &= \int_0^t [(L + M + Ns_0T)\|v_n - v_{n-1}\| + L\|w_n - w_{n-1}\| \\ &\quad + (M + Ns_0T)\|v_{n+1} - v_n\|] d\tau. \end{aligned}$$

By Fatou lemma, taking limit, we have

$$\begin{aligned} m_1(t) &\leq \int_0^t [(L + M + Ns_0T)m_1(\tau) + Lm_2(\tau) + (M + Ns_0T)m_1(\tau)] d\tau \\ &= \int_0^t [(L + 2M + 2Ns_0T)m_1(\tau) + Lm_2(\tau)] d\tau. \end{aligned}$$

Similarly, we can obtain

$$m_2(t) \leq \int_0^t [(L + 2M + 2Ns_0T)m_2(\tau) + Lm_1(\tau)] d\tau.$$

Therefore,

$$m(t) = m_1(t) + m_2(t) \leq 2 \int_0^t (L + M + Ns_0T)m(\tau) d\tau.$$

Notice that $m(0) = 0$, so $m(t) \equiv 0$ on I . The proof of Lemma 2.4 is complete. $\square$

Proof of theorem 2.1

By Lemma 2.3, the sequences $\{v_n\}, \{w_n\}$ have uniformly convergent subsequences $\{v_{n_k}\}, \{w_{n_k}\}$.

We let $\lim_{k \rightarrow \infty} v_{n_k} = v$, $\lim_{k \rightarrow \infty} w_{n_k} = w$ and notice that

$$\begin{aligned} v_{n_k}(t) &= u_0 + \int_0^t [-Mv_{n_k}(\tau) - N(Sv_{n_k})(\tau) + H_0(\tau, v_{n_k-1}(\tau), (Sv_{n_k-1})(\tau)) \\ &\quad + H_1(\tau, v_{n_k-1}(\tau), (Sv_{n_k-1})(\tau)) + H_2(\tau, w_{n_k-1}(\tau), (Sw_{n_k-1})(\tau)) + Mv_{n_k-1}(\tau) + N(Sv_{n_k-1})(\tau)] d\tau, \end{aligned}$$

and Lemma 2.4 implies

$$\lim_{k \rightarrow \infty} v_{n_k-1} = \lim_{k \rightarrow \infty} v_{n_k} = v, \lim_{k \rightarrow \infty} w_{n_k-1} = \lim_{k \rightarrow \infty} w_{n_k} = w \text{ uniformly on } I.$$

Therefore, letting $k \rightarrow \infty$, we get

$$v(t) = u_0 + \int_0^t [H_0(\tau, v(\tau), (Sv)(\tau)) + H_1(\tau, v(\tau), (Sv)(\tau)) + H_2(\tau, w(\tau), (Sw)(\tau))] d\tau.$$

Similarly, we have

$$w(t) = u_0 + \int_0^t [H_0(\tau, w(\tau), (Sw)(\tau)) + H_1(\tau, w(\tau), (Sw)(\tau)) + H_2(\tau, v(\tau), (Sv)(\tau))] d\tau.$$

Thus v, w are the coupled quasi-solutions of (1.1). By (A'_1) , we obtain

$$\|v(t) - w(t)\| \leq 2L \int_0^t \|v(\tau) - w(\tau)\| d\tau.$$

It then follows that $\|v(t) - w(t)\| \equiv 0$ on I since $\|v(0) - w(0)\| = 0$. That is, $v = w$ is a solution of (1.1). It is easy to prove that the solution of (1.1) is unique by (A'_1) . The proof of theorem 2.1 is therefore complete. $\square$

Remark When $H_1 = H_2 = 0$, theorem 2.1 in this paper is just theorem 2.1 in [1].

References

- [1] Y. Chen and W. Zhuang, Existence via method of upper and lower solutions for nonlinear integro-differential equations in a Banach space, Journal of Maths. and Phy. Sciences, 20(2)(1986), 181-191.
- [2] K. Deimling, Ordinary differential equations in Banach spaces. Berlin-Heidelberg, New York, 1977.
- [3] S.J. Daher, On a fixed point principle of Sadovskii, Nonlinear Anal. , (1978), 643-645.
- [4] H. Mönch and von Harten, On the Cauchy problem for O.D.E in Banach spaces, Arch. Math., 39(1982),153-160.
- [5] Y. Chen and W. Zhuang, The existence of maximal and minimal solutions of the nonlinear integro-differential equations in Banach spaces, Applicable analysis A., 22(2)(1986), 139-147.
- [6] V. Lakshmikantham and S. Leela, Nonlinear differential equations in abstract spaces, Pergamon, Oxford, 1981.

- [7] B.N. Sadvskii, Limit-compact and condensing operators, Russian Math. Surveys 27(1972),85-155.
- [8] Y. Chen, W. Zhuang, Extremal solutions and comparison principle for nonlinear integro-differential equations in a Banach space, Acta Mathematics Sinica, 2(1993),152-159.