

SOME APPLICATIONS OF CELEBRATED MASTER THEOREM OF RAMANUJAN

M. I. Qureshi¹, Kaleem A. Quraishi^{2,*} and Ram Pal³

¹Department of Applied Sciences and Humanities, Faculty of Engineering and Technology, Jamia Millia Islamia (A Central University), New Delhi-110025 (India)

²Mathematics Section, Mewat Engineering College (Wakf), Palla, Nuh, Mewat-122107, Haryana (India)

³Department of Applied Sciences and Humanities, Aryabhat Polytechnic, G. T. Karnal Road, Delhi-110033 (India)

E-mails: miqureshi_delhi@yahoo.co.in; kaleemspn@yahoo.co.in;
rampal1966@rediffmail.com

ABSTRACT

If $F(x)$ is expanded in the form of Maclaurin's series

$$F(x) = \sum_{n=0}^{\infty} \left\{ (-1)^n \frac{d^n F(x)}{dx^n} \right\}_{x=0} \frac{(-x)^n}{n!}$$

then Ramanujan asserts that the value of $I = \int_0^{\infty} x^{s-1} F(x) dx$ can be found from the coefficient of $\frac{(-x)^n}{n!}$ in the expansion of $F(x)$. Conversely Ramanujan claims that if the value of I is known, then the Maclaurin's coefficient of $F(x)$ can be found.

In this paper, we obtain Maclaurin's expansions and Mellin transforms of some composite functions, using *Ramanujan's Master theorem*.

2010 Mathematics Subject Classification: Primary 44A99; Secondary 44A05, 44A20.

Keywords and Phrases: Ramanujan Master Theorem; Gauss hypergeometric function; Pochhammer's symbol; Legendre's duplication formula; Mellin transform; Inverse Mellin transform; Maclaurin's expansion; Beta and Gamma functions.

1. INTRODUCTION, DEFINITIONS AND PRELIMINARIES

Gamma Function is defined by[11,p.347(7.3.1,7.3.2)]

$$\Gamma(s) = \begin{cases} \int_0^{\infty} e^{-x} x^{s-1} dx & ; \text{ if } \Re(s) > 0 \\ \int_0^{\infty} (e^{-x} - 1) x^{s-1} dx & ; \text{ if } -1 < \Re(s) < 0 \end{cases} \quad (1.1)$$

Pochhammer's symbol or generalized factorial function $(b)_k$ is defined by[22,p.21(14); see also 18]

$$(b)_k = \frac{\Gamma(b+k)}{\Gamma(b)} = \begin{cases} b(b+1)(b+2) \cdots (b+k-1); & \text{if } k \in \{1, 2, 3, \dots\} \\ 1 & ; \text{if } k = 0 \\ k! & ; \text{if } b = 1, k \in \{1, 2, 3, \dots\} \end{cases} \quad (1.2)$$

The generalized Gaussian hypergeometric function ${}_A F_B$ of one variable is defined by [22,p.42(1); see also 18]

$${}_A F_B \left[\begin{matrix} a_1, a_2, \dots, a_A & ; \\ b_1, b_2, \dots, b_B & ; \end{matrix} \quad z \right] = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k \dots (a_A)_k z^k}{(b_1)_k (b_2)_k \dots (b_B)_k k!} \quad (1.3)$$

where denominator parameters $b_1, b_2, \dots, b_B$ are neither zero nor negative integers and A, B are non-negative integers.

If $A \leq B$, then series ${}_A F_B$ is always convergent for all finite values of z (real or complex).

If $A = B + 1$, then series ${}_A F_B$ is convergent for $|z| < 1$.

We recall here one of the gems in Ramanujan's notebooks[19;20;21], which is popularly known as the Ramanujan's Master theorem.

Ramanujan's Master-Theorem[1,p.186;4,p.298(1.1); see also 13]

Suppose that, in some neighbourhood of $x = 0$,

$$F(x) = \sum_{n=0}^{\infty} \frac{\Phi(n) (-x)^n}{n!} \quad (1.4)$$

then, Mellin transform of $F(x)$ is given by

$$\mathcal{M}\{F(x) : x \rightarrow s\} = \int_0^{\infty} x^{s-1} F(x) dx = \Gamma(s) \Phi(-s) \quad (1.5)$$

where s is not necessarily a positive integer and $F(x)$ is called inverse Mellin transform of $\Gamma(s) \Phi(-s)$ and it is given by

$$\mathcal{M}^{-1}\{\Gamma(s) \Phi(-s) : s \rightarrow x\} = F(x) = \frac{1}{2\pi i} \int_{h-i\infty}^{h+i\infty} x^{-s} \Gamma(s) \Phi(-s) ds \quad (1.6)$$

$$(0 < h < \delta, 0 < \delta < 1; x > 0)$$

provided that above integral exist.

Mellin transforms[9;16;17] is treated as a sum of two-one sided Laplace transforms.

Above theorems were reported by Ramanujan in his first quarterly report submitted to the university of Madras sometimes in August 1913. There are two other quarterly reports the second and the third submitted later by him. These reports have not been published as such. They are reported in Berndt's note book [4,pp.295-336]. All the three reports contain various applications of Master theorem for the evaluation of a variety of integrals and expansion formulae.

In excellent five "parts"[4;5;6;7;8; see also 14;15] B. C. Berndt has examined 3254 results from the various notebooks left by S. Ramanujan(1887-1920). Also, R. P. Agarwal has

made a comprehensive study of Ramanujan's work in his remarkable three "volumes" [1;2;3]. These works also contain complete references of the contributions made by other eminent mathematicians on Ramanujan's mathematics. Undoubtedly, some of Ramanujan's work has embedded in it unparalleled motivation, wisdom, depth, imagination and enough scope to pursue further research.

2. EXPANSIONS OF SOME FUNCTIONS WHEN MELLIN TRANSFORMS ARE KNOWN

[11,p.310(6.2.19)]

Since

$$\int_0^\infty x^{s-1} (1 + \alpha x)^{-\nu} dx = \alpha^{-s} B(s, \nu - s) = \alpha^{-s} \frac{\Gamma(s)\Gamma(\nu - s)}{\Gamma(\nu)} \quad (2.1)$$

$$\left(|\arg \alpha| < \pi; 0 < \Re(s) < \Re(\nu) \right)$$

then by the application of Ramanujan's Master theorem (1.4) and (1.5), we get

$$(1 + \alpha x)^{-\nu} = \sum_{n=0}^{\infty} \alpha^n \frac{\Gamma(\nu + n)}{\Gamma(\nu)} \frac{(-x)^n}{n!} \quad (2.2)$$

[11,p.311(6.2.28)]

Since

$$\int_0^\infty x^{s-1} (a^2 + x^2)^{\frac{-1}{2}} [x + \sqrt{a^2 + x^2}]^\nu dx = \frac{a^{\nu+s-1}}{2^s} B\left(s, \frac{1-s-\nu}{2}\right) = \frac{a^{\nu+s-1} \Gamma(s) \Gamma(\frac{1-s-\nu}{2})}{2^s \Gamma(\frac{1+s-\nu}{2})} \quad (2.3)$$

$$\left(\Re(a) > 0; 0 < \Re(s) < -\Re(\nu) + 1 \right)$$

then

$$\frac{[x + \sqrt{a^2 + x^2}]^\nu}{\sqrt{a^2 + x^2}} = a^{\nu-1} \sum_{n=0}^{\infty} \frac{\Gamma(\frac{1+n-\nu}{2}) \left(\frac{-2x}{a}\right)^n}{\Gamma(\frac{1-n-\nu}{2}) n!} \quad (2.4)$$

[11,p.324(6.6.14)]

Since

$$\int_0^\infty x^{s-1} \frac{\sinh(\nu \sinh^{-1} x)}{\sqrt{1+x^2}} dx = \frac{\Gamma(s) \sin(\frac{\pi s}{2}) \sin(\frac{\pi \nu}{2}) \Gamma(\frac{1-s+\nu}{2}) \Gamma(\frac{1-s-\nu}{2})}{2^s \pi} \quad (2.5)$$

$$\left(-1 < \Re(s) < 1 - |\Re(\nu)| \right)$$

then

$$\frac{\sinh(\nu \sinh^{-1} x)}{\sqrt{1+x^2}} = \sum_{n=0}^{\infty} \frac{2^n \sin(\frac{-\pi n}{2}) \sin(\frac{\pi \nu}{2}) \Gamma(\frac{1+n+\nu}{2}) \Gamma(\frac{1+n-\nu}{2})}{\pi} \frac{(-x)^n}{n!} \quad (2.6)$$

[11,p.324(6.6.15)]

Since

$$\int_0^\infty x^{s-1} \frac{\cosh(\nu \sinh^{-1} x)}{\sqrt{1+x^2}} dx = \frac{\Gamma(s) \cos(\frac{\pi s}{2}) \cos(\frac{\pi \nu}{2}) \Gamma(\frac{1-s+\nu}{2}) \Gamma(\frac{1-s-\nu}{2})}{2^s \pi} \quad (2.7)$$

$$\left(0 < \Re(s) < 1 - |\Re(\nu)|\right)$$

then

$$\frac{\cosh(\nu \sinh^{-1} x)}{\sqrt{1+x^2}} = \sum_{n=0}^{\infty} \frac{2^n \cos(\frac{\pi n}{2}) \cos(\frac{\pi \nu}{2}) \Gamma(\frac{1+n+\nu}{2}) \Gamma(\frac{1+n-\nu}{2}) (-x)^n}{\pi n!} \quad (2.8)$$

[11,p.348(7.3.9)]

Since

$$\int_0^{\infty} x^{s-1} e^{-x \cos \alpha} \sin(x \sin \alpha) dx = \Gamma(s) \sin(\alpha s) \quad (2.9)$$

$$\left(\Re(s) > -1; \quad -\frac{\pi}{2} < \Re(\alpha) < \frac{\pi}{2}\right)$$

then

$$e^{-x \cos \alpha} \sin(x \sin \alpha) = \sum_{n=1}^{\infty} (-1)^{n+1} \sin(\alpha n) \frac{x^n}{n!} \quad (2.10)$$

[11,p.348(7.3.11)]

Since

$$\int_0^{\infty} x^{s-1} e^{-x \cos \alpha} \cos(x \sin \alpha) dx = \Gamma(s) \cos(\alpha s) \quad (2.11)$$

$$\left(\Re(s) > 0; \quad -\frac{\pi}{2} < \Re(\alpha) < \frac{\pi}{2}\right)$$

then

$$e^{-x \cos \alpha} \cos(x \sin \alpha) = \sum_{n=0}^{\infty} (-1)^n \cos(\alpha n) \frac{x^n}{n!} \quad (2.12)$$

[11,p.316(6.4.27),p.347(7.3.25)]

Since

$$\int_0^{\infty} x^{s-1} \ln(1 + 2x \cos \theta + x^2) dx = \frac{2\pi \operatorname{cosec}(\pi s) \cos(\theta s)}{s} \quad (2.13)$$

$$\left(-1 < \Re(s) < 0; \quad -\pi < \theta < \pi\right)$$

then

$$\ln(1 + 2x \cos \theta + x^2) = 2 \sum_{n=1}^{\infty} (-1)^{n+1} \cos(\theta n) \frac{x^n}{n} \quad (2.14)$$

$$\left(|2x \cos \theta + x^2| < 1\right)$$

[11,p.317(6.4.28)]

Since

$$\int_0^{\infty} x^{s-1} \ln(1 - 2\alpha e^{-x} \cos \theta + \alpha^2 e^{-2x}) dx = -2\Gamma(s) \sum_{m=1}^{\infty} \frac{\alpha^m \cos(m\theta)}{(m+1)} \quad (2.15)$$

$$\left(0 < \alpha < 1; \quad -\pi < \theta < \pi; \quad \Re(s) > 0\right)$$

then

$$\ln(1 - 2\alpha e^{-x} \cos \theta + \alpha^2 e^{-2x}) = -2 \left(\sum_{m=1}^{\infty} \frac{\alpha^m \cos(m\theta)}{(m+1)} \right) e^{-x} \quad (2.16)$$

[11,p.322(6.5.46)]

Since

$$\int_0^{\infty} x^{s-1} \tan^{-1} x \, dx = -\frac{\pi}{2s} \sec\left(\frac{\pi}{2}\right) \quad (2.17)$$

$$\left(-1 < \Re(s) < 0 \right)$$

then

$$\tan^{-1} x = \sum_{n=1}^{\infty} (-1)^{n+1} \sin\left(\frac{n\pi}{2}\right) \frac{x^n}{n}; \quad |x| < 1 \quad (2.18)$$

[11,p.336(6.9.3)]

Since

$$\int_0^{\infty} x^{s-1} {}_2F_1 \left[\begin{matrix} \alpha, \beta & ; \\ \gamma & ; \end{matrix} -x \right] dx = \frac{B(s, \alpha-s) B(s, \beta-s)}{B(s, \gamma-s)} \quad (2.19)$$

$$\left(0 < \Re(s) < \min\{\Re(\alpha), \Re(\beta)\} \right)$$

then

$${}_2F_1 \left[\begin{matrix} \alpha, \beta & ; \\ \gamma & ; \end{matrix} -x \right] = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\beta)} \sum_{n=0}^{\infty} \frac{\Gamma(\alpha+n) \Gamma(\beta+n)}{\Gamma(\gamma+n)} \frac{(-x)^n}{n!} \quad (2.20)$$

[10,p.285(6.15.2.10)]

Since

$$\int_0^{\infty} x^{s-1} {}_1F_1 \left[\begin{matrix} \alpha & ; \\ \gamma & ; \end{matrix} -x \right] dx = \frac{\Gamma(s) \Gamma(\gamma) \Gamma(\alpha-s)}{\Gamma(\alpha) \Gamma(\gamma-s)} \quad (2.21)$$

$$\left(0 < \Re(s) < \Re(\alpha) \right)$$

then

$${}_1F_1 \left[\begin{matrix} \alpha & ; \\ \gamma & ; \end{matrix} -x \right] = \frac{\Gamma(\gamma)}{\Gamma(\alpha)} \sum_{n=0}^{\infty} \frac{\Gamma(\alpha+n)}{\Gamma(\gamma+n)} \frac{(-x)^n}{n!} \quad (2.22)$$

[12,p.229(A.8.1.1)]

Since

$$\int_0^{\infty} x^{s-1} {}_A F_B \left[\begin{matrix} (a_A) & ; \\ (b_B) & ; \end{matrix} -kx \right] dx$$

$$= \frac{\Gamma(b_1) \Gamma(b_2) \cdots \Gamma(b_B) \Gamma(a_1-s) \Gamma(a_2-s) \cdots \Gamma(a_A-s) \Gamma(s)}{k^s \Gamma(a_1) \Gamma(a_2) \cdots \Gamma(a_A) \Gamma(b_1-s) \Gamma(b_2-s) \cdots \Gamma(b_B-s)} \quad (2.23)$$

$$\left(0 < \Re(s) < \min\{\Re(a_j)\}, j \in \{1, 2, 3, \dots, A\}\right)$$

then

$$\begin{aligned} & {}_A F_B \left[\begin{matrix} (a_A) & ; \\ (b_B) & ; \end{matrix} \begin{matrix} -kx \end{matrix} \right] \\ &= \frac{\Gamma(b_1) \Gamma(b_2) \cdots \Gamma(b_B)}{\Gamma(a_1) \Gamma(a_2) \cdots \Gamma(a_A)} \sum_{n=0}^{\infty} \frac{\Gamma(a_1 + n) \Gamma(a_2 + n) \cdots \Gamma(a_A + n)}{\Gamma(b_1 + n) \Gamma(b_2 + n) \cdots \Gamma(b_B + n)} \frac{(-kx)^n}{n!} \end{aligned} \quad (2.24)$$

[17,p.22(Entry 1.2(2.47));23,p.117(17)]

Since

$$\int_0^{\infty} x^{s-1} [(x+a)+\sqrt{(x^2+2ax)}]^{-\nu} dx = \frac{2\nu a^{-\nu} \left(\frac{a}{2}\right)^s \Gamma(2s) \Gamma(\nu-s)}{\Gamma(1+\nu+s)} = \frac{\nu \Gamma(s) \Gamma(\nu-s) 2^s \Gamma(s+\frac{1}{2})}{\sqrt{\pi} a^{\nu-s} \Gamma(\nu+s+1)} \quad (2.25)$$

$$\left(0 < \Re(s) < \Re(\nu)\right)$$

then

$$[(x+a)+\sqrt{(x^2+2ax)}]^{-\nu} = \sum_{n=0}^{\infty} \frac{\nu \Gamma(\nu+n) 2^{-n} \Gamma(\frac{1}{2}-n) (-x)^n}{\sqrt{\pi} a^{\nu+n} \Gamma(\nu+1-n) n!} = a^{-\nu} {}_2F_1 \left[\begin{matrix} \nu, -\nu & ; \\ \frac{1}{2} & ; \end{matrix} -\frac{x}{2a} \right] \quad (2.26)$$

[1,p.191(21)]

Since

$$\int_0^{\infty} x^{s-1} \left(\frac{2}{1+\sqrt{1+4x}} \right)^{\mu} dx = \frac{\mu \Gamma(s) \Gamma(\mu-2s)}{\Gamma(\mu-s+1)} \quad (2.27)$$

$$\left(0 < \Re(s) < \Re\left(\frac{\mu}{2}\right)\right)$$

then

$$\left(\frac{2}{1+\sqrt{1+4x}} \right)^{\mu} = \mu \sum_{n=0}^{\infty} \frac{\Gamma(\mu+2n)}{\Gamma(\mu+n+1)} \frac{(-x)^n}{n!} = {}_2F_1 \left[\begin{matrix} \frac{\mu}{2}, \frac{\mu+1}{2} & ; \\ \mu+1 & ; \end{matrix} -4x \right] \quad (2.28)$$

[11,p.350(7.3.23)]

Since

$$\int_0^{\infty} x^{s-1} x^{-\frac{\nu}{2}} J_{\nu}(2\sqrt{x}) dx = \frac{\Gamma(s)}{\Gamma(\nu-s+1)} \quad (2.29)$$

$$\left(0 < \Re(s) < \Re\left(\frac{\nu}{2}\right) + \frac{1}{4}\right)$$

then

$$x^{-\frac{\nu}{2}} J_{\nu}(2\sqrt{x}) = \sum_{n=0}^{\infty} \frac{1}{\Gamma(\nu+n+1)} \frac{(-x)^n}{n!} = \frac{1}{\Gamma(\nu+1)} {}_0F_1 \left[\begin{matrix} - & ; \\ \nu+1 & ; \end{matrix} -x \right] \quad (2.30)$$

[11,p.351(7.3.31)]

Since

$$\int_0^\infty x^{s-1} \frac{[\sqrt{x} + \sqrt{4+x}]^{1-2\nu}}{2^{1-2\nu} \sqrt{4+x}} dx = \frac{\Gamma(2s) \Gamma(\nu-s)}{\Gamma(\nu+s)} = \frac{\Gamma(s) 2^{2s-1} \Gamma(s+\frac{1}{2}) \Gamma(\nu-s)}{\sqrt{\pi} \Gamma(\nu+s)} \quad (2.31)$$

$$(0 < \Re(s) < \Re(\nu))$$

then

$$\frac{[\sqrt{x} + \sqrt{4+x}]^{1-2\nu}}{\sqrt{(4+x)}} = \frac{1}{4^\nu} {}_2F_1 \left[\begin{matrix} \nu, 1-\nu & ; \\ \frac{1}{2} & ; \end{matrix} -\frac{x}{4} \right] \quad (2.32)$$

[11,p.322(6.5.46)]

Since

$$\int_0^\infty x^{s-1} \tan^{-1} x dx = -\frac{\pi}{2s} \sec\left(\frac{\pi s}{2}\right) \quad (2.33)$$

$$(-1 < \Re(s) < 0)$$

then

$$\tan^{-1} x = \sum_{n=1}^\infty (-1)^{n+1} \sin\left(\frac{n\pi}{2}\right) \frac{x^n}{n}; \quad |x| < 1 \quad (2.34)$$

[11,p.336(6.9.1)]

Since

$$\int_0^\infty x^{s-1} \exp\left(\frac{-x^2}{4}\right) D_{-\nu}(x) dx = \frac{\sqrt{\pi} \Gamma(s)}{2^{\frac{s+\nu}{2}} \Gamma(\frac{s+\nu+1}{2})} \quad (2.35)$$

$$(\Re(s) > 0)$$

then

$$\exp\left(\frac{-x^2}{4}\right) D_{-\nu}(x) = \sqrt{\frac{\pi}{2^\nu}} \sum_{n=0}^\infty \frac{1}{\Gamma(\frac{\nu-n+1}{2})} \frac{(-\sqrt{2}x)^n}{n!} \quad (2.36)$$

where $D_{-\nu}(x)$ is parabolic cylinder function.

[11,p.318(6.5.7)]

Since

$$\int_0^\infty x^{s-1} e^{-\alpha x} \sin(\beta x) dx = \frac{\Gamma(s) \sin[s \tan^{-1}(\frac{\beta}{\alpha})]}{(\sqrt{\alpha^2 + \beta^2})^s} \quad (2.37)$$

$$(\Re(\alpha) > |\Im(\beta)|; \Re(s) > -1)$$

then

$$e^{-\alpha x} \sin(\beta x) = \sum_{n=0}^\infty (\alpha^2 + \beta^2)^{\frac{n}{2}} (-1)^{n+1} \sin\left(n \tan^{-1} \frac{\beta}{\alpha}\right) \frac{x^n}{n!} \quad (2.38)$$

[11,p.320(6.5.27)]

Since

$$\int_0^\infty x^{s-1} e^{-\alpha x} \cos(\beta x) dx = \frac{\Gamma(s) \cos[s \tan^{-1}(\frac{\beta}{\alpha})]}{(\sqrt{\alpha^2 + \beta^2})^s} \quad (2.39)$$

$$\left(\Re(\alpha) > |\Im(\beta)|; \Re(s) > 0 \right)$$

then

$$e^{-\alpha x} \cos(\beta x) = \sum_{n=0}^{\infty} (\alpha^2 + \beta^2)^{\frac{n}{2}} (-1)^n \cos\left(n \tan^{-1} \frac{\beta}{\alpha}\right) \frac{x^n}{n!} \quad (2.40)$$

[11,p.315(6.4.15)]

Since

$$\int_0^\infty x^{s-1} \ln(1 + \alpha x) dx = \frac{\Gamma(s) \Gamma(1-s)}{s \alpha^s} \quad (2.41)$$

$$\left(|\arg(\alpha)| < \pi; -1 < \Re(s) < 0 \right)$$

then

$$\ln(1 + \alpha x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(\alpha x)^n}{n!} \quad (2.42)$$

3. MELLIN TRANSFORMS OF SOME FUNCTIONS WHEN SERIES EXPANSIONS ARE KNOWN

By means of Leibnitz theorem for successive differentiation, Binomial expansion, decomposition of a fraction into its partial fractions, Demoivre's theorem, Euler's formula, properties of Gamma functions, after long systematic calculation, we can find the Maclaurin's expansion of required functions. Using Ramanujan's Master theorem (1.4) and (1.5), we get Mellin transforms.

Since we know that

$$\sin(ax) = \sum_{n=1}^{\infty} (-a)^n \sin\left(\frac{n\pi}{2}\right) \frac{(-x)^n}{n!} \quad (3.1)$$

then

$$\int_0^\infty x^{s-1} \sin(ax) dx = \Gamma(s) (a)^{-s} \sin\left(\frac{s\pi}{2}\right) \quad (3.2)$$

Similarly we can obtain

$$\int_0^\infty x^{s-1} \cos(ax) dx = \Gamma(s) (-a)^{-s} \cos\left(\frac{s\pi}{2}\right)$$

Since

$$\sin^2(ax) = -\frac{1}{2} \sum_{n=0}^{\infty} (-2a)^n \cos\left(\frac{n\pi}{2}\right) \frac{(-x)^n}{n!} \quad (3.3)$$

then

$$\int_0^\infty x^{s-1} \sin^2(ax) dx = \Gamma(s) (-2)^{-s-1} a^{-s} \cos\left(\frac{s\pi}{2}\right) \quad (3.4)$$

$$\left(-2 < \Re(s) < 0\right)$$

Similarly we can obtain

$$\int_0^\infty x^{s-1} \cos^2(ax) dx = \Gamma(s) (2)^{-s-1} (-a)^{-s} \cos\left(\frac{s\pi}{2}\right)$$

Since

$$\frac{1}{(\alpha+x)(\beta+x)} = \frac{1}{(\beta-\alpha)} \sum_{n=0}^\infty (\alpha^{-n-1} - \beta^{-n-1}) (-x)^n \quad (3.5)$$

then

$$\int_0^\infty x^{s-1} (\alpha+x)^{-1} (\beta+x)^{-1} dx = \frac{\pi \operatorname{cosec}(\pi s) (\alpha^{s-1} - \beta^{s-1})}{(\beta-\alpha)} \quad (3.6)$$

$$\left(|\arg(\alpha)| < \pi; |\arg(\beta)| < \pi; 0 < \Re(s) < 2; \alpha \neq \beta\right)$$

In (3.5) and (3.6), put $\alpha = ae^{i\theta}$ and $\beta = ae^{-i\theta}$, then

$$\int_0^\infty x^{s-1} (x^2 + 2ax \cos \theta + a^2)^{-1} dx = -\Gamma(s) \Gamma(1-s) a^{s-2} \operatorname{cosec} \theta \sin[(s-1)\theta] \quad (3.7)$$

$$\left(a > 0; -\pi < \theta < \pi; 0 < \Re(s) < 2\right)$$

Since

$$\frac{x+\alpha}{(\beta+x)(\gamma+x)} = \sum_{n=0}^\infty \Gamma(1+n) \left[\left(\frac{\beta-\alpha}{\beta-\gamma}\right) \beta^{-1-n} + \left(\frac{\gamma-\alpha}{\gamma-\beta}\right) \gamma^{-1-n} \right] \frac{(-x)^n}{n!} \quad (3.8)$$

then

$$\int_0^\infty \frac{x^{s-1} (x+\alpha)}{(\beta+x)(\gamma+x)} dx = \Gamma(s) \Gamma(1-s) \left[\left(\frac{\beta-\alpha}{\beta-\gamma}\right) \beta^{s-1} + \left(\frac{\gamma-\alpha}{\gamma-\beta}\right) \gamma^{s-1} \right] \quad (3.9)$$

$$\left(|\arg(\beta)| < \pi; |\arg(\gamma)| < \pi; 0 < \Re(s) < 1\right)$$

Since

$$(x^2 + 2ax \cos \theta + a^2)^{-\nu} = (x + ae^{i\theta})^{-\nu} (x + ae^{-i\theta})^{-\nu} = a^{-2\nu} \sum_{n=0}^\infty \sum_{m=0}^\infty \frac{(\nu)_n (\nu)_m (-1)^{n+m} x^{n+m}}{n! m! (ae^{i\theta})^n (ae^{-i\theta})^m}$$

$$= \sum_{n=0}^\infty \frac{(\nu)_n a^{-2\nu}}{(ae^{i\theta})^n} {}_2F_1 \left[\begin{matrix} -n, \nu & ; \\ 1 - \nu - n & ; \end{matrix} e^{2i\theta} \right] \frac{(-x)^n}{n!} \quad (3.10)$$

then

$$\int_0^\infty x^{s-1} (x^2 + 2ax \cos \theta + a^2)^{-\nu} dx = \frac{\Gamma(s)\Gamma(\nu-s)a^{-2\nu}}{\Gamma(\nu)(ae^{i\theta})^{-s}} {}_2F_1 \left[\begin{matrix} s, \nu & ; \\ 1-\nu+s & ; \end{matrix} e^{2i\theta} \right] \quad (3.11)$$

$$\left(-\pi < \theta < \pi \right)$$

ACKNOWLEDGEMENTS

Authors are thankful to the referees for their valuable suggestions and comments in the improvement of this paper.

REFERENCES

1. Agarwal, R. P.; Resonance of Ramanujan's Mathematics. Vol. I, New Age International (P) Ltd., New Delhi, 1996.
2. Agarwal, R. P.; Resonance of Ramanujan's Mathematics. Vol. II, New Age International (P) Ltd., New Delhi, 1996.
3. Agarwal, R. P.; Resonance of Ramanujan's Mathematics. Vol. III, New Age International (P) Ltd., New Delhi, 1999.
4. Berndt, B. C.; Ramanujan's Notebooks. Part I, Springer-Verlag, New York, 1985.
5. Berndt, B. C.; Ramanujan's Notebooks. Part II, Springer-Verlag, New York, 1989.
6. Berndt, B. C.; Ramanujan's Notebooks. Part III, Springer-Verlag, New York, 1991.
7. Berndt, B. C.; Ramanujan's Notebooks. Part IV, Springer-Verlag, New York, 1994.
8. Berndt, B. C.; Ramanujan's Notebooks. Part V, Springer-Verlag, New York, 1998.
9. Ditkin, V. A., Prudnikov, A. P.; Integral Transforms and Operational Calculus, Pergamon Press, Oxford, London, Frankfurt, 1965.
10. Erdélyi, A., Magnus, W., Oberhettinger, F., Tricomi, F. G.; Higher Transcendental Functions, Vol. I (Bateman Manuscript Project), McGraw-Hill Book Co. Inc., New York, Toronto and London, 1953.
11. Erdélyi, A., Magnus, W., Oberhettinger, F., Tricomi, F. G.; Table of Integral Transforms, Vol. I (Bateman Manuscript Project), McGraw-Hill Book Co. Inc., New York, Toronto and London, 1954.
12. Exton, H.; Handbook of Hypergeometric Integrals, Theory, Applications, Tables and Computer Programs, Halsted Press (Ellis Horwood Ltd., Chichester, U. K.), John Wiley and Sons, New York, Chichester, Sussex, England, 1978.

13. Garg, M., Mittal, S.; Ramanujan-A Tribute and an Application of his Master Theorem, *J. Raj. Acad. Phys. Sci.* 2004, 3, 139-142.
14. Hardy, G. H.; Ramanujan: Twelve Lectures on Subjects Suggested by his Life and Work, AMS, Chelsea Publishing Company, Providence, Rhode Island, 1999.
15. Hardy, G. H., Aiyer, P. V. Seshu, Wilson, B. M.; Collected Papers of Srinivasa Ramanujan, First Published by Cambridge University Press, Cambridge, 1927; Reprinted by Chelsea, New York, 1962; Reprinted by the American Mathematical Society, Providence, Rhode Island, 2000.
16. Marichev, O. I.; Handbook of Integral Transforms of Higher Transcendental Functions, Theory and Algorithmic Tables, Halsted Press (Ellis Horwood, Chichester, U.K) John Wiley and Sons, New York, Brisbane, Chichester and Toronto, 1983.
17. Oberhettinger, F.; Tables on Mellin Transforms, Springer-Verlag, New York, Heidelberg, Berlin, 1974.
18. Rainville, E. D.; Special Functions, The Macmillan Co. Inc., New York 1960; Reprinted by Chelsea Publ. Co. Bronx, New York, 1971.
19. Ramanujan, S.; Notebooks of Srinivasa Ramanujan, Vol. I, Tata Institute of Fundamental Research, Bombay, 1957; Reprinted by Narosa Publishing House, New Delhi, 1984.
20. Ramanujan, S.; Notebooks of Srinivasa Ramanujan, Vol. II, Tata Institute of Fundamental Research, Bombay, 1957; Reprinted by Narosa Publishing House, New Delhi, 1984. .
21. Ramanujan, S.; The Lost Notebook and Other Unpublished Papers, Narosa Publishing House, New Delhi, 1988.
22. Srivastava, H. M., Manocha, H. L.; A Treatise on Generating Functions, Halsted Press (Ellis Horwood Limited, Chichester, U.K.) John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1984.
23. Srivastava, H. M., Qureshi, M. I., Singh, R., Arora, A.; A Family of Hypergeometric Integrals Associated with Ramanujan's Integral Formula, *Advanced Studies in Contemporary Mathematics*, 2009, 18(2), 113-125.