

2-tuple linguistic Bonferroni mean operators and their application to multiple attribute group decision making

Zhiming Zhang^{1,2*}, Chong Wu²

¹*School of Management, Harbin Institute of Technology, Harbin 150001, Heilongjiang Province, PR China*

²*College of Mathematics and Computer Science, Hebei University, Baoding 071002, Hebei Province, PR China*

ABSTRACT

Aims: The aim of this paper is to develop the 2-tuple linguistic Bonferroni mean and the weighted 2-tuple linguistic Bonferroni mean.

Study design: Some desirable properties and special cases of the developed operators are discussed. The geometric Bonferroni mean (GBM) is a generalization of the Bonferroni mean and geometric mean. In this paper, we also investigate the GBM under 2-tuple linguistic environments. We develop the 2-tuple linguistic geometric Bonferroni mean and the weighted 2-tuple linguistic geometric Bonferroni mean. We investigate some fundamental properties and special cases of them.

Place and Duration of Study: The Bonferroni Mean (BM) operator is a traditional mean type aggregation operator, which can capture the expressed interrelationship of the individual arguments and which is only suitable to aggregate crisp data.

Methodology: This paper extends the BM operator to 2-tuple linguistic environments.

Results: Based on these operators, we develop two approaches for multiple attribute group decision making with 2-tuple linguistic information.

Conclusion: Two numerical examples are provided to illustrate the effectiveness and practicality of the proposed approaches.

Keywords: Multiple attribute group decision making; 2-tuple linguistic information; 2-tuple linguistic Bonferroni mean; 2-tuple linguistic geometric Bonferroni mean.

1. INTRODUCTION

In many multiple attribute group decision making (MAGDM) problems, the decision information about alternatives is usually uncertain or fuzzy due to the increasing complexity of the socio-economic environment and the vagueness of inherent subjective nature of human thinking [1,2,3]; thus, it may be appropriate and sufficient to assess the decision information in a qualitative form rather than a quantitative form. For example, when evaluating a house's cost, linguistic terms such as "high", "medium", and "low" are usually used, and when evaluating a house's design, linguistic terms like "good", "medium", and "bad" can be frequently used. To date, many methods have been developed for dealing with linguistic information [4-13]. Herrera and Martinez [14] introduced a 2-tuple fuzzy linguistic representation model that represents the linguistic information by means of 2-tuples, which are composed by a linguistic term and a number [14,15]. The main advantage of this representation is to allow a continuous representation of the linguistic information on its domain; therefore, it can represent any counting of information obtained in a aggregation

* Tel.: +86 312 5079660; fax: +86 312 5079660.

E-mail address: zhimingzhang@ymail.com.

process without any loss of information [14,15]. In the past few decades, a variety of 2-tuple linguistic aggregation operators [15-24] have been developed for aggregating 2-tuple linguistic information. However, these 2-tuple linguistic aggregation operators only emphasize the importance of each data or ordered position and they cannot reflect the interrelationships of individual data.

The Bonferroni mean, originally introduced by Bonferroni [25], is a mean-type aggregation operator and it can provide for the aggregation lying between the max and min operators and the logical “or” and “and” operators [26]. The desirable characteristic of the BM is its capability to capture the interrelationship between input arguments [26]. Recently, Yager [27] gave a detailed studied of the BM and proposed some generalizations of the BM. However, the BM [25] and its these generalizations [27] only can accommodate the situations where the input arguments take the form of crisp numbers. In many group decision makings, the attribute values are given in the form of the other types of domains which are not suitable to be aggregated by the BM, such as interval numbers [28], intuitionistic fuzzy numbers [29], interval-valued intuitionistic fuzzy numbers [30], hesitant fuzzy elements [31], uncertain linguistic variables [32], and **triangular** fuzzy linguistic variables [33]. To address this issue, some authors have suggested some new generalizations of the BM, including the uncertain Bonferroni mean operator [34], the weighted uncertain Bonferroni mean operator [34], the intuitionistic fuzzy BM (IFBM) [26], the weighted intuitionistic fuzzy Bonferroni mean (WIFBM) [26], the Atanassov's intuitionistic fuzzy geometric Bonferroni mean [35], the weighted Atanassov's intuitionistic fuzzy geometric Bonferroni mean [35], the interval-valued intuitionistic fuzzy Bonferroni mean [36], the weighted interval-valued intuitionistic fuzzy Bonferroni mean [36], the hesitant fuzzy geometric Bonferroni mean (HFCGBM) [37], the hesitant fuzzy Choquet geometric Bonferroni mean (HFCGBM) [37], the weighted hesitant fuzzy geometric Bonferroni mean (WHFCGBM) [37], the weighted hesitant fuzzy Choquet geometric Bonferroni mean (WHFCGBM) [37], the uncertain linguistic Bonferroni mean (ULBM) operator [38], the uncertain linguistic weighted Bonferroni mean (ULWBM) operator [38], the uncertain linguistic geometric Bonferroni mean (ULGBM) operator [38], the uncertain linguistic weighted geometric Bonferroni mean (ULWGBM) operator [38], the trapezoid fuzzy linguistic Bonferroni mean (TFLBM) operator [39], the trapezoid fuzzy linguistic weighted Bonferroni mean (TFLWBM) operator [39], the trapezoid fuzzy linguistic Bonferroni OWA (TFLBOWA) operator [39], the trapezoid fuzzy linguistic weighted Bonferroni OWA (TFLWBOWA) operator [39], the generalized intuitionistic fuzzy weighted Bonferroni mean (GIFWBM) [40], and the generalized weighted Bonferroni geometric mean (GWBGM) [40]. However, these Bonferroni mean operators cannot accommodate the situations where the input arguments take the form of 2-tuples.

Based on the aforementioned analysis, we can conclude that the existing 2-tuple linguistic aggregation operators do not consider the interrelationship of the individual arguments, while the existing Bonferroni mean operators cannot accommodate the situations in which the input arguments take the form of 2-tuples. To overcome this drawback, it is therefore necessary to develop some new aggregation operators that not only accommodate 2-tuple linguistic information but also consider the interrelationship of the individual arguments. To do this, we extend the BM operator to 2-tuple linguistic environments and then develop two 2-tuple linguistic aggregation operators, including the 2-tuple linguistic Bonferroni mean and the weighted 2-tuple linguistic Bonferroni mean. The desirable characteristic of these two operators is that they not only accommodate the input arguments in the form of 2-tuples but also reflect the interrelationship of the input arguments. Xia et al. [35] introduced a new Bonferroni mean called the geometric Bonferroni mean based on the BM and the geometric mean (GM). We further extend the GBM operator to 2-tuple linguistic environments and develop the 2-tuple linguistic geometric Bonferroni mean and the weighted 2-tuple linguistic geometric Bonferroni mean. Finally, we utilize the proposed operators to develop two approaches for multiple attribute group decision making with 2-tuple linguistic information and then apply both the developed approaches to two practical examples.

The remainder of this paper is organized as follows. In Section 2, we briefly review some basic concepts and operations related to the 2-tuple fuzzy linguistic representation model, the BM, and the GBM. In Section 3, the 2-tuple linguistic Bonferroni mean and the weighted 2-tuple linguistic Bonferroni mean are developed, some desirable properties of these operators are studied, and some special cases are discussed. Section 4 develops the 2-tuple linguistic geometric Bonferroni mean and the weighted 2-tuple linguistic geometric Bonferroni mean. Section 5 introduces two approaches based on these operators for multiple attribute group decision making with 2-tuple linguistic information. In Section 6, we give two practical examples to illustrate the group decision making steps based on the proposed approaches. Section 7 ends this paper with some concluding remarks.

2. PRELIMINARIES

In this section, we will introduce the basic notions of the 2-tuple fuzzy linguistic approach, Bonferroni mean, and geometric Bonferroni mean.

2.1. The 2-tuple fuzzy linguistic representation model

Let $S = \{s_i | i = 0, 1, 2, \dots, g\}$ be a finite and totally ordered discrete linguistic term set with odd cardinality, where s_i represents a possible value for a linguistic variable, and it should satisfy the following characteristics [14,41,42].

- (1) The set is ordered: $s_i \geq s_j$ if $i \geq j$;
- (2) There is the negation operator: $neg(s_i) = s_j$ such that $j = g - i$;
- (3) Max operator: $\max(s_i, s_j) = s_i$ if $s_i \geq s_j$;
- (4) Min operator: $\min(s_i, s_j) = s_i$ if $s_i \leq s_j$.

For example, a set of seven terms S could be given as follows [43-47]:

$$S = \{s_0 = \text{nothing}, s_1 = \text{very low}, s_2 = \text{low}, s_3 = \text{medium}, s_4 = \text{high}, s_5 = \text{very high}, s_6 = \text{perfect}\}.$$

Based on the concept of symbolic translation, Herrera and Martinez [14,41] introduced a 2-tuple fuzzy linguistic representation model for dealing with linguistic information. This model represents the linguistic assessment information by means of a 2-tuple (s_i, α) , where $s_i \in S$ represents a linguistic label from the predefined linguistic term set S and $\alpha \in [-0.5, 0.5]$ is the value of symbolic translation.

Definition 2.1 [14,41]. Let β be the result of an aggregation of the indexes of a set of labels assessed in a linguistic term set S , i.e., the result of a symbolic aggregation operation. $\beta \in [0, g]$, being $g+1$ the cardinality of S . Let $i = \text{round}(\beta)$ and $\alpha = \beta - i$ be two values such that $i \in [0, g]$ and $\alpha \in [-0.5, 0.5]$ then α is called a symbolic translation, where $\text{round}(\cdot)$ is the usual round operation.

Definition 2.2 [14,41]. Let $S = \{s_i | i = 0, 1, 2, \dots, g\}$ be a linguistic term set and $\beta \in [0, g]$ a value representing the result of a symbolic aggregation operation. Then, the 2-tuple that expresses the equivalent information to β is obtained with the following function:

$$\Delta: [0, g] \rightarrow S \times [-0.5, 0.5] \quad (1)$$

$$\Delta(\beta) = (s_i, \alpha), \quad \text{with} \begin{cases} s_i, & i = \text{round}(\beta) \\ \alpha = \beta - i, & \alpha \in [-0.5, 0.5] \end{cases} \quad (2)$$

where s_i has the closest index label to β and α is the value of the symbolic translation.

Theorem 2.1 [14,41]. Let $S = \{s_i | i = 0, 1, 2, \dots, g\}$ be a linguistic term set and (s_i, α) be a 2-tuple. There is always a Δ^{-1} function such that from a 2-tuple it returns its equivalent numerical value $\beta \in [0, g] \subset R$, where

$$\Delta^{-1} : S \times [-0.5, 0.5) \rightarrow [0, g] \quad (3)$$

$$\Delta^{-1}(s_i, \alpha) = i + \alpha = \beta. \quad (4)$$

It is obvious that the conversion of a linguistic term into a 2-tuple consists of adding a value zero as symbolic translation

$$s_i \in S \Rightarrow (s_i, 0).$$

Definition 2.3 [14,41]. The comparison of linguistic information represented by 2-tuples is carried out according to an ordinary lexicographic order. Let (s_k, α_k) and (s_l, α_l) be two 2-tuples, with each one representing a counting of information as follows.

(1) If $k < l$ then (s_k, α_k) is smaller than (s_l, α_l) .

(2) If $k = l$ then

- if $\alpha_k = \alpha_l$ then (s_k, α_k) , (s_l, α_l) represents the same information;
- if $\alpha_k < \alpha_l$ then (s_k, α_k) is smaller than (s_l, α_l) ;
- if $\alpha_k > \alpha_l$ then (s_k, α_k) is bigger than (s_l, α_l) .

Theorem 2.2. Let (s_k, α_k) and (s_l, α_l) be two 2-tuples, $\beta_k = \Delta^{-1}(s_k, \alpha_k)$, and $\beta_l = \Delta^{-1}(s_l, \alpha_l)$. Then, $(s_k, \alpha_k) < (s_l, \alpha_l)$ if and only if $\beta_k < \beta_l$, and $(s_k, \alpha_k) = (s_l, \alpha_l)$ if and only if $\beta_k = \beta_l$.

Proof. (1) We first prove that $(s_k, \alpha_k) < (s_l, \alpha_l)$ if and only if $\beta_k < \beta_l$. Assume that $(s_k, \alpha_k) < (s_l, \alpha_l)$. Then, $k < l$, or $k = l$ and $\alpha_k < \alpha_l$. If $k < l$, then we have $\beta_k = k + \alpha_k < k + 0.5 \leq l - 0.5 \leq l + \alpha_l = \beta_l$. If $k = l$ and $\alpha_k < \alpha_l$, then we have $\beta_k = k + \alpha_k < l + \alpha_l = \beta_l$.

Assume that $\beta_k < \beta_l$. Then, $k < l$, or $k = l$ and $\alpha_k < \alpha_l$. If $k < l$, then we have $(s_k, \alpha_k) < (s_l, \alpha_l)$. If $k = l$ and $\alpha_k < \alpha_l$, then we have $(s_k, \alpha_k) < (s_l, \alpha_l)$.

(2) We next prove that $(s_k, \alpha_k) = (s_l, \alpha_l)$ if and only if $\beta_k = \beta_l$. If $(s_k, \alpha_k) = (s_l, \alpha_l)$, then $k = l$ and $\alpha_k = \alpha_l$, which implies that $\beta_k = k + \alpha_k = l + \alpha_l = \beta_l$. If $\beta_k = \beta_l$, then $k = l$ and $\alpha_k = \alpha_l$, which implies that $(s_k, \alpha_k) = (s_l, \alpha_l)$.

2.2. Bonferroni mean and geometric Bonferroni mean

Bonferroni [25] originally introduced a mean type aggregation operator, called Bonferroni mean, which can provide for aggregation lying between the max, min operators and the logical "or" and "and" operators.

Definition 2.4 [25]. Let $p, q \geq 0$, and let $a_i (i = 1, 2, \dots, n)$ be a collection of non-negative real numbers. Then, the aggregation function:

$$B^{p,q}(a_1, a_2, \dots, a_n) = \left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n a_i^p a_j^q \right)^{\frac{1}{p+q}} \quad (5)$$

is called a Bonferroni mean (BM).

Based on the usual geometric mean (GM) and the BM, Xia et al. [35] introduced the geometric Bonferroni mean, which was defined as follows:

Definition 2.5 [35]. Let $p, q \geq 0$, and $a_i (i=1, 2, \dots, n)$ be a collection of non-negative numbers. If

$$GB^{p,q}(a_1, a_2, \dots, a_n) = \frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (pa_i + qa_j)^{\frac{1}{n(n-1)}} \quad (6)$$

then we call $GB^{p,q}$ the geometric Bonferroni mean (GBM).

3. 2-TUPLE LINGUISTIC BONFERRONI MEAN AND WEIGHTED 2-TUPLE LINGUISTIC BONFERRONI MEAN

In this section, we first extend the Bonferroni mean operator (Eq. (5)) to 2-tuple linguistic environment, i.e., develop a 2-tuple linguistic Bonferroni mean operator and its weight form.

Definition 3.1. Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ ($r_i \in S, \alpha_i \in [-0.5, 0.5], i=1, 2, \dots, n$) be a collection of 2-tuples. If

$$2TLB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n ((\Delta^{-1}(r_i, \alpha_i))^p \cdot (\Delta^{-1}(r_j, \alpha_j))^q) \right)^{\frac{1}{p+q}} \right), \quad (7)$$

then $2TLB^{p,q}$ is called the 2-tuple linguistic Bonferroni mean (2TLBM).

In what follows, we investigate some desirable properties of the 2TLBM:

Theorem 3.1. Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let

$\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ ($r_i \in S, \alpha_i \in [-0.5, 0.5], i=1, 2, \dots, n$) be a collection of 2-tuples.

Then, the following properties hold.

(1) Commutativity: If $\{(r'_1, \alpha'_1), (r'_2, \alpha'_2), \dots, (r'_n, \alpha'_n)\}$ is any permutation of $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$, then

$$2TLB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = 2TLB^{p,q}((r'_1, \alpha'_1), (r'_2, \alpha'_2), \dots, (r'_n, \alpha'_n)). \quad (8)$$

(2) Idempotency: If $(r_i, \alpha_i) = (r, \alpha)$ for all i , then

$$2TLB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = (r, \alpha). \quad (9)$$

(3) Boundedness:

$$\min_{1 \leq i \leq n} \{(r_i, \alpha_i)\} \leq 2TLB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \leq \max_{1 \leq i \leq n} \{(r_i, \alpha_i)\}. \quad (10)$$

(4) Monotonicity: Let $\{(\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n)\}$ and $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ be two collections of 2-tuples, if $(r_i, \alpha_i) \leq (\bar{r}_i, \bar{\alpha}_i)$, for all i , then

$$2TLB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \leq 2TLB^{p,q}((\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n)). \quad (11)$$

Proof. (1) Since $\{(r'_1, \alpha'_1), (r'_2, \alpha'_2), \dots, (r'_n, \alpha'_n)\}$ is any permutation of $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$, we have

$$\begin{aligned}
& 2\text{TLB}^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\
198 \quad &= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r_i, \alpha_i))^p \cdot (\Delta^{-1}(r_j, \alpha_j))^q \right) \right)^{\frac{1}{p+q}} \right) \\
&= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r'_i, \alpha'_i))^p \cdot (\Delta^{-1}(r'_j, \alpha'_j))^q \right) \right)^{\frac{1}{p+q}} \right) \\
&= 2\text{TLB}^{p,q}((r'_1, \alpha'_1), (r'_2, \alpha'_2), \dots, (r'_n, \alpha'_n))
\end{aligned}$$

199 (2) If $(r_i, \alpha_i) = (r, \alpha)$ for all i , then

$$\begin{aligned}
& 2\text{TLB}^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\
&= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r_i, \alpha_i))^p \cdot (\Delta^{-1}(r_j, \alpha_j))^q \right) \right)^{\frac{1}{p+q}} \right) \\
200 \quad &= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r, \alpha))^p \cdot (\Delta^{-1}(r, \alpha))^q \right) \right)^{\frac{1}{p+q}} \right) \\
&= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r, \alpha))^{p+q} \right) \right)^{\frac{1}{p+q}} \right) = (r, \alpha)
\end{aligned}$$

201 (3) Because $\min_{1 \leq i \leq n} \{(r_i, \alpha_i)\} \leq 2\text{TLB}^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \leq \max_{1 \leq i \leq n} \{(r_i, \alpha_i)\}$, we have

$$\begin{aligned}
\min_{1 \leq i \leq n} \{(r_i, \alpha_i)\} &= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(\min_{1 \leq i \leq n} \{(r_i, \alpha_i)\}))^p \cdot (\Delta^{-1}(\min_{1 \leq i \leq n} \{(r_i, \alpha_i)\}))^q \right) \right)^{\frac{1}{p+q}} \right) \\
202 \quad &\leq \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r_i, \alpha_i))^p \cdot (\Delta^{-1}(r_j, \alpha_j))^q \right) \right)^{\frac{1}{p+q}} \right) \\
&\leq \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(\max_{1 \leq i \leq n} \{(r_i, \alpha_i)\}))^p \cdot (\Delta^{-1}(\max_{1 \leq i \leq n} \{(r_i, \alpha_i)\}))^q \right) \right)^{\frac{1}{p+q}} \right) \\
&= \max_{1 \leq i \leq n} \{(r_i, \alpha_i)\}.
\end{aligned}$$

203 (4)

$$\begin{aligned}
2\text{TLB}^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) &= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r_i, \alpha_i))^p \cdot (\Delta^{-1}(r_j, \alpha_j))^q \right) \right)^{\frac{1}{p+q}} \right) \\
&\leq \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(\bar{r}_i, \bar{\alpha}_i))^p \cdot (\Delta^{-1}(\bar{r}_j, \bar{\alpha}_j))^q \right) \right)^{\frac{1}{p+q}} \right) \\
&= 2\text{TLB}^{p,q}((\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n))
\end{aligned}$$

The proof of Theorem 3.1 is complete.

Theorem 3.2. Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ ($r_i \in S, \alpha_i \in [-0.5, 0.5], i = 1, 2, \dots, n$) be a collection of 2-tuples. Then, we have

$$2\text{TLB}^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = 2\text{TLB}^{q,p}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)). \quad (12)$$

Proof.

$$\begin{aligned}
2\text{TLB}^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) &= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r_i, \alpha_i))^p \cdot (\Delta^{-1}(r_j, \alpha_j))^q \right) \right)^{\frac{1}{p+q}} \right) \\
&= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{j,i=1 \\ j \neq i}}^n \left((\Delta^{-1}(r_j, \alpha_j))^q \cdot (\Delta^{-1}(r_i, \alpha_i))^p \right) \right)^{\frac{1}{q+p}} \right) \\
&= 2\text{TLB}^{q,p}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n))
\end{aligned} \quad (13)$$

The proof of Theorem 3.2 is complete.

In the following, let us consider some special cases of the 2TLBM operator by taking different values of the parameters p and q .

Case1. If $q \rightarrow 0$, then, by Eq. (7), we have

$$\begin{aligned}
& \lim_{q \rightarrow 0} 2TLB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\
&= \lim_{q \rightarrow 0} \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r_i, \alpha_i))^p \cdot (\Delta^{-1}(r_j, \alpha_j))^q \right) \right)^{\frac{1}{p+q}} \right) \\
216 \quad &= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r_i, \alpha_i))^p \right)^{\frac{1}{p}} \right)^{\frac{1}{p}} \right) = \Delta \left(\left(\frac{1}{n(n-1)} \sum_{i=1}^n \left((n-1) (\Delta^{-1}(r_i, \alpha_i))^p \right)^{\frac{1}{p}} \right)^{\frac{1}{p}} \right) \quad (14) \\
&= \Delta \left(\left(\frac{1}{n} \sum_{i=1}^n (\Delta^{-1}(r_i, \alpha_i))^p \right)^{\frac{1}{p}} \right) = 2TLB^{p,0}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n))
\end{aligned}$$

217 which we call the generalized 2-tuple linguistic mean [21].

218 **Case 2.** If $p = 2$ and $q \rightarrow 0$, then Eq. (7) is transformed as:

$$\begin{aligned}
219 \quad & 2TLB^{2,0}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r_i, \alpha_i))^2 \right)^{\frac{1}{2}} \right)^{\frac{1}{2}} \right) \\
&= \Delta \left(\left(\frac{1}{n} \sum_{i=1}^n (\Delta^{-1}(r_i, \alpha_i))^2 \right)^{\frac{1}{2}} \right) \quad (15)
\end{aligned}$$

220 which we call the 2-tuple linguistic square mean [21].

221 **Case 3.** If $p = 1$ and $q \rightarrow 0$, then Eq. (7) reduces to the 2-tuple linguistic average [21]:

$$\begin{aligned}
222 \quad & 2TLB^{1,0}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r_i, \alpha_i))^1 \right)^1 \right)^1 \right) \\
&= \Delta \left(\frac{1}{n} \sum_{i=1}^n \Delta^{-1}(r_i, \alpha_i) \right) \quad (16)
\end{aligned}$$

223 **Case 4.** If $p = q = 1$, then Eq. (7) reduces to the following:

$$224 \quad 2TLB^{1,1}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left(\Delta^{-1}(r_i, \alpha_i) \cdot \Delta^{-1}(r_j, \alpha_j) \right) \right)^{\frac{1}{2}} \right) \quad (17)$$

225 which we call the 2-tuple linguistic interrelated square mean.

226 **Case 5.** If $p \rightarrow +\infty$ and $q \rightarrow 0$, then Eq. (7) reduces to the 2-tuple linguistic maximum
227 operator:

228

$$\begin{aligned}
& \lim_{\substack{p \rightarrow +\infty \\ q \rightarrow 0}} 2TLB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\
&= \lim_{p \rightarrow +\infty} \left(\lim_{q \rightarrow 0} \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((\Delta^{-1}(r_i, \alpha_i))^p \cdot (\Delta^{-1}(r_j, \alpha_j))^q \right) \right)^{\frac{1}{p+q}} \right) \right) \\
&= \lim_{p \rightarrow +\infty} \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n (\Delta^{-1}(r_i, \alpha_i))^p \right)^{\frac{1}{p}} \right) \tag{18} \\
&= \lim_{p \rightarrow +\infty} \Delta \left(\left(\frac{1}{n(n-1)} \sum_{i=1}^n ((n-1)(\Delta^{-1}(r_i, \alpha_i))^p) \right)^{\frac{1}{p}} \right) = \lim_{p \rightarrow +\infty} \Delta \left(\left(\frac{1}{n} \sum_{i=1}^n (\Delta^{-1}(r_i, \alpha_i))^p \right)^{\frac{1}{p}} \right) \\
&= \lim_{p \rightarrow 0} \Delta \left(\max_{1 \leq i \leq n} \{ \Delta^{-1}(r_i, \alpha_i) \} \right) = 2TLB^{+\infty, 0}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n))
\end{aligned}$$

230 In the 2TLBM operator, the importance of the input arguments is not emphasized.
 231 Nevertheless, in many practical situations, the weights of the input arguments should be
 232 taken into account. If we allow the arguments to have different weights, then the weighted 2-
 233 tuple linguistic Bonferroni mean (W2TLBM) operator can be defined as follows:

234 **Definition 3.2.** Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let
 235 $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ ($r_i \in S, \alpha_i \in [-0.5, 0.5], i = 1, 2, \dots, n$) be a collection of 2-tuples.
 236 $w = (w_1, w_2, \dots, w_n)^T$ is the weight vector of $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$, where w_i indicates
 237 the importance degree of (r_i, α_i) , satisfying $w_i \in [0, 1]$ ($i = 1, 2, \dots, n$) and $\sum_{i=1}^n w_i = 1$. If

$$\begin{aligned}
& W2TLB_w^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\
&= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((w_i \cdot \Delta^{-1}(r_i, \alpha_i))^p \cdot (w_j \cdot \Delta^{-1}(r_j, \alpha_j))^q \right) \right)^{\frac{1}{p+q}} \right), \tag{19}
\end{aligned}$$

239 then $W2TLB_w^{p,q}$ is called the weighted 2-tuple linguistic Bonferroni mean (W2TLBM).

240 **Theorem 3.3.** Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let
 241 $\{(\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n)\}$ and $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ be two collections of 2-tuples,
 242 if $(r_i, \alpha_i) \leq (\bar{r}_i, \bar{\alpha}_i)$, for all i , then

$$243 W2TLB_w^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \leq W2TLB_w^{p,q}((\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n)). \tag{20}$$

244 **Proof.**

$$\begin{aligned}
& \text{W2TLB}_w^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\
&= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((w_i \cdot \Delta^{-1}(r_i, \alpha_i))^p \cdot (w_j \cdot \Delta^{-1}(r_j, \alpha_j))^q \right) \right)^{\frac{1}{p+q}} \right) \\
&\leq \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((w_i \cdot \Delta^{-1}(\bar{r}_i, \bar{\alpha}_i))^p \cdot (w_j \cdot \Delta^{-1}(\bar{r}_j, \bar{\alpha}_j))^q \right) \right)^{\frac{1}{p+q}} \right) \\
&= \text{W2TLB}_w^{p,q}((\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n))
\end{aligned}$$

246 The proof of Theorem 3.3 is complete.

247 The W2TLBM operator is neither idempotent, bounded, nor commutative.

248 **Theorem 3.4.** Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let

249 $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ ($r_i \in S, \alpha_i \in [-0.5, 0.5], i = 1, 2, \dots, n$) be a collection of 2-tuples.

250 $w = (w_1, w_2, \dots, w_n)^T$ is the weight vector of $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$, satisfying $w_i \in [0, 1]$

251 ($i = 1, 2, \dots, n$) and $\sum_{i=1}^n w_i = 1$. Then, we have

$$252 \quad \text{W2TLB}_w^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = \text{W2TLB}_w^{q,p}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)). \quad (21)$$

253 **Proof.**

$$\begin{aligned}
& \text{W2TLB}_w^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\
&= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \left((w_i \cdot \Delta^{-1}(r_i, \alpha_i))^p \cdot (w_j \cdot \Delta^{-1}(r_j, \alpha_j))^q \right) \right)^{\frac{1}{p+q}} \right) \\
&= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{j,i=1 \\ j \neq i}}^n \left((w_j \cdot \Delta^{-1}(r_j, \alpha_j))^q \cdot (w_i \cdot \Delta^{-1}(r_i, \alpha_i))^p \right) \right)^{\frac{1}{q+p}} \right) \\
&= \text{W2TLB}_w^{q,p}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n))
\end{aligned}$$

255 The proof of Theorem 3.4 is complete.

256

257 4. 2-TUPLE LINGUISTIC GEOMETRIC BONFERRONI MEAN AND WEIGHTED 2- 258 TUPLE LINGUISTIC GEOMETRIC BONFERRONI MEAN

259

260 In this section, we shall investigate the geometric Bonferroni mean under 2-tuple linguistic
261 environments, i.e., develop a 2-tuple linguistic geometric Bonferroni mean operator and its
262 weight form.

Definition 4.1. Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ ($r_i \in S, \alpha_i \in [-0.5, 0.5], i = 1, 2, \dots, n$) be a collection of 2-tuples. If

$$2TLGB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(r_i, \alpha_i) + q \cdot \Delta^{-1}(r_j, \alpha_j))^{\frac{1}{n(n-1)}} \right), \quad (22)$$

then $2TLGB^{p,q}$ is called the 2-tuple linguistic geometric Bonferroni mean (2TLGBM).

In what follows, we investigate some desirable properties of the 2TLGBM:

Theorem 4.1. Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ ($r_i \in S, \alpha_i \in [-0.5, 0.5], i = 1, 2, \dots, n$) be a collection of 2-tuples. Then, the following properties hold.

(1) Commutativity: If $\{(r'_1, \alpha'_1), (r'_2, \alpha'_2), \dots, (r'_n, \alpha'_n)\}$ is any permutation of $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$, then

$$2TLGB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = 2TLGB^{p,q}((r'_1, \alpha'_1), (r'_2, \alpha'_2), \dots, (r'_n, \alpha'_n)). \quad (23)$$

(2) Idempotency: If $(r_i, \alpha_i) = (r, \alpha)$ for all i , then

$$2TLGB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = (r, \alpha). \quad (24)$$

(3) Boundedness:

$$\min_{1 \leq i \leq n} \{(r_i, \alpha_i)\} \leq 2TLGB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \leq \max_{1 \leq i \leq n} \{(r_i, \alpha_i)\}. \quad (25)$$

(4) Monotonicity: Let $\{(\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n)\}$ and $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ be two collections of 2-tuples, if $(r_i, \alpha_i) \leq (\bar{r}_i, \bar{\alpha}_i)$, for all i , then

$$2TLGB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \leq 2TLGB^{p,q}((\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n)). \quad (26)$$

Proof. (1) Since $\{(r'_1, \alpha'_1), (r'_2, \alpha'_2), \dots, (r'_n, \alpha'_n)\}$ is any permutation of $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$, then we have

$$\begin{aligned} & 2TLGB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\ &= \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(r_i, \alpha_i) + q \cdot \Delta^{-1}(r_j, \alpha_j))^{\frac{1}{n(n-1)}} \right) \\ &= \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(r'_i, \alpha'_i) + q \cdot \Delta^{-1}(r'_j, \alpha'_j))^{\frac{1}{n(n-1)}} \right) \\ &= 2TLGB^{p,q}((r'_1, \alpha'_1), (r'_2, \alpha'_2), \dots, (r'_n, \alpha'_n)) \end{aligned}$$

(2) If $(r_i, \alpha_i) = (r, \alpha)$ for all i , then

285

$$\begin{aligned}
2\text{TLGB}^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) &= \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(r_i, \alpha_i) + q \cdot \Delta^{-1}(r_j, \alpha_j))^{\frac{1}{n(n-1)}} \right) \\
&= \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(r, \alpha) + q \cdot \Delta^{-1}(r, \alpha))^{\frac{1}{n(n-1)}} \right) \\
&= (r, \alpha)
\end{aligned}$$

286

(3) Because $\min_{1 \leq i \leq n} \{(r_i, \alpha_i)\} \leq 2\text{TLPG}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \leq \max_{1 \leq i \leq n} \{(r_i, \alpha_i)\}$, we have

287

$$\begin{aligned}
\min_{1 \leq i \leq n} \{(r_i, \alpha_i)\} &= \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(\min_{1 \leq i \leq n} \{(r_i, \alpha_i)\}) + q \cdot \Delta^{-1}(\min_{1 \leq i \leq n} \{(r_i, \alpha_i)\}))^{\frac{1}{n(n-1)}} \right) \\
&\leq \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(r_i, \alpha_i) + q \cdot \Delta^{-1}(r_j, \alpha_j))^{\frac{1}{n(n-1)}} \right) \tag{4} \\
&\leq \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(\max_{1 \leq i \leq n} \{(r_i, \alpha_i)\}) + q \cdot \Delta^{-1}(\max_{1 \leq i \leq n} \{(r_i, \alpha_i)\}))^{\frac{1}{n(n-1)}} \right) \\
&= \max_{1 \leq i \leq n} \{(r_i, \alpha_i)\}.
\end{aligned}$$

288

289

$$\begin{aligned}
2\text{TLGB}^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) &= \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(r_i, \alpha_i) + q \cdot \Delta^{-1}(r_j, \alpha_j))^{\frac{1}{n(n-1)}} \right) \\
&\leq \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(\bar{r}_i, \bar{\alpha}_i) + q \cdot \Delta^{-1}(\bar{r}_j, \bar{\alpha}_j))^{\frac{1}{n(n-1)}} \right) \\
&= 2\text{TLGB}^{p,q}((\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n))
\end{aligned}$$

290

The proof of Theorem 4.1 is complete.

291

Theorem 4.2. Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let

292

$\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ ($r_i \in S, \alpha_i \in [-0.5, 0.5], i = 1, 2, \dots, n$) be a collection of 2-tuples.

293

Then, we have

294

$$2\text{TLGB}^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = 2\text{TLGB}^{q,p}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)). \tag{27}$$

295

Proof.

296

$$\begin{aligned}
2\text{TLGB}^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) &= \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(r_i, \alpha_i) + q \cdot \Delta^{-1}(r_j, \alpha_j))^{\frac{1}{n(n-1)}} \right) \\
&= \Delta \left(\frac{1}{q+p} \prod_{\substack{j,i=1 \\ j \neq i}}^n (q \cdot \Delta^{-1}(r_j, \alpha_j) + p \cdot \Delta^{-1}(r_i, \alpha_i))^{\frac{1}{n(n-1)}} \right) \\
&= 2\text{TLGB}^{q,p}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n))
\end{aligned}$$

297 The proof of Theorem 4.2 is complete.

298 In the following, let us consider some special cases of the 2TLGBM operator by taking
299 different values of the parameters p and q .

300 **Case1.** If $q \rightarrow 0$, then, by Eq. (22), we have

$$\begin{aligned}
 & \lim_{q \rightarrow 0} 2TLGB^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\
 301 &= \lim_{q \rightarrow 0} \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(r_i, \alpha_i) + q \cdot \Delta^{-1}(r_j, \alpha_j))^{\frac{1}{n(n-1)}} \right) \quad (28) \\
 &= \Delta \left(\frac{1}{p} \prod_{\substack{i,j=1 \\ i \neq j}}^n (p \cdot \Delta^{-1}(r_i, \alpha_i))^{\frac{1}{n(n-1)}} \right) = \Delta \left(\frac{1}{p} \prod_{i=1}^n (p \cdot \Delta^{-1}(r_i, \alpha_i))^{\frac{1}{n}} \right) \\
 &= 2TLGB^{p,0}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n))
 \end{aligned}$$

302 which we call the generalized 2-tuple linguistic geometric mean [21].

303 **Case 2.** If $p = 2$ and $q \rightarrow 0$, then Eq. (22) is transformed as:

$$\begin{aligned}
 304 \quad 2TLGB^{2,0}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) &= \Delta \left(\frac{1}{2} \prod_{\substack{i,j=1 \\ i \neq j}}^n (2 \cdot \Delta^{-1}(r_i, \alpha_i))^{\frac{1}{n(n-1)}} \right) \quad (29) \\
 &= \Delta \left(\frac{1}{2} \prod_{i=1}^n (2 \cdot \Delta^{-1}(r_i, \alpha_i))^{\frac{1}{n}} \right)
 \end{aligned}$$

305 which we call the 2-tuple linguistic square geometric mean [21].

306 **Case 3.** If $p = 1$ and $q \rightarrow 0$, then Eq. (22) reduces to the 2-tuple linguistic geometric
307 average [25]:

$$\begin{aligned}
 308 \quad 2TLGB^{1,0}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) &= \Delta \left(\prod_{\substack{i,j=1 \\ i \neq j}}^n (\Delta^{-1}(r_i, \alpha_i))^{\frac{1}{n(n-1)}} \right) \quad (30) \\
 &= \Delta \left(\prod_{i=1}^n (\Delta^{-1}(r_i, \alpha_i))^{\frac{1}{n}} \right)
 \end{aligned}$$

309 **Case 4.** If $p = q = 1$, then Eq. (22) reduces to the following:

$$310 \quad 2TLGB^{1,1}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = \Delta \left(\frac{1}{2} \prod_{\substack{i,j=1 \\ i \neq j}}^n (\Delta^{-1}(r_i, \alpha_i) + \Delta^{-1}(r_j, \alpha_j))^{\frac{1}{n(n-1)}} \right) \quad (31)$$

312 which we call the 2-tuple linguistic interrelated square geometric mean.

313 It should be noted that the 2TLGBM operator does not consider the importance of the
314 aggregated arguments, but in many practical problem, especially in some group decision
315 makings, the aggregated arguments have different weights, to overcome this drawback, we
316 introduce the following definition:

317 **Definition 4.2.** Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let

318 $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ ($r_i \in S, \alpha_i \in [-0.5, 0.5], i = 1, 2, \dots, n$) be a collection of 2-tuples.

319 $w = (w_1, w_2, \dots, w_n)^T$ is the weight vector of $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$, where w_i indicates
 320 the importance degree of (r_i, α_i) , satisfying $w_i \in [0, 1]$ ($i = 1, 2, \dots, n$) and $\sum_{i=1}^n w_i = 1$. If

$$\begin{aligned} & \text{W2TLGB}_w^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\ 321 &= \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n \left(p \cdot (\Delta^{-1}(r_i, \alpha_i))^{w_i} + q \cdot (\Delta^{-1}(r_j, \alpha_j))^{w_j} \right)^{\frac{1}{n(n-1)}} \right), \end{aligned} \quad (32)$$

322 then $\text{W2TLGB}_w^{p,q}$ is called the weighted 2-tuple linguistic geometric Bonferroni mean
 323 (W2TLGBM).

324 **Theorem 4.3.** Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let
 325 $\{(\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n)\}$ and $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ be two collections of 2-tuples,
 326 if $(r_n, \alpha_n) \leq (\bar{r}_n, \bar{\alpha}_n)$, for all i , then

$$327 \quad \text{W2TLGB}_w^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \leq \text{W2TLGB}_w^{p,q}((\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n)). \quad (33)$$

328 **Proof.**

$$\begin{aligned} & \text{W2TLGB}_w^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\ &= \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n \left(p \cdot (\Delta^{-1}(r_i, \alpha_i))^{w_i} + q \cdot (\Delta^{-1}(r_j, \alpha_j))^{w_j} \right)^{\frac{1}{n(n-1)}} \right) \\ 329 &\leq \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n \left(p \cdot (\Delta^{-1}(\bar{r}_i, \bar{\alpha}_i))^{w_i} + q \cdot (\Delta^{-1}(\bar{r}_j, \bar{\alpha}_j))^{w_j} \right)^{\frac{1}{n(n-1)}} \right) \\ &= \text{W2TLGB}_w^{p,q}((\bar{r}_1, \bar{\alpha}_1), (\bar{r}_2, \bar{\alpha}_2), \dots, (\bar{r}_n, \bar{\alpha}_n)) \end{aligned}$$

330 The proof of Theorem 4.3 is complete.

331 The W2TLGBM operator is neither idempotent, bounded, nor commutative.

332 **Theorem 4.4.** Let $p \geq 0$, $q \geq 0$, and p, q do not take the value 0 simultaneously. Let
 333 $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$ ($r_i \in S, \alpha_i \in [-0.5, 0.5], i = 1, 2, \dots, n$) be a collection of 2-tuples.

334 $w = (w_1, w_2, \dots, w_n)^T$ is the weight vector of $\{(r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)\}$, satisfying $w_i \in [0, 1]$

335 ($i = 1, 2, \dots, n$) and $\sum_{i=1}^n w_i = 1$. Then, we have

$$336 \quad \text{W2TLGB}_w^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = \text{W2TLGB}_w^{q,p}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)). \quad (34)$$

337 **Proof.**

$$\begin{aligned}
& \text{W2TLGB}_w^{p,q}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) \\
&= \Delta \left(\frac{1}{p+q} \prod_{\substack{i,j=1 \\ i \neq j}}^n \left(p \cdot (\Delta^{-1}(r_i, \alpha_i))^{w_i} + q \cdot (\Delta^{-1}(r_j, \alpha_j))^{w_j} \right)^{\frac{1}{n(n-1)}} \right) \\
&= \Delta \left(\frac{1}{q+p} \prod_{\substack{j,i=1 \\ j \neq i}}^n \left(q \cdot (\Delta^{-1}(r_j, \alpha_j))^{w_j} + p \cdot (\Delta^{-1}(r_i, \alpha_i))^{w_i} \right)^{\frac{1}{n(n-1)}} \right) \\
&= \text{W2TLGB}_w^{q,p}((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n))
\end{aligned}$$

338

339 The proof of Theorem 4.4 is complete.

340

341 5. APPROACHES FOR MULTIPLE ATTRIBUTE GROUP DECISION MAKING 342 WITH 2-TUPLE LINGUISTIC INFORMATION

343

344 In this section, we utilize the proposed aggregation operators to develop some approaches
345 for multiple attribute group decision making with 2-tuple linguistic information.346 The multiple attribute group decision making (MAGDM) with 2-tuple linguistic information can
347 be formulated as follows: Let $X = \{x_1, x_2, \dots, x_m\}$ be a set of m alternatives, and let348 $C = \{c_1, c_2, \dots, c_n\}$ be a collection of n attributes, whose weight vector is349 $w = (w_1, w_2, \dots, w_n)^T$, with $w_j \in [0, 1]$, $j = 1, 2, \dots, n$, and $\sum_{j=1}^n w_j = 1$, where w_j denotes the350 importance degree of the attribute c_j , and let $D = \{d_1, d_2, \dots, d_l\}$ be a set of l decision351 makers, whose weight vector is $\omega = (\omega_1, \omega_2, \dots, \omega_l)^T$ with $\omega_k \in [0, 1]$, $k = 1, 2, \dots, l$, and352 $\sum_{k=1}^l \omega_k = 1$, where ω_k denotes the importance degree of the decision maker d_k . Each353 decision maker provides his/her own linguistic decision matrix $A^{(k)} = (a_{ij}^{(k)})_{m \times n}$ ($k = 1, 2, \dots, l$),354 where $a_{ij}^{(k)} \in S$ is a performance value, which takes the form of linguistic variable, given by355 the decision maker $d_k \in D$, for the alternative $x_i \in X$ with respect to the attribute $c_j \in C$.356 If all the attributes c_j ($j = 1, 2, \dots, n$) are of the same type, then the performance values do

357 not need normalization. Whereas there are, generally, benefit attributes (the bigger the

358 performance values the better) and cost attributes (the smaller the performance values the

359 better) in multiple attribute group decision making, in such cases, we may transform the

360 performance values of the cost type into the performance values of the benefit type. Then,

361 we can transform $A^{(k)} = (a_{ij}^{(k)})_{m \times n}$ into the matrix $R^{(k)} = (r_{ij}^{(k)})_{m \times n}$, where

$$r_{ij}^{(k)} = \begin{cases} a_{ij}^{(k)}, & \text{for benefit attribute } c_j \\ \text{neg}(a_{ij}^{(k)}), & \text{for cost attribute } c_j \end{cases}, \quad i = 1, 2, \dots, m, \quad j = 1, 2, \dots, n, \quad k = 1, 2, \dots, l \quad (35)$$

363 where $\text{neg}(a_{ij}^{(k)})$ is the complement of $a_{ij}^{(k)}$.

In the following, we utilize the W2TLBM (or W2TLGBM) operator to develop an approach for multiple attribute group decision making under 2-tuple linguistic environments. The algorithm involves the following steps.

Approach I

Step 1. Transform the linguistic decision matrix $A^{(k)} = (a_{ij}^{(k)})_{m \times n}$ into a normalized linguistic decision matrix $R^{(k)} = (r_{ij}^{(k)})_{m \times n}$ using Eq. (35).

Step 2. Transform the linguistic decision matrix $R^{(k)} = (r_{ij}^{(k)})_{m \times n}$ ($k=1,2,\dots,l$) into 2-tuple linguistic decision matrix $\bar{R}^{(k)} = ((r_{ij}^{(k)}, 0))_{m \times n}$ ($k=1,2,\dots,l$).

Step 3. Utilize the W2TLBM operator (Eq. (19)),

$$\begin{aligned} \bar{r}_{ij} &= (r_{ij}, \alpha_{ij}) = \text{W2TLB}_{\omega}^{p,q} \left((r_{ij}^{(1)}, 0), (r_{ij}^{(2)}, 0), \dots, (r_{ij}^{(l)}, 0) \right) \\ &= \Delta \left(\left(\frac{1}{l(l-1)} \sum_{\substack{s,t=1 \\ s \neq t}}^l \left((\omega_s \cdot \Delta^{-1}(r_{ij}^{(s)}, 0))^p \cdot (\omega_t \cdot \Delta^{-1}(r_{ij}^{(t)}, 0))^q \right) \right)^{\frac{1}{p+q}} \right) \end{aligned} \quad (36)$$

or the W2TLGBM operator (Eq. (32)),

$$\begin{aligned} \bar{r}_{ij} &= (r_{ij}, \alpha_{ij}) = \text{W2TLGB}_{\omega}^{p,q} \left((r_{ij}^{(1)}, 0), (r_{ij}^{(2)}, 0), \dots, (r_{ij}^{(l)}, 0) \right) \\ &= \Delta \left(\frac{1}{p+q} \prod_{\substack{s,t=1 \\ s \neq t}}^l \left(p \cdot (\Delta^{-1}(r_{ij}^{(s)}, 0))^{\omega_s} + q \cdot (\Delta^{-1}(r_{ij}^{(t)}, 0))^{\omega_t} \right)^{\frac{1}{l(l-1)}} \right) \end{aligned} \quad (37)$$

to aggregate all of the individual 2-tuple linguistic decision matrices $\bar{R}^{(k)} = ((r_{ij}^{(k)}, 0))_{m \times n}$ ($k=1,2,\dots,l$) into the collective 2-tuple linguistic decision matrix $\bar{R} = (\bar{r}_{ij})_{m \times n} = ((r_{ij}, \alpha_{ij}))_{m \times n}$.

Step 4. Utilize the W2TLBM operator (Eq. (19)),

$$\begin{aligned} \bar{r}_i &= (r_i, \alpha_i) = \text{W2TLB}_w^{p,q} \left((r_{i1}, \alpha_{i1}), (r_{i2}, \alpha_{i2}), \dots, (r_{in}, \alpha_{in}) \right) \\ &= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{x,y=1 \\ x \neq y}}^n \left((w_x \cdot \Delta^{-1}(r_{ix}, \alpha_{ix}))^p \cdot (w_y \cdot \Delta^{-1}(r_{iy}, \alpha_{iy}))^q \right) \right)^{\frac{1}{p+q}} \right) \end{aligned} \quad (38)$$

or the W2TLGBM operator (Eq. (32)),

$$\begin{aligned} \bar{r}_i &= (r_i, \alpha_i) = \text{W2TLGB}_w^{p,q} \left((r_{i1}, \alpha_{i1}), (r_{i2}, \alpha_{i2}), \dots, (r_{in}, \alpha_{in}) \right) \\ &= \Delta \left(\frac{1}{p+q} \prod_{\substack{x,y=1 \\ x \neq y}}^n \left(p \cdot (\Delta^{-1}(r_{ix}, \alpha_{ix}))^{w_x} + q \cdot (\Delta^{-1}(r_{iy}, \alpha_{iy}))^{w_y} \right)^{\frac{1}{n(n-1)}} \right) \end{aligned} \quad (39)$$

to aggregate all of the preference values $\bar{r}_{ij}$ ($j=1,2,\dots,n$) in the i th line of $\bar{R}$, and then derive the collective overall preference value $\bar{r}_i = (r_i, \alpha_i)$ ($i=1,2,\dots,m$) of alternative x_i ($i=1,2,\dots,m$).

385 **Step 5.** Rank the $\bar{r}_i = (r_i, \alpha_i)$ ($i=1,2,\dots,m$) in descending order using Definition 2.3.

386 **Step 6.** Rank all of the alternatives x_i ($i=1,2,\dots,m$), and then select the best one(s) in
 387 accordance with the collective overall preference values $\bar{r}_i = (r_i, \alpha_i)$ ($i=1,2,\dots,m$).

388 **Step 7.** End.

389 If the information regarding the weights of the decision makers and attributes is unknown,
 390 then we utilize the 2TLBM (or 2TLGBM) operator to develop an alternative approach for the
 391 MAGDM problem with 2-tuple linguistic information, which is described below.

392 **Approach II**

393 **Step 1.** Transform the linguistic decision matrix $A^{(k)} = (a_{ij}^{(k)})_{m \times n}$ into a normalized linguistic
 394 decision matrix $R^{(k)} = (r_{ij}^{(k)})_{m \times n}$ using Eq. (35).

395 **Step 2.** Transform the linguistic decision matrix $R^{(k)} = (r_{ij}^{(k)})_{m \times n}$ ($k=1,2,\dots,l$) into 2-tuple
 396 linguistic decision matrix $\bar{R}^{(k)} = ((r_{ij}^{(k)}, 0))_{m \times n}$ ($k=1,2,\dots,l$).

397 **Step 3.** Utilize the 2TLBM operator (Eq. (7)),

$$\begin{aligned} \bar{r}_{ij} = (r_{ij}, \alpha_{ij}) &= 2TLB_{\omega}^{p,q} \left((r_{ij}^{(1)}, 0), (r_{ij}^{(2)}, 0), \dots, (r_{ij}^{(l)}, 0) \right) \\ &= \Delta \left(\left(\frac{1}{l(l-1)} \sum_{\substack{s,t=1 \\ s \neq t}}^l \left((\Delta^{-1}(r_{ij}^{(s)}, 0))^p \cdot (\Delta^{-1}(r_{ij}^{(t)}, 0))^q \right) \right)^{\frac{1}{p+q}} \right) \end{aligned} \quad (40)$$

399 or the 2TLGBM operator (Eq. (22)),

$$\begin{aligned} \bar{r}_{ij} = (r_{ij}, \alpha_{ij}) &= 2TLGB_{\omega}^{p,q} \left((r_{ij}^{(1)}, 0), (r_{ij}^{(2)}, 0), \dots, (r_{ij}^{(l)}, 0) \right) \\ &= \Delta \left(\frac{1}{p+q} \prod_{\substack{s,t=1 \\ s \neq t}}^l \left(p \cdot \Delta^{-1}(r_{ij}^{(s)}, 0) + q \cdot \Delta^{-1}(r_{ij}^{(t)}, 0) \right)^{\frac{1}{l(l-1)}} \right) \end{aligned} \quad (41)$$

401 to aggregate all of the individual 2-tuple linguistic decision matrices $\bar{R}^{(k)} = ((r_{ij}^{(k)}, 0))_{m \times n}$
 402 ($k=1,2,\dots,l$) into the collective 2-tuple linguistic decision matrix $\bar{R} = (\bar{r}_{ij})_{m \times n} = ((r_{ij}, \alpha_{ij}))_{m \times n}$.

403 **Step 4.** Utilize the 2TLBM operator (Eq. (7)),

$$\begin{aligned} \bar{r}_i = (r_i, \alpha_i) &= 2TLB_w^{p,q} \left((r_{i1}, \alpha_{i1}), (r_{i2}, \alpha_{i2}), \dots, (r_{in}, \alpha_{in}) \right) \\ &= \Delta \left(\left(\frac{1}{n(n-1)} \sum_{\substack{x,y=1 \\ x \neq y}}^n \left((\Delta^{-1}(r_{ix}, \alpha_{ix}))^p \cdot (\Delta^{-1}(r_{iy}, \alpha_{iy}))^q \right) \right)^{\frac{1}{p+q}} \right) \end{aligned} \quad (42)$$

405 or the 2TLGBM operator (Eq. (22)),

$$\begin{aligned} \bar{r}_i = (r_i, \alpha_i) &= 2TLGB_w^{p,q} \left((r_{i1}, \alpha_{i1}), (r_{i2}, \alpha_{i2}), \dots, (r_{in}, \alpha_{in}) \right) \\ &= \Delta \left(\frac{1}{p+q} \prod_{\substack{x,y=1 \\ x \neq y}}^n \left(p \cdot \Delta^{-1}(r_{ix}, \alpha_{ix}) + q \cdot \Delta^{-1}(r_{iy}, \alpha_{iy}) \right)^{\frac{1}{n(n-1)}} \right) \end{aligned} \quad (43)$$

to aggregate all of the preference values $\bar{r}_{ij}$ ($j=1,2,\dots,n$) in the i th line of $\bar{R}$, and then derive the collective overall preference value $\bar{r}_i=(r_i, \alpha_i)$ ($i=1,2,\dots,m$) of alternative x_i ($i=1,2,\dots,m$).

Step 5. Rank the $\bar{r}_i=(r_i, \alpha_i)$ ($i=1,2,\dots,m$) in descending order using Definition 2.3.

Step 6. Rank all of the alternatives x_i ($i=1,2,\dots,m$), and then select the best one(s) in accordance with the collective overall preference values $\bar{r}_i=(r_i, \alpha_i)$ ($i=1,2,\dots,m$).

Step 7. End.

Remark 5.1. Approach I is designed for situations where the weights of the decision makers and attributes can be predefined and it utilizes the W2TLBM (or W2TLGBM) operator to aggregate all of the individual 2-tuple linguistic decision matrices into the collective 2-tuple linguistic decision matrix. Approach II is designed for situations where the information regarding the weights of the decision makers and attributes is unknown and it utilizes the 2TLBM (or 2TLGBM) operator to aggregate all of the individual 2-tuple linguistic decision matrices into the collective 2-tuple linguistic decision matrix.

6. ILLUSTRATIVE EXAMPLES

In this subsection, let us consider a numerical example adapted from Herrera et al. [48] and Herrera and Martínez [41].

Example 6.1 [41,48]. Suppose that an investment company wants to invest a sum of money in the best option. There is a panel with four possible alternatives in which to invest the money: (1) x_1 is a car industry; (2) x_2 is a food company; (3) x_3 is a computer company; and (4) x_4 is an arms industry. The investment company must make a decision according to the following four attributes: (1) c_1 is the risk analysis; (2) c_2 is the growth analysis; (3) c_3 is the social-political impact analysis; and (4) c_4 is the environmental impact analysis. Among the considered attributes, c_2 is the benefit attribute, and c_j ($j=1,3,4$) are the cost attributes. The weight vector of attributes c_j ($j=1,2,3,4$) is $w=(0.3,0.25,0.25,0.2)^T$. The four possible alternatives x_i ($i=1,2,3,4$) are to be evaluated using the linguistic term set

$$S = \left\{ \begin{array}{l} s_0 = \text{extremely poor}, s_1 = \text{very poor}, s_2 = \text{poor}, s_3 = \text{slightly poor}, s_4 = \text{fair}, \\ s_5 = \text{slightly good}, s_6 = \text{good}, s_7 = \text{very good}, s_8 = \text{extremely good} \end{array} \right\}$$

by three decision makers d_k ($k=1,2,3$) (suppose that the weight vector of three decision makers is $\omega=(0.2,0.5,0.3)^T$) under the above four attributes, and construct, respectively, the linguistic decision matrices $A^{(k)}=(a_{ij}^{(k)})_{4 \times 4}$ ($k=1,2,3$) as shown in Tables 1-3.

1	c_1	c_2	c_3	c_4
x_1	s_4	s_3	s_7	s_3
x_2	s_5	s_6	s_2	s_0
x_3	s_5	s_2	s_1	s_3
x_4	s_0	s_1	s_5	s_2

Table 1. Linguistic decision matrix $A^{(1)}$ provided by d_1 .

2	c_1	c_2	c_3	c_4
x_1	s_3	s_2	s_1	s_5
x_2	s_1	s_4	s_0	s_2
x_3	s_1	s_8	s_3	s_2
x_4	s_0	s_6	s_3	s_5

Table 2. Linguistic decision matrix $A^{(2)}$ provided by d_2 .

3	c_1	c_2	c_3	c_4
x_1	s_3	s_1	s_6	s_0
x_2	s_1	s_8	s_2	s_3
x_3	s_3	s_6	s_5	s_4
x_4	s_2	s_8	s_3	s_1

Table 3. Linguistic decision matrix $A^{(3)}$ provided by d_3 .

Assume that the weights of the decision makers and attributes are known. We use Approach I to find the decision result.

Step 1. Using Eq. (35), we transform the linguistic decision matrix $A^{(k)} = (a_{ij}^{(k)})_{4 \times 4}$ into a normalized linguistic matrix $R^{(k)} = (r_{ij}^{(k)})_{4 \times 4}$ (see Table 4, 5, and 6).

4	c_1	c_2	c_3	c_4
x_1	s_4	s_3	s_1	s_5
x_2	s_3	s_6	s_5	s_8
x_3	s_3	s_2	s_7	s_5
x_4	s_8	s_1	s_3	s_6

Table 4. Linguistic decision matrix $R^{(1)}$ provided by d_1 .

5	c_1	c_2	c_3	c_4
x_1	s_5	s_2	s_7	s_3
x_2	s_7	s_4	s_8	s_6
x_3	s_7	s_8	s_5	s_6
x_4	s_8	s_6	s_5	s_3

Table 5. Linguistic decision matrix $R^{(2)}$ provided by d_2 .

6	c_1	c_2	c_3	c_4
x_1	s_5	s_1	s_2	s_8
x_2	s_7	s_8	s_6	s_5
x_3	s_5	s_6	s_3	s_4
x_4	s_6	s_8	s_5	s_7

Table 6. Linguistic decision matrix $R^{(3)}$ provided by d_3 .

Step 2. Transform the linguistic decision matrices $R^{(k)} = (r_{ij}^{(k)})_{4 \times 4}$ ($k=1,2,3$) given in Tables 4-6 into 2-tuple linguistic decision matrices $\bar{R}^{(k)} = ((r_{ij}^{(k)}, 0))_{4 \times 4}$ ($k=1,2,3$) which are given in Tables 7-9.

7	c_1	c_2	c_3	c_4
x_1	$(s_4, 0)$	$(s_3, 0)$	$(s_1, 0)$	$(s_5, 0)$
x_2	$(s_3, 0)$	$(s_6, 0)$	$(s_5, 0)$	$(s_8, 0)$
x_3	$(s_3, 0)$	$(s_2, 0)$	$(s_7, 0)$	$(s_5, 0)$
x_4	$(s_8, 0)$	$(s_1, 0)$	$(s_3, 0)$	$(s_6, 0)$

Table 7. 2-tuple linguistic decision matrix $\bar{R}^{(1)}$.

8	c_1	c_2	c_3	c_4
x_1	$(s_5, 0)$	$(s_2, 0)$	$(s_7, 0)$	$(s_3, 0)$
x_2	$(s_7, 0)$	$(s_4, 0)$	$(s_8, 0)$	$(s_6, 0)$
x_3	$(s_7, 0)$	$(s_8, 0)$	$(s_5, 0)$	$(s_6, 0)$
x_4	$(s_8, 0)$	$(s_6, 0)$	$(s_5, 0)$	$(s_3, 0)$

Table 8. 2-tuple linguistic decision matrix $\bar{R}^{(2)}$.

9	c_1	c_2	c_3	c_4
x_1	$(s_5, 0)$	$(s_1, 0)$	$(s_2, 0)$	$(s_8, 0)$
x_2	$(s_7, 0)$	$(s_8, 0)$	$(s_6, 0)$	$(s_5, 0)$
x_3	$(s_5, 0)$	$(s_6, 0)$	$(s_3, 0)$	$(s_4, 0)$
x_4	$(s_6, 0)$	$(s_8, 0)$	$(s_5, 0)$	$(s_7, 0)$

Table 9. 2-tuple linguistic decision matrix $\bar{R}^{(3)}$.

Step 3. Use the W2TLBM operator (Eq. (36)) (here, we take $p = q = 1$) to aggregate all of the individual 2-tuple linguistic decision matrices $\bar{R}^{(k)} = \left((r_{ij}^{(k)}, 0) \right)_{4 \times 4}$ ($k = 1, 2, 3$) into the collective 2-tuple linguistic decision matrix $\bar{R} = \left(\bar{r}_{ij} \right)_{4 \times 4} = \left((r_{ij}, \alpha_{ij}) \right)_{4 \times 4}$ (see Table 10).

10	c_1	c_2	c_3	c_4
x_1	$(s_2, -0.4779)$	$(s_1, -0.4000)$	$(s_1, -0.0134)$	$(s_2, -0.4169)$
x_2	$(s_2, -0.1106)$	$(s_2, -0.1670)$	$(s_2, 0.0817)$	$(s_2, -0.0232)$
x_3	$(s_2, -0.3417)$	$(s_2, -0.2186)$	$(s_2, -0.4714)$	$(s_2, -0.3875)$
x_4	$(s_2, 0.3438)$	$(s_2, -0.3387)$	$(s_1, 0.4318)$	$(s_2, -0.4220)$

559

Table 10. The collective 2-tuple linguistic decision matrix $\bar{R}$.

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

Step 4. Use the W2TLBM operator (Eq. (38)) to aggregate all of the preference values (r_{ij}, α_{ij}) ($j = 1, 2, 3, 4$) in the i th line of $\bar{R}$ and then derive the collective overall preference value $\bar{r}_i = (r_i, \alpha_i)$ ($i = 1, 2, 3, 4$) of the alternative x_i ($i = 1, 2, 3, 4$).

$$\bar{r}_1 = (s_0, 0.2852), \quad \bar{r}_2 = (s_0, 0.4837), \quad \bar{r}_3 = (s_0, 0.4101), \quad \bar{r}_4 = (s_0, 0.4394).$$

Using Definition 2.3, we then rank the $\bar{r}_i$ ($i = 1, 2, 3, 4$) in descending order:

$$\bar{r}_2 > \bar{r}_4 > \bar{r}_3 > \bar{r}_1.$$

Step 5. Rank all of the alternatives x_i ($i = 1, 2, 3, 4$) as follows:

$$x_2 \succ x_4 \succ x_3 \succ x_1.$$

The best alternative is x_2 .

We can find that as the values of the parameters p and q change according to the decision makers' subjective preferences, we may obtain different rankings of the alternatives, which can reflect the decision makers' risk preferences. As the values of the parameters p and q change, the collective overall preference value $\bar{r}_i = (r_i, \alpha_i)$ ($i = 1, 2, 3, 4$) and the ranking of alternatives can be obtained and shown in Table 11.

578

11	$p = 0, q = 20$	$p = 0.5, q = 15$	$p = q = 5$	$p = 15, q = 0.1$	$p = 20, q = 0.05$
x_1	$(s_1, -0.2255)$	$(s_1, -0.3110)$	$(s_0, 0.3779)$	$(s_1, -0.2668)$	$(s_1, -0.2308)$
x_2	$(s_1, -0.0577)$	$(s_1, -0.1233)$	$(s_1, -0.4132)$	$(s_1, -0.0948)$	$(s_1, -0.0610)$
x_3	$(s_1, -0.0577)$	$(s_1, -0.1302)$	$(s_1, -0.4712)$	$(s_1, -0.0963)$	$(s_1, -0.0615)$
x_4	$(s_1, 0.0598)$	$(s_1, -0.0280)$	$(s_1, -0.4210)$	$(s_1, 0.0074)$	$(s_1, 0.0560)$
Ranking	$x_4 \succ x_2 \succ x_3 \succ x_1$	$x_4 \succ x_2 \succ x_3 \succ x_1$	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_4 \succ x_2 \succ x_3 \succ x_1$	$x_4 \succ x_2 \succ x_3 \succ x_1$

579

580

581

582

583

584

585

586


587

588

589


Table 11. The collective overall preference values obtained with the W2TLBM operator and rankings of the alternatives.

Furthermore, it is possible to analyze how the different attitudinal characters p and q play a role in the aggregation results. As the values of the parameters p and q change between 0 and 20, different results of a symbolic aggregation operation $\beta_i = \Delta^{-1}(\bar{r}_i)$ ($i=1,2,3,4$) of the collective overall preference values $\bar{r}_i$ ($i=1,2,3,4$) of the four alternatives x_i ($i=1,2,3,4$) can be obtained. Figs. 1-4 illustrate the values $\beta_i = \Delta^{-1}(\bar{r}_i)$ ($i=1,2,3,4$) of the four alternatives x_i ($i=1,2,3,4$) obtained by the W2TLBM operator in detail.




590


591 **Fig. 1.** The values β for alternative x_1 obtained by the W2TLBM operator ($p \in (0, 20]$,
 592 $q \in (0, 20]$).
 593
 594



595 **Fig. 2.** The values β for alternative x_2 obtained by the W2TLBM operator ($p \in (0, 20]$,
 596 $q \in (0, 20]$).
 597



598
599 **Fig. 3.** The values β for alternative x_3 obtained by the W2TLBM operator ($p \in (0,20]$,
600 $q \in (0,20]$).
601
602




603

604 **Fig. 4.** The values β for alternative x_4 obtained by the W2TLBM operator ($p \in (0, 20]$,
 605 $q \in (0, 20]$).

606

607 If we let the parameter p fixed, different values β_i and rankings of the alternatives can be
 608 obtained as the parameter q changed which was shown in Fig. 5.



609

610 **Fig. 5.** Variation of β obtained with the W2TLBM operator ($p = 1$, $q \in (0, 20]$).

611 From Fig. 5, we can find that,

612 1) when $q \in (0, 4.6560]$, the ranking of the four alternatives is $x_2 \succ x_4 \succ x_3 \succ x_1$ and the best
 613 choice is x_2 .

614 2) when $q \in (4.6560, 20]$, the ranking of the four alternatives is $x_4 \succ x_2 \succ x_3 \succ x_1$ and the best
 615 choice is x_4 .

616 From Fig. 5, we also find that the collective overall preference values obtained by the
 617 W2TLBM aggregation operators become bigger as parameter p and q increase for the
 618 same aggregation arguments. Therefore, it plays a crucial part in decision making. For
 619 example, in the real group decision making problems, the decision makers who takes a
 620 gloomy view of the prospects could choose the smaller values of the parameters p and q
 621 while the decision makers who are optimistic could choose the bigger values of the
 622 parameters p and q .

623 If the W2TLGBM operator is used in place of the W2TLBM operator to aggregate the values
 624 of the alternatives in steps 3 and 4, then the collective overall preference values and the
 625 rankings of the alternatives are listed in Table 12.

626

627

628

629

630

631

632


633

634


12	$p = 0, q = 20$	$p = 0.5, q = 15$	$p = q = 5$	$p = 15, q = 0.1$	$p = 20, q = 0.05$
x_1	$(s_1, 0.1057)$	$(s_1, 0.1066)$	$(s_1, 0.1130)$	$(s_1, 0.1059)$	$(s_1, 0.1057)$
x_2	$(s_1, 0.1604)$	$(s_1, 0.1614)$	$(s_1, 0.1677)$	$(s_1, 0.1606)$	$(s_1, 0.1605)$
x_3	$(s_1, 0.1464)$	$(s_1, 0.1476)$	$(s_1, 0.1558)$	$(s_1, 0.1466)$	$(s_1, 0.1465)$
x_4	$(s_1, 0.1477)$	$(s_1, 0.1488)$	$(s_1, 0.1560)$	$(s_1, 0.1479)$	$(s_1, 0.1478)$
Ranking	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_4 \succ x_2 \succ x_3 \succ x_1$	$x_4 \succ x_2 \succ x_3 \succ x_1$

Table 12. The collective overall preference values obtained with the W2TLGBM operator and rankings of the alternatives.


Furthermore, it is possible to analyze how the different attitudinal characters p and q play a role in the aggregation results. As the values of the parameters p and q change between 0 and 20, different results of a symbolic aggregation operation $\beta_i = \Delta^{-1}(\bar{r}_i)$ ($i=1,2,3,4$) of the collective overall preference values $\bar{r}_i$ ($i=1,2,3,4$) of the four alternatives x_i ($i=1,2,3,4$) can be obtained. Figs. 6-9 illustrate the values $\beta_i = \Delta^{-1}(\bar{r}_i)$ ($i=1,2,3,4$) of the four alternatives x_i ($i=1,2,3,4$) obtained by the W2TLGBM operator in detail.




647 **Fig. 6.** The values β for alternative x_1 obtained by the W2TLGBM operator ($p \in (0, 20]$,
 648 $q \in (0, 20]$).
 649



650 **Fig. 7.** The values β for alternative x_2 obtained by the W2TLGBM operator ($p \in (0, 20]$,
 651 $q \in (0, 20]$).
 652
 653




654 **Fig. 8.** The values β for alternative x_3 obtained by the W2TLGBM operator ($p \in (0, 20]$,
 655 $q \in (0, 20]$).
 656
 657



658 **Fig. 9.** The values β for alternative x_4 obtained by the W2TLGBM operator ($p \in (0, 20]$,
 659 $q \in (0, 20]$).
 660

661
 662 If we let the parameter p fixed, different values β_i and rankings of the alternatives can be
 663 obtained as the parameter q changed which was shown in Fig. 10.




664
 665 **Fig. 10.** Variation of β obtained with the W2TLGBM operator ($p = 1, q \in (0, 20]$).
 666

667 From Fig. 10, we can find that when $q \in (0, 20]$, the ranking of the four alternatives is
 668 $x_2 \succ x_4 \succ x_3 \succ x_1$ and the best choice is x_2 .


669 From Fig. 10, we find that the collective overall preference values obtained by the
 670 W2TLGBM aggregation operators become smaller as parameter p and q increase for the
 671 same aggregation arguments. Therefore, it plays a crucial part in decision making. For
 672 example, in the real group decision making problems, the decision makers who take a
 673 gloomy view of the prospects could choose the bigger values of the parameters p and q
 674 while the decision makers who are optimistic could choose the smaller values of the
 675 parameters p and q .

676 It should be noted that the beta obtained by the W2TLBM operator are smaller than the
 677 beta obtained by the W2TLGBM operator, which indicates that the W2TLBM operator can
 678 obtain more unfavorable (or pessimistic) expectations, while the W2TLGBM operator has
 679 more favorable (or optimistic) expectations. Therefore, we can conclude that the W2TLBM
 680 operator can be considered as the pessimistic operator, while the W2TLGBM operator can be
 681 considered as the optimistic operator and the values of the parameters can be considered
 682 as the optimistic or pessimistic levels. By Figs. 1-10, we can conclude that the decision
 683 makers who take a gloomy view of the prospects could use the W2TLBM operator and
 684 choose the smaller values of the parameters p and q , while the decision makers who are
 685 optimistic could use the W2TLGBM operator and choose the smaller values of the
 686 parameters p and q .

687 In order to obtain the more neutral results, we can use the arithmetic averages of the
 688 optimistic and pessimistic results, which can be found in Figs. 11-14.
 689




690
 691 **Fig. 11.** The values β for alternative x_1 obtained by the W2TLBM and W2TLGBM operators
 692 ($p \in (0, 20]$, $q \in (0, 20]$).
 693



694

695 **Fig. 12.** The values β for alternative x_2 obtained by the W2TLBM and W2TLGBM
 696 operators ($p \in (0, 20]$, $q \in (0, 20]$).
 697



698 **Fig. 13.** The values β for alternative x_3 obtained by the W2TLBM and W2TLGBM
 699 operators ($p \in (0, 20]$, $q \in (0, 20]$).
 700
 701




Fig. 14. The values β for alternative x_4 obtained by the W2TLBM and W2TLGBM operators ($p \in (0, 20]$, $q \in (0, 20]$).

In the following, we compare our operators and approaches with the existing 2-tuple linguistic aggregation operators and approaches so as to demonstrate the advantages of the operators and approaches proposed here. In Example 5.1, we utilize the 2-tuple linguistic weighted average (2TLWA) operator [14,41]:

$$2TLWA((r_1, \alpha_1), (r_2, \alpha_2), \dots, (r_n, \alpha_n)) = \Delta \left(\sum_{i=1}^n w_i \cdot \Delta^{-1}(r_i, \alpha_i) \right) \quad (44)$$

to replace the W2TLBM operator; then the collective overall preference value $\bar{r}_i = (r_i, \alpha_i)$ ($i=1, 2, 3, 4$) of alternative x_i ($i=1, 2, 3, 4$) are shown as follows:

$$\bar{r}_1 = (s_4, -0.0300), \quad \bar{r}_2 = (s_6, 0.1800), \quad \bar{r}_3 = (s_5, 0.4700), \quad \bar{r}_4 = (s_6, -0.2700).$$

Using Definition 2.3, we then rank the $\bar{r}_i$ ($i=1, 2, 3, 4$) in descending order:

$$\bar{r}_2 > \bar{r}_4 > \bar{r}_3 > \bar{r}_1.$$

According to the ranking of the $\bar{r}_i$ ($i=1, 2, 3, 4$), rank all of the alternatives x_i ($i=1, 2, 3, 4$) as follows:

$$x_2 \succ x_4 \succ x_3 \succ x_1.$$

Thus, the best alternative is x_2 .

Based on the aforementioned analysis, we can see that the 2TLWA operator is simpler than the W2TLBM and W2TLGBM operators from the computational point of view, but the W2TLBM and W2TLGBM operators can capture the interrelationship of the aggregated arguments [35] and can provide the decision makers more choices by changing the values of the parameters determined by the preferences of the decision makers [35].

Example 6.2. Let us reconsider Example 6.1. Assume that $A^{(k)} = (a_{ij}^{(k)})_{4 \times 4}$ ($k=1,2,3$) are three linguistic decision matrices shown in Tables 1-3. $R^{(k)} = (r_{ij}^{(k)})_{4 \times 4}$ ($k=1,2,3$) are three normalized linguistic decision matrices given in Tables 4-6. $\bar{R}^{(k)} = ((r_{ij}^{(k)}, 0))_{4 \times 4}$ ($k=1,2,3$) are three corresponding 2-tuple linguistic decision matrices given in Tables 7-9. Suppose that the weights of the decision makers and the attributes are unknown; then, we use Approach II to determine the decision. The collective 2-tuple linguistic decision matrix $\bar{R}' = (\bar{r}'_{ij})_{4 \times 4} = ((r'_{ij}, \alpha'_{ij}))_{4 \times 4}$ is given in Table 13. The best alternative is x_2 .

13	c_1	c_2	c_3	c_4
x_1	$(s_5, -0.3453)$	$(s_2, -0.0851)$	$(s_3, -0.2311)$	$(s_5, 0.1316)$
x_2	$(s_6, -0.4924)$	$(s_6, -0.1122)$	$(s_6, 0.2716)$	$(s_6, 0.2716)$
x_3	$(s_5, -0.1352)$	$(s_5, 0.0332)$	$(s_5, -0.1352)$	$(s_5, -0.0332)$
x_4	$(s_7, 0.3030)$	$(s_5, -0.4539)$	$(s_4, 0.2817)$	$(s_5, 0.1962)$

Table 13. The collective 2-tuple linguistic decision matrix $\bar{R}'$.


We can find that as the values of the parameters p and q change according to the decision makers' subjective preferences, we may obtain different rankings of the alternatives, which can reflect the decision makers' risk preferences. As the values of the parameters p and q change, the collective overall preference value $\bar{r}'_i = (r'_i, \alpha'_i)$ ($i=1,2,3,4$) and the ranking of alternatives can be obtained and shown in Table 14.

14	$p = 0, q = 20$	$p = 0.5, q = 15$	$p = q = 5$	$p = 15, q = 0.1$	$p = 20, q = 0.05$
x_1	$(s_7, 0.0891)$	$(s_7, -0.4043)$	$(s_4, 0.4483)$	$(s_7, -0.2153)$	$(s_7, 0.0684)$
x_2	$(s_7, 0.4823)$	$(s_7, 0.2594)$	$(s_6, 0.3807)$	$(s_7, 0.3246)$	$(s_7, 0.4752)$
x_3	$(s_7, 0.1133)$	$(s_7, -0.2625)$	$(s_6, -0.4656)$	$(s_7, -0.1379)$	$(s_7, 0.0981)$
x_4	$(s_7, 0.4740)$	$(s_7, 0.2032)$	$(s_6, 0.2012)$	$(s_7, 0.2934)$	$(s_7, 0.4625)$
Ranking	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_2 \succ x_4 \succ x_3 \succ x_1$


Table 14. The collective overall preference values obtained with the 2TLBM operator and rankings of the alternatives.

As the values of the parameters p and q change between 0 and 20, different results of a symbolic aggregation operation $\beta_i = \Delta^{-1}(\bar{r}'_i)$ ($i=1,2,3,4$) of the collective overall


758 preference values $\bar{r}_i$ ($i=1,2,3,4$) of the four alternatives x_i ($i=1,2,3,4$) can be obtained.
 759 Figs. 15-18 illustrate the values $\beta_i = \Delta^{-1}(\bar{r}_i')$ ($i=1,2,3,4$) of the four alternatives x_i
 760 ($i=1,2,3,4$) obtained by the 2TLBM operator in detail.
 761



762 **Fig. 15.** The values β for alternative x_1 obtained by the 2TLBM operator ($p \in (0,20]$,
 763 $q \in (0,20]$).
 764
 765
 766




767
 768 **Fig. 16.** The values β for alternative x_2 obtained by the 2TLBM operator ($p \in (0, 20]$,
 769 $q \in (0, 20]$).
 770




771

772 **Fig. 17.** The values β for alternative x_3 obtained by the 2TLBM operator ($p \in (0, 20]$,
 773 $q \in (0, 20]$).
 774



775 **Fig. 18.** The values β for alternative x_4 obtained by the 2TLBM operator ($p \in (0, 20]$,
 776 $q \in (0, 20]$).
 777
 778

779 If we let the parameter p fixed, different values β_i and rankings of the alternatives can be
 780 obtained as the parameter q changed which was shown in Fig. 19.



781

782 **Fig. 19.** Variation of β obtained with the 2TLBM operator ($p=1, q \in (0, 20]$).

783 From Fig. 19, we can find that when $q \in (0, 20]$, the ranking of the four alternatives is
784 $x_2 \succ x_4 \succ x_3 \succ x_1$ and the best choice is x_2 .

785 From Fig. 19, we find that the collective overall preference values obtained by the 2TLBM
786 aggregation operators become bigger as parameter p and q increase for the same
787 aggregation arguments. Therefore, it plays a crucial part in decision making. For example, in
788 the real group decision making problems, the decision makers who take a gloomy view of
789 the prospects could choose the smaller values of the parameters p and q while the
790 decision makers who are optimistic could choose the bigger values of the parameters p and
791 q .

792 If the 2TLGBM operator is used in place of the 2TLBM operator to aggregate the values of
793 the alternatives in steps 1 and 2, **then** the collective overall preference values and the
794 rankings of the alternatives are listed in Table 15.
795

15	$p = 0, q = 20$	$p = 0.5, q = 15$	$p = q = 5$	$p = 15, q = 0.1$	$p = 20, q = 0.05$
x_1	$(s_3, 0.1644)$	$(s_3, 0.2330)$	$(s_4, -0.3675)$	$(s_3, 0.1792)$	$(s_3, 0.1700)$
x_2	$(s_6, -0.1442)$	$(s_6, -0.1208)$	$(s_6, 0.0328)$	$(s_6, -0.1392)$	$(s_6, -0.1423)$
x_3	$(s_5, -0.2650)$	$(s_5, -0.2273)$	$(s_5, 0.0098)$	$(s_5, -0.2570)$	$(s_5, -0.2620)$
x_4	$(s_5, -0.1387)$	$(s_5, -0.0524)$	$(s_5, 0.3822)$	$(s_5, -0.1196)$	$(s_5, -0.1314)$
Ranking	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_2 \succ x_4 \succ x_3 \succ x_1$	$x_2 \succ x_4 \succ x_3 \succ x_1$

796
797 **Table 15.** The collective overall preference values obtained with the 2TLGBM operator and
798 rankings of the alternatives.
799

800 As the values of the parameters p and q change between 0 and 20, different results of a
801 symbolic aggregation operation $\beta_i = \Delta^{-1}(\bar{r}_i)$ ($i=1,2,3,4$) of the collective overall
802 preference values $\bar{r}_i$ ($i=1,2,3,4$) of the four alternatives x_i ($i=1,2,3,4$) can be obtained.
803 Figs. 20-23 illustrate the values $\beta_i = \Delta^{-1}(\bar{r}_i)$ ($i=1,2,3,4$) of the four alternatives x_i
804 ($i=1,2,3,4$) obtained by the 2TLGBM operator in detail.
805
806




Fig. 20. The values β for alternative x_1 obtained by the 2TLGBM operator ($p \in (0, 20]$, $q \in (0, 20]$).





Fig. 21. The values β for alternative x_2 obtained by the 2TLGBM operator ($p \in (0, 20]$, $q \in (0, 20]$).


815



816

817 **Fig. 22.** The values β for alternative x_3 obtained by the 2TLGBM operator ($p \in (0,20]$,
 818 $q \in (0,20]$).

819




820

821 **Fig. 23.** The values β for alternative x_4 obtained by the 2TLGBM operator ($p \in (0,20]$,
 822 $q \in (0,20]$).

823

824 If we let the parameter p fixed, then different values β_i and rankings of the alternatives can
 825 be obtained as the parameter q changed, which was shown in Fig. 24.



826

827 **Fig. 24.** Variation of β obtained with the 2TLGBM operator ($p=1$, $q \in (0,20]$).

828


829 From Fig. 24, we can find that when $q \in (0,20]$, the ranking of the four alternatives is
 830 $x_2 \succ x_4 \succ x_3 \succ x_1$ and the best choice is x_2 .

831 From Fig. 24, we find that the collective overall preference values obtained by the 2TLGBM
 832 aggregation operators become smaller as parameter p and q increase for the same
 833 aggregation arguments. Therefore, it plays a crucial part in decision making. For example, in
 834 the real group decision making problems, the decision makers who take a gloomy view of
 835 the prospects could choose the bigger values of the parameters p and q while the decision
 836 makers who are optimistic could choose the smaller values of the parameters p and q .


837 It is worth noting that most of the beta obtained by the 2TLBM operator are bigger than most
 838 of the beta obtained by the 2TLGBM operator, which indicates that the 2TLBM operator can
 839 obtain more favorable (or optimistic) expectations, while the 2TLGBM operator has more
 840 unfavorable (or pessimistic) expectations. Therefore, we can conclude that the 2TLBM
 841 operator can be considered as the optimistic operator, while the 2TLGBM operator can be
 842 considered as the pessimistic operator and the values of the parameters can be considered
 843 as the optimistic or pessimistic levels. By Figs. 15-24, we can conclude that the decision
 844 makers who take a gloomy view of the prospects could use the 2TLGBM operator and
 845 choose the bigger values of the parameters p and q , while the decision makers who are
 846 optimistic could use the 2TLBM operator and choose the bigger values of the parameters p
 847 and q .

848 In order to obtain the more neutral results, we can use the arithmetic averages of the
 849 optimistic and pessimistic results, which can be found in Figs. 25-28.

850



851
852 **Fig. 25.** The values β for alternative x_1 obtained by the 2TLBM and 2TLGBM operators
853 $(p \in (0, 20], q \in (0, 20])$.
854



855
856 **Fig. 26.** The values β for alternative x_2 obtained by the 2TLBM and 2TLGBM operators
857 $(p \in (0, 20], q \in (0, 20])$.
858




Fig. 27. The values β for alternative x_3 obtained by the 2TLBM and 2TLGBM operators ($p \in (0, 20]$, $q \in (0, 20]$).




Fig. 28. The values β for alternative x_4 obtained by the 2TLBM and 2TLGBM operators ($p \in (0, 20]$, $q \in (0, 20]$).

By examples 1 and 2, we can see that the decision results obtained with Approach I may be different from the decision results obtained with Approach II. Approach I utilizes the W2TLBM (or W2TLGBM) operator to aggregate all of the individual 2-tuple linguistic decision

matrices into a collective 2-tuple linguistic decision matrix and then utilize the W2TLBM (or W2TLGBM) operator to derive the collective overall preference values of each alternative. Approach II utilizes the 2TLBM (or 2TLGBM) operator to aggregate all of the individual 2-tuple linguistic decision matrices into a collective 2-tuple linguistic decision matrix and then utilize the 2TLBM (or 2TLGBM) operator to derive the collective overall preference values of each alternative. The 2TLBM and 2TLGBM operators only involve the input data and their interrelationships, but the importance of each datum is not emphasized. The W2TLBM and W2TLGBM operators not only involve the input data and their interrelationships but also take the importance of each datum into account.

7. CONCLUSIONS

In this paper, we have developed several new 2-tuple linguistic aggregation operators, including the 2TLBM, W2TLBM, 2TLGBM, and W2TLGBM operators. We have studied some fundamental properties of the developed operators, such as commutativity, idempotency, boundedness, and monotonicity. We also discuss some special cases of the proposed operators. Compared with the existing 2-tuple linguistic aggregation operators, the primary advantage of these operators is that they capture the interrelationship of the input arguments and thus consider the decision information as much as possible. Furthermore, we have used the proposed operators to develop two approaches for multiple attribute group decision making with 2-tuple linguistic information. Finally, two numerical examples are provided to illustrate the developed approaches and to compare the developed approaches with the existing ones. In the future, we will study the Bonferroni mean and the geometric Bonferroni mean under interval-valued 2-tuple linguistic environments [49,50].

ACKNOWLEDGEMENTS

The authors thank the anonymous referees for their valuable suggestions in improving this paper. This work is supported by the National Natural Science Foundation of China (Grant Nos. 71271070 and 61375075) and the Natural Science Foundation of Hebei Province of China (Grant Nos. F2012201020 and A2012201033).

COMPETING INTERESTS

AUTHORS' CONTRIBUTIONS

'Zhiming Zhang' designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. 'Chong Wu' managed the analyses of the study and the literature searches. All authors read and approved the final manuscript.

REFERENCES

1. Cebi S, Kahraman C. Developing a group decision support system based on fuzzy information axiom. *Knowledge-Based Systems*. 2010;23:3-16.
2. Gao CY, Peng DH. SWOT analysis with nonhomogeneous uncertain preference information. *Knowledge-Based Systems*. 2011;24:796-808.
3. Noor-E-Alam M, Lipi TF, Hasin MAA, Ullah AMMS. Algorithms for fuzzy multi expert multi criteria decision making (ME-MCDM). *Knowledge-Based Systems*. 2011;24:367-377.
4. Degani R, Bortolan G. The problem of linguistic approximation in clinical decision making. *International Journal of Approximate Reasoning*. 1988;2:143-162.
5. Delgado M, Verdegay JL, Vila MA. On aggregation operations of linguistic labels. *International Journal of Intelligent Systems*. 1993;8:351-370.
6. Martin O, Klir GJ. On the problem of retranslation in computing with perceptions. *International Journal of General Systems*. 2006;35 (6):655-674.

- 922 7. Pedrycz W, Ekel P, Parreiras R. Fuzzy Multicriteria Decision-Making: Models, Methods
923 and Applications, John Wiley & Sons, Ltd., Chichester, UK, 2010.
- 924 8. Xu ZS. A method based on linguistic aggregation operators for group decision making
925 with linguistic preference relations. *Information Sciences*. 2004;166(1-4):19-30.
- 926 9. Xu ZS. EOWA and EOWG operators for aggregating linguistic labels based on linguistic
927 preference relations. *International Journal of Uncertainty, Fuzziness and Knowledge-Based
928 Systems*. 2004;12:791-810.
- 929 10. Xu ZS. Deviation measures of linguistic preference relations in group decision making.
930 *Omega*. 2005;33:249-254.
- 931 11. Xu ZS. On generalized induced linguistic aggregation operators. *International Journal of
932 General Systems*. 2006;35:17-28.
- 933 12. Xu ZS. Induced uncertain linguistic OWA operators applied to group decision making.
934 *Information Fusion*. 2006;7:231-238.
- 935 13. Xu ZS. An approach based on the uncertain LOWG and induced uncertain LOWG
936 operators to group decision making with uncertain multiplicative linguistic preference
937 relations. *Decision Support Systems*. 2006;41:488-499.
- 938 14. Herrera F, Martínez L. A 2-tuple fuzzy linguistic representation model for computing with
939 words. *IEEE Transactions on Fuzzy Systems*. 2000;8:746-752.
- 940 15. Martínez L, Herrera F. An overview on the 2-tuple linguistic model for computing with
941 words in decision making: Extensions, applications and challenges. *Information Sciences*.
942 2012;207:1-18.
- 943 16. Chang KH, Wen TC. A novel efficient approach for DFMEA combining 2-tuple and the
944 OWA operator. *Expert Systems with Applications*. 2010;37:2362-2370.
- 945 17. Jiang YP, Fan ZP. Property analysis of the aggregation operators for 2-tuple linguistic
946 information. *Control and Decision*. 2003;18(6):754-757.
- 947 18. Ju YB, Wang AH, Liu XY. Evaluating emergency response capacity by fuzzy AHP and 2-
948 tuple fuzzy linguistic approach. *Expert Systems with Applications*. 2012;39:6972-6981.
- 949 19. Wei GW. A method for multiple attribute group decision making based on the ET-WG
950 and ET-OWG operators with 2-tuple linguistic information. *Expert Systems with Applications*.
951 2010;37(12):7895-7900.
- 952 20. Wei GW. Grey relational analysis method for 2-tuple linguistic multiple attribute group
953 decision making with incomplete weight information. *Expert Systems with Applications*.
954 2011;38:4824-4828.
- 955 21. Wei GW. Some generalized aggregating operators with linguistic information and their
956 application to multiple attribute group decision making. *Computers & Industrial Engineering*.
957 2011;61:32-38.
- 958 22. Wei GW, Zhao XF. Some dependent aggregation operators with 2-tuple linguistic
959 information and their application to multiple attribute group decision making. *Expert Systems
960 with Applications*. 2012;39:5881-5886.
- 961 23. Xu YJ, Huang L. An approach to group decision making problems based on 2-tuple
962 linguistic aggregation operators, in: *ISECS International Colloquium on Computing,
963 Communication, Control, and Management*, IEEE Computer Society, Guangzhou, China,
964 2008, pp. 73-77.
- 965 24. Yang W, Chen ZP. New aggregation operators based on the Choquet integral and 2-
966 tuple linguistic information. *Expert Systems with Applications*. 2012;39:2662-2668.
- 967 25. Bonferroni C. Sulle medie multiple di potenze. *Bolletino Matematica Italiana*. 1950;5:267-
968 270.
- 969 26. Xu ZS, Yager RR. Intuitionistic fuzzy Bonferroni means. *IEEE Transactions on Systems,
970 Man and Cybernetics*. 2011;41:568-578.
- 971 27. Yager RR. On generalized Bonferroni mean operators for multi-criteria aggregation.
972 *International Journal of Approximate Reasoning*. 2009;50:1279-1286.
- 973 28. Xu RN, Zhai XY. Extensions of the analytic hierarchy process in fuzzy environment.
974 *Fuzzy Sets and Systems*. 1992;52:251-257.

- 975 29. Xu ZS, Yager RR. Some geometric aggregation operators based on intuitionistic fuzzy
976 sets. *International Journal of General Systems*. 2006;35:417-433.
- 977 30. Xu ZS. Methods for aggregating interval-valued intuitionistic fuzzy information and their
978 application to decision making. *Control and Decision*. 2007;22:215-219.
- 979 31. Xia MM, Xu ZS. Hesitant fuzzy information aggregation in decision making. *International*
980 *Journal of Approximate Reasoning*. 2011;52:395-407.
- 981 32. Xu ZS. Uncertain linguistic aggregation operators based approach to multiple attribute
982 group decision making under uncertain linguistic environment. *Information Science*.
983 2004;168:171-184.
- 984 33. Xu ZS. *Group Decision Making with Triangular Fuzzy Linguistic Variables*, Springer,
985 Berlin, Heidelberg, pp. 17-26, 2007.
- 986 34. Xu ZS. Uncertain Bonferroni mean operators. *International Journal of Computational*
987 *Intelligence Systems*. 2010;3(6):761-769.
- 988 35. Xia MM, Xu ZS, Zhu B. Geometric Bonferroni means with their application in multi-
989 criteria decision making. *Knowledge-Based Systems*. 2013;40:88-100.
- 990 36. Xu ZS, Chen Q. A multi-criteria decision making procedure based on interval-valued
991 intuitionistic fuzzy bonferroni means. *Journal of Systems Science and Systems Engineering*.
992 2011;20:217-228.
- 993 37. Zhu B, Xu ZS, Xia MM. Hesitant fuzzy geometric Bonferroni means. *Information*
994 *Sciences*. 2010;205(1):72-85.
- 995 38. Wei GW, Zhao XF, Lin R, Wang HJ. Uncertain linguistic Bonferroni mean operators and
996 their application to multiple attribute decision making. *Applied Mathematical Modelling*.
997 2013;37:5277-5285.
- 998 39. Liu PD, Jin F. The trapezoid fuzzy linguistic Bonferroni mean operators and their
999 application to multiple attribute decision making. *Scientia Iranica*. 2012;19:1947-1959.
- 1000 40. Xia MM, Xu ZS, Zhu B. Generalized intuitionistic fuzzy Bonferroni means. *International*
1001 *Journal of Intelligent systems*. 2012;27:23-47.
- 1002 41. Herrera F, Martínez L. An approach for combining linguistic and numerical information
1003 based on 2-tuple fuzzy linguistic representation model in decision-making. *International*
1004 *Journal of Uncertainty, Fuzziness, Knowledge-Based Systems*. 2000;8:539-562.
- 1005 42. Herrera F, Martínez L. A model based on linguistic 2-tuples for dealing with multigranular
1006 hierarchical linguistic contexts in multi-expert decision making. *IEEE Transactions on*
1007 *Systems, Man, and Cybernetics-Part B: Cybernetics*. 2001;31:227-234.
- 1008 43. Bonissone PP, Decker KS. Selecting uncertainty calculi and granularity: an experiment in
1009 trading-off precision and complexity, in: L.H. Kanal, J.F. Lemmer (Eds.), *Uncertainty in*
1010 *Artificial Intelligence*, North-Holland, Amsterdam, 1986, pp. 217-247.
- 1011 44. Delgado M, Herrera F, Herrera-Viedma E, Martín-Bautista MJ, Martínez L, Vila MA. A
1012 communication model based on the 2-tuple fuzzy linguistic representation for a distributed
1013 intelligent agent system on internet. *Soft Computing*. 2002;6:320-328.
- 1014 45. Dong YC, Xu YF, Li HY, Feng B. The OWA-based consensus operator under linguistic
1015 representation models using position indexes. *European Journal of Operational Research*.
1016 2010;203:455-463.
- 1017 46. Herrera F, Herrera-Viedma E. Linguistic decision analysis: steps for solving decision
1018 problems under linguistic information. *Fuzzy Sets and Systems*. 2000;115:67-82.
- 1019 47. Herrera F, Herrera-Viedma E, Martínez L. A fusion approach for managing
1020 multigranularity linguistic term sets in decision making. *Fuzzy Sets and Systems*.
1021 2000;114:43-58.
- 1022 48. Herrera F, Martínez L, Sanchez PJ. Managing non-homogeneous information in group
1023 decision-making. *European Journal of Operational Research*. 2005;166:115-132.
- 1024 49. Zhang HM. The multi attribute group decision making method based on aggregation
1025 operators with interval-valued 2-tuple linguistic information. *Mathematical and Computer*
1026 *Modelling*. 2012;56:27-35.

- 1027 50. Zhang HM. Some interval-valued 2-tuple linguistic aggregation operators and application
1028 in multiattribute group decision making. *Applied Mathematical Modelling*. 2013;37:4269-4282.