

Original Research Article

The law of large numbers. A multicriterial analysis

ABSTRACT

The paper presents the law of large numbers followed by a multicriterial analysis of the six theorems that compose the weak form. This group of theorems work on issues of sequences limit laws of sums of random variables in the sense of convergence almost surely of such sequences

Keywords: Bernoulli, Cebâşev, the law of large numbers, Markov, multicriterial analysis, Poisson

1. INTRODUCTION

A type of boundary problems, based on the notion of convergence in probability, the paramount significance of the link between the basic concepts of probability theory and empirical terms, is known as the law of large numbers. This is a result set looking the convergence to 1 of the probabilities of a series of events depending on an increasing number of random factors.

2. THE LAW OF LARGE NUMBERS (THE WEAK FORM)

Theorem1.¹ *Let $(Y_n)_{n \in \mathbb{N}^*}$ be a string of random variables and mean values and finite variances*

$$M[Y_n] = M_n < \infty, \quad D^2[Y_n] = \sigma_n^2 < \infty.$$

If there exist a constant M such that

$$\lim_{n \rightarrow \infty} M_n = M$$

and

$$\lim_{n \rightarrow \infty} \sigma_n^2 = 0$$

(1)

then $\{Y_n\} \xrightarrow{p} M$.

Demonstration

We observe that for any $\varepsilon > 0$

$$\{|Y_n - M| > \varepsilon\} \subseteq \{|Y_n - M_n| > \frac{\varepsilon}{2}\} \cup \{|M_n - M| > \frac{\varepsilon}{2}\}.$$

we can write

$$P(\{|Y_n - M| > \varepsilon\}) \leq P\left(\{|Y_n - M_n| > \frac{\varepsilon}{2}\}\right) + P\left(\{|M_n - M| > \frac{\varepsilon}{2}\}\right). \quad (2)$$

Applying random variable Y_n on Cebâşev's inequality we obtain

$$P(\{|Y_n - M| > \varepsilon\}) \leq \frac{D^2[Y_n]}{\varepsilon^2} = \frac{\sigma_n^2}{\varepsilon^2}. \quad (3)$$

Since

¹ [9]

$$\lim_{n \rightarrow \infty} M_n = M$$

It results that $\forall \varepsilon > 0, \exists N(\varepsilon) \in \mathbb{N}$ such that $\forall n \geq N(\varepsilon)$ the inequality $|M_n - M| > \frac{\varepsilon}{2}$ becomes impossible, ie

$$\forall n \geq N(\varepsilon): P(\{|M_n - M| > \varepsilon\}) = 0.$$

Passing to the limit in (2), taking into account of (3) and the ipohesis (1) it results that

$$\lim_{n \rightarrow \infty} P(\{|Y_n - M| > \varepsilon\}) = 0.$$

Theorem 2. (the weak form of the Jacob Bernoulli² theorem)

The first formulation³: If A is an event caracterized by $P(A) = p \neq 0$ and $\frac{S_n}{n}$ of the relative frequency achievement of the event A in the first n samples of an infinite number of independent samples, then

$$\forall \varepsilon > 0 : \lim_{n \rightarrow \infty} P\left(\left|\frac{S_n}{n} - p\right| > \varepsilon\right) = 0$$

or, equivalent,

$$\forall \varepsilon > 0 : \lim_{n \rightarrow \infty} P\left(\left|\frac{S_n}{n} - p\right| \leq \varepsilon\right) = 1.$$

The second formulation: If $\{X_n\}_{n \in \mathbb{N}}$ is a set of independent random variables such that $X_n: B_i(n, p), n \in \mathbb{N}$, then the string of random variables

$$\left\{\frac{1}{n} \sum_{k=1}^n X_k\right\}_{n \in \mathbb{N}}$$

converges in probabiliy to p .

Demonstration

We demonstrate Bernoulli's theorem in the second form. We note

$$Y_n = \frac{1}{n} \sum_{k=1}^n X_k$$

and considering that

$$M[Y_n] = \frac{1}{n} \sum_{k=1}^n M[X_k] = \frac{1}{n} np = p,$$

$$D^2[Y_n] = \frac{1}{n^2} \sum_{k=1}^n D^2[X_k] = \frac{npq}{n^2} = \frac{pq}{n}.$$

So

$$\lim_{n \rightarrow \infty} M[Y_n] = p$$

and

$$\lim_{n \rightarrow \infty} D^2[Y_n] = 0$$

which indicates that the hypotheses of the theorem 1 are satisfied, then

$$\left\{\frac{1}{n} \sum_{k=1}^n X_k\right\} \xrightarrow{p} p.$$

Remark 1. The first statement of the Bernoulli theorem approach is theoretical justification made between the concepts of relative frequency and probability of an event. It expresses the fact that for very large strings evidence any significant difference between the relative frequency and probability of an event is very improbable. So convergence of relative frequencies to probability of the event it does not to be understood as a common numerical convergence; not apparent from the above theorem and is not acceptable intuitive idea that with increasing number of samples relative frequency values are necessarily closer probability.

² Jakob Bernoulli (1654-1705), *Analysis problematis ante hac propositi, de inventione linea descensus a corpore gravi percurrendae uniformiter, sic ut temporibus aequales altitudines emetiatur: & alterius cujusdam problematis propositio*", *Acta Eruditorum* , 1690

³ [9], [8]

Remark 2. The two formulations under one of the other if we consider the absolute frequency of realization of A in n samples. Let X_n be a random variable distributed $B_i(n, p)$. If we note

$$S_n = \sum_{k=1}^n X_k$$

then

$$\frac{S_n}{n} = \frac{1}{n} \sum_{k=1}^n X_k$$

and

$$\lim_{n \rightarrow \infty} P \left(\left| \frac{S_n}{n} - p \right| > \varepsilon \right) = 0$$

which means that the

$$\left\{ \frac{1}{n} \sum_{k=1}^n X_k \right\} \xrightarrow{p} p$$

Remark 3. The theorem result is the theoretical justification of the definition of a parameter estimation of a distribution.

Theorem 3.⁴(Cebâșev⁵) If $(X_n)_{n \in \mathbb{N}^*}$ is a set of independent random variables two by two having dispersions bounded by the same constant

$$D^2[X_n] \leq c^2, \quad n \in \mathbb{N}$$

then

$$\forall \varepsilon > 0 : \lim_{n \rightarrow \infty} P \left(\left| \frac{1}{n} \sum_{k=1}^n X_k - \frac{1}{n} \sum_{k=1}^n M[X_k] \right| > \varepsilon \right) = 0$$

or, equivalent,

$$\forall \varepsilon > 0 : \lim_{n \rightarrow \infty} P \left(\left| \frac{1}{n} \sum_{k=1}^n X_k - \frac{1}{n} \sum_{k=1}^n M[X_k] \right| \leq \varepsilon \right) = 1,$$

ie occurs the convergence in probability string of random variables,

$$\left\{ \frac{1}{n} \sum_{k=1}^n (X_k - M[X_k]) \right\} \xrightarrow{p} 0$$

Demonstration

If

$$Y_n = \frac{1}{n} \sum_{k=1}^n (X_k - M[X_k])$$

then

$$M[Y_n] = M \left[\frac{1}{n} \sum_{k=1}^n (X_k - M[X_k]) \right] = \frac{1}{n} \sum_{k=1}^n M[X_k] - \frac{1}{n} \sum_{k=1}^n M[X_k] = 0$$

and because the random variables X_n are independent two by two, it results that $(X_k - M[X_k])$ are independent two by two and therefore

$$D^2[Y_n] = D^2 \left[\frac{1}{n} \sum_{k=1}^n (X_k - M[X_k]) \right] = \frac{1}{n^2} \sum_{k=1}^n D^2(X_k - M[X_k]) = \frac{1}{n} \sum_{k=1}^n D^2[X_k].$$

Applying to random variable Y_n the Cebâșev's inequality and taking into account the assumptions of the problem, we can write

$$P \left(\left| \frac{1}{n} \sum_{k=1}^n X_k - \frac{1}{n} \sum_{k=1}^n M[X_k] \right| > \varepsilon \right) \leq \frac{1}{n^2 \varepsilon^2} \sum_{k=1}^n D^2[X_k] < \frac{nc^2}{n^2 \varepsilon^2} = \frac{c^2}{n \varepsilon^2}$$

⁴ [9], [6]

⁵ *Paŋnuti Lvovici Cebîșev (1821 – 1894), Experiența elementară în analiza teoriei probabilităților*, 1845

and so

$$\lim_{n \rightarrow \infty} P \left(\left\{ \left| \frac{1}{n} \sum_{k=1}^n X_k - \frac{1}{n} \sum_{k=1}^n M[X_k] \right| > \varepsilon \right\} \right) \leq \lim_{n \rightarrow \infty} \frac{c^2}{n\varepsilon^2} = 0$$

Theorem 4.⁶(Markov⁷) *If the random variables $X_1, X_2, \dots, X_n, \dots$ have the property that:*

$$\frac{D^2[\sum_{i=1}^n X_i]}{n^2} \rightarrow 0, \quad n \rightarrow \infty$$

then we have the law of large numbers.

Demonstration

It must be demonstrated that the string $\{X_n\}_{n \in \mathbb{N}}$ verifies the relation:

$$\forall \varepsilon > 0 : \lim_{n \rightarrow \infty} P \left(\left\{ \left| \frac{1}{n} \sum_{k=1}^n X_k - \frac{1}{n} \sum_{k=1}^n M[X_k] \right| \leq \varepsilon \right\} \right) = 1.$$

Applying Cebâşev's inequality for the random variables:

$$Y = \sum_{i=1}^n \frac{X_i}{n}, D^2[Y] = D^2 \left[\sum_{i=1}^n X_i/n \right] = \frac{1}{n^2} D^2 \left[\sum_{i=1}^n X_i \right] \rightarrow 0, \quad n \rightarrow \infty$$

Therefore $D^2[Y] < \infty$ and then Cebâşev's inequality holds:

$$P(|Y - M[Y]| \leq \varepsilon) \geq 1 - \frac{1}{\varepsilon^2} \frac{1}{n^2} D^2 \left[\sum_{i=1}^n X_i \right].$$

But

$$\frac{1}{n^2} D^2 \left[\sum_{i=1}^n X_i \right] \rightarrow 0, \quad n \rightarrow \infty$$

Passing to the limit in the previous inequality, we obtain the assertion of the theorem.

Remark 4. In the particular case when the random variables are independent, provided the statement is written:

$$\frac{D^2[\sum_{i=1}^n X_i]}{n} \rightarrow 0, \quad n \rightarrow \infty$$

Theorem 5.⁸(Hincin) *If $\{X_n\}_{n \in \mathbb{N}}$ is a set of independent random variables with the same distribution two by two and having finite dispersions, then*

$$\lim_{n \rightarrow \infty} P \left(\left\{ \left| \frac{1}{n} \sum_{k=1}^n X_k - M \right| > \varepsilon \right\} \right) = 0$$

where $M = M[X_n], n \in \mathbb{N}$.

Demonstration

If the random variable $\{X_n\}_{n \in \mathbb{N}}$ follow the same distribution, it results that $D^2[X_n] = \sigma^2, n \in \mathbb{N}$, which implies satisfying the assumptions of Cebâşev's theorem.

Remark 5. This result substantiates the usual theory of approximation of the average of statistical characteristic of a population studied by the arithmetic mean of the observed values. Indeed, if $X_1, X_2, \dots, X_n, \dots$ are seen as potential values of a random variable X obtained by a number of independent samples, the theorem shows that their average converges to $M = M[X]$.

As a special case of Cebâşev's theorem we obtain Poisson's theorem.

Theorem 6. (Poisson⁹) *Let A an event of which the probability to output varies during a series of independent samples so that the sample of rank $k, P(A) = p_k, k = 1, 2, \dots$ and let $\frac{S_n}{n}$ the frequency of execution of the event A in the first n samples. Then*

$$\lim_{n \rightarrow \infty} P \left(\left\{ \left| \frac{S_n}{n} - \frac{p_1 + p_2 + \dots + p_n}{n} \right| > \varepsilon \right\} \right) = 0$$

⁶ [9], [1]

⁷ Andrey Andreyevich Markov (1856 - 1922), *Extension of the limit theorems of probability theory to a sum of variables connected in a chain*, reprinted in Appendix B of: R.

Howard. *Dynamic Probabilistic Systems, volume 1: Markov Chains*. John Wiley and Sons, 1971

⁸ [9], [2]

⁹ Siméon Denis Poisson (1781-1840), “*The Poisson law*”

Demonstration

We define for each $k = 1, 2, \dots, n, \dots$ the random variable X_k takes the values 1 or 0 as A is realized or not in the sample with the rank k . Hereupon, X_k is defined

$$X_k: \begin{pmatrix} 0 & 1 \\ 1-p_k & p_k \end{pmatrix}$$

Noting with S_n the number of realisations of A in the first n samples, we denote that

$$f_n(A) = \frac{S_n}{n} = \frac{1}{n} \sum_{k=1}^n X_k$$

Because $X_1, X_2, \dots, X_n, \dots$ are independent random variables and because

$$M[X_k] = p_k, D^2[X_k] = M[X_k^2] - M[X_k]^2 = p_k - p_k^2 = p_k(1 - p_k) \leq \frac{1}{4}$$

it results that the string $\{X_n\}_{n \in \mathbb{N}}$ verify the hypotheses of the Cebâşev's theorem and so

$$\lim_{n \rightarrow \infty} P \left(\left| \frac{S_n}{n} - \frac{p_1 + p_2 + \dots + p_n}{n} \right| > \varepsilon \right) = \lim_{n \rightarrow \infty} P \left(\left| \frac{1}{n} \sum_{k=1}^n X_k - \frac{1}{n} \sum_{k=1}^n M[X_k] \right| > \varepsilon \right) = 0.$$

3. THE MULTICRITERIAL ANALYSIS OF THE DYNAMICS THAT MAKE UP THE APPLICATION FORM WEAK THEOREMS OF LAW OF LARGE NUMBERS

We have the following variants:

- variant (a): the extent of which one applies more frequently theorems
- variant (b): the hardness
- variant (c): reality

6 criteria have been chosen:

1.	T. Bernoulli 1
2.	T. Bernoulli 2
3.	T. Cebâşev
4.	T. Markov
5.	T. Hincin
6.	T. Poisson

Fig. 1

Based bet on score, weighting of the criteria resulted as follows:

	T. Bernoulli 1	T. Bernoulli 2	T. Cebâşev	T. Markov	T. Hincin	T. Poisson	Point s	Leve l	γ_i
T. Bernoulli 1	1/2	1	0	0	0	1/2	2	6	0,5
T. Bernoulli 2	0	1/2	0	1	1/2	1/2	2,5	3,5	1
T. Cebâşev	1	1	1/2	1	0	0	3,5	2,5	2,4
T. Markov	1	0	0	1/2	0	1	2,5	3,5	1
T. Hincin	1	1/2	1	1	1/2	0	4	1	3,8
T. Poisson	1/2	1/2	1	0	1	1/2	3,5	2,5	2,4

Fig. 2

By the following formula,

$$\gamma_i = \frac{p + 4p + m + 0,5}{-\Delta p' + \frac{N_{crt}}{2}}$$

we have

$$\gamma_{Bernoulli 1} = \frac{2 + (2 - 2) + 0 + 0,5}{-(2 - 4) + \frac{6}{2}} = \frac{2,5}{5} = 0,5$$

$$\gamma_{Bernoulli 2} = \frac{2,5 + (2,5 - 2) + 1 + 0,5}{-(2,5 - 4) + \frac{6}{2}} = \frac{4,5}{4,5} = 1 = \gamma_{Markov}$$

$$\gamma_{Poisson} = \gamma_{Cebăşev} = \frac{3,5 + (3,5 - 2) + 3 + 0,5}{-(3,5 - 4) + \frac{6}{2}} = \frac{8,5}{3,5} = 2,4$$

$$\gamma_{Hincin} = \frac{4 + (4 - 2) + 5 + 0,5}{-(4 - 4) + \frac{6}{2}} = \frac{11,5}{3} = 3,8$$

It is noted that the main diagonal of the array contains only quadratic criteria for scoring 1/2 values because no criteria may be more important or less important than the criteria itself.

According to the criteria were the following notes for each variant N_i .

	Variant (a)	Variant (b)	Variant (c)
Criteria	N_i	N_i	N_i
T. Bernoulli 1	7	6	8
T. Bernoulli 2	9	9	9
T. Cebăşev	10	10	7
T. Markov	6	5	4
T. Hincin	9	4	5
T. Poisson	8	7	7

Fig. 3

It may take into account different weight now and consequence of each criterion, complementing and enhancing the table above notes (lines) with the coefficient of importance:

		Variant (a)		Variant (b)		Variant (c)	
Criteria	γ_i	N_i	$N_i \times \gamma_i$	N_i	$N_i \times \gamma_i$	N_i	$N_i \times \gamma_i$
T. Bernoulli 1	0,5	7	3,5	6	3	8	4
T. Bernoulli 2	1	9	9	9	9	9	9
T. Cebăşev	2,4	10	24	10	24	7	16,8
T. Markov	1	6	6	5	5	4	4
T. Hincin	3,8	9	34,2	4	15,2	5	19
T. Poisson	2,4	8	19,2	7	16,8	7	16,8
Final clasament			95,9		73		69,6

Fig. 4

4. CONCLUSIONS

It is noted that after the ranking did, the theorems of the weak large numbers law are preferred to be taken in the variant (a).

We want to emphasize that point III is our own creation. Efforts have been made to develop this multicriterial analysis applied to the law of large numbers, hoping that we will develop it in the future.

REFERENCES

- [1] M. Capinski, E. Kopp, Measure, Integral and Probability, (Springer-Verlag, 2003)
- [2] G. Cenuşa, V. Burlacu, I. Sacuiu, Teoria probabilităţilor şi statistică matematică, (Editura ASE, 1999)
- [3] W. Feller, An introduction to probability theory and its applications, Third Edition, Vol. I, (John Wiley & Sons, Inc., New York, 1968), ISBN 0-471-25711-7
- [4] C. Geiss, S. Geiss, An Introduction to Probability Theory, (University of Jyväskylä, 2004)
- [5] Klebaner, C. Fima, Introduction to Stochastic Calculus with Applications, (Imperial College Press, London, 2004)
- [6] M. Loève, Fundamental limit theorems of probability theory, The Annals of Mathematical Statistics, Vol. 21, No. 3 (Sep., 1950), pp. 321-338

- [7] B. Øksendal, Stochastic differential equations – An introduction with applications, (Springer, Berlin, 2007), ISBN-10:3-540-04758-1, ISBN-13:978-3-540-04758-2, p.113-p.137
- [8] G.V. Orman, Handbook of limit theorems and stochastic approximation, (Transilvania University Press, Braşov, 2003), ISBN 973-635-227-7
- [9] A. Pletea, L. Popa, Teoria probabilităţilor, (Ed. Universităţii Tehnice Gh. Asachi, Iaşi, 1999)