Mathematical model of formation aerosol particles fractal storm cloud

Tembulat Kumykov[*1] **and Roman Parovik**[2,3]

[1] *Department of Mathematical modeling of geophysical processes, Institute of Applied Mathematics and Automation, Shortanova Str. 89A, 360000 Nalchik, RUSSIA*

[2] *Laboratory of Mathematical modeling of physical processes, Institute of Cosmophysical Research and Radio Wave Propagation Far Eastern Branch of Russian Academy of Science, Mirnaya Str. 7, 684034 Kamchatka region, Paratunka, RUSSIA*

[3] *Faculty of Physics and Mathematics, Vitus Bering Kamchatka State University, Pogranichnaya Str., 4, 683032 Kamchatka region, Petropavlovsk-Kamchatsky, RUSSIA*

Short Research Article

Abstract

The paper proposes a new educational model of aerosol particles in thunderclouds. The model takes into account the fractal properties of the storm clouds, and its solution was obtained by numerical methods of fractional calculus. Built new profiles calculated curves that are consistent with the classical Lifshitz-Slezov-Wagner theory.

Keywords: Coalescence, Lifshitz-Slezov-Wagner theory, Fractal dimension, the mathematical model

2010 Mathematics Subject Classification: 93A30; 35Q86

1 Introduction

Formation aerosol particles is an important process in the formation of thunderstorm electricity and studied by the theory of Lifshitz-Slyozov-Wagner (LSW) [Lifshic (1958); Wagner (1961)]. This process is referred to as coalescence. According to the theory LSV distribution of aerosols in size depending on the time of power law: $R(t) \sim t^{\frac{1}{3}}$ in the case of the diffusion mechanism of formation [Lifshic (1958)], according to the law $R(t) \sim t^{\frac{1}{2}}$ in the case of the reaction on the surface of aerosol [Lifshic (1958)].

It is known that power laws describe well the fractal objects and processes [Feder, (1988)]. In [Ernst (1986)], the authors, using the experimental data, was found fractal dimension hail clouds, which has a value of $d_f = 1,36 \pm 0,1$. Therefore it can be concluded that the storm clouds have fractal

**Corresponding author: E-mail: macist20@mail.ru*

properties that can be described by the mathematical apparatus of fractional calculus [Nahushev (2003)].

2 Statement of the problem

Consider the scheme coalescence LSW when fall grains (aerosols) are fixed and grow by diffusion from the surrounding solution Lifshic (1979). In determining the concentration of the bulk amount of the substance dissolved in a unit volume of solution, the diffusion flux is determined by the formula

$$q\left(r,t\right)=\lambda\frac{\partial u\left(r,t\right)}{\partial r}, \tag{2.1}$$

when $\lambda>0$ – the diffusion coefficient of solute, $u\left(r,t\right)$ – concentration of the saturated solution, r – the aerosol radius, $t>0$ – time.

Diffusion flux (2.1) at the surface of the aerosol ($r=R$) coincides with the rate of change of its radius:

$$\frac{dR\left(t\right)}{dt}=\lambda\frac{\partial u\left(r,t\right)}{\partial r}|_{r=R}. \tag{2.2}$$

In [Lifshic (1979)], taking into account the critical radius $R_c\left(t\right)$ and the relation (2.2), we obtain the equation:

$$\frac{dR\left(t\right)}{dt}=\frac{R_c^3\left(0\right)}{R\left(t\right)}\left(\frac{1}{R_c\left(t\right)}-\frac{1}{R_c\left(t\right)}\right), \tag{2.3}$$

when $R_c^3\left(0\right)$ – the value of critical radius, it is responsible for the beginning stages of coalescence. In the formula (2.3) with values $R\left(t\right)>R_c\left(t\right)$ of the aerosol is increased, and with - $R\left(t\right)<R_c\left(t\right)$ – dissolved.

In the case of the fractal structure of a thundercloud diffusion flux (2.1) can be generalized by the formula [Nahushev (2003)]

$$q\left(r,t\right)=\lambda\partial_{0t}^{\alpha}u\left(r,\tau\right),0<\alpha<1, \tag{2.4}$$

when $\partial_{0t}^{\alpha}u\left(r,\tau\right)=\frac{1}{\Gamma\left(1-\alpha\right)}\int\limits_0^t\frac{u_\tau'\left(r,\tau\right)d\tau}{\left(t-\tau\right)^\alpha}$ – fractional differentiation operator Gerasimov-Caputo order [Nahushev (2012)]; $\Gamma\left(x\right)$ – Euler gamma-function. Taking into account (2.4), in [Kumykov (2014)] proposed a generalization of equation (2.3), which can be written as:

$$\partial_{0t}^{\alpha}R\left(\tau\right)=\frac{R_c^3\left(0\right)}{R\left(t\right)R_c\left(t\right)}\left(1-\frac{R_c\left(t\right)}{R\left(t\right)}\right), \tag{2.5}$$

We solve equation (2.5) using an explicit finite-difference scheme. We introduce a uniform grid with a constant pitch τ in the time interval from 0 to T , where T – any positive integer. Then we have the following approximation [Sweilam (2011)]

$$\partial_{0t}^{\alpha}R\left(\tau\right)\approx B\sum_{k=0}^{j-1}b_k\left(R_{j-k+1}-R_{j-k}\right),j=1,..,N \tag{2.6}$$

when $B=\frac{\tau^{-\alpha}}{\Gamma\left(2-\alpha\right)}$, N - number of points and $b_k=\left(k+1\right)^{1-\alpha}-k^{1-\alpha}$. Using (2.6) of the formula (2.5), we obtain

$$R_{j+1}=R_j-\sum_{k=0}^{j-1}b_k\left(R_{j-k+1}-R_{j-k}\right)+\frac{R_c^3}{BR_{cj}R_j}\left(1-\frac{R_{cj}}{R_j}\right). \tag{2.7}$$

Equation (2.7) is a numerical solution of the equation formation - dissolution of aerosol in the generalized theory of coalescence LSW fractal thundercloud.

3 Results and Discussion

Numerical modeling was performed using Maple [D'yakonov (2006)], and the values of the parameters in the model were chosen according to [Michael (1998)].



Figure 1: Calculated curves the aerosol radius, constructed according to the decision (2.7) of the generalized model LSW depending on the parameter values α

Fig. 1 shows the calculated curves found by the formula (2.7), which describe the variation the aerosol radius according to the parameter values In the framework of the generalized theory of Lifshitz-Slyozov-Wagner.

Note that when $\alpha = 1$ generalized theory LSW is the classical theory of LSW (red line in Fig. 1). In the general case of Fig. 1 shows that with decreasing the parameter is the rapid growth of the aerosol (the theory of Lifshitz-Slezov), and its dissolution (Wagner theory) on the contrary slows down. This effect is related to the fractal properties of the medium, namely with a power law distribution of aerosols in size depending on the time [Ernst (1986)] and the effects of "memory" in the medium.

The calculated curves of Fig. 1, to some extent, are similar to the calculated curves of the radius of crystals in the framework of LSW (fig. 2a), and especially on the calculated curves in the theory of "communicating neighbors" (fig. 2b), presented in [Michael (1998)].



Figure 2: The calculated curves of the radius of crystals, presented in [Michael (1998)] *a* – classical theory LSW; *b* - the theory of "communicating neighbors"

4 CONCLUSIONS

Indeed, in the theory of ”communicating neighbors” in the generalized theory of LSV accounted for nonlocal effects in time or ”memory” effect, so the calculated curves behave similarly. On this basis, we can conclude that it makes sense to continue the further development and study of the properties of the generalized theory of LSW.

Acknowledgment

The work was performed as part of the project 12-I-DPS-16 ”Fundamental Problems of impact-power radio waves on the atmosphere and the Earth plasmosphere” and Ministry of Education and Science of the Russian Federation on the strategic development ”Kamchatka State University by Vitus Bering ” for 2012-2016 .

References

Lifshic, I.M. and Slezov, V.V. (1958). O kinetike diffuzionnogo raspad peresyschennyh tverdyh rastvorov. Zh‘ETF, 35, 479–492.

Wagner, C.Z. (1961). Theorie der alterung von niederschlgen durch umlsen (ostwaldreifung). Z. Elektrochem, 65, 581-611.

Feder, J. (1988). Fractals. Plenum.

Ernst, M. H. and Pietronero, L. and Tosatti, E. (1986). Fractals in Physics. North-Holland, Amsterdam.

Nahushev, A.M. (2003). Drobnoe ischislenie i ego primenenie. Russia, Moscow.

Lifshic, E.M. and Pitaevskij L.P., (1979). Fizicheskaya kinetika. Russia, Moscow.

Nahushev, A.M. (2003). Nagruzhennye uravneniya i ih primenenie. Russia, Moscow.

Kumykov, T.S.(2014). Primenenie drobnogo ischisleniya v teorii Lifshica-Slezova. Sovremennye voprosy matematicheskoj fiziki, matematicheskoj biologii i informatiki. Russia, Nal’chik, 72–75.

Sweilam, N.H. and Khader, M.M. and Mahdy, A.M. (2011). Numerical studies for solving fractional-order logistic equation. International Journal of Pure and Applied Mathematics, 78, No 8, 1199–1210.

D’yakonov, V.P. (2006). Maple 9.5/10 v matematike, fizike i obrazovanii. Russia, Moscow.

Michael, D.H. (1998). Origin of Anorthosite by Textural Coarsening: Quantitative Measurements of a Natural Sequence of Textural Development. Journal of Petrology, 39, No 7, 1307–1323.