

# Changes in Some Testicular Biometric Parameters and Testicular Function in Cadmium Chloride Administered Wistar Rats

Ekhoje, E. I.<sup>1,\*</sup>, Aloamaka, C.P.<sup>1</sup>, Nwangwa, E. K<sup>1</sup>.

<sup>1</sup>Department of Physiology, Faculty of Basic Medical Science, College of Health Science, Delta State University, P.M.B 001, Abraka, Delta State, Nigeria.

## ABSTRACT

**Introduction:** Recently there has been an increased association between environmental factors and male infertility.

**Aims:** In the present study, the effect of changes in testicular biometric parameters (weight and volume) and testicular function (Sperm count, morphology, testosterone level) in Cadmium chloride administered Wistar rats was studied.

**Methodology:** Twenty male albino Wistar rats were randomized into four groups (n=5). Group A (control) received rat chow and water, while Group B, C and D received 15mg/L, 20mg/L and 25mg/L of Cadmium chloride respectively for 6 weeks.

**Result:** There was a significant ( $P=.05$ ) and dose dependent decrease in testicular function parameters in the rats and a significant ( $P=.05$ ) and positive correlation between the biometric parameters and testicular function.

**Conclusion:** The findings showed that Cadmium chloride has a deleterious effect on testicular function and biometric parameters of the testes may be important in the assessment of testicular function.

*Keyword: Cadmium chloride, male infertility, testicular weight, testicular volume*

## 1. INTRODUCTION

Over the years, infertility has been on the increase in both males and females. The increase in male infertility has become a source of global concern [1]. According to the World Health Organization, 'infertility is the inability of a sexually active, non-contracepting couple to achieve pregnancy in one year [2].

Currently, the etiology of suboptimal semen quality and male reproductive hormone is poorly understood, and many physiological and environmental factors have been implicated [3,4], factors such as smoking, use of restricted drugs, stress, and lack of exercise specifically have been implicated [1].

Cadmium is one of environmental pollutants arising from electroplating, fertilizers, pigment and plastic manufactures. Therefore it easily contaminates the soil, plants, air and water [5]. Cadmium in tobacco represents a contributory source to the total body burden of the smoker [6,7]. Humans and animals can easily be exposed to cadmium toxicity by consuming plants, water and air. Cadmium is absorbed and accumulates in various tissues [8,9] even red blood cells [10], the heart [11] and the skeletal muscle of rats [12].

Most animals with scrotal testes are susceptible to cadmium-induced testicular toxicity [13]. Although, only about 1–2 % of acute cadmium dose is usually taken up by the testes, testicular toxicity is almost

\*Tel: +2347038648797

Email: tareonline@yahoo.com

invariably evident. It has been reported that as low as 1-2 mg Cd/kg body weight can cause testicular damage without pathological changes to other organs [14]. Exposure to cadmium has been reported to reduce male fertility in both humans and rodents [15], but the mechanism is still unknown.

Exposure to cadmium can negatively affect the male reproductive system via degenerative changes in testes, epididymis, and seminal vesicles [16]. Recently, Azoospermic persons were found to have higher serum and seminal plasma cadmium level compared with oligospermic ones [17]. Also positive relationship was found between cadmium exposure and asthenozoospermia in a rat model [15].

The aim of the study is to establish an association between changes in some testicular biometric parameters and testicular function in Wistar rats induced with Cadmium chloride.

## **2. MATERIALS AND METHODS**

### **2.1 Animals and Housing:**

Twenty male albino Wistar rats weighing between 160- 180g and aged 12-14 weeks were used in this study. Animals were raised in the Animal House Unit in Faculty of Basic Medical Science, Delta State University, Abraka, Nigeria. They were maintained in wooden cages with stainless steel wire lids (bedded with wood shavings). Rat chow and water were supplied *ad libitum*. Rats were housed at a temperature of  $28 \pm 1^\circ\text{C}$ , 60 % humidity and under a 12 -hour - light: 12- hour- dark schedule.

### **2.2 Administration of Cadmium:**

Cadmium Chloride ( $\text{CdCl}_2$ ) in crystalline form was obtained from the Petroleum Training Institute (PTI), Warri, Nigeria, and was dissolved in distilled water. The concentrations were 15mg/L, 20mg/L and 25mg/L of the solution [18]. Male rats were randomly assigned to four groups of 5 animals each (control and experimental groups). Experimental male rats were provided access to drinking water containing  $\text{CdCl}_2$  for 6 weeks. The control group received tap water only.

### **2.3 Experimental Design**

A total of twenty (20) male albino Wistar rats were randomly divided into four groups of five rats each.

Group A: Control, fed with rat chow and water daily (n=5)

Group B: 15mg/L Cadmium chloride solution treated rats (n=5)

Group C: 20mg/L Cadmium chloride solution treated rats (n=5)

Group D: 25mg/L Cadmium chloride solution treated rats (n=5)

### **2.4 Male fertility Assessment:**

#### **Sex Organ Weight and Volume**

The rats were sacrificed from each group by decapitation at the end of experiment. The testes were dissected and weighed using a digital electronic weighing balance (EW, Germany) and recorded. The volume of the right testes was measured by first drying the surface of the right testes (using a filter paper), and then immersing it into a  $10 \pm 0.2$  ml measuring cylinder containing 5ml physiological saline, and the volume of displaced fluid in the cylinder recorded.

\*Tel: +2347038648797

Email: tareonline@yahoo.com

### 2.4.1 Semen Quality Analysis:

Seminal content of epididymis was obtained by cutting of the cauda epididymis using surgical blades and squeezed in a sterile clean watch glass. This content was diluted 10 times with 2.9% sodium citrate dihydrate solution and thoroughly mixed to estimate the progressive motility and sperm count [19]. One drop of the suspension was smeared on a glass slide and stained by Eosin – nigrosin stain to determine the percentage of sperm cell morphology [20].

### 2.4.2 Testosterone Assay

Blood samples were collected directly from the heart, by cardiac puncture, in plain non-heparinized bottles and the serum was separated and frozen until hormonal assay was done. Testosterone was estimated in the serum as described by Ismail [21].

### 2.4.3 Histological Section


The portion of testis was fixed in Bouins fluid, dehydrated in ascending grades of alcohol, Embedded in paraffin wax and sectioned at 5 $\mu$  and stained with haematoxyline and eosin. The micrometric measurements such as diameter of seminiferous tubule and diameter of the Leydig cell were measured from randomly selected twenty round sections from each group.

## 2.5 Statistical Analysis

Values were expressed in mean  $\pm$  SEM. SPSS version 17 was used for statistical analysis. Differences between groups were assessed by Student *t*-test and Pearson's product moment correlation.  $P < 0.05$  was taken as significant.


## 3. RESULTS

Result showed the effect of cadmium chloride on testicular function and some testicular biometric parameters, it showed a dose dependent significant ( $P = .05$ ) reduction in the semen quality, serum testosterone, testicular weight/volume.




**Fig 1: Effect of Cadmium Chloride on Testicular Weight**

\*  $p < 0.05$  compared with the Group 1 (control); (n=5)




**Fig 2: Effect of Cadmium Chloride on Testicular Volume**

\*  $p < 0.05$  compared with the Group 1(control); (n=5)




**Fig 3: Effect of Cadmium Chloride on Sperm Count**

\*  $p < 0.05$  compared with the Group 1(control); (n=5)




**Fig 4: Effect of Cadmium Chloride on Sperm motility**

\*  $p < 0.05$  compared with the Group 1(control); (n=5)



**Fig 5: Effect of Cadmium Chloride on Sperm morphology**


\*  $p < 0.05$  compared with the Group 1(control); (n=5)




**Fig 6: Effect of Cadmium Chloride on Testosterone**

\*  $p < 0.05$  compared with the Group 1 (control); (n=5)

#### Histological sections




**Plate 1:** Normal Testis Showing Seminiferous tubules (A), tubules that are lined by stratified layers of cells that are of different stages of maturation (B). Blood vessels (C) and Leydig cells (D). (H & E x100)




**Plate 2:** Testis of Wistar rats treated with 15mg/L of Cadmium chloride treated rats showing mild interstitial fibrosis (A). (H & E, x100).





**Plate 3:** Testis of Wistar rats treated with 20mg/L of cadmium chloride, showing interstitial fibrosis (A), degeneration of seminiferous tubules (B). (H & E, x100).



**Plate 4:** Testis of Wistar rat treated with 25mg/L of cadmium chloride, showing severe Interstitial edema (A) and Focal area of calcification at the tunica albuginea (B). Also present was Leydig cell hypertrophy (C) (H & E, x100).

The histology of the control rats (**Plate 1**) showed normal morphology of the testes. Severity in testicular morphological derangement was dose dependent as the effect of 15mg/L cadmium chloride on the histology of rat testes (**Plate 2**) caused a mild interstitial fibrosis, 20mg/L cadmium chloride solution on the testes (**Plate 3**) induced interstitial fibrosis and degeneration of the seminiferous tubules; and 25mg/L of cadmium chloride (**Plate 4**) caused severe interstitial edema, calcification of the tunica albuginea and hypertrophy of the leydig cells

The relationship between the testicular weight and/or volume with the semen quality and serum testosterone level, following cadmium chloride treatment was ascertained on the basis of correlation coefficient. This was done to establish a possible relationship between the testicular weight or volume with semen quality and serum testosterone level.

The values in the table below indicate the correlation coefficient, a positive value signify a positive correlation.

**Table 1: Relationship between the Testicular Weight and Volume with Semen Quality and Serum Testosterone Level after Cadmium Chloride Treatment.**

	Correlation	
	Testicular Weight (g/kg.bw)	Testicular Volume (ml)
<b>Sperm count (<math>\times 10^6</math> cells/ml)</b>	0.67*	0.60*
<b>Sperm motility (%)</b>	0.56*	0.69*
<b>Sperm morphology (%)</b>	0.75*	0.68*
<b>Testosterone (ng/ml)</b>	0.58*	0.61*

\* Correlation is significant at the  $p < 0.05$  level (2-tailed); (n=20)

In the above table (Table 1), the testicular volume was positively and significantly correlated with the sperm count, sperm motility, percentage of sperm with normal morphology and serum testosterone levels.

\*Tel: +2347038648797

Email: tareonline@yahoo.com

Similarly, the testicular weights of the rats were positively and significantly correlated with the sperm count, progressive sperm motility, percentage of sperm with normal morphology and serum testosterone level.

#### 4. DISCUSSION

In this study, effects of administration of CdCl<sub>2</sub> in the drinking water of adult male rat on semen quality, testosterone and some testicular biometric parameters were investigated. Cadmium has been found to produce wide range of biochemical and physiological dysfunctions in humans and laboratory animals [22] and such target organs as the testes have been affected [23].

From the study there are evidences that exposure to cadmium chloride can interfere with male fertility. At the various dose of administration, Cadmium Chloride adversely affected ( $P=.05$ ) the semen quality parameters and testosterone, to the extent that it may impair fertility in the male Wistar rat used for the study. The histological results in CdCl<sub>2</sub> treated groups showed damage and degeneration of seminiferous tubules in testes. Cadmium, is very dangerous to testicular function [24]. According to Waisberg et al. [25], through increase in oxidative stress, cadmium chloride causes testicular damage, and impairs male fertility. Testicular damage caused by CdCl<sub>2</sub> is attributed to compete of CdCl<sub>2</sub> with Zinc in Zinc-containing enzymes and decreased activity of Testis-specific enzymes [26, 27]. In line with the above, Neeven et al. [28], Benoff et al. [15], Bench et al. [29] and Babara et al. [30] have also shown that cadmium chloride administration would interfere with male fertility, especially by its detrimental effect on testicular function. The decrease in testicular function from this study due to CdCl<sub>2</sub> treatment proves that cadmium impairs spermatogenesis. This result is similar to the observation of [31, 32]. The decrease in the testicular function could also be ascribed to the effect of cadmium chloride at the hypothalamic-pituitary-gonadal axis as suggested by Lafuente et al. [33] whose study revealed that cadmium affects the hypothalamic-pituitary-testicular axis function by acting at the three levels.

Different studies have shown that cadmium affects plasma gonadotropin levels [34]. Reduction in Testosterone hormone as shown in this study was manifested by lowered FSH and LH plasma levels [35, 36], as well as, CdCl<sub>2</sub> decrease the total testosterone hydroxylase activity [37] whose caused a dramatic decreasing in testosterone hormone. In this study the reduced testosterone level can also be attributed to interstitial fibrosis and interstitial edema caused by cadmium chloride which was in line with Laskey et al [38], who found that CdCl<sub>2</sub> caused necrosis in Leydig cells, these cells count primary synthesis site to testosterone hormone [39].

Cadmium chloride also significantly ( $P=.05$ ) decreased the testicular weight and volume, this could be understandable after reports from Rekha et al. [40] show derangement in some testicular structures such as seminiferous tubules and leydig cells that consist 70% to 80% of testicular mass [41].

According to Bailey *et al.* [42], testicular volume and weight are highly correlated. Also, Arai *et al.* [43] have suggested a possible correlation between testicular volume and testicular function which was also observed in this present study.

#### 5. CONCLUSION

The exposure of rats to cadmium chloride induces biochemical effects in the testes. Cadmium chloride decreased some testicular biometric parameters, semen quality and testosterone and these changes in the testicular function and testicular biometric parameters are positively correlated.



## REFERENCES

- [1] Berkley M. Infertility: Men and Sperm Health. Biology (3rd Ed.) St. Louis: Mosby (2004).
- [2] World Health Organisation. Manual for the Standardised Investigation and Diagnosis of the Infertile Couple. Cambridge: Cambridge University Press. (2000).
- [3] Sharlip, I.D., Jarow, J.P., and Belker, A.M. Best practice policies for male infertility. *Fertil. Steril.*; (2002): 77: 873–882.
- [4] Kenkel S, Rolf C, Nieschlag E. Occupational risks for male fertility: An analysis of patients attending a tertiary referral centre. *Int. J. Androl.*; (2001): 24: 318–26.
- [5] Ognjanovic, B.I., Markovic, S.D., Pavlovic, S.Z., Zikic, R.V., Stajn, A.S., and Saicic Z.S. Effect of chronic cadmium exposure on antioxidant defense system in some tissues of rats: Protective effect of selenium. *Physiol. Res.* (2008). 57: 403 - 411.
- [6] Schroeder, H. A., Balassa, J. J., Hogencamp, J. C Abnormal trace metals in man. Cadmium. *J. Chronic Dis.* (1961).14, 236-58.
- [7] Lewis, G. P., Jusko, W. J., Coughlin, L. L., Hartz, S., Contribution of cigarette smoking to Cadmium accumulation in man. *Lancet.*(1972). 1(7745): 291 – 292.
- [8] Casalino, E., Calzaretti, G., Sblano, C., and Landriscina C. Molecular inhibitory mechanisms of antioxidant enzymes in rat liver and kidney by cadmium. *Toxicology.* (2002). 179. 37-50.
- [9] Weisberg, E. Smoking and reproductive health. *Clin. Reprod. Afr. J. Biotechnol. Fertil.* (1985). 3: 175-186.
- [10] Kostic M.M., Ognjanovic B., Dimitrijevic S., Zikic R.V., Stajn A., Rosic G.L. Cadmium – induced changes of antioxidant and metyabolic status in red blood cells of rat: In vivo effects. *Eur. J. Haematol.* (1993). 51: 86 - 92.
- [11] Zikic, R.V., Stajn, A.S., Ognjanovic, B.I., Saicic, Z.S., Kostic, M.M., Pavlovic, S.Z., and Petrovic, V.M. The effect of cadmium and selenium on the antioxidant enzyme activities in rat heart. *J. Environ Pathol. Toxicol. Oncol.*, (1998). 17: 259 – 264.
- [12] Pavlovic S.Z., Ognjanovic B.I., Stajn A.S., Zikic R.V., Saicic Z.S., Petrovic V.M. Antioxidant defense system in skeletal muscle of rats treated with cadmium. A possible protective role of coenzyme Q10. *Jugoslav . Med. Biochem.*, (2001). 20: 229 - 235.
- [13] King LM, Banks AW, George WJ Differences in Cadmium Transport to the Testis, Epididymis and Brain in Cd-sensitive and Resistant Murine Strains *Pharmacol. Exp. Ther.*, (1999). 2: 825-830.
- [14] Prozialeck, W. C.; Edwards, J. R.; Woods, J. M. The Vascular Endothelium as a Target of Cadmium Toxicity. *Life Sci.*, (2006). 79 (16): 1493-1506.
- [15] Benoff, S., Auburn, K., Marmar, L. J., Hurley, R. I. Link between low-dose environmentally relevant cadmium exposures and asthenozoospermia in a rat model. *Fertil. Steril.*, (2008). 89 ,2 : 73 – 79.
- [16] Ibrahim M. E.; Sameh A. Y. The Antagonistic Effect of Chlorpromazine on Cadmium Toxicity . *Toxicology and Applied Pharmacology* (2002). 161. 34-3.

\*Tel: +2347038648797

Email: tareonline@yahoo.com

- [17] Oluyemi A., Ayodele O. Arowojolu, O., Shittu B. and John I A. Cadmium toxicity: a possible cause of male infertility in Nigeria. *Reprod. Biol.* (2006). 6 (1): 17 - 30.
- [18] Mervat, M .K., Abeer, H. A , Kawkab, A . A and Gehan, M. K. Exposure of Adult Male Rats to Cadmium: Assessment of Sexual Behaviour, Fertility, Aggression as well as Anxiety like Behaviour with Special Reference to Biochemical and Pathological Alterations. *Life Sci. J.* (2011). 8 (2): 106-119.
- [19] Bearden J.H.; Fluquary J.W. *Appl Anim Reprod.* Reston, VA: Rreston Publishers, (1980). p. 157 – 169.
- [20] Miller P.G.; Pass N.P. *Manual of infertility and artificial insemination in cattle*. Biolliere Tindall and Cox, London, U.K., (1952). p. 29 - 35.
- [21] Ismail A.A. The role of testosterone measurement in the investigation of androgen disorders. *Ann. Clin. Biochem.* (1986), 23: 113 - 134.
- [22] Santos FW, Oro T, Zeni G, Rocha JBT. Do Nascimento PC & Nogueira CW. Cadmium induced testicular damage and its response to administration of succimer and diphenyl discelenide in mice. *Toxicol Lett*; . (2004). 152: 255-63
- [23] Oteiza PI, Adonaylo VN & Keen CL. Cadmium induced testes oxidative damage in rats can be influenced by Zinc intake. *Toxicol*; (1999). 137 (1): 13-22.
- [24] Li LH, Heindel JJ. Sertoli cell toxicants. In: Korach KS, Ed. *Reproductive and Developmental Toxicology*. New York: Marcel Dekker, (1998). 655–691.
- [25] Waisberg, M., Joseph, P., Hale, B., and Beyersmann, D. Molecular and cellular mechanisms of cadmium carcinogenesis: a review. *Toxicology*. (2003). 192: 95 – 117.
- [26] Verboost, P . M., Flik, G., Look, R. A. and Wendelaar, B. S. Cadmium inhibits plasma membrane calcium transport. *J. Membr . Biol .* (1988). 102:97-104
- [27] Caslino, E., Sblano, C. and Landriscina, C. Enzyme activity alteration by cadmium administration to rats: the possibility of iron involvement in lipid peroxidation. *Arch . Biochem . Biophys.* (1997). 346: 171-179.
- [28] Neveen , T. , Elbetieha , A. , Darmani, H. , and Khamas, W. Effects of long term exposure to Cadmium chloride on fertility in adult male mice. *Vet. Res.* ( 2007 ). 1 (2): 40 -48.
- [29] Bench, G., Corzett, M.H., Martinelli,R., and Balhorn, R. Cadmium concentrations in the testes, sperm and spermatids of mice subjected to long – term Cadmium chloride exposure. *Cytometry* (1999). 35: 30 – 36.
- [30] Barbara P., Lukaszuk, A., Senczuk, W. The anti – reproductive effects of long term oral administration of Cadmium on the adult male rat. *Int. J. Androl.* (2008).2: 150-161.
- [31] Xu, B., Chin, S. E., Tsakak, M. and Ong, C.N. Trace elements in blood and seminal plasma and their relationship to sperm quality. *Reprod . Toxicol .* (1993) 7:613-618.
- [32] Chia, S. E., Xu, B., Ong, C.N., TSAKOK, F. M. H. and Lee, S.T. Effect of cadmium and cigarette smoking on human semen quality. *Int . J .fertil.* (1994). 39:292-298.
- [33] Lafuente A., Márquez N., Pérez-Lorenzo M., Pazo D. and Esquifino A.I. Cadmium Effects on Hypothalamic-Pituitary-Testicular Axis in Male Rats. *Exp Biol Med Vol.* (2001). 226(6):605–611.

\*Tel: +2347038648797

Email: tareonline@yahoo.com

- [34] Lafuente A, Blanco A, Ma´rquez N, Alvarez-Demanuel E, Esquifino AI. Effects of acute and subchronic cadmium administration on pituitary hormone secretion in rat. *Rev Esp Fisiol.* (1997). 53:265–270.
- [35] Paksy, K., Varga, B. and Lazar, P.. Zinc protection against cadmium-induced infertility in female rats. Effect of zinc and cadmium on the progesterone production of cultured granulose cells. *Bio . Metals.* (1996). 10: 27-35.
- [36] Lafuente, A., Marquez , N., Perez-Lorenzo, M., Pazo, D. and Esquifino, A., I. Pubertal and postpubertal cadmium exposure differentially affects the hypothalamic-pituitary- testicular axis function in the rat. *Food and Chemical Toxicology.* (2000). 38: 913-923.
- [37] Clark, J. I., Jimenez, B., Evans, S. L., Barrow, R., Winfree, M. and Mrotek, J. J. Cadmium-induced sexual dysfunction does not in vive increased hepatic metabolism of testosterone nor increased circulating levels of corticosterone .*Physiol . Behav .* (1994). 56: 975-981
- [38] Laskey, J. W.,. Rehnberg , G. L., Laws, S. C. and Hein , J.F. Reproductive effects of low acute doses of cadmium chloride in adult male rats. *Toxicol. Appl . Pharmacol.* (1984). 73:250- 255
- [39] Guyton, A. C. and Hall, J. E. Textbook of medical physiology. 9th ed. W.B. Saunders company. Philadelphia. (1996). 998 - 1000
- [40] Rekha D K, Tripathi Y, Raghuveer CV, Sheila R P, Ramaswamy C, Priya K. Role of vitamin c as an antioxidant in cadmium chloride Induced testicular damage. *Int J. App. Biol. & Pharm Tech.* (2011). 2 (3): 484-488.
- [41] Setchell, B.P., and Brooks, D.E. Anatomy, vasculature, innervations and fluids of the male reproductive tract. In: Knobil, E., and Neil, J.D., eds *The physiology of Reproduction.* New York, NY; Raven, (1998). 753 – 836.
- [42] Bailey, J.L., Hudson, R.S., Rowe, T.A., Riddell, M.G., Wolfe, D.F., and Carson, R.L. Caliper and Ultrasonographic measurements of bovine testicles and mathematical formula for determining testicular volume and weight in vivo. *Radiology.* (1998). 222: 114–119.
- [43] Arai, T., Kitahara, S., Horiuchi, S., Sumi, S., and Yoshida, K. Relationship of testicular volume to semen profiles and serum hormone concentrations in infertile Japanese males. *Int J Fertil.* (1998). 43:40–47.