

**Original Research Article**

**Screening for Vascular Disease in Pulmonary Rehabilitation**

**ABSTRACT**

**Background:** Several of the comorbidities and risk factors associated with COPD and other pulmonary disorders are also similar to those for lower extremity peripheral artery disease (PAD) and cerebrovascular disease. The later condition is often manifested as cognitive impairment, not only from large vessel associated stroke, but also from vascular cognitive impairment (VCI). We hypothesized that pulmonary rehabilitation (PR) would be an appropriate site to discern the prevalence of both PAD and mild cognitive impairment (MCI). Individuals in PR with severe pulmonary disease may have impaired cognition, which could impact learning and treatment adherence while PAD may contribute to functional limitation.

**Study Design:** In this pilot study of 14 patients enrolled in phase II PR, we describe the prevalence of PAD and cognitive impairment, the utility of screening tools, and the potential relationship between PAD, depression, and cognitive impairment.

**Results:** Twenty-eight percent of patients had an abnormal ABI with two subjects manifesting low ( $< 0.90$ ) and two subjects with abnormally high ( $> 1.40$ ) ABI. The Edinburgh Claudication Questionnaire was negative in all subjects and was not useful in the detection of PAD (sensitivity 0%, specificity 11%, PPV 0%, NPV 78.57%). The Walking Impairment Questionnaire did not differentiate patients with or without abnormal ABIs with respect to walking distance (40.02 vs. 64.46 meters,  $P = .154$ ), speed (27.15 vs. 25.92 miles per hour,  $P = .545$ ), or symptoms (46.88 vs. 63.75,  $P = .152$ ). Furthermore, eleven of the fourteen patients (79%) had an overall score of less than 26 on the Montreal Cognitive Assessment, indicating cognitive impairment. Scores indicating mild to moderate depression or anxiety measured by the PHQ-9 and STAI questionnaires revealed a concurrence between depressive and anxiety symptoms.

**Keywords:** [Pulmonary Rehabilitation, PAD, COPD, Mild Cognitive Impairment]

**1. INTRODUCTION**

Pulmonary disorders are an increasing cause of disability and functional impairment, followed by high rates of mortality. [1] Patients enrolled in a pulmonary rehabilitation (PR)

program are provided with individually tailored therapy to minimize their disability, to improve functional impairment and disease specific knowledge, and to decrease the psychological burden of their disorder. A majority of patients enrolled in PR are diagnosed with chronic obstructive pulmonary disease (COPD), the fourth-leading cause of death in the United States. [1] Several of the co-morbidities and risk factors associated with this and other pulmonary disorders are notably similar to those for lower extremity peripheral artery disease (PAD) and cerebrovascular disease. Symptomatic and functional consequences of PAD emerge from intermittent or chronic leg ischemia leading to claudication. Cerebrovascular disease is often manifested as cognitive impairment, not only from large vessel associated strokes, but also from vascular cognitive impairment (VCI). This latter condition is a heterogeneous group of disorders including VCI-no dementia, vascular dementia, and cognitive impairment of mixed (Alzheimer's disease and vascular dementia) origin. [2] Additionally, patients with VCI-no dementia have a high risk of progression to dementia, mixed primary neurodegenerative dementia, and vascular dementia, particularly if they have recurrent strokes. [2] Furthermore, structured lower extremity exercise is the fundamental component of treatment for claudication, and there is an increasing amount of evidence affirming that it may result in improvement of cognitive dysfunction. [3] Therefore, we hypothesized that PR would be an appropriate site to discern the prevalence of both PAD and mild cognitive impairment (MCI), as well as potentially to administer therapeutic interventions targeted to improve both vascular maladies.

PAD is often undetected and untreated, despite the high prevalence of this disease and its significant clinical consequences. [3] The ankle-brachial index (ABI) is a widely used non-invasive and cost effective test for screening, which compares the higher of the brachial artery systolic blood pressures with both the left and right lower extremities (either the *dorsalispedis* or *posterior tibial*) systolic blood pressure in a simple ratio. [4] Standardized questionnaires, validated in settings other than PR have been used as well. The Edinburgh Claudication Questionnaire is one such questionnaire that has been validated as a PAD screening tool with reported 91% sensitivity and 99% specificity for diagnosis in the general populations. [5] Additionally, the Walking Impairment Questionnaire is another validated method that assesses functional limitations by helping identify the degree of walking ability in the PAD populations in comparison to graded treadmill tests. [6]

Cognitive impairment is increasingly recognized as common in many medical conditions and may relate to the ability of the cardiovascular system to provide adequate brain perfusion. Reduced cerebral blood flow and structural brain changes including greater global and region-specific atrophy are often implicated, as past work in patients with COPD or depression has demonstrated a strong relationship between cognitive dysfunction and decreased fitness in patients with COPD or depression [7]. Specifically, the recent Cardiovascular Risk Factors, Aging and Dementia (CAIDE) study concluded that midlife COPD and asthma were associated with an almost two-fold risk of mild cognitive impairment and dementia later in life [8]. Therefore, it may be hypothesized that individuals in PR with severe pulmonary disease may have impaired cognition, and this especially may be the case with those diagnosed with vascular diseases such as PAD.

## 2. MATERIALS AND METHODS

### 2.1 Goals

The primary goal of this study was to determine the validity of screening tools in detecting PAD in comparison to ABI status in a representative population of patients in pulmonary rehabilitation. Secondary goals included the detection of cognitive impairment (defined by standardized neuropsychological testing with the Montreal Cognitive Assessment, MoCA)

and of depression. Thus, we sought to find a quantitative relationship between these important medical conditions.

## 2.2 Protocol

This was a prospective observational pilot study of 25 consecutive subjects enrolled in phase II PR at University Hospitals Harrington Heart & Vascular Institute's Pulmonary Rehabilitation Program in Ohio. All patients enrolled in PR between June 21, 2013 and August 27, 2013 were considered for this study. Exclusion criteria included individuals with required use of an assistive device (walker, cane, or wheelchair) for ambulation (n=2), documented dementia (n=0), or failure to provide informed consent (n=10).

The research assistant screened subjects for the study during the course of the patients' rehabilitation sessions. If the subject met the selection criteria, informed consent was obtained the day of screening or at the following session.

The study was conducted with minimal interruption to the course of pulmonary rehabilitation. After consent was obtained, the Walking Impairment Questionnaire, the Edinburgh Claudication Questionnaire, and the Seton Medical Center Pulmonary Knowledge Test were administered. Following the first set of questionnaires, the research assistant obtained ABI measurements according to the American Heart Association recommendations. [9]At the next scheduled rehabilitation session, the following surveys were administered: the Montreal Cognitive Assessment (MoCA), the State Trait Anxiety Inventory, and the Patient Health Questionnaire 9 (PHQ-9), which were used to screen for anxiety and depression symptoms respectively.

## 2.3 Data Analysis

The prevalence of peripheral artery disease, defined as an ABI  $<0.9$  or  $>1.4$  was described for the overall study population. Baseline characteristics of subjects with abnormal ABI were compared to those subjects with normal ABI. Furthermore, continuous variables were reported with standard descriptive statistics including mean and standard deviation. Between-group differences were tested for statistical significance with an unpaired t-test, and categorical variables were described as frequency and percentage. The test characteristics of the Edinburgh Claudication Questionnaire for the detection of abnormal ABI in this population were described in terms of sensitivity, specificity, and negative and positive predictive value. Unpaired t-tests were used to compare the walking ability scores on the Walking Impairment Questionnaire of patients with normal ABI to those with abnormal ABI. Additionally, the prevalence of depressive symptoms, assessed by PHQ-9, was described as percentages. Similarly, the overall scores on the MoCA were compared between both groups by percentages. The pulmonary knowledge of the rehabilitation patients was related with standard descriptive statistics. Lastly, the results of the MoCA were correlated with those of the PHQ-9 and STAI questionnaires using chi-square testing and two sample t-tests. All statistical tests were two-sided with a 0.05 significance level. Analyses were performed with SPSS version 20 (IBM SPSS Statistics for Windows, Armonk, NY).


## 3. RESULTS

### 3.1 Patient Population

Twenty-five consecutive patients enrolled in phase II pulmonary rehabilitation were assessed for eligibility and all met criteria; 8 were not consented within the first six exercise sessions

123 and 17 agreed to participate however 3 did not complete testing due to early withdrawal from  
124 the program.

125 The prevalence of PAD in this population of PR patients was 28% (4/14), with two subjects  
126 manifesting low ( $< 0.90$ ) and two subjects with abnormally high ( $> 1.40$ ) ABI.(Figure 1)



127

128 **Fig. 1.Prevalence of PAD in PR**  
129 *PAD= Peripheral Artery Disease; PR= Pulmonary Rehabilitation*

130

131 Table 1 contains the baseline characteristics of subjects with normal ABI compared to those  
132 with abnormal ABI as well as risk factors for PAD.

133 **Table 1.Baseline Characteristics and Risk Factors for PAD in Both ABI Groups**  
134 *DM=; Diabetes Mellitus HTN= Hypertension; DLD= Dyslipidemia; COPD= Chronic Obstructive*  
135 *Pulmonary Disorder; OA= Osteoarthritis*

136

Demographic	Normal ABI	Abnormal ABI
Number	10 (72%)	4 (28%)
Age	66 (+/- 7.5)	66.8 (+/- 12.8)
Sex		
<i>Male</i>	4 (28.6%)	2 (14.3%)
<i>Female</i>	6 (42.8%)	2 (14.3%)
Race		
<i>African American</i>	5 (35.7%)	2 (14.3%)

<i>White</i>	5 (35.7%)	2 (14.3%)
BMI (kg/m <sup>2</sup> )	32.3 (+/- 9.2)	30.4 (+/-12.6)
Past Medical History		
<i>PAD</i>	0	0
<i>DM</i>	2 (20%)	0
<i>HTN</i>	4 (40%)	3 (75%)
<i>DLD</i>	1 (10%)	1 (25%)
<i>COPD</i>	10 (100%)	2 (50%)
<i>Asthma</i>	2 (20%)	1 (25%)
<i>OA</i>	6 (60%)	2 (20%)
Smoking History		
<i>Current/Former</i>	9 (90%)	4 (100%)
<i>Never</i>	1 (10%)	0

137

138

### 3.2 Risk Factors

139

140

141

142

143

144

The total number of traditional risk factors for PAD was assessed in both groups, including the presence of smoking history, diabetes, obesity (BMI>30), hypertension, dyslipidemia, and age >50. Most of the patients in each PR group displayed at least three risk factors, however there was no statistically significant difference between the average numbers of risk factors when comparing subjects with normal ABI to those with abnormal ABI ( $P=.465$ ).

145

### 3.3 Edinburg Claudication Questionnaire (ECQ)

146

147

148

149

The Edinburg Claudication Questionnaire performed poorly in this population of PR patients, as it was negative in all subjects. Therefore, it was not useful in the detection of PAD in this population. (sensitivity 0%, specificity 11%, PPV 0%, NPV 78.57%).

150

### 3.4 Walking Impairment Questionnaire (WIQ)

151

152

153

154

155


The WIQ also performed poorly in this population, as it did not correlate with the presence of PAD. The questionnaire did not reveal any differences between patients with abnormal compared to normal ABIs with regard to walking distance, speed, or symptoms. (Table 2;  $P=.154$ ,  $P=.545$ ,  $P=.152$ )

**Table 2. Average Results on the WIQ between Patients with Abnormal and Normal ABIs**  
ABI= Ankle-Brachial Index

Section	Abnormal ABI (average score)	Normal ABI (average score)	P-value
Walking Distance (m)	40.015	64.456	0.154
Walking Speed (mph)	27.15	25.92	0.545
Walking Symptoms	46.875	63.75	0.152

### 3.5 Montreal Cognitive Assessment (MoCA)


Eleven of the fourteen patients (79%) had an overall score of less than 26, indicating cognitive impairment. One of the three patients who scored >26 had an abnormal ABI. Among the sub-scores, lowest performances were found in abstraction, delayed recall, and visuospatial/executive skills, in which 9 of 14 patients completed the alternating trail making incorrectly. Even with a high proportion of patients scoring less than 26, there was not statistically significant evidence of a relationship between low scores and patients with abnormal ABIs ( $P=.837$ ). (Figure 2)



**Fig. 2. MoCA Results of Patients with Abnormal and Normal ABIs**  
Scores >26 considered normal

### 3.6 Seton Knowledge Test

The mean score on the Seton Knowledge test in this population was a 19/25 (76%), which would be considered average performance. Six patients scored 19 points or below, whereas 8 patients scored 20 or above. There was a large range of scores with a minimum of 11 and a maximum of 24. Of the six patients with scores 19 or less, all scored <26 on the MoCA, five scored minimal to mild depression on the PHQ-9, and five scored low to moderate state anxiety on the STAI. Moreover, half of the six patients had abnormal ABIs. (Figure 3)



**Fig. 3. Performance on the MoCA& Seton Knowledge Test Amongst Both ABI Groups**

### 3.7 STAI/PHQ-9

Most patients' scores correlated to moderate state anxiety on the STAI questionnaire and similarly, those of the PHQ-9 corresponded with none, minimal, or mild depression. Although, one patient scored moderate to severe state anxiety and another scored moderate depression. Three of the four patients with abnormal ABIs that scored low to moderate on the STAI, and minimal to mild depression on the PHQ-9, had an overall score of <26 on the MoCA. Similarly among patients with normal ABIs, 9 of 10 patients that scored none to moderate on the PHQ-9, scored <26 on the MoCA. Despite this trend, there was not a statistically significant correlation with depression and anxiety levels and results of the MoCA among both patient groups ( $P=.466$ ,  $P=.075$ ).

## 4. DISCUSSION

### 4.1 Screening for Occult Peripheral Artery Disease

PAD is prevalent in patients with known cardiovascular disease as reported in previous studies, [2] however little is known about its relation to pulmonary disease and its prevalence in the pulmonary rehabilitation population. This study confirms and updates prior studies indicating that lower extremity PAD is prevalent in the pulmonary rehabilitation population. Despite the ability of the Edinburgh Claudication Questionnaire and other screening modalities to detect PAD in the general population, they are less sensitive in the PR population. These patients are likely to have other functionally limiting co-morbid conditions that may mask the presence of typical claudication symptoms, confounding the clinical detection of PAD. Additionally, selective screening based on risk factor stratification is of limited value in the PR population. As a result, this supports the application of ABI as a simple screening test for all patients enrolled in PR to detect PAD, as many patients are asymptomatic or have atypical symptoms.



217

## 218 **4.2 Prevalence of Mild Cognitive Impairment**

219

220 Cognitive impairment is prevalent in our representative PR population. It has been found that  
 221 patients with Vascular Cognitive Impairment (VCI) even if they do not have lesions on  
 222 neuroimaging can have a notably poor prognosis.[2] Additionally, lower scores on the Seton  
 223 Knowledge test could be related to below normal scores on the MoCA, as all six patients that  
 224 answered 19 or fewer questions correctly out of 25 on the Seton Knowledge test also scored  
 225 lower than 26 on the MoCA. Half of these patients had abnormal ABIs. This may reflect an  
 226 inability to retain information learned in rehabilitation sessions as a consequence of MCI. As  
 227 a result, there is an emerging concurrence between PAD and cognitive impairment.

## 228 **4.3 Prevalence of Depressive and Anxiety Symptoms**

229

230 Past prospective studies have found that depressive symptoms are associated with cognitive  
 231 decline and that the effect of anxiety on cognitive function is indirectly mediated by  
 232 depression [10]. Thus, the established relationship between cognitive impairment and  
 233 depression can be taken into consideration amongst patients with PAD, who have markedly  
 234 reduced health-related quality of life and higher prevalence of depression. [11] The  
 235 association between depressive symptoms and reduced cognitive function can be explained  
 236 by the ruminative processing of negative information that may reduce effort and attention to  
 237 demanding cognitive tasks. [10] Furthermore, depression in patients with cognitive  
 238 impairment is marked by abnormalities on neuropsychological tests including impairments in  
 239 memory, attention, and executive function [12], all of which are assessed in the MoCA.  
 240 Consequently, our results affirm the existing relationship between depressive and anxiety  
 241 symptoms and cognitive impairment, and also allude to a relationship between PAD and  
 242 depression and anxiety.

243

## 244 **5. STUDY LIMITATIONS AND IMPLICATIONS**

245

246 PAD was defined in our study by abnormal ABI. Other less common but more sophisticated,  
 247 costly, and often higher risk diagnostic methods such as pulse volume recording, computed  
 248 tomography angiography, and magnetic resonance angiography were not utilized given  
 249 financial and logistical restrictions. Furthermore, this pilot study was limited by the small  
 250 sample size, making it difficult to correlate MoCA scores with results of the PHQ-9 and the  
 251 STAI among patients with abnormal and normal ABIs.

252 As the prevalence of PAD is common in both pulmonary and cardiac rehabilitation [13],our  
 253 study extends other work for future studies assessing MCI in a larger PR population.  
 254 Furthermore, inquiry into the prevalence of MCI in patients with PAD would be of great  
 255 significance as well considering both of their clinical consequences. In a population- based  
 256 study, progression from VCI-no dementia to incident dementia occurred at the highest rate in  
 257 patients with both memory and functional impairment at baseline. Equivalently, up to a third  
 258 of people change diagnostic category (no cognitive impairment; cognitive impairment-no  
 259 dementia; dementia) within one year after stroke. [2] Therefore, cognitive impairment has  
 260 become increasingly common and debilitating, and should be a target for further study in the  
 261 proposed populations.

## 262 **6. CONCLUSIONS**

263

264 PAD is common in patients in phase II pulmonary rehabilitation and is often clinically occult.  
 265 It is infrequently detected by standard screening questionnaires in this population,



266 suggesting a need for a broad-based screening strategy including ABI measurements. In  
 267 fact, unappreciated PAD may correlate with mild cognitive impairment, which was observed  
 268 in a significant percentage of patients. Overall, our results have affirmed the substantial  
 269 prevalence of both PAD and cognitive impairment in PR. We found a positive correlation  
 270 between low Seton Knowledge test scores and low MoCA results, indicating the lack of  
 271 acquisition of disease related knowledge, which is one of the primary goals of PR. The  
 272 concurrence between depressive and anxiety symptoms and cognitive impairment was  
 273 affirmed as well. Individuals with depression have a greater tendency to submit to unhealthy  
 274 habits and risks, which may lead to increasing the risks of cardiovascular disease, cognitive  
 275 impairment, or worsening existing conditions. [12] Therefore, our conclusions have led us  
 276 to postulate a possible relationship between PAD and MCI, along with that of PAD and  
 277 depression and anxiety. Future studies, with a large sample size, will provide more insight on  
 278 ameliorating care for patients whose vascular disease affects cognitive function and mental  
 279 health.

280  
 281

## 282 **CONSENT**

283

284 All authors declare that 'written informed consent was obtained from the patient (or other  
 285 approved parties) for publication of this case report and accompanying images. A copy of  
 286 the written consent is available for review by the Editorial office/Chief Editor/Editorial Board  
 287 members of this journal.

288

## 289 **ETHICAL APPROVAL**

290

291 All authors hereby declare that all experiments have been examined and approved by the  
 292 appropriate ethics committee and have therefore been performed in accordance with the  
 293 ethical standards laid down in the 1964 Declaration of Helsinki.

294

## 295 **REFERENCES**

296

297 1. Pulmonary Rehabilitation Executive Summary\*: Joint American College Of Chest  
 298 Physicians/American Association Of Cardiovascular And Pulmonary Rehabilitation  
 299 Evidence-Based Clinical Practice Guidelines Ries AL, Bauldoff GS, Carlin BW, et al. *Chest*.  
 300 2007;131(5\_suppl):1S-3S.

301

302 2. Moorhouse, P., & Rockwood, K. (2008). Vascular cognitive impairment: Current  
 concepts and clinical developments. *Lancet Neurology*, 7, 246-255.

303

304 3. Hamburg NM, Balady GJ. Exercise rehabilitation in peripheral artery disease: functional  
 impact and mechanisms of benefits. *Circulation*. 2011 Jan 4; 123(1): 87-89.

305

306 4. Hirsch AT, Haskal ZJ, Hertzner NR, Bakal CW, Creager MA, Halperin JL, et al. ACC/AHA  
 307 2005 Practice Guidelines for the management of patients with peripheral arterial disease  
 308 (lower extremity, renal, mesenteric, and abdominal aortic): a collaborative report from the  
 309 American Association for Vascular Surgery/Society for Vascular Surgery, Society for  
 310 Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology,  
 311 Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines  
 312 (Writing Committee to Develop Guidelines for the Management of Patients With Peripheral  
 313 Arterial Disease): endorsed by the American Association of Cardiovascular and Pulmonary  
 Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing;

- 314 TransAtlantic Inter-Society Consensus; and Vascular Disease Foundation.Circulation. 2006  
315 Mar 21;113(11):e463–654.
- 316 5. Leng GC, Fowkes FG. The Edinburgh Claudication Questionnaire: an improved version  
317 of the WHO/Rose Questionnaire for use in epidemiological surveys. J. Clin. Epidemiol.1992  
318 Oct; 45(10): 1101-9.
- 319 6. Sagar SP, Brown PM, Zelt DT, Pickett WL, Tranmer JE. Further clinical validation of the  
320 walking impairment questionnaire for classification of walking performance in patients with  
321 peripheral artery disease. Int. J. Vasc. Med. 2012;2012:190641.
- 322 7. Garcia S, Alosco M-L, Sptiznagel M, et al. Cardiovascular fitness associated with  
323 cognitive performance in heart failure patients enrolled in cardiac rehabilitation. BMC  
324 CardiovascDisord. 2013; 13: 29.
- 325 8. Rusanen N, Ngandu T, Laatikainen T, Tuomilehto J, Soininen H, Kivipelto M. Chronic  
326 Obstructive Pulmonary Disease and Asthma and the Risk of Mild Cognitive Impairment and  
327 Dementia: A Population Based CAIDE Study. Current Alzheimer Research. 2013; 10:  
328 549-555.
- 329 9. Measurement and Interpretation of the Ankle-Brachial Index: A Scientific Statement From  
330 the American Heart Association. Circulation.2012; 126: 2890-2909.
- 331 10. Pietrzak RH, Maruff P, Woodward, M, Fredrickson, JB, Fredrickson AB, Krystal, JH,  
332 Southwick, SM, Darby, D. Mild Worry Symptoms Predict Decline in Learning and Memory in  
333 Healthy Older Adults: A 2-Year Prospective Cohort Study. American Journal of Geriatric  
334 Psychiatry.Vol. 20(3), p.266-275, Mar. 2012.  
335
- 336 11. McDermott MM, Guralnik JM, Ferrucci L, Tian L, Liu K, Liao Y, et al. Asymptomatic  
337 peripheral arterial disease is associated with more adverse lower extremity characteristics  
338 than intermittent claudication. Circulation. 2008 May 13;117(19):2484–91.
- 339 12. Sohani ZN, Samaan Z. Does Depression Impact Cognitive Impairment in Patients with  
340 Heart Failure? Cardiology Research and Practice. Vol. 2012.
- 341 13. Marty C. Tam, Richard Sukeena, Chen H. Chow, Chris T. Longenecker, Marianne Vest,  
342 Sri Krishnan Madan Mohan, Teresa Carman, Sahil A. Parikh, Richard A. Josephson,  
343 Peripheral Arterial Disease Worsens Outcomes for Patients in Cardiac Rehabilitation.  
344 Journal of Cardiopulmonary Rehabilitation and Prevention, Vol. 33, No. 5., p. 339, Sept.  
345 2013.  
346  
347  
348  
349