**Original Research Article** 1 2 Screening for Vascular Disease in Pulmonary 3 Rehabilitation 4 5 Sofia Arruda<sup>\*</sup>, Marty Tam<sup>1</sup>, Marianne Vest<sup>1</sup>, 6 Joel Hughes<sup>2</sup>, John Gunstad<sup>2</sup>, Mary Dolansky<sup>1, 3</sup>, Richard Sukeena,<sup>1</sup> 7 Richard Josephson<sup>1</sup> 8 9 10 <sup>1</sup>University Hospitals Case Medical Center, Harrington Heart and Vascular Institute 11 <sup>2</sup>Kent State University, Kent, Ohio <sup>3</sup>Payne Bolton School of Nursing, Case Western Reserve University, 12 13 Cleveland, Ohio 1<del>5</del> 16

#### ABSTRACT

17 18

**Background:** Several of the comorbidities and risk factors associated with COPD and other pulmonary disorders are also similar to those for lower extremity peripheral artery disease (PAD) and cerebrovascular disease. The later condition is often manifested as cognitive impairment, not only from large vessel associated stroke, but also from vascular cognitive impairment (VCI). We hypothesized that pulmonary rehabilitation (PR) would be an appropriate site to discern the prevalence of both PAD and mild cognitive impairment (MCI).Individuals in PR with severe pulmonary disease may have impaired cognition, which could impact learning and treatment adherence while PAD may contribute to functional limitation.

**Study Design**: In this pilot study of 14 patients enrolled in phase II PR, we describe the prevalence of PAD and cognitive impairment, the utility of screening tools, and the potential relationship between PAD, depression, and cognitive impairment.

**Results:** Twenty-eight percent of patients had an abnormal ABI with two subjects manifesting low (< 0.90) and two subjects with abnormally high (>1.40) ABI. The Edinburgh Claudication Questionnaire was negative in all subjects and was not useful in the detection of PAD (sensitivity 0%, specificity 11%, PPV 0%, NPV 78.57%.) The Walking Impairment Questionnaire did not differentiate patients with or without abnormal ABIs with respect to walking distance (40.02 vs. 64.46 meters, P= .154), speed (27.15 vs. 25.92 miles per hour, P=.545), or symptoms (46.88 vs. 63.75, P= .152). Furthermore, eleven of the fourteen patients (79%) had an overall score of less than 26 on the Montreal Cognitive Assessment, indicating cognitive impairment. Scores indicating mild to moderate depression or anxiety measured by the PHQ-9 and STAI questionnaires revealed a concurrence between depressive and anxiety symptoms.

19

20 Keywords: [Pulmonary Rehabilitation, PAD, COPD, Mild Cognitive Impairment]

- 21
- 22 23

#### 24 1. INTRODUCTION

25 26 Pulmonary disorders are an increasing cause of disability and functional impairment, 27 followed by high rates of mortality. [1] Patients enrolled in a pulmonary rehabilitation (PR) program are provided with individually tailored therapy to minimize their disability, to improve 28 functional impairment and disease specific knowledge, and to decrease the psychological 29 30 burden of their disorder. A majority of patients enrolled in PR are diagnosed with chronic 31 obstructive pulmonary disease (COPD), the fourth-leading cause of death in the United States. [1] Several of the co-morbidities and risk factors associated with this and other 32 33 pulmonary disorders are notably similar to those for lower extremity peripheral artery disease 34 (PAD) and cerebrovascular disease. Symptomatic and functional consequences of PAD 35 emerge from intermittent or chronic leg ischemia leading to claudication. Cerebrovascular 36 disease is often manifested as cognitive impairment, not only from large vessel associated 37 strokes, but also from vascular cognitive impairment (VCI). This latter condition is a heterogeneous group of disorders including VCI-no dementia, vascular dementia, and 38 cognitive impairment of mixed (Alzheimer's disease and vascular dementia) origin. [2] 39 Additionally, patients with VCI-no dementia have a high risk of progression to dementia, 40 41 mixed primary neurodegenerative dementia, and vascular dementia, particularly if they have recurrent strokes. [2] Furthermore, structured lower extremity exercise is the fundamental 42 43 component of treatment for claudication, and there is an increasing amount of evidence affirming that it may result in improvement of cognitive dysfunction. [3] Therefore, we 44 45 hypothesized that PR would be an appropriate site to discern the prevalence of both PAD and mild cognitive impairment (MCI), as well as potentially to administer therapeutic 46 47 interventions targeted to improve both vascular maladies.

48

49 PAD is often undetected and untreated, despite the high prevalence of this disease and its 50 significant clinical consequences. [3] The ankle-brachial index (ABI) is a widely used non-51 invasive and cost effective test for screening, which compares the higher of the brachial artery systolic blood pressures with both the left and right lower extremities (either the 52 dorsalispedis or posterior tibial) systolic blood pressure in a simple ratio. [4] Standardized 53 questionnaires, validated in settings other than PR have been used as well. The Edinburgh 54 55 Claudication Questionnaire is one such questionnaire that has been validated as a PAD screening tool with reported 91% sensitivity and 99% specificity for diagnosis in the general 56 populations. [5] Additionally, the Walking Impairment Questionnaire is another validated 57 58 method that assesses functional limitations by helping identify the degree of walking ability in 59 the PAD populations in comparison to graded treadmill tests. [6]

60

61 Cognitive impairment is increasingly recognized as common in many medical conditions and 62 may relate to the ability of the cardiovascular system to provide adequate brain perfusion. 63 Reduced cerebral blood flow and structural brain changes including greater global and 64 region-specific atrophy are often implicated, as past work in patients with COPD or 65 depression has demonstrated a strong relationship between cognitive dysfunction and decreased fitness in patients with COPD or depression [7]. Specifically, the recent 66 Cardiovascular Risk Factors, Aging and Dementia (CAIDE) study concluded that midlife 67 68 COPD and asthma were associated with an almost two-fold risk of mild cognitive impairment 69 and dementia later in life [8]. Therefore, it may be hypothesized that individuals in PR with 70 severe pulmonary disease may have impaired cognition, and this especially may be the case 71 with those diagnosed with vascular diseases such as PAD.

72

#### 73 2. MATERIALS AND METHODS

74

#### 75 2.1 Goals

The primary goal of this study was to determine the validity of screening tools in detecting PAD in comparison to ABI status in a representative population of patients in pulmonary rehabilitation. Secondary goals included the detection of cognitive impairment (defined by standardized neuropsychological testing with the Montreal Cognitive Assessment, MoCA) and of depression. Thus, we sought to find a quantitative relationship between these important medical conditions.

#### 83 **2.2 Protocol**

84

82

This was a prospective observational pilot study of 25 consecutive subjects enrolled in phase II PR at University Hospitals Harrington Heart & Vascular Institute's Pulmonary Rehabilitation Program in Ohio. All patients enrolled in PR between June 21, 2013 and August 27, 2013 were considered for this study. Exclusion criteria included individuals with required use of an assistive device (walker, cane, or wheelchair) for ambulation (n=2), documented dementia (n=0), or failure to provide informed consent (n=10).

91 The research assistant screened subjects for the study during the course of the patients' 92 rehabilitation sessions. If the subject met the selection criteria, informed consent was 93 obtained the day of screening or at the following session.

94 The study was conducted with minimal interruption to the course of pulmonary rehabilitation. 95 After consent was obtained, the Walking Impairment Questionnaire, the Edinburgh 96 Claudication Questionnaire, and the Seton Medical Center Pulmonary Knowledge Test were 97 administered. Following the first set of questionnaires, the research assistant obtained ABI 98 measurements according to the American Heart Association recommendations. [9]At the 99 next scheduled rehabilitation session, the following surveys were administered: the Montreal 100 Cognitive Assessment (MoCA), the State Trait Anxiety Inventory, and the Patient Health 101 Questionnaire 9 (PHQ-9), which were used to screen for anxiety and depression symptoms 102 respectively.

103

#### 104 2.3 Data Analysis

105

106 The prevalence of peripheral artery disease, defined as an ABI <0.9 or >1.4 was described 107 for the overall study population. Baseline characteristics of subjects with abnormal ABI were compared to those subjects with normal ABI. Furthermore, continuous variables were 108 109 reported with standard descriptive statistics including mean and standard deviation. 110 Between-group differences were tested for statistical significance with an unpaired t-test, 111 and categorical variables were described as frequency and percentage. The test 112 characteristics of the Edinburgh Claudication Questionnaire for the detection of abnormal 113 ABI in this population were described in terms of sensitivity, specificity, and negative and 114 positive predictive value. Unpaired t-tests were used to compare the walking ability scores 115 on the Walking Impairment Questionnaire of patients with normal ABI to those with abnormal 116 ABI. Additionally, the prevalence of depressive symptoms, assessed by PHQ-9, was 117 described as percentages. Similarly, the overall scores on the MoCA were compared between both groups by percentages. The pulmonary knowledge of the rehabilitation 118 patients was related with standard descriptive statistics. Lastly, the results of the MoCA 119 were correlated with those of the PHQ-9 and STAI questionnaires using chi-square testing 120 121 and two sample t-tests. All statistical tests were two-sided with a 0.05 significance level. 122 Analyses were performed with SPSS version 20 (IBM SPSS Statistics for Windows, Armonk, 123 NY).

#### 3. RESULTS 124

125

#### **3.1 Patient Population** 126

127

128 Twenty-five consecutive patients enrolled in phase II pulmonary rehabilitation were assessed 129 for eligibility and all met criteria; 8 were not consented within the first six exercise sessions 130 and 17 agreed to participate however 3 did not complete testing due to early withdrawal from 131 the program.

132 The prevalence of PAD in this population of PR patients was 28% (4/14), with two subjects 133 manifesting low (< 0.90) and two subjects with abnormally high (>1.40) ABI.(Figure 1)



#### 135 Fig. 1. Prevalence of PAD in PR

- 136 PAD= Peripheral Artery Disease; PR= Pulmonary Rehabilitation
- 137

134

138 Table 1 contains the baseline characteristics of subjects with normal ABI compared to those 139 with abnormal ABI as well as risk factors for PAD.

#### 140 Table 1.Baseline Characteristics and Risk Factors for PAD in Both ABI Groups

141 DM=; Diabetes Mellitus HTN= Hypertension; DLD= Dyslipidemia; COPD= Chronic Obstructive 142 Pulmonary Disorder; OA= Osteoarthritis 143

Demographic	Normal ABI	Abnormal ABI	
Number	10 (72%)	4 (28%)	
Age	66 (+/- 7.5)	66.8 (+/- 12.8)	
Sex			
Male	4 (28.6%)	2 (14.3%)	

Female	6 (42.8%)	2 (14.3%)	
Race			
African American	5 (35.7%)	2 (14.3%)	
White	5 (35.7%)	2 (14.3%)	
BMI (kg/m <sup>2</sup> )	32.3 (+/- 9.2)	30.4 (+/-12.6)	
Past Medical History			
PAD	0	0	
DM	2 (20%)	0	
HTN	4 (40%)	3 (75%)	
DLD	1 (10%)	1 (25%)	
COPD	10 (100%)	2 (50%)	
Asthma	2 (20%)	1 (25%)	
OA	6 (60%)	2 (20%)	
Smoking History			
Current/Former	9 (90%)	4 (100%)	
Never	1 (10%)	0	

#### 144

#### 145 3.2 Risk Factors

146

147 The total number of traditional risk factors for PAD was assessed in both groups, including 148 the presence of smoking history, diabetes, obesity (BMI>30), hypertension, dyslipidemia, 149 and age >50. Most of the patients in each PR group displayed at least three risk factors, 150 however there was no statistically significant difference between the average numbers of risk 151 factors when comparing subjects with normal ABI to those with abnormal ABI (P=.465).

#### 152 **3.3 Edinburg Claudication Questionnaire (ECQ)**

153

The Edinburgh Claudication Questionnaire performed poorly in this population of PR patients, as it was negative in all subjects. Therefore, it was not useful in the detection of PAD in this population. (sensitivity 0%, specificity 11%, PPV 0%, NPV 78.57%.)

#### 157 3.4 Walking Impairment Questionnaire (WIQ)

158

The WIQ also performed poorly in this population, as it did not correlate with the presence of PAD. The questionnaire did not reveal any differences between patients with abnormal compared to normal ABIs with regard to walking distance, speed, or symptoms. (Table 2; P=.154, P=.545, P=.152)

## Table 2.Average Results on the WIQ between Patients withAbnormal and Normal ABIs ABI= Ankle-Brachial Index

165

Section	Abnormal ABI	Normal ABI	P-value
	(average score)	(average score)	
Walking Distance (m)	40.015	64.456	0.154
Walking Speed (mph)	27.15	25.92	0.545
Walking Symptoms	46.875	63.75	0.152

166

### 167 3.5 Montreal Cognitive Assessment (MoCA)

168

Eleven of the fourteen patients (79%) had an overall score of less than 26, indicating cognitive impairment. One of the three patients who scored >26 had an abnormal ABI. Among the sub-scores, lowest performances were found in abstraction, delayed recall, and visuospatial/executive skills, in which 9 of 14 patients completed the alternating trail making incorrectly. Even with a high proportion of patients scoring less than 26, there was not statistically significant evidence of a relationship between low scores and patients with abnormal ABIs (*P*=.837). (Figure 2)

176



- 177 178
- 179

Fig. 2.MoCA Results of Patients with Abnormal and Normal ABIs Scores >26 considered normal

- 180 181
- 182 **3.6 Seton Knowledge Test**

The mean score on the Seton Knowledge test in this population was a 19/25 (76%), which would be considered average performance. Six patients scored 19 points or below, whereas 8 patients scored 20 or above. There was a large range of scores with a minimum of 11 and a maximum of 24. Of the six patients with scores 19 or less, all scored <26 on the MoCA, five scored minimal to mild depression on the PHQ-9, and five scored low to moderate state anxiety on the STAI. Moreover, half of the six patients had abnormal ABIs. (Figure 3)





190 191 192

Fig. 3.Performance on the MoCA& Seton Knowledge Test Amongst Both ABI Groups

# 193194 **3.7 STAI/PHQ-9**

195

196 Most patients' scores correlated to moderate state anxiety on the STAI questionnaire and 197 similarly, those of the PHQ-9 corresponded with none, minimal, or mild depression. 198 Although, one patient scored moderate to severe state anxiety and another scored moderate 199 depression. Three of the four patients with abnormal ABIs that scored low to moderate on 200 the STAI, and minimal to mild depression on the PHQ-9, had an overall score of <26 on the 201 MoCA. Similarly among patients with normal ABIs, 9 of 10 patients that scored none to moderate on the PHQ-9, scored <26 on the MoCA. Despite this trend, there was not a 202 statistically significant correlation with depression and anxiety levels and results of the MoCA 203 204 among both patient groups (P=.466, P=.075).

### 205 4. DISCUSSION

206

## 207 4.1 Screening for Occult Peripheral Artery Disease

208

209 PAD is prevalent in patients with known cardiovascular disease as reported in previous 210 studies, [2] however little is known about its relation to pulmonary disease and its prevalence in the pulmonary rehabilitation population. This study confirms and updates prior studies 211 212 indicating that lower extremity PAD is prevalent in the pulmonary rehabilitation population. 213 Despite the ability of the Edinburgh Claudication Questionnaire and other screening 214 modalities to detect PAD in the general population, they are less sensitive in the PR 215 population. These patients are likely to have other functionally limiting co-morbid conditions 216 that may mask the presence of typical claudication symptoms, confounding the clinical detection of PAD. Additionally, selective screening based on risk factor stratification is of limited value in the PR population. As a result, this supports the application of ABI as a simple screening test for all patients enrolled in PR to detect PAD, as many patients are asymptomatic or have atypical symptoms.

221

222

223 224

### 225 4.2 Prevalence of Mild Cognitive Impairment

226

227 Cognitive impairment is prevalent in our representative PR population. It has been found that 228 patients with Vascular Cognitive Impairment (VCI) even if they do not have lesions on 229 neuroimaging can have a notably poor prognosis.[2] Additionally, lower scores on the Seton 230 Knowledge test could be related to below normal scores on the MoCA, as all six patients that 231 answered 19 or fewer questions correctly out of 25 on the Seton Knowedge test also scored 232 lower than 26 on the MoCA. Half of these patients had abnormal ABIs. This may reflect an 233 inability to retain information learned in rehabilitation sessions as a consequence of MCI. As 234 a result, there is an emerging concurrence between PAD and cognitive impairment.

### 235 **4.3 Prevalence of Depressive and Anxiety Symptoms**

236

237 Past prospective studies have found that depressive symptoms are associated with cognitive 238 decline and that the effect of anxiety on cognitive function is indirectly mediated by 239 depression [10]. Thus, the established relationship between cognitive impairment and 240 depression can be taken into consideration amongst patients with PAD, who have markedly 241 reduced health-related quality of life and higher prevalence of depression. [11] The 242 association between depressive symptoms and reduced cognitive function can be explained 243 by the ruminative processing of negative information that may reduce effort and attention to 244 demanding cognitive tasks. [10] Furthermore, depression in patients with cognitive 245 impairment is marked by abnormalities on neuropsychological tests including impairments in memory, attention, and executive function [12], all of which are assessed in the MoCA. 246 247 Consequently, our results affirm the existing relationship between depressive and anxiety 248 symptoms and cognitive impairment, and also allude to a relationship between PAD and 249 depression and anxiety.

250 251

## 5. STUDY LIMITATIONS AND IMPLICATIONS

252

PAD was defined in our study by abnormal ABI. Other less common but more sophisticated, costly, and often higher risk diagnostic methods such as pulse volume recording, computed tomography angiography, and magnetic resonance angiography were not utilized given financial and logistical restrictions. This pilot study was limited by the small sample size, precluding robust multivariate analysis or other analyses such as net reclassification. Future studies, with larger populations, could utilize these techniques to provide additional insight.

259 As the prevalence of PAD is common in both pulmonary and cardiac rehabilitation [13], our 260 study extends other work for future studies assessing MCI in a larger PR population. 261 Furthermore, inquiry into the prevalence of MCI in patients with PAD would be of great 262 significance as well considering both of their clinical consequences. In a population- based 263 study, progression from VCI-no dementia to incident dementia occurred at the highest rate in 264 patients with both memory and functional impairment at baseline. Equivalently, up to a third 265 of people change diagnostic category (no cognitive impairment; cognitive impairment-no 266 dementia; dementia) within one year after stroke. [2] Therefore, cognitive impairment has 267 become increasingly common and debilitating, and should be a target for further study in the 268 proposed populations.

Overall, pulmonary rehabilitation serves a broad spectrum of patients, even amongst those with a single index diagnosis such as COPD. Pulmonary therapies such as continuous positive air pressure (CPAP) and corticosteroids along with comorbidities, namely hyperlipidemia, diabetes mellitus, and hypertension, may influence vascular and muscle structure and function. [14] As a result, the listed therapies and comorbidities could potentially influence the development and clinical manifestations of PAD in patients enrolled in pulmonary rehabilitation and could be explored in future work.

#### 276 6. CONCLUSIONS

277

278 PAD is common in patients in phase II pulmonary rehabilitation and is often clinically occult. 279 It is infrequently detected by standard screening questionnaires in this population, 280 suggesting a need for a broad-based screening strategy including ABI measurements. In 281 fact, unappreciated PAD may correlate with mild cognitive impairment, which was observed 282 in a significant percentage of patients. Overall, our results have affirmed the substantial 283 prevalence of both PAD and cognitive impairment in PR. We found a positive correlation 284 between low Seton Knowledge test scores and low MoCA results, indicating the lack of 285 acquisition of disease related knowledge, which is one of the primary goals of PR. The 286 concurrence between depressive and anxiety symptoms and cognitive impairment was 287 affirmed as well. Individuals with depression have a greater tendency to submit to unhealthy 288 habits and risks, which may lead to increasing the risks of cardiovascular disease, cognitive 289 impairment, or worsening existing conditions. [12] Therefore, our conclusions have leaded us to postulate a possible relationship between PAD and MCI, along with that of PAD and 290 291 depression and anxiety. Future studies, with a large sample size, will provide more insight on 292 ameliorating care for patients whose vascular disease affects cognitive function and mental 293 health

#### 294 AUTHORS' CONTRIBUTIONS

Dr. Richard Josephson was responsible for the coordination of the overall study, including: the study design, collaboration amongst investigators, data analysis, and manuscript preparation. Sofia Arruda wrote the first draft of the manuscript. Marty Tam and Sofia Arruda managed the analyses of the study and the literature searches. Dolanksy, Hughes, and Gunstad contributed to the study design and selection of cognitive biobehavioral instruments. Vest and Sukeena assisted with staff training, manuscript review, and study design. 302

#### 303 CONSENT

304

All authors declare that 'written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal.

#### 310 ETHICAL APPROVAL

311

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

- 316 **REFERENCES**
- 317

315

Pulmonary Rehabilitation Executive Summary\*: Joint American College Of Chest
 Physicians/American Association Of Cardiovascular And Pulmonary Rehabilitation
 Evidence-Based Clinical Practice GuidelinesRies AL, Bauldoff GS, Carlin BW, et al. *Chest.* 2007;131(5\_suppl):1S-3S.

- 322 2. Moorhouse, P., & Rockwood, K. (2008). Vascular cognitive impairment: Current 323 concepts and clinical developments. *Lancet Neurology*, 7, 246-255.
- 324 3. Hamburg NM, Balady GJ. Exercise rehabilitation in peripheral artery disease: functional 325 impact and mechanisms of benefits. Circulation. 2011 Jan 4; 123(1): 87-89.

326 4. Hirsch AT, Haskal ZJ, Hertzer NR, Bakal CW, Creager MA, Halperin JL, et al. ACC/AHA 327 2005 Practice Guidelines for the management of patients with peripheral arterial disease 328 (lower extremity, renal, mesenteric, and abdominal aortic): a collaborative report from the 329 American Association for Vascular Surgery/Society for Vascular Surgery, Society for 330 Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines 331 332 (Writing Committee to Develop Guidelines for the Management of Patients With Peripheral 333 Arterial Disease): endorsed by the American Association of Cardiovascular and Pulmonary 334 Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; 335 TransAtlantic Inter-Society Consensus; and Vascular Disease Foundation. Circulation. 2006 336 Mar 21;113(11):e463-654.

- 5. Leng GC, Fowkes FG. The Edinburgh Claudication Questionnaire: an improved version
  of the WHO/Rose Questionnaire for use in epidemiological surveys. J. Clin. Epidemiol.1992
  Oct; 45(10): 1101-9.
- Sagar SP, Brown PM, Zelt DT, Pickett WL, Tranmer JE. Further clinical validation of the
  walking impairment questionnaire for classification of walking performance in patients with
  peripheral artery disease. Int. J. Vasc. Med. 2012;2012:190641.
- 343 7. Garcia S, Alosco M-L, Sptiznagel M, et al. Cardiovascular fitness associated with
   344 cognitive performance in heart failure patients enrolled in cardiac rehabilitation. BMC
   345 CardiovascDisord. 2013; 13: 29.
- 8. Rusanen N, Ngandu T, Laatikainen T, Tuomilehto J, Soininen H, Kivipelto M. Chronic
   Obstructive Pulmonary Disease and Asthma and the Risk of Mild Cognitive Impairment and

348 Dementia: A Population Based CAIDE Study. Current Altzheimer Research. 2013; 10: 349 549-555.

9. Measurement and Interpretation of the Ankle-Brachial Index: A Scientific Statement From
 the American Heart Association. Circulation.2012; 126: 2890-2909.

10. Pietrzak RH, Maruff P, Woodward, M, Fredrickson, JB, Fredrickson AB, Krystal, JH,
Southwick, SM, Darby, D. Mild Worry Symptoms Predict Decline in Learning and Memory in
Healthy Older Adults: A 2-Year Prospective Cohort Study. American Journal of Geriatric
Psychiatry.Vol. 20(3), p.266-275, Mar. 2012.

356

McDermott MM, Guralnik JM, Ferrucci L, Tian L, Liu K, Liao Y, et al. Asymptomatic
 peripheral arterial disease is associated with more adverse lower extremity characteristics
 than intermittent claudication. Circulation. 2008 May 13;117(19):2484–91.

360 12. Sohani ZN, Samaan Z. Does Depression Impact Cognitive Impairment in Patients with
 361 Heart Failure? Cardiology Research and Practice. Vol. 2012.
 362

13. Marty C. Tam, Richard Sukeena, Chen H. Chow, Chris T. Longenecker, Marianne Vest,
Sri Krishnan Madan Mohan, Teresa Carman, Sahil A. Parikh, Richard A. Josephson,
Peripheral Arterial Disease Worsens Outcomes for Patients in Cardiac Rehabilitation.
Journal of Cardiopulmonary Rehabilitation and Prevention, Vol. 33, No. 5., p. 339, Sept.
2013.

369 14. Is there a correlation between OSAS duration/severity and carotid intima-media
 370 thickness? Ciccone, Marco Matteo et al. Respiratory Medicine, Volume 106, Issue 5, 740 –
 371 746.

- 372
- 373 374
- 375
- 376

377