

Unappreciated Cognitive Dysfunction is Common Among Patients Hospitalized with Cardiovascular Disease

ABSTRACT

Background Better understanding of the prevalence of unappreciated cognitive dysfunction among patients with cardiovascular disease during hospitalization is integral to patient-centered care. We tested the hypothesis that there is a substantial prevalence of debilitating cognitive dysfunction in cardiac patients which health care providers do not appreciate.

Methods This observational prospective study evaluated 51 patients on admission to a cardiac intensive care unit (CICU) who did not have a history of conditions known to adversely affect cognition and appeared cognitively intact to treating physicians and nursing staff. Patients underwent neuropsychological testing to assess the domains of global cognition, attention, memory, and executive function at important points during hospitalization.

Results Twenty-six percent of participants were impaired globally based upon the Modified Mini Mental Status Examination. Of the study's participants, 46% were impaired in memory on short recall and 40% were impaired on long recall based upon Rey Auditory Verbal Learning, while 49% were impaired based upon Digit Span. Further, 38% of participants were impaired in attention based upon Trail Making Test Part A. Twenty-five percent of participants were impaired in executive function based upon Trail Making Test Part B, and 18% were impaired based upon Frontal Assessment Battery.

Conclusions In patients hospitalized with cardiovascular disease, unappreciated cognitive dysfunction is common. The dysfunction involves multiple domains and likely impedes patient participation in longitudinal care and their comprehension of health education which ultimately hinders the transition from hospital to home care. This process creates a setting of poor self-management with significantly increased potential for rehospitalizations.

Keywords: cardiovascular disease, cognition, executive function, memory

1. INTRODUCTION

Cardiovascular disease (CVD) risk factors have been associated with cognitive decline in older populations.^{4, 5} These findings suggest that patients are in the middle of the vascular cognitive impairment continuum, performing worse than healthy age-matched peers, but substantially better than persons with dementia or other significant neurological disorders. This cognitive dysfunction can be subtle, and can go unrecognized by health care providers.

Patients with heart failure have been found to manifest difficulties with memory and concentration that are associated with poor health outcomes distinct from age-related memory and concentration abnormalities. Depending on the population under investigation, including inpatients and outpatients and the methodology used, it has been reported that 30% to 80% of patients with heart failure exhibit cognitive dysfunction⁶, particularly on tests of memory and executive function. Several mechanisms have been implicated in this cognitive dysfunction including reduced cerebral perfusion⁷ and cerebrovascular pathology.^{8, 9, 10, 11, 12}

Recent studies suggest that baseline cognitive dysfunction is associated with a five times greater risk of mortality and six times greater risk of functional decline in the future.^{13, 14, 15} Cognitive dysfunction is associated with dependency in activities of daily living¹⁶, which is

further linked to disability and frequent rehospitalization.^{17, 18} One study found cognitive dysfunction to be associated with in-hospital mortality and decreased 1-year survival among older heart failure patients.¹³ Cognitive dysfunction in attention, executive function, and memory can negatively impact patients' self-care behaviors such as taking medications, dietary compliance, and exercise. Such dysfunction also prevents patients from providing health care providers with accurate histories, which further reduces optimal medical care.

In hospitals, patient histories are often obtained on the first morning after admission and the patients' cognition plays a key role in providing an accurate history to health care providers. Furthermore, the majority of patient education regarding self-management occurs prior to discharge. As more and more patients are discharged from intensive care units, it is important to understand their cognitive status in order to ensure effective teaching related to discharge instructions.

We tested the hypothesis that there is a substantial prevalence of cognitive dysfunction in cardiac patients that is unappreciated by health care providers throughout the hospitalization course.

2. METHODOLOGY

Participants

A convenience sample of 51 patients being treated for acute cardiovascular conditions was recruited following their admissions to a cardiac intensive care unit at an academic tertiary care hospital. Patients were 18 years of age or older and were awake and conversant at the time of enrollment.

Criteria were chosen to maximize generalizability to other samples. Patients with conditions known to present cognitive dysfunction independent of their cardiac disease or its related conditions were excluded. Patients were excluded for a history of neurological disorder or injury (Alzheimer's disease, dementia, stroke, seizures) as well as moderate or severe head injury (defined as >10 minutes loss of consciousness). Patients with a past or current history of severe psychiatric illness, including psychotic disorders (schizophrenia and bipolar disorder), were excluded, but not those individuals with treated depression or anxiety disorders. Patients were also excluded if they had a history of alcohol or drug abuse within the last 5 years or a history of learning disorder or developmental disability. Patients were excluded for renal failure requiring dialysis, diagnosed untreated sleep apnea, and severely ill individuals, including those who were sedated, on a ventilator, or exhibiting acute hemodynamic changes. Furthermore, attending physicians and nurses were asked to exclude patients deemed inappropriate for the study due to clear cognitive dysfunction noted upon routine clinical examination. Thus, the study group represents a patient population that most health care providers would consider to have "normal" cognitive function.

Procedures

The local Human Subjects Institutional Review Board approved all study methods prior to conducting the study. Newly admitted patients were referred by nurse managers and physicians on the unit. Prior to approaching the patient, the patient's nurse was asked if he/she noted any cognitive dysfunction. If not, the patient was approached and informed of the study, and written informed consent was obtained. A series of neuropsychological tests were administered to consented participants, and when logistically feasible, data collection was repeated prior to discharge. Testing sessions required approximately 1 hour.

Measures

Neuropsychological Test Battery. Participants completed a battery of well-established and validated neuropsychological measures assessing multiple domains. Researchers examined

global cognitive functioning and cognitive performance in the domains of memory, language, and attention/executive function^{35, 36}. The test battery was administered on the first morning following admission (Time 1) and again within three hours of anticipated discharge (Time 2). The order of neuropsychological testing was not random but rather the Modified Mini Mental Status Examination first, Rey Auditory Verbal Learning, Trails testing, followed by Digit Span Backwards, Rey Auditory Verbal Learning Long Delay and Recognition, and then finally Frontal Assessment Battery. Repeat testing prior to discharge was intended to confirm that the cognitive dysfunction noted at Time 1 was not transitory or unique to admission. The following neuropsychological tests were administered:

Global Cognitive Functioning

Modified Mini Mental Status Examination (3MS; Teng & Chui, 1987) provides a screening measure of global cognitive function. It is comprised of several short tasks, including orientation, similarities, animal fluency, learning, and brief and delayed recall of a short list of target words, and a copying a simple geometric figure. It is sensitive to a range of cognitive dysfunction including Alzheimer's disease and other forms of dementia.

Attention/Executive Function

Trail Making Test Parts A and B (Reitan, 1958)⁴². The Trail Making Test Part A requires participants to connect a series of 25 numbered dots in ascending order from 1 to 25 as quickly as possible. Trail Making Test Part B adds a set-shifting component, requiring participants to alternate between numbers and letters in ascending order (eg, 1, A, 2, B). The first part of this test assesses psychomotor speed and visual scanning, whereas the second part evaluates the ability to respond to shifting demands.

*Frontal Assessment Battery*⁴³. The test includes subtests: (a) conceptualization and abstract reasoning (similarities test); (b) mental flexibility (verbal fluency test); (c) motor programming and executive control of action (Luris motor sequences); (d) resistance to interference (conflicting instructions); (e) inhibitory control (go-no go test); and (f) environmental autonomy (prehension behavior).

Memory

Rey Auditory Verbal Learning Test. Participants read a list of 15 words a total of 5 times followed by recalling as many words as possible. Following presentation and recall of a distraction list, participants are again asked to recall words from the original list. After a 30-minute delay, participants are again asked to recall target words from the original list. Finally, a recognition trial comprised of target words and foil (words not initially asked to learn) is completed.

Digit Span Backwards test. Participants hear a series of numerical digits read with one second intervals between digits. The participants are then asked to repeat the digits aloud in reverse order. Comprising each sequence gradually increases from 3 to 9 digits.

Neuropsychiatric Normative Data and Validity

All neuropsychiatric assessments utilized for the study were well validated in age, socioeconomic, and education matched control populations alleviating the need to recruit normative data from a control population. The Modified Mini Mental Status Examination, Rey Auditory Verbal Learning Test, Digit Span Backward testing, and Trail Making testing have been validated across a variety of ages and educational backgrounds, thereby providing norms on the composite scores.^{58, 59, 60, 61, 62} Furthermore, tests such as the Frontal Assessment Battery have demonstrated good interrater reliability, internal consistency, and discriminant validity.⁶³ The well-defined normative data allow the selected neuropsychiatric assessments to detect clinically relevant cognitive dysfunction correlated to impairment in the tasks of daily living.

138 **Analyses**

139 Raw scores from each cognitive test were standardized to t- or z-scores using
140 existing normative data based on age. Cutoffs for impairment were then created using these
141 data, with scores falling greater than 1.5 standard deviations below normative values.

142
143
144 **3. RESULTS AND DISCUSSION**

145
146 **Demographic Information**

147 Table 1 shows the demographic characteristics of the study population. The
148 median age was 62 with an even distribution of males and females. The majority of
149 participants were Caucasian (82%) and 17.6% were African-American. The family structure
150 and education backgrounds of the study's participants were comparable to CVD patients in a
151 typical urban CICU in the United States.

152
153 **Medical Conditions**

154 Figure 1 shows the primary admission diagnosis of participants, which was
155 predominantly myocardial infarction followed by heart failure exacerbation and arrhythmia. A
156 review of patient comorbidities was used to calculate the Charlson Comorbidity Index, with
157 28.6% having a score of 0; 45.2% having a score of 1-2; 21.5% having a score of 3-4; and
158 4.8% having a score of 5 or greater. The patient sample selected is comparable to national
159 CCU trends.⁶⁴

160
161 **Invasive Procedures Prior to Time 1 Testing**

162 Patients underwent many invasive procedures between Time 1 and Time 2. 60.3%
163 of participants underwent diagnostic cardiac catheterizations, 22.2% underwent
164 percutaneous coronary intervention, and 4.7% underwent an electrophysiology study or
165 pacemaker placement. . To ensure that medications were not influencing cognitive testing,
166 we collected data on the medication administration of all narcotic and sedatives and invasive
167 procedures. Fewer than 3.1% of patients received opioids, benzodiazepines, antihistamines,
168 or anti-cholinergic medications and no participants received medications during the 4 hours
169 prior to cognitive testing. Therefore, medication side effects were unlikely to account for
170 substantial cognitive dysfunction in this population.

171
172 **Cognitive Dysfunction Prevalence**

173 Figure 2 and Supplemental Tables 1 and 2 indicate the prevalence of cognitive
174 dysfunction in the CICU setting according to cognitive test and domains at Time 1. Among
175 the participants at Time 1, 26 % had global impairment 45.5% demonstrated impairment
176 on the short recall and 40.4% demonstrated impairment on the long recall in memory based
177 upon Rey Auditory Verbal Learning Test, 49% were impaired in memory based upon Digit
178 Span testing , 38% were impaired in attention based upon Trail Making Test A, 25%
179 demonstrated impairment in executive function based upon Trail Making Test B and 18%
180 had impairment in executive function based upon Frontal Assessment Battery.

181
182 **Cognitive Dysfunction remains Prevalent prior to Discharge.**

183 Figure 2 reveals the prevalence of cognitive dysfunction in the CICU setting at Time
184 2. Among participants, 33% demonstrated impairment globally based upon the Modified Mini
185 Mental Status Examination , 16.7% demonstrated impairment on the short recall and 43.8%
186 demonstrated impairment on the long recall in memory based upon Rey Auditory Verbal
187 Learning Test, 61% demonstrated impairment in memory based upon Digit Span testing,
188 18% demonstrated impairment in attention based upon Trail Making Test A, 26% of
189 demonstrated impairment in executive function , and 29% demonstrated impairment in
190 executive function based upon Frontal Assessment Battery testing.

191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242

Discussion

In this study, the prevalence of unappreciated cognitive dysfunction for patients hospitalized in the CICU, as evidenced by neurocognitive testing, was substantial. Nearly half of all patients manifested significant impairment in at least one measured domain. The most prevalent dysfunction appeared to occur in global cognition and memory. However, significant impairment in attention and executive function was also detected.

When logistically feasible, testing was repeated prior to discharge to confirm that prior findings of impairment at Time 1 were not an isolated or transient finding. In addition, the timing of the second test battery administration represents an important point when clinical staff often provides patient teaching. Reassessment before discharge indicated a high prevalence of cognitive dysfunction across multiple domains.

Strengths of the study include that the sample was chosen to represent the general CICU population. The distribution of demographics, admission diagnoses, comorbid conditions, medications administered, and invasive procedures performed are consistent with the majority of CICUs nationwide. Furthermore, patients were enrolled on the first morning of hospitalization, which is a critical time in patient care because history is often obtained by care providers at that time, particularly in a teaching environment. Moreover, both the first morning of hospitalization and the brief period of time prior to discharge are key points for patient disease orientation and self-care education. By conducting assessments at two time points during the hospitalization, this study indicates that the detected cognitive dysfunction is neither transient nor unique to a specific time point in hospitalization.

This study was designed to exclude participants with conditions that have been previously correlated to cognitive dysfunction, such as stroke and obstructive sleep apnea. We used care providers' clinical examinations to further eliminate from participating patients with clear cognitive dysfunction. Therefore, the study group represented what most physicians and nurses would consider "normal" cognitive function. This study also reveals the clear weaknesses of conventional clinical assessments currently being used, in which a large portion of cognitive dysfunction often passes unnoticed.

The cognitive function testing instruments used in this study have been validated in prior studies to measure other types of clinically significant dysfunctions. While these neurocognitive testing tools have been used extensively in other scientific fields, this study demonstrated the merits of these instruments in detecting cognitive dysfunction as well. As the public health costs of continuing to ignore cognitive dysfunction become more apparent, so too will the value of administration of these sensitive neurocognitive screening tests.

A notable barrier to implementation of neurocognitive testing in the broader health care system is the time resources required to appropriately perform these assessments. The study was impacted by the difficulty in completing the full battery of testing, which often required an uninterrupted hour. The majority of subjects were able to complete the Modified Mini Mental Status Examination ($n = 51$), but only 32 were able to complete the Trail Making Test Part B testing at Time 1. Patient refusal due to fatigue or anxiety, and interruption of medical procedures, sometimes prohibited the administration of the complete testing battery.

Research staff was not always available to perform Time 2 testing during evenings, weekends, or holidays. This resulted in fewer ($n=24$) patients undergoing testing at both time points. Although longitudinal reassessment of all patients would have been ideal, this was not feasible due to staff availability and a number of patient-related factors.

243 4. CONCLUSION

244
245 Early in the hospital stay and prior to discharge, dysfunction in cognition is predictive of
246 adversely affecting patient care by impairing the accuracy of the history provided and
247 reducing a patient's ability to learn how to care for their illness, whether it is in the form of
248 lifestyle modification, dietary changes, or medication compliance.^{56, 57} Further understanding
249 of cognitive dysfunction in cardiac patients may reveal the underlying etiology of many of the
250 difficulties and gaps in care associated with transitions from hospital to home.

251 Unappreciated cognitive dysfunction is common even in hospitalized CICU patients, who
252 staff physicians and nurses considered to be cognitively intact. In this study's sample group,
253 while cognitive dysfunction was more pronounced in the domains of memory and attention,
254 significant impairment was also noted in global cognition and executive function.

255
256 Cognitive dysfunction likely impedes patients from participating in longitudinal care and
257 comprehension of health education, with significantly increased potential for
258 rehospitalizations. Future studies will address the determinants of cognitive dysfunction and
259 better clarify the time course of the dysfunction, both in acute hospitalization and transition to
260 outpatient care. Such studies will contribute to tailoring the delivery of education to optimize
261 patient understanding, as well as offering a means of improving patient satisfaction. The
262 implications for individual patients, caregivers, and the public health system are promising.

264 CONSENT (WHERE EVER APPLICABLE)

265
266 All authors declare that 'written informed consent was obtained from the patient (or other
267 approved parties) for publication of this case report and accompanying images. A copy of
268 the written consent is available for review by the Editorial office/Chief Editor/Editorial Board
269 members of this journal.

270 271 272 ETHICAL APPROVAL (WHERE EVER APPLICABLE)

273
274 All authors hereby declare that all experiments have been examined and approved by the
275 appropriate ethics committee and have therefore been performed in accordance with the
276 ethical standards laid down in the 1964 Declaration of Helsinki.

277 278 REFERENCES

- 279
280 (1) Lopez AD, Mathers CD, Ezzati M, Jaimson DT, Murray CJL. Global and regional burden
281 of disease and risk factors, 2001: systemic analysis of population health data. *The*
282 *Lancet* 2006;367(9524):1747-57.
- 283 (2) American Heart Association. Heart Disease and Stroke Statistics -- 2009 Update.
284 Dallas, TX; 2009.
- 285 (3) Trogon JG, Finkelstein EA, Nwaise IA, Tangka FK, Orenstein D. The Economic
286 Burden of Chronic Cardiovascular Disease for Major Insurers. *Health Promotion*
287 *Practice* 2007 July 1;8(3):234-42.
- 288 (4) Knopman D, Boland LL, Mosley T et al. Cardiovascular risk factors and cognitive
289 decline in middle-aged adults. *Neurology* 2001 January 9;56(1):42-8.

- 290 (5) Breteler MMB, Bots ML, Ott A, Hofman A. Risk factors for vascular disease and
291 dementia. *Haemostasis* 1998;28:167-73.
- 292 (6) Bennett SJ, Sauve MJ. Cognitive deficits in patients with heart failure: a review of
293 the literature. *Journal of Cardiovascular Nursing* 2003;18:219-42.
- 294 (7) Alves TCTF, Rays J, Fraguas R et al. Localized cerebral blood flow reductions in
295 patients with heart failure: A study using Tc-99m-HMPAO SPECT. *Journal of*
296 *Neuroimaging* 2005 April;15(2):150-6.
- 297 (8) Woo MA, Macey PM, Fonarow GC, Hamilton MA, Harper RM. Regional brain gray
298 matter loss in heart failure. *J Appl Physiol* 2003 August 1;95(2):677-84.
- 299 (9) Zuccala G, Cattell C, Manes-Gravina E, Di Niro MG, Cocchi A, Bernabei R. Left
300 ventricular dysfunction: a clue to cognitive impairment in older patients with heart
301 failure. *J Neurol Neurosurg Psychiatry* 1997 October 1;63(4):509-12.
- 302 (10) Almeida OP, Flicker L. The mind of a failing heart: a systematic review of the
303 association between congestive heart failure and cognitive function. *Internal*
304 *Medicine Journal* 2001;31:290-5.
- 305 (11) Breteler MMB, Claus JJ, Grobbee DE, Hofman A. Cardiovascular disease and
306 distribution of cognitive function in elderly people: the Rotterdam study. *BMJ* 1994
307 June 18;308(6944):1604-8.
- 308 (12) Singh-Manoux A, Sabia S, Lajnef M et al. History of coronary heart disease and
309 cognitive performance in midlife: the Whitehall II study. *Eur Heart J* 2008 September
310 1;29(17):2100-7.
- 311 (13) Zuccala G, Pedone C, Cesari M et al. The effects of cognitive impairment on
312 mortality among hospitalized patients with heart failure. *The American Journal of*
313 *Medicine* 2003 August 1;115(2):97-103.
- 314 (14) Barclay LL, Weiss EM, Mattis S, Bond O, Blass JP. Unrecognized cognitive
315 impairment in cardiac rehabilitation patients. *Journal of the American Geriatrics*
316 *Society* 1988;36:22-8.
- 317 (15) Konstam MA. Improving clinical outcomes with drug treatment in heart failure: what
318 have trials taught? *The American Journal of Cardiology* 2003 March 20;91(6,
319 Supplement 1):9-14.
- 320 (16) Zuccala G, Onder G, Pedone C et al. Cognitive dysfunction as a major determinant
321 of disability in patients with heart failure: results from a multicentre survey. *J Neurol*
322 *Neurosurg Psychiatry* 2001 January 1;70(1):109-12.
- 323 (17) Thom T, Haase N, Rosamond W et al. Heart Disease and Stroke Statistics--2006
324 Update: A Report From the American Heart Association Statistics Committee and
325 Stroke Statistics Subcommittee. *Circulation* 2006 February 14;113(6):e85-151.
- 326 (18) Walsh JT, Charlesworth A, Andrews R, Hawkins M, Cowley AJ. Relation of daily
327 activity levels in patients with chronic heart failure to long-term prognosis. *The*
328 *American Journal of Cardiology* 1997;79:1364-9.

- 329 (19) Gorkin L, Norvell NK, Rosen RC et al. Assessment of quality of life as observed from
330 the baseline data of the Studies of Left Ventricular Dysfunction (SOLVD) trial quality-
331 of-life substudy. *American Journal of Cardiology* 1993;71:1069-73.
- 332 (20) Schmidt R, Fazekas F, Offenbacher H, Dusleag J, Lechner H. Brain magnetic
333 resonance imaging and neuropsychologic evaluation of patients with idiopathic
334 dilated cardiomyopathy. *Stroke* 1991 February 1;22(2):195-9.
- 335 (21) Trojano L, Antonelli RI, Acanfora D et al. Cognitive impairment: a key feature of
336 congestive heart failure in the elderly. *Journal of Neurology* 2003;250(12):1456-63.
- 337 (22) Cline CMJ, Bjorck-Linne AK, Israelsson BYA, Willenheimer RB, Erhardt LR. Non-
338 compliance and knowledge of prescribed medication in elderly patients with heart
339 failure. *Eur J Heart Fail* 1999 June 1;1(2):145-9.
- 340 (23) FREEDMAN NEIL, KOTZER NATA, SCHWAB RICH. Patient Perception of Sleep
341 Quality and Etiology of Sleep Disruption in the Intensive Care Unit. *Am J Respir Crit*
342 *Care Med* 1999 April 1;159(4):1155-62.
- 343 (24) Harrison Y, Horne JA. Sleep deprivation affects speech. *Sleep* 1997
344 October;20(10):871-7.
- 345 (25) Harrison Y, Horne JA. Sleep loss impairs short and novel language tasks having a
346 prefrontal focus. *Journal of Sleep Research* 1998;7(2):95-100.
- 347 (26) Drummond SPA, Brown GG, Gillin JC, Stricker JL, Wong EC, Buxton RB. Altered
348 brain response to verbal learning following sleep deprivation. *Nature* 2000 February
349 10;403(6770):655-7.
- 350 (27) Chapman E, Gavrin J. The contributions of persistent pain. *The Lancet* 1999 June
351 26;353:2233-7.
- 352 (28) Brown SC, Glass JM, Park DC. The relationship of pain and depression to cognitive
353 function in rheumatoid arthritis patients. *Pain* 2002 April;96(3):279-84.
- 354 (29) Benton D, Sargent J. Breakfast, blood glucose and memory. *Biological Psychology*
355 1992 July;33(2-3):207-10.
- 356 (30) Smith AP, Kendrick AM, Maben AL. Effects of breakfast and caffeine on
357 performance and mood in the late morning and after lunch. *Neuropsychobiology*
358 1992;26(4):198-204.
- 359 (31) Conners CK, Blouin AG. Nutritional effects on behavior of children. *Journal of*
360 *Psychiatric Research* 1983;17(2):193-201.
- 361 (32) Agostini JV, Leo-Summers LS, Inouye SK. Cognitive and Other Adverse Effects of
362 Diphenhydramine Use in Hospitalized Older Patients. *Arch Intern Med* 2001
363 September 24;161(17):2091-7.
- 364 (33) Canet J, Raeder J, Rasmussen LS et al. Cognitive dysfunction after minor surgery in
365 the elderly. *Acta Anaesthesiologica Scandinavica* 2003;47(10):1204-10.

- 366 (34) Newman MF, Kirchner JL, Phillips-Bute B et al. Longitudinal Assessment of
367 Neurocognitive Function after Coronary-Artery Bypass Surgery. *N Engl J Med* 2001
368 February 8;344(6):395-402.
- 369 (35) Paul RH, Lawrence J, Williams LM, Richard CC, Cooper N, Gordon E. Preliminary
370 Validity of "Integneuro TM": A New Computerized Battery of Neurocognitive Tests.
371 *International Journal of Neuroscience* 2005;115(11):1549-67.
- 372 (36) Clark CR, Paul RH, Williams LM et al. Standardized assessment of cognitive
373 functioning during development and aging using an automated touchscreen battery.
374 *Archives of Clinical Neuropsychology* 2006 August;21(5):449-67.
- 375 (37) Teng EL, Chui HC. The Modified Mini-Mental State (3MS) Examination. *Journal of*
376 *Clinical Psychiatry* 1987 August;48(8):314-8.
- 377 (38) Rosenberg SJ, Ryan JJ, Prifitera A. Rey auditory-verbal learning test performance of
378 patients with and without memory impairment. *Journal of Clinical Psychology* 2006
379 February 21;40(3):785-7.
- 380 (39) Conn HO. Trailmaking and number-connection tests in the assessment of mental
381 state in portal systemic encephalopathy. *Digestive Diseases and Sciences* 1977
382 June 1;22(6):541-50.
- 383 (40) Slachevsky A, Villalpando JM, Sarazin M, Hahn-Barma V, Pillon B, Dubois B.
384 Frontal Assessment Battery and Differential Diagnosis of Frontotemporal Dementia
385 and Alzheimer Disease. *Archives of Neurology* 2004 July 1;61(7):1104-7.
- 386 (41) Torgesen JK, Houck DG. Processing deficiencies of learning-disabled children who
387 perform poorly on the digit span test. *Journal of Educational Psychology* 1980
388 April;72(2):141-60.
- 389 (42) Reitan RM. Validity of the Trail Making Test as an indicator of organic brain damage.
390 *Perceptual and Motor Skills* 1958;8:271-6.
- 391 (43) Slachevsky A, Villalpando JM, et al. Frontal assessment battery and differential
392 diagnosis of frontotemporal dementia and Alzheimer disease. *Arch Neurol*
393 2004;61:1104-7.
- 394 (44) Callahan LF, DeVellis RF, DeWalt DA. PROMIS (Patient-Reported Outcomes
395 Measurement Information System): Review of the first 3 years of progress and
396 implications for arthritis outcomes. *Arthritis and Rheumatism* 2008;58(9):S1162.
- 397 (45) Garcia SF, Cella D, Clauser SB et al. Standardizing patient-reported outcomes
398 assessment in cancer clinical trials: A patient-reported outcomes measurement
399 information system initiative. *Journal of Clinical Oncology* 2007 November
400 10;25(32):5106-12.
- 401 (46) Smid HG, Trumper BG, Pottag G et al. Differentiation of hypoglycaemia induced
402 cognitive impairments. An electrophysiological approach. *Brain* 1997 June
403 1;120(6):1041-56.

- 404 (47) Alaimo K, Olson CM, Frongillo Jr EA. Food Insufficiency and American School-Aged
405 Children's Cognitive, Academic, and Psychosocial Development. *Pediatrics* 2001
406 July 1;108(1):44-53.
- 407 (48) Simeon DT, Grantham-McGregor S. Effects of missing breakfast on the cognitive
408 functions of school children of differing nutritional status. *Am J Clin Nutr* 1989 April
409 1;49(4):646-53.
- 410 (49) Green MW, Elliman NA, Rogers PJ. Lack of effect of short-term fasting on cognitive
411 function. *Journal of Psychiatric Research* 1995;29(3):245-53.
- 412 (50) Omichinski L. *You Count, Calories Don't*. Winnipeg: Hyperion Press; 1992.
- 413 (51) Lorr M, McNair DM, Droppleman LF. *Manual: Profile of Mood States*. San Diego,
414 CA: Educational and Industrial Testing Service; 1971.
- 415 (52) Curran SL, Andrykowski MA, Studts JL. Short form of the Profile of Mood States
416 (POMS-SF): psychometric information. *Psychological Assessment* 1995;7(1):80-3.
- 417 (53) Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying
418 prognostic comorbidity in longitudinal studies: development and validation. *Journal*
419 *of Chronic Disease* 1987;40:373-83.
- 420 (54) Gortner SR, Jaeger AA, Harr J, Miller T. Elders' expected and realized benefits from
421 cardiac surgery. *Cardio-vascular Nursing* 21994;30:9-14.
- 422 (55) Safeer RS, Keenan J. Health Literacy: The gap between physicians and patients.
423 *American Family Physician* 2005;72:463-8.
424
- 425 (56) Kakos LS, Szabo AJ, Gunstad J, Stanek KM, Waechter D, Hughes J, Luyster F,
426 Josephson R, Rosneck J. Reduced executive functioning is associated with poorer
427 outcome in cardiac rehabilitation. *Preventive Cardiology* 2010 Summer;13(3):100-
428
- 429 (57) Stanek KM, Gunstad J, Spitznagel MB, Waechter D, Hughes JW, Luyster F,
430 Josephson
431 R, Rosneck J. Improvements in cognitive function following cardiac rehabilitation for
432 older adults with cardiovascular disease. *International Journal of Neuroscience*
433 2011 Feb;121(2):86-93. Epub 2010 Nov 10.
434
- 435 (58) Lancu I, Olmer A. The minimental state examination an up-to-date review.
436 *Harefuah*. 2006 Sep;145(9):687-90, 701.
- 437 (59) Tschanz J, Welsh-Bohmer K, Plassman B, Norton M, Wyse B, Breitner J; Cache
438 County Study Group. An adaptation of the modified mini-mental state examination:
439 analysis of demographic influences and normative data: the cache county study.
440 *Neuropsychiatry Neuropsychol Behav Neurol*. 2002 Mar;15(1):28-38.
- 441 (60) Vakil E, Greenstein Y, Blachstein H. Normative data for composite scores for
442 children and adults derived from the Rey Auditory Verbal Learning Test. *Clin*
443 *Neuropsychol*. 2010 May;24(4):662-77. Epub 2010 Feb 11.
- 444 (61) Miller G. The Magical Number Seven, Plus or Minus Two: Some Limits on Our
445 Capacity for Processing Information. *Psychological Review* 63 (2): 81–97. PMID
446 13310704.
- 447 (62) Mitrushina M, Boone K, Razani J, D'Elia L. *Handbook of Normative Data for*
448 *Neuropsychological Assessment*. Feb 10, 2005.

449 (63) Dubois B, Slachevsky A, Litvan I, Pillon B. The FAB: a Frontal Assessment Battery
 450 at bedside. *Neurology*. 2000;55:1621-1626.
 451 (64) Groeger J, Guntupalli K, Strosberg M, Halpern N, Raphaely R, Cerra F, Kaye W.
 452 Descriptive analysis of critical care units in the United States: patient characteristics
 453 and intensive care unit utilization. *Critical Care Med*. 1993 Feb; 21(2):279-91.
 454
 455
 456 **APPENDIX**

Table 1. Patient Demographics (N=51)

Variable	Total Sample N(%)	
Age	Mean= 61.3 STD +/- 13.3	
Gender	Women/ Men	23 (48.4)/ 28 (51.9)
Current Living Arrangements	Live Alone	12 (25.5)
	With Spouse, Family, Friend	35 (74.5)
Marital Status	Single	7 (14)
	Married	27 (54)
	Widowed, Separated, Divorced	16 (32)
Race	Black	10 (19)
	Caucasian	41 (81)
Highest Education Level	Less than High School	6 (14.6)
	High School	15 (36.6)
	Advanced Degree or Training	20 (48.7)
Employment Status	Retired	21 (45.7)
	Work	18 (39.1)
	Unemployed	7 (15.2)

Figure 1. Primary admission diagnosis

Unappreciated Cognitive Dysfunction is Common 3/20/2015





Figure 2. Prevalence of Cognitive dysfunction



**Global
Cognition**

Memory

Attention

**Executive
Function**

Supplemental Table 1. Cognitive testing on admission

Global

3MS

Impaired (< 90) 26% N=51

Memory

Ray Auditory Verbal Learning Test

Short recall-scaled 6.7 + 3.1 N= 37

Impaired (< 7) 46%

Long recall-scaled 6.8 + 3.1 N= 37

Impaired (< 7) 40%

Digit Span Backward 3.7 + 1.3 N= 43

Impaired (< 4) 49%

Attention

Unappreciated Cognitive Dysfunction is Common 3/20/2015

Trails A	8.7 + 2.8	N= 34
Impaired (< 7)	38%	
<i>Executive Function</i>		
Trails-B	8.8 + 3.4	N= 32
Impaired (< 7)	25%	
Frontal Assessment Battery	51.5+8.9	N= 34
Impaired (< 40)	18%	

Supplemental Table 2. Cognitive testing prior to discharge

<i>Global</i>		
3MS	90.5+7.4	N= 24
Impaired (< 90)	33%	
<i>Memory</i>		
Ray Auditory Verbal Learning Test		
Short recall-scaled	8.2 + 3.3	N= 16
Impaired (< 7)	17%	
Long recall-scaled	7.4 + 3.9	N= 16
Impaired (< 7)	44%	
Digit Span Backward	3.3 + 1.0	N= 18
Impaired (< 4)	61%	
<i>Attention</i>		
Trails A	8.1 + 3.7	N= 19
Impaired (< 7)	18%	
<i>Executive Function</i>		
Trails-B	7.9+ 3.0	N= 19
Impaired (< 7)	26%	
Frontal Assessment Battery	48.4+11.8	N= 21
Impaired (< 40)	29%	