SCIENCEDOMAIN international

www.sciencedomain.org



SDI FINAL EVALUATION FORM 1.1

PART 1:

Journal Name:	British Journal of Pharmaceutical Research
Manuscript Number:	2013 BJPR 3295
Title of the Manuscript:	Free radical scavenging activities of Nyctanthes arbor-tristis. L on adjuvant induced arthritic rats

PART 2:

Regarding explanation to Fig. 2 (the levels of lipid peroxides and activities of antioxidant enzymes in the tissue homogenates were increased in arthritic rats and significantly decreased by ANT treatments), authors describe that over production of oxygen free radicals causes lipid peroxidation and the increase in catalase/GPs activities signifies over production of hydrogen peroxide. Thus, free radical scavenging action of ANT mainly concerns to anti-arthritic activity. In this story, it is reasonable. Also, followings are needed to recheck. 1) In the Methods, tissue homogenate was prepared in the presence of 5% TCA followed by centrifugation(reference No.10). I wonder enzyme activities such as SOD, catalase or GPX, were tost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein 3) Title of Table 3 should be "Activities of membrane marker enzymes in the serum (not tissues) and also sentence on the discussion "A marked increase in were observed in the serum (not joint tissues) of arthritic rats". These enzymes are marker for liver damage in serum. 4) The pictures of histological examinations need the title as Figure 4. Histopathological slides are titled as Figure 4	PART 2:		
enzymes in the tissue homogenates were increased in arthritic rats and significantly decreased by ANT treatments), authors describe that over production of oxygen free radicals causes lipid peroxidation and the increase in catalase/GPx activities signifies over production of hydrogen peroxide. Thus, free radical scavenging action of ANT mainly concerns to anti-arthritic activity. In this story, it is reasonable. Also, followings are needed to recheck. 1) In the Methods, tissue homogenate was prepared in the presence of 5% TCA followed by centrifugation(reference No.10). I wonder enzyme activities such as SOD, catalase or GpX, were lost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein The unit of SOD has been changed in the Figure 2 The title has been changed.		Authors' response to final evaluator's comments	
decreased by ANT treatments), authors describe that over production of oxygen free radicals causes lipid peroxidation and the increase in catalase/GPx activities signifies over production of hydrogen peroxide. Thus, free radical scavenging action of ANT mainly concerns to anti-arthritic activity. In this story, it is reasonable. Also, followings are needed to recheck. 1) In the Methods, tissue homogenate was prepared in the presence of 5% TCA followed by centrifugation(reference No.10). I wonder enzyme activities such as SOD, catalase or GpX, were lost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein The unit of SOD has been changed in the Figure 2 The title has been changed.			
radicals causes lipid peroxidation and the increase in catalase/GPx activities signifies over production of hydrogen peroxide. Thus, free radical scavenging action of ANT mainly concerns to anti-arthritic activity. In this story, it is reasonable. Also, followings are needed to recheck. 1) In the Methods, tissue homogenate was prepared in the presence of 5% TCA followed by centrifugation(reference No.10). I wonder enzyme activities such as SOD, catalase or GpX, were lost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig. 2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein The unit of SOD has been changed in the Figure 2 3) Title of Table 3 should be "Activities of membrane marker enzymes in the serum (not tissues) and also sentence on the discussion "A marked increase in "" were observed in the serum (not joint tissues) of arthritic rats". These enzymes are marker for liver damage in serum.			
production of hydrogen peroxide. Thus, free radical scavenging action of ANT mainly concerns to anti-arthritic activity. In this story, it is reasonable. Also, followings are needed to recheck. 1) In the Methods, tissue homogenate was prepared in the presence of 5% TCA followed by centrifugation(reference No.10). I wonder enzyme activities such as SOD, catalase or GpX, were lost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein The unit of SOD has been changed in the Figure 2 The title has been changed. The title has been changed.	decreased by ANT treatments), authors describe that over production of oxygen free		
Also, followings are needed to recheck. 1) In the Methods, tissue homogenate was prepared in the presence of 5% TCA followed by centrifugation(reference No.10). I wonder enzyme activities such as SOD, catalase or GpX, were lost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein The unit of SOD has been changed in the Figure 2 The title has been changed. The title has been changed.	radicals causes lipid peroxidation and the increase in catalase/GPx activities signifies over		
Also, followings are needed to recheck. 1) In the Methods, tissue homogenate was prepared in the presence of 5% TCA followed by centrifugation(reference No.10). I wonder enzyme activities such as SOD, catalase or GpX, were lost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein 3) Title of Table 3 should be "Activities of membrane marker enzymes in the serum (not tissues) and also sentence on the discussion "A marked increase in were observed in the serum (not joint tissues) of arthritic rats". These enzymes are marker for liver damage in serum.	production of hydrogen peroxide. Thus, free radical scavenging action of ANT mainly		
1)In the Methods, tissue homogenate was prepared in the presence of 5% TCA followed by centrifugation(reference No.10). I wonder enzyme activities such as SOD, catalase or GpX, were lost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein The typographical error has been corrected in the manuscript The typographical error has been corrected in the typ	concerns to anti-arthritic activity. In this story, it is reasonable.		
1)In the Methods, tissue homogenate was prepared in the presence of 5% TCA followed by centrifugation(reference No.10). I wonder enzyme activities such as SOD, catalase or GpX, were lost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein The typographical error has been corrected in the manuscript The typographical error has been corrected in the manuscript The unit of SOD has been changed in the Figure 2 The unit of SOD has been changed in the Figure 2 The title has been changed.			
centrifugation(reference No.10). I wonder enzyme activities such as SOD, catalase or GpX, were lost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein The unit of SOD has been changed in the Figure 2 The title has been changed. The title has been changed.	Also, followings are needed to recheck.		
centrifugation(reference No.10). I wonder enzyme activities such as SOD, catalase or GpX, were lost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein The unit of SOD has been changed in the Figure 2 The title has been changed. The title has been changed.			
GpX, were lost (inactivated) by 5% TCA. In the No.10, TCA was used in only GSH assay. Correct homogenizing conditions. #Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein The unit of SOD has been changed in the Figure 2 The title has been changed. The title has been changed.		The typographical error has been corrected in the manuscript	
#Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein 3) Title of Table 3 should be "Activities of membrane marker enzymes in the serum (not tissues) and also sentence on the discussion "A marked increase in were observed in the serum (not joint tissues) of arthritic rats". These enzymes are marker for liver damage in serum.			
#Fig.2; Concerning the unit of SOD activity, it seems to be microgram (or nanogram) of SOD for 50% inhibition of epinephrine autooxidation/mg protein The unit of SOD has been changed in the Figure 2 The unit of SOD has been changed in the Figure 2 The title has been changed. The title has been changed.			
SOD for 50% inhibition of epinephrine autooxidation/mg protein 3) Title of Table 3 should be "Activities of membrane marker enzymes in the serum (not tissues) and also sentence on the discussion "A marked increase in were observed in the serum (not joint tissues) of arthritic rats". These enzymes are marker for liver damage in serum. The unit of SOD has been changed in the Figure 2 The title has been changed.	Correct homogenizing conditions.		
SOD for 50% inhibition of epinephrine autooxidation/mg protein 3) Title of Table 3 should be "Activities of membrane marker enzymes in the serum (not tissues) and also sentence on the discussion "A marked increase in were observed in the serum (not joint tissues) of arthritic rats". These enzymes are marker for liver damage in serum. The unit of SOD has been changed in the Figure 2 The title has been changed.			
3) Title of Table 3 should be "Activities of membrane marker enzymes in the serum (not tissues) and also sentence on the discussion "A marked increase in were observed in the serum (not joint tissues) of arthritic rats". These enzymes are marker for liver damage in serum.		The unit of COD has been shound in the Figure 2	
tissues) and also sentence on the discussion "A marked increase in were observed in the serum (not joint tissues) of arthritic rats". These enzymes are marker for liver damage in serum.	SOD for 50% inhibition of epinephrine autooxidation/mg protein	The unit of SOD has been changed in the Figure 2	
tissues) and also sentence on the discussion "A marked increase in were observed in the serum (not joint tissues) of arthritic rats". These enzymes are marker for liver damage in serum.			
in the serum (not joint tissues) of arthritic rats". These enzymes are marker for liver damage in serum.	,	The title has been changed.	
damage in serum.	tissues) and also sentence on the discussion "A marked increase in were observed	The vive has seen enanged.	
	in the serum (not joint tissues) of arthritic rats". These enzymes are marker for liver		
4) The pictures of histological examinations need the title as Figure 4. Histopathological slides are titled as Figure 4	damage in serum.		
4) The pictures of histological examinations need the title as Figure 4. Histopathological slides are titled as Figure 4.			
	4) The pictures of histological examinations need the title as Figure 4.	Histopathological slides are titled as Figure 4	

Created by: EA Checked by: ME Approved by: CEO Version: 1.5 (4th August, 2012)