

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20

Research paper

Hepatoprotective effect of *Drynaria quercifolia* fronds hydroalcoholic extract and isolated constituent against CCl₄-induced hepatocellular damage

Pradeep Kamboj^{1*}, Ajudhia Nath Kalia²

¹Assistant Professor, Department of Pharmacognosy,
JCDM College of Pharmacy,

Barnala Road, Sirsa 125055 (Haryana), India

²Professor & Head, Department of Pharmacognosy,
ISF College of Pharmacy,

GT-Road, Moga 142001(Punjab), India

ABSTRACT

Aims: The present study was conducted to evaluate the hepatoprotective effect of hydroalcoholic extract of *Drynaria quercifolia* fronds (Dq), its fractions and isolated compound (Dq-4) from ethyl acetate (EA) fraction.

Place and Duration of Study: Department of Pharmacognosy and Department of Herbal Drug Research, ISF College of Pharmacy, Moga, between June 2010 and May 2012.

Methodology: The toxicant CCl₄ (1ml/kg) was administered on 4th and 5th day to induce hepatotoxicity in Wistar rats (*in-vivo*) and the *in-vitro* hepatoprotection was evaluated against CCl₄ (1%) induced toxicity in HepG2 cellines.

Results: The pre-treatment of rats with Dq extract, EA fraction and Dq-4 for 7 days produced a significant dose dependent hepatoprotective action by decreased levels of hepatic enzymes, total bilirubin and TBARS and increased levels of total proteins, albumin, and reduced glutathione. The histological examination provided the supportive evidences. Additionally, Dq extract, EA fraction and Dq-4 significantly decreased the CCl₄-induced *in-vitro* toxicity in HepG2 cellines evident by MTT reduction assay and trypan blue method.

Conclusion: The study scientifically validated the traditional use of *D. quercifolia* for liver disorders and strongly demonstrates antioxidative effect on hepatocytes in restoring their normal architecture and functional ability.

21
22
23
24
25
26
27
28
29
30
31
32

Keywords: Hepatoprotective, HepG2, *Drynaria quercifolia*, Naringin

1. INTRODUCTION

The liver as a vital organ in the body is primarily responsible for the metabolism of endogenous and exogenous agents [1]. The pathogenesis of liver injury is initiated by the participation of toxic agents or by their bio-activation to chemically reactive metabolites [2, 3]. These metabolites can be electrophilic chemicals or free radicals, that either elicits an immune response or directly affects the biochemistry of the cells by interacting with cellular macromolecules *viz.* proteins, lipids and nucleic acids leading to protein dysfunction, lipid peroxidation, DNA damage, oxidative stress and depletion of natural antioxidants [4, 5]. Hepatocellular damage is known to be associated with impaired hepatic drug metabolizing

* Tel.: +91 7357000002.

E-mail address: dr.pradeepkamboj@gmail.com.

33 capacity and impaired activity of hepatic enzymes [6, 7]. In absence of reliable conventional
34 and synthetic drugs for alleviation of hepatic diseases, traditional medicines are
35 recommended for the treatment of liver diseases in India [8, 9]. Therefore, many folk
36 remedies from plant origin are scientifically evaluated for their possible hepatoprotective
37 potential against experimental induced hepatotoxicity.

38 *Drynaria quercifolia* J. Smith (Polypodiaceae) is locally known as Attukalkizhangu and Gurar
39 [10, 11]. Traditionally, the fronds of plant are reported to be used by tribal communities of
40 Tamil Nadu and Kerala in treatment of diverse ailments including typhoid fever [12], chronic
41 jaundice, anti-inflammatory agent [13], as a poultice and antifertility agent [14, 15], and
42 antipyretic agent [16]. The whole plant is used to treat chest and skin diseases, and is also
43 anthelmintic, expectorant and tonic [17]. Various phytoconstituents like 3,4-dihydroxybenzoic
44 acid friedelin, epifriedelinol, β -amyrin, β -sitosterol and β -sitosterol 3- β -D-glucopyranoside
45 has been isolated from the plant [16]. Although the plant is widely used for remission of
46 several ailments related to liver disorders, there are no systematic scientific reports in the
47 modern literature regarding the usefulness of the plant and its phytoconstituents as a
48 hepatoprotective agent. Hence, to scientifically validate this ethnopharmacological
49 relevance, hepatoprotective potential of *Drynaria quercifolia* fronds was studied in CCl₄-
50 intoxicated both *in-vivo* and *in-vitro* experimental models of hepatocellular damage.

51

52 2. MATERIAL AND METHODS

53

54 2.1. Plant material

55 The fronds of *Drynaria quercifolia* were collected from the forest of Mudumalai National Park,
56 district Udhagamandalam, Tamil Nadu (India) in the month of December 2009. Botanical
57 identification and authentication was done by Dr. H.B. Singh, Scientist F & Head, Raw
58 Material Herbarium & Museum, National Institute of Science Communication and Information
59 Resources (NISCAIR), New Delhi, India under references and authoritative voucher
60 specimen number: NISCAIR/RHMD/Consult/-2010-11/1646/244.

61 2.2. Extraction and fractionation procedures

62 The fronds were cleaned and air dried for a week and pulverized in electric grinder. The
63 dried and powdered fronds (1200 g) were extracted to exhaustion by triple maceration with
64 50% hydroalcohol (2000 ml \times 3) at ambient temperature with constant stirring. The combined
65 filtrate was concentrated under reduced pressure below 40^oC to afford hydroalcoholic extract
66 of *Drynaria quercifolia* fronds (Dq extract: 9.76% w/w). Dq extract was further suspended in
67 distilled water and fractionated through successive extractions with chloroform (1000 ml \times 6),
68 ethyl acetate (1000 ml \times 9) and *n*-butanol (1000 ml \times 12). Each fraction was concentrated to
69 dryness under reduced pressure to give CHCl₃ (19.86% w/w), EA (27.13% w/w) and *n*-
70 BuOH (12.46% w/w) fractions, respectively. The extract and fractions were preserved under
71 refrigeration till further use.

72 2.3. Phytochemical screening

73 The extract and fractions obtained were screened phytochemically for the presence of
74 alkaloids, steroids, terpenoids, glycosides, flavonoids, saponins, proteins, tannins and
75 phenolic compounds, as previously described by [18, 19].

76 2.4. Chemicals and reagents

77 Carbon tetrachloride (CCl₄) and silymarin were purchased from S.D. Fine-Chem. Ltd.,
78 Mumbai and Micro Labs Ltd., Baddi, respectively. All other chemicals and solvents used
79 were of analytical grade and obtained from Sigma Chemicals Co., USA and Merck India Ltd.,
80 Mumbai. Biochemical enzymatic kits were procured from ERBA, Diagnostics Mannheim
81 GmbH, Germany.

* Tel.: +91 7357000002.

E-mail address: dr.pradeepkamboj@gmail.com.

82 **2.5. Experimental animals**

83 Wistar albino rats of either sex weighing between 180-220g were employed in the present
84 study *vide* approval no ISF/CPCSEA/IAEC/2010/39. The experiments were conducted
85 according to the ethical norms approved by Institutional Animal Ethics Committee (IAEC)
86 guide lines for animal care and were adhered to as recommended by CPCSEA guidelines
87 for the use and care of experimental animals. Animals were housed in environmentally
88 controlled (25 ± 2 °C, 12 h light & dark cycle) small cages, with free access to standard
89 laboratory chow diet and water *ad-libitum*.

90 **2.6. Acute toxicity study**

91 Acute oral toxicity studies were performed as per revised OECD guideline No. 423 in the
92 albino mice [20]. The animals were fasted overnight with water *ad-libitum* and administered
93 with single dose of 2000mg/kg test drug. Animals were observed individually at
94 predetermined time intervals during the first 24 h, with special attention given during the first
95 4 h followed by daily observation for a total of 14 days. The animals were observed for toxic
96 symptoms such as behavioral changes, locomotion and mortality [21].

97 **2.7. In-vivo experimental design**

98 The rats were divided into eight groups comprising of six animals each (n=6). The test and
99 standard drug silymarin were suspended in 0.5% w/v carboxymethyl cellulose (CMC) for oral
100 administration. The toxicant 50% CCl₄ in olive oil (1 ml/kg, s.c.) was given on 4th and 5th day,
101 2 hrs after the test and standard drug administration [22]. The doses of the fractions (CHCl₃,
102 EA and *n*-BuOH) were calculated according to their percentage yields. The treatment
103 protocol is summarized and given below.

104 **Group 1:** Normal control; rats received 0.5% CMC for 7 days

105 **Group 2:** CCl₄ control; rats administered with 0.5% CMC for 7 days and received
106 toxicant CCl₄ on 4th and 5th day

107 **Group 3:** Silymarin; rats treated with silymarin suspension 50 mg/kg for 7 days and
108 received toxicant CCl₄ on 4th and 5th day

109 **Group 4 & 5:** Dq extract; rats treated with Dq extract 200 & 400 mg/kg, respectively
110 for 7 days and received toxicant CCl₄ on 4th and 5th day

111 **Group 6:** CHCl₃ fraction; rats treated with CHCl₃ fraction 72.40 mg/kg for 7 days and
112 received toxicant CCl₄ on 4th and 5th day

113 **Group 7:** EA fraction; rats treated with EA fraction 74.55 mg/kg for 7 days and
114 received toxicant CCl₄ on 4th and 5th day

115 **Group 8:** *n*-BuOH fraction; rats treated with *n*-BuOH fraction 45.40 mg/kg for 7 days
116 and received toxicant CCl₄ on 4th and 5th day

117 **2.8. Analysis of hepatic injury**

118 The blood samples were withdrawn from the orbital sinus on 7th day to obtain haemolysis
119 free serum for biochemical estimations. The serum ALT- alanine transaminase, AST-
120 aspartate transaminase [23], ALP- alkaline phosphatase [24], TB- total bilirubin [25], TP-
121 total proteins [26] and ALB- albumin [27] were estimated using commercial enzymatic
122 biochemical diagnostic kits.

123 **2.9. Analysis of in-vivo GSH level and oxidative stress**

124 All the animals were sacrificed by an overdose of ketamine and xylazine mixture and liver
125 was quickly excised, free from any adhering tissues, washed and perfused with chilled
126 normal saline and blotted dry. Perfused liver samples were minced and homogenized in
127 chilled 10mM Tris-HCl buffer (pH 7.4) to obtain 10% whole liver homogenate for the
128 estimation of GSH [28]. The malondialdehyde (MDA) content, a measure of lipid

129 peroxidation, was assayed in the form of thiobarbituric acid reacting substances (TBARS)
130 [29].

131 **2.10. Histopathological studies**

132 The livers were immediately removed and the tissues were fixed in 10% formalin,
133 dehydrated in ethanol (50–100%), cleared in xylene and embedded in paraffin wax. These
134 were then cut into 4–5µm thick sections in rotary microtome and stained with haematoxylin-
135 eosin for photomicroscopic assessment [30, 31].

136 **2.11. Isolation of compounds**

137 The EA fraction showed significant *in-vivo* hepatoprotective activity and 5.02g of fraction was
138 charged in column. The column was initially eluted using chloroform and ethyl acetate of
139 increasing polarity. 220 fractions, each of 45-50 ml were collected and fractions with similar
140 TLC profile were pooled. The pooled fractions A1(6-24), A2(40-55) and A3(69-86) resulted in
141 the isolation of compounds Dq-1 (15.24mg), Dq-2 (13.87mg) and Dq-3 (14.43mg),
142 respectively (data not shown). The pooled fraction (198-220), obtained through elution with
143 100% ethyl acetate, when concentrated and left overnight in deep freezer resulted in a pale
144 yellow coloured compound, which on repeated crystallization yielded pure compound Dq-4
145 (17.91mg).

146 **2.12. Characterization of compounds**

147 The identification of compounds was done by physical, chemical and spectral analysis. The
148 melting points were determined in open-glass capillaries on Stuart SMP10 melting point
149 apparatus. The IR spectra (ν , cm^{-1}) were obtained with a Nicolet 380 FTIR spectrometer
150 (Thermo Scientific) in KBr pellets. $^1\text{H-NMR}$ spectra (δ , ppm) were recorded in DMSO- d_6
151 solutions on a Varian-Mercury 400 MHz spectrometer using tetramethylsilane (TMS) as the
152 internal reference. Mass spectra were recorded on a Shimadzu GCMS-QP 1000 EX.

153 **2.13. *In-vitro* hepatoprotective activity**

154 The *in-vitro* hepatoprotective activity was performed on human HepG2 cellines, obtained
155 from the National Centre for Cell Science (NCCS) Pune, India. The cells were cultured in 96-
156 well plates at density of 1.0×10^5 cells/well over night in DMEM containing 10% FBS
157 maintained at 5% CO_2 at 37 °C [32]. After 24 hours, when partial monolayer was formed, the
158 supernatant was flicked off and the monolayer was washed once. The hepatocytes were
159 exposed to fresh medium containing CCl_4 (1%) along with various concentrations of Dq
160 extract, fractions (CHCl_3 , EA & *n*-BuOH) and isolated compound Dq4. 60 min after the CCl_4
161 intoxication, cytotoxicity was assessed by estimating the percentage viability of HepG2 cells
162 by MTT reduction assay [33]. In addition, morphological changes in HepG2 cells and loss of
163 membrane integrity during the later stages of cell death were determined by trypan blue dye
164 membrane integrity assay [34].

165 **2.14. Statistical analysis**

166 The data of *in-vivo* and *in-vitro* studies were expressed as mean \pm SD and mean \pm SEM of
167 triplicate experiments, respectively. The data was analysed by one-way ANOVA followed by
168 Tukey's multiple comparison analysis as *post-hoc* test using GraphPad Prism 4 (GraphPad
169 Software Inc., CA, USA). The $p < 0.05$ was considered to be statistically significant.

170
171
172
173

174 **3. RESULTS**

175

176 **3.1. Phytochemical study**

177 Preliminary phytochemical screening revealed presence of carbohydrates, proteins,
178 saponins, steroids, triterpenes, phenolic compounds and flavonoids in Dq extract. The EA
179 fraction showed the presence of terpenoids, phenolic compounds and flavonoids.

180 **3.2. Acute toxicity study**

181 The Dq extract and fractions (CHCl₃, EA & *n*-BuOH) did not show any sign and symptoms of
182 behavioral changes, toxicity and mortality up to 2000 mg/kg; b.w.

183 **3.3. Effect of Dq extract and fractions (CHCl₃, EA & *n*-BuOH) on serum
184 biochemical parameters**

185 Compared to the vehicle (normal control) group, CCl₄ at sub lethal dose caused increases in
186 serum AST (280.80%), ALT (300.57%), ALP (70.61%) and TB (151.72%) levels. CCl₄ also
187 caused marked reductions in serum TP (51.32%) and ALB (53.07%) levels. The pre-
188 treatment of groups with Dq extract and EA fraction for 7 days reversed the toxicity affect
189 and exhibited hepatocellular protection as manifested by the reduction in serum ALT, AST,
190 ALP, TB and decreased levels of serum TP and ALB. The EA fraction exhibited the highest
191 percent of protection among all the tested samples. The EA fraction (74.55 mg/kg) markedly
192 reduced the levels of serum AST (40.88%), ALT (41.49%), ALP (19.72%) and TB (28.76%)
193 along with increased levels of serum TP (60.38%) and ALB (28.57%) as compared to CCl₄
194 control group. However, CHCl₃ (72.40 mg/kg) and *n*-BuOH (45.40 mg/kg) fractions did not
195 show any protection against CCl₄-induced hepatocellular injury (Tables 1).

196 **3.4. Effect of Dq extract and fractions (CHCl₃, EA & *n*-BuOH) on tissue GSH
197 and TBARS levels**

198 As shown in Table 1, the GSH level was decreased and TBARS contents were increased in
199 the liver homogenate in CCl₄ control group by 51.34% and 163.67%, respectively as
200 compared to normal control group. The pre-treatment of rats for 7 days with Dq extract and
201 EA fraction markedly reversed these toxic effects and restored the altered levels of GSH and
202 TBARS. The Dq extract (400 mg/kg) and EA fraction (74.55 mg/kg) increased the level of
203 reduced GSH by 22.09% and 22.46% and inhibited the levels of TBARS by 26.23% and
204 22.38%, respectively as compared to CCl₄ control group.

205 **3.5. Histopathological studies**

206 The histological examination of CCl₄-toxicated liver section showed various degree of
207 architecture damage with de-arrangement of normal hepatic cells, centrilobular necrosis and
208 fibrosis, ventral vein enlargement, sinusoidal dilation, fatty vacuolization, ballooning
209 degeneration and broad infiltration of lymphocytes. The Dq extract and EA pre-treated rats
210 caused degenerative changes and retained the structural integrity of hepatic cells, which
211 closely resembles to the liver histology of the normal control group with less vacuole
212 formation, absence of necrosis and overall less visible parenchymal injury (Figure 1).

213 **3.6. Characterization of compound Dq-4**

214 The isolated compound Dq-4 was characterized as 7-[[2-O-(6-Deoxy- α -L-monopyranoxyl)- β -
215 D-glucopyranoxyl] oxy]-2,3-dehydro-5,7-dihydroxy-2-(4 hydroxyphenyl)-4H-1-benzopyran-4-
216 one (flavanone glycoside; naringin) Figure 2. The yield was 17.91mg; m.p. 169-172 °C; IR
217 (KBr; cm⁻¹): 3468 (OH stretch), 2935 (CH stretch), 2865, 2843 (CH₂ stretch), 1685 (C=O), 1606
218 (C=C Ar), 1235 (C-O). ¹H-NMR (400 MHz, DMSO-*d*₆, δ ppm): 11.99 (s, 1H, OH, C-5,
219 chromone), 9.38 (s, 1H, OH, chromone, C-7), 7.93 (s, 1H, OH, C-4', phenyl), 7.29 (d, 2H, C-

* Tel.: +91 7357000002.

E-mail address: dr.pradeepkamboj@gmail.com.

220 2'-C-6', phenyl, $J = 8.4$ Hz), 6.38 (*d*, 2H, C-3'-C-5', phenyl, $J = 8.4$ Hz), 6.11 (*s*, 1H, C-8,
221 chromone), 5.39 (*m*, 1H, C-2, chromone), 5.19 (*m*, 3H, OH, C-3'',C-4'',C-5'', pyranoside),
222 4.99 (*t*, 1H, C-3''), 4.23 (*m*, 1H, C-2''), 3.83 (*m*, 2H, CH₂-OH, pyranoside), 3.63 (*m*, 1H, C-
223

* Tel.: +91 7357000002.

E-mail address: dr.pradeepkamboj@gmail.com.

224 **Table 1. Effect of Dq extract and fractions (CHCl₃, EA & n-BuOH) on serum biochemical parameters, tissue GSH and TBARS**
 225 **levels in CCl₄-induced hepatic injury in rats**

Groups	Serum parameters						Tissue parameters	
	AST (U/L)	ALT (U/L)	ALP (U/L)	TB (mg/dl)	TP (g/dl)	ALB (g/dl)	TBARS (nM/mg protein)	GSH (μ M/mg protein)
Normal control	47.78 \pm 7.06	41.66 \pm 9.12	160.31 \pm 14.94	0.29 \pm 0.06	6.43 \pm 0.53	3.58 \pm 0.41	21.31 \pm 5.41	108.96 \pm 14.92
CCl ₄ control (1 ml/kg; s.c.)	177.17 \pm 14.92 (270.80%) ^a \uparrow	166.88 \pm 13.92 (300.57%) ^a \uparrow	273.51 \pm 21.10 (70.61%) ^a \uparrow	0.98 \pm 0.14 (237.93%) ^a \uparrow	3.13 \pm 0.51 (51.32%) ^a \downarrow	1.68 \pm 0.25 (53.07%) ^a \downarrow	56.19 \pm 8.63 (163.67%) ^a \uparrow	53.01 \pm 9.41 (51.34%) ^a \downarrow
Silymarin (50 mg/kg; p.o.)	65.51 \pm 9.83 (63.02%) ^b \downarrow	54.69 \pm 11.63 (67.22%) ^b \downarrow	190.74 \pm 18.35 (30.26%) ^b \downarrow	0.39 \pm 0.09 (60.20%) ^b \downarrow	6.18 \pm 0.55 (97.44%) ^b \uparrow	3.14 \pm 0.27 (86.90%) ^b \uparrow	31.59 \pm 6.19 (43.78%) ^b \downarrow	84.83 \pm 10.29 (60.02%) ^b \uparrow
Dq extract (200 mg/kg; p.o.)	127.98 \pm 13.78 (27.76%) ^b \downarrow	123.43 \pm 13.66 (26.03%) ^b \downarrow	245.23 \pm 16.73 (10.33%) \downarrow	0.57 \pm 0.08 (41.83%) ^b \downarrow	4.75 \pm 0.41 (51.75%) ^b \uparrow	2.17 \pm 0.30 (29.16%) \uparrow	47.68 \pm 8.42 (15.14%) \downarrow	67.14 \pm 8.26 (26.65%) \uparrow
Dq extract (400 mg/kg; p.o.)	96.59 \pm 11.73 (45.48%) ^b \downarrow	90.24 \pm 13.28 (45.92%) ^b \downarrow	217.40 \pm 16.60 (20.51%) ^b \downarrow	0.48 \pm 0.08 (51.02%) ^b \downarrow	5.61 \pm 0.43 (79.23%) ^b \uparrow	2.66 \pm 0.38 (58.33%) ^b \uparrow	41.45 \pm 8.28 (26.23%) ^b \downarrow	74.72 \pm 7.38 (40.95%) ^b \uparrow
CHCl ₃ fraction (72.40 mg/kg; p.o.)	167.18 \pm 14.18 (5.63%) \downarrow	153.41 \pm 14.62 (8.07%) \downarrow	255.23 \pm 17.71 (6.68%) \downarrow	0.87 \pm 0.09 (11.22%) \downarrow	3.55 \pm 0.31 (13.41%) \uparrow	1.79 \pm 0.23 (6.54%) \uparrow	52.63 \pm 8.13 (6.33%) \downarrow	56.11 \pm 7.29 (5.84%) \uparrow
EA fraction (74.55 mg/kg; p.o.)	104.73 \pm 12.13 (40.88%) ^{b,c,d} \downarrow	97.64 \pm 14.17 (41.49%) ^{b,c,d} \downarrow	201.57 \pm 17.61 (26.30%) ^{b,c,d} \downarrow	0.52 \pm 0.07 (46.93%) ^{b,c,d} \downarrow	5.42 \pm 0.45 (73.16%) ^{b,c,d} \uparrow	2.66 \pm 0.27 (58.33%) ^{b,c} \uparrow	43.61 \pm 7.76 (22.38%) ^{b,c,d} \downarrow	64.92 \pm 6.12 (22.46%) ^{b,c,d} \uparrow
n-BuOH fraction (45.40mg/kg; p.o.)	175.89 \pm 13.21 (0.72%) \downarrow	165.28 \pm 14.93 (0.95%) \downarrow	270.47 \pm 16.92 (1.12%) \downarrow	0.90 \pm 0.09 (8.16%) \downarrow	3.24 \pm 0.54 (3.51%) \uparrow	1.72 \pm 0.31 (2.38%) \uparrow	55.23 \pm 7.75 (1.70%) \downarrow	54.25 \pm 6.94 (2.33%) \uparrow

226

227

**The results are expressed as the Mean \pm SD of six rats/group; One way ANOVA followed by Tukey's multiple test*

228

a = p < 0.05 vs normal group; b = p < 0.05 vs CCl₄ control; c = p < 0.05 vs CHCl₃ (72.40 mg/kg); d = p < 0.05 vs n-BuOH (45.40 mg/kg)

229

230

231

* Tel.: +91 7357000002.

E-mail address: dr.pradeepkamboj@gmail.com.

232 4"), 3.53 (*m*, 2H, C-3'-chromone), a doublet at 1.157, *J* = 6 Hz of 3H for CH₃ of rhamnose .
 233 MS (ESI) *m/z* = 580.53 (M⁺) analysed for C₂₇H₃₂O₁₄.
 234 The glycoside was hydrolysed with 5% HCl, which resulted into a crystalline product and
 235 was separated by filtration. The solid was crystallized from methanol to give a crystalline
 236 product (m.p. 250-251 °C, λ_{max} 226 & 292 nm) and was therefore identified as a glycone
 237 (naringenin). The aqueous part was concentrated after neutralization and subjected to paper
 238 chromatography using isopropanol: 5% boric acid (7:3), which resulted in the identification of
 239 two sugars as rhamnose and glucose at R_f 0.43 and 0.18, respectively. Therefore the
 240 compound Dq-4 was confirmed as naringin.


241 **3.7. *In-vitro* hepatoprotective effect in HepG2 cell line**

242 The *in-vitro* hepatoprotective activity of Dq extract, fractions (CHCl₃, EA & *n*-BuOH) and
 243 isolated compound (Dq-4) at dose levels 3, 6, 12.5, 25, 50 & 100 µg/ml were evaluated. The
 244 hepatocytes exposed to CCl₄ (1%) showed a decrease percentage of cells viability (41.25%)
 245 as compared to normal control, indicating the HepG2 cells injury caused by CCl₄ toxicant.
 246 The isolated compounds Dq-4 at dose level 50µg/ml markedly protected the viability of
 247 HepG2 cells against CCl₄-induced cytotoxicity by 91.66%. Similar kinds of results were
 248 observed in case of Dq extract and EA fraction. The percentage protection of Dq extract and
 249 EA fraction at dose level 100µg/ml was 71.62% and 84.33%, respectively. However, CHCl₃
 250 and *n*-BuOH fractions did not show any hepatoprotection in both the assays (Figures 3, 4, 5
 251 & 6).

252 **3.8. Effect on morphology of the HepG2 cells**



253 Normal HepG2 cells (in clumps adherent to the walls) were of spindle shape, clear cell
 254 borders and nuclei with darker cytoplasm. When exposed to toxicant CCl₄ (1%), morphology
 255 of HepG2 cells changed to round shape (swollen), showed irregular and bleeding plasma
 256 membrane, the ability of cells to adhere to walls was decreased and debris emitted
 257 increased around the cells. The changes in the cells were largely prevented with the
 258 increasing concentration of Dq extract, EA fraction and isolated compound (Dq-4) (Figures 3,
 259 4, 5 & 6).
 260
 261

262 **Figure 1. Effect of Dq extract & EA fraction on histological characteristics in CCl₄-**
 263 **induced hepatic injury in rats**





264

265




266



267 **Fv**- fatty vacuoles, **Cn**- centrilobular necrosis, **Fb**- hepatic fibrosis, **Hbd**- hepatocyte
 268 ballooning degeneration and broad infiltration of lymphocytes. Arrows shows the loss of
 269 cellular boundaries
 270
 271


Figure 2: Structure of isolated compound (Dq-4)



272
 273
 274


7-[[2-O-(6-Deoxy- α -L-monopyranosyl)- β -D-glucopyranosyl]oxy]-2,3-dehydro-5,7-dihydroxy-2-(4 hydroxyphenyl)-4H-1-benzopyran-4-one

275 **Figure 3. MTT assay and cytoprotection of Dq extract, fractions (CHCl₃, EA & n-BuOH)**
 276 **and Dq-4 in HepG2 cells**




277 Values are Mean ± SEM of three separate experiments; a = $p < 0.05$ vs normal control; b =
 278 $p < 0.05$ vs CCl₄ control
 279
 280
 281
 282

Figure 4. Face contrast images of treated and untreated HepG2 cells; at 100X




283




Silymarin (25 µg/ml)

284




Dq extract (100 µg/ml)




EA fraction (100 µg/ml) of Dq extract

285
286




CHCl₃ fraction (100 µg/ml) of Dq extract



n-BuOH fraction (100 µg/ml) of Dq extract

287
288
289
290
291
292
293




Dq-4 (50µg/ml); isolated compound

* Tel.: +91 7357000002.


E-mail address: dr.pradeepkamboj@gmail.com.

294 **Figure 5. Trypan blue assay; showing the viability of treated and untreated HepG2**
 295 **cells**
 296



297 Values are Mean ± SEM of three separate experiments; a = $p < 0.05$ vs normal control; b =
 298 $p < 0.05$ vs CCl_4 control
 299
 300
 301
 302


303 **Figure 6. Face contrast images of trypan blue assay: treated and untreated HepG2**
 304 **cells; 40X**



305




Silymarin (25 µg/ml)




EA fraction (100 µg/ml) of Dq extract

306
307



CHCl₃ fraction (100 µg/ml) of Dq extract



Dq-4 (50 µg/ml); isolated compound

308
309
310
311

4. DISCUSSION

312 The present study was undertaken to study the possible hepatoprotective role of *Drynaria*
313 *quercifolia* fronds, a hepatoprotective agent, used by tribals in India, against CCl₄-induced
314 hepatocellular damage.

315 Hepatic damage induced by CCl₄ results from its metabolic bioactivation, primarily through
316 the activity of CYP2E1, to the free radicals CCl₃[•] and CCl₃OO[•] [2]. The free radicals bind
317 covalently to macromolecules by abstracting a hydrogen atom from the polyunsaturated fatty
318 acids of phospholipids and induce peroxidative degradation of the membrane lipids of
319 endoplasmic reticulum [35, 36]. This initiates the formation of lipid peroxides followed by
320 pathological changes such as depression of protein synthesis [37], elevated levels of serum
321 marker enzymes, alkaline phosphatase, total bilirubin, lipid peroxidation [38] and depletion of
322 glutathione content [39]. The antioxidant activity or the inhibition of the generation of free
323 radicals is important in providing protection against hepatic damage [40].

324 Serum transaminases and alkaline phosphatase have long been considered as sensitive
325 indicators of hepatic injury [41]. Injury to the hepatocytes alters enzymes transport function
326 and membrane permeability, leading to leakage of enzymes from the cells; this leakage
327 causes increased levels of AST and ALT in the blood [42]. ALP activity is related to the
328 functioning of hepatocytes, increase in its activity being due to increased synthesis in the

* Tel.: +91 7357000002.

E-mail address: dr.pradeepkamboj@gmail.com.

329 presence of increased biliary pressure [43]. The induced elevation of this enzymatic activity
330 in the blood is associated with high level of bilirubin content [44]. This may explain the
331 increased levels of serum AST, ALT, ALP and bilirubin in CCl₄-intoxicated rats in the present
332 study. The pre-treated groups, Dq extract & EA fraction, induced significant ($p<0.05$)
333 suppression of the increased serum AST, ALT and ALP activities with the significant
334 ($p<0.05$) depletion of raised serum bilirubin. The reduction in the levels of AST and ALT
335 suggests the ability of the test drugs to stabilize the plasma membrane as well as repair of
336 hepatic tissue during CCl₄-induced liver injury. The depletion of increased ALP activity with
337 simultaneous suppression of raised bilirubin level indicates the stabilization of biliary
338 dysfunction in rat liver during the hepatic injury [45].

339 CCl₄ induces fatty liver and cell necrosis, which plays a significant role in diminution of
340 serum protein and albumin, depletion of GSH and increased lipid peroxidation [39, 46]. This
341 expected decline in serum TP, ALB and enhanced lipid peroxidation can be deemed as a
342 useful index for the severity of hepatocellular dysfunction and liver injury [7, 47, 48]. The pre-
343 treated groups, Dq extract & EA fraction, demonstrated hepatoprotective activity by
344 significantly ($p<0.05$) increasing the CCl₄-induced reduction of serum TP and ALB.
345 Scavenging of free radicals is one of the major antioxidation mechanisms to inhibit the chain
346 reaction of lipid peroxidation. A major defense mechanism involves the antioxidant enzymes
347 as well as GSH (non-enzymatic biological antioxidant), which convert active oxygen
348 molecules into non-toxic compounds [46]. Increase in TBARS levels in the CCl₄-intoxicated
349 rats, suggests enhanced lipid peroxidation leading to tissue damage and failure of
350 antioxidant defense mechanisms to prevent formation of excessive free radicals [49]. These
351 effects were significantly ($p<0.05$) reversed in pre-treated groups, attenuation of these
352 hepatotoxic effects might be either through decreased production of free radical derivatives
353 or due to the abilities of Dq extract and EA fraction to act as radical scavengers that might
354 lessen oxidative damage to the liver tissue [50].

355 The histopathological observations of liver sample provided the supportive evidence for the
356 biochemical analysis and explain the hepatoprotective potential of tested plant [51, 52]. The
357 liver of intoxicated rats manifested massive fatty changes, gross necrosis and broad
358 infiltration of lymphocytes and kupffer cells around the central vein and loss of cellular
359 boundaries. The Dq extract and EA fraction pre-treated rats showed a more or less normal
360 architecture, having reversed to a large extent, the hepatic lesions produced by the toxin,
361 thus protecting the histostructural integrity of the liver cells.

362 The human HepG2 cells have proven to be a valuable tool to study *in-vitro* hepatotoxicity of
363 different chemicals or drugs as these cells retain many of the morphological and biochemical
364 characteristics of normal hepatocytes [53, 54]. The percent cell viability, determined using
365 MTT assay and Trypan blue dye exclusion method is useful to predict the cell injury that
366 affects cell attachment or progress to cell death [55]. CCl₄-induced hepatic cell damage
367 causes instability of cell metabolism, inducing triacylglycerol accumulation, increased lipid
368 peroxidation and membrane damage [56]. The Dq extract, EA fraction and Dq-4 exhibited
369 significant ($p<0.05$) restoration of the cell viability and altered morphological changes
370 towards normal in CCl₄ intoxicated HepG2 cells. The reversed hepatotoxic effects is due to
371 the presence of flavonoids in test drugs, which could accelerates the excretion of free radical
372 derivatives and inhibit lipid peroxidation that leads to decrease in severity of oxidative
373 damage in the HepG2 cells [40, 57].

374 5. CONCLUSION

375
376 The present study concludes that the plant *Drynaria quercifolia* exhibited hepatoprotective
377 potential due to the presence of flavonoids. The hepatocellular protection might be due to

* Tel.: +91 7357000002.

E-mail address: dr.pradeepkamboj@gmail.com.

378 the antioxidant & antifibrotic properties and/or due to the membrane stabilizing cascades for
379 the prevention of progression in liver injury. The results of present study support and justify
380 the traditional and folklore medicinal claims attributed to this plant in the treatment of liver
381 ailment. However, in addition to its free radical scavenging potential, further investigations
382 are in process to ascertain the precise cellular/molecular mechanism(s) of hepatoprotective
383 effect.

384

385 **ACKNOWLEDGEMENTS**

386

387 The authors extend their sincere thanks to Sh. Parveen Garg (Chairman, ISF College of
388 Pharmacy) for his full co-operation and for providing the required institutional facilities.

389 **REFERENCES**

390

391 1. Bacon BR, Bisceglie AD. Liver disease: diagnosis and management. Philadelphia:
392 Churchill Livingstone; 1999.

393 2. Gonzalez FJ. The molecular biology of cytochrome P450s. *Pharmacol Rev.*
394 1998;40(4):243-88.

395 3. Guengerich FP. Analysis and characterization of enzymes. In: Hayes AW, editor.
396 Principles and methods of toxicology. Raven Press: New York; 1994.

397 4. Zimmerman HJ. Drug-induced liver disease. In: Zimmerman HJ, editor. Hepatotoxicity: the
398 adverse effects of drugs and other chemicals on the liver. Lippincott Williams & Wilkins:
399 Philadelphia, PA; 1999.

400 5. DeLeve LD, et al. Characterization of a reproducible rat model of hepatic veno-occlusive
401 disease. *Hepatology.* 1999;29(6):1779-91.

402 6. Amin A, Hamza AA. Hepatoprotective effects of Hibiscus, Rosmarinus and Salvia on
403 azathioprine-induced toxicity in rats. *Life Sci.* 2005;77(3):266-78.

404 7. Navarro VJ, Senior JR. Drug-related hepatotoxicity. *New Engl J Med.* 2006;354(7):731-39.

405 8. Handa SS, Sharma A, Chakraborti KK. Natural products and plants as liver protecting
406 drugs. *Fitoterapia.* 1986;57(5):307-51.

407 9. Chatterjee TK. Medicinal plants with hepatoprotective properties. In: Herbal Options.,
408 Books and Allied (P) Ltd., Kolkata; 2000.

409 10. Kirtikar KR, Basu BD. Indian Medicinal Plants. Dehra Dun Publisher Ltd., Dehra Dun,
410 India; 1994.

411 11. Bhattacharya S. In: Chrinjib Banoushadi. Anand Publishing Ltd., Calcutta, India; 1990.

412 12. Dixit RD, Vohra JN. A dictionary of pteridophytes of India. In: Botanical survey of India.
413 Howrah: India; 1984.

* Tel.: +91 7357000002.

E-mail address: dr.pradeepkamboj@gmail.com.

- 414 13. Irudayaraj V, Senthamarai R. Pharmacognostical studies on a medicinal fern, *Drynaria*
415 *quercifolia* (L.) J. Sm. (Polypodiaceae: Pteridophyta). Phytomorphology. 2004;54(3-4):193-
416 200.
- 417 14. Shokeen P, et al. Preliminary studies on activity of *Ocimum sanctum*, *Drynaria*
418 *quercifolia*, and *Annona squamosa* against *Neisseria gonorrhoeae*. Sex Transm Dis.
419 2005;32(2):106-11.
- 420 15. Rajendran A, Rajan S. *Drynaria quercifolia*: an antifertility agent. Ancient Sci Life.
421 1996;15(4):286-87.
- 422 16. Khan A, et al. Isolation of antibacterial constituent from rhizome of *Drynaria quercifolia*
423 and its sub-acute toxicological studies. DARU. 2007;15(4):205-11.
- 424 17. Koushik P, Dhiman AK. Common medicinal pteridophytes. Indian Fern Journal.
425 1995;12(1-2):139-45.
- 426 18. Harborne JB. Phytochemical methods: A guide to modern techniques of plant analysis.
427 Chapman and Hall, London; 1984.
- 428 19. Farnsworth NR. Biological and phytochemical screening of plants. J Pharmaceut Sci.
429 1966;55(3):225-76.
- 430 20. OECD, In: Guidelines for the testing of chemicals revised draft guideline 423: acute oral
431 toxicity. 2000.
- 432 21. Botham PA. Acute systemic toxicity prospects for tiered testing strategies. Toxicol Vitro.
433 2004;18:227-30.
- 434 22. Rosa MPG, Solis RV. Hepatoprotective and inhibition of oxidative stress in liver of
435 *Prostechea michuacana*. Record Nat Prod. 2009;3(1):46-51.
- 436 23. Bradley DW, et al. Transaminase activities in serum of long-term hemodialysis patients.
437 Clin Chem. 1972;18:1442.
- 438 24. Lieberman D, Phillips D. Isolated elevation of alkaline phosphatase: significance in
439 hospitalized patients. J Clin Gastroenterol. 1990;12(4):415-19.
- 440 25. Jendrassik L, Grof P. Determination of total and direct bilirubin in serum or plasma on
441 manual system. Biochemische Zeitschrift. 1938;297:81-9.
- 442 26. Doumas BT. Standards for total serum protein assays-a collaborative study. Clin Chem.
443 1975;21:1159-66.
- 444 27. Ingwersen S, Raabo E. Improved and more specific bromcresol green methods for the
445 manual and automatic determination of serum albumin. Clin Chim Acta. 1978;88(3):545-50.
- 446 28. Moron MJ, Dipierre JW, Mannerv KB. Levels of glutathione, glutathione reductase and
447 glutathione-s-transferase activities in rat lungs and liver. Biochim Biophys Acta.
448 1979;582:67-71.

* Tel.: +91 7357000002.

E-mail address: dr.pradeepkamboj@gmail.com.

- 449 29. Wills ED. Mechanisms of lipid peroxide formation in animal tissues. *Biochem J.*
450 1996;99(3):667.
- 451 30. Galigher AE, Kozloff EN. *Essentials of Practical Microtechniques.* Lea and Febiger:
452 Philadelphia; 1971.
- 453 31. Krajian AA. Tissue cutting and staining. In: Frankel S, Reitman S, editors. *Gradwohl's*
454 *Clinical Laboratory methods and diagnosis. The CV.* Mosby Co.: Saint Louis, USA; 1963.
- 455 32. Mitra SK, et al. Liv.52 regulates ethanol induced PPAR c and TNF a expression in
456 HepG2 cells. *Mol Cell Biochem.* 2008;315(1-2):9-15.
- 457 33. Ke H, et al. Antineoplastic agents III: steroidal glycosides from *Solanum nigrum.* *Planta*
458 *Medica.* 1999;65(1):35-38.
- 459 34. Freshney I. Cytotoxicity, culture of animal cells. In: *A manual of basic technique.* Wiley-
460 Liss: New York; 2000.
- 461 35. Cesaratto L, et al. The importance of redoxstate in liver damage. *Ann Hepatol.*
462 2004;3(3):86-92.
- 463 36. Weber LW, Boll M, Stampfl A. Hepatotoxicity and mechanism of action of haloalkanes:
464 carbon tetrachloride as a toxicological model. *Crit Rev Toxicol.* 2003;33(2):105-36.
- 465 37. Faroon O, De Rosa CT, Smith L. Carbon tetrachloride: Health affects toxicokinetics,
466 human exposure and environment fate. *Toxicol Ind Health.* 1994;10:4-20.
- 467 38. Zimmerman HJ, Seeff LB. Enzymes in hepatic disease. In: Goodly EE, editor. *Diagnostic*
468 *enzymology.* Lea and Febiger: Philadelphia; 1970.
- 469 39. Kamiyama T, Sato C, Liu J. Role of lipid peroxidation in acetaminophen induced
470 hepatotoxicity; comparision with carbontetrachloride. *Toxicol Lett.* 1993;66:7-12.
- 471 40. Ferguson LR. Role of plant polyphenols in genomic stability. *Mutat Res.* 2001;475(1-
472 2):89-111.
- 473 41. Kew MC. Serum aminotransferase concentration as evidence of hepatocellular damage.
474 *Lancet.* 2000;355:591-92.
- 475 42. Nkosi CZ, Opoku AR, Terblanche SE. Effect of pumpkind seed (*Cucurbita pepo*) protein
476 isolate on the activity levels of certain plasma enzymes in CCl₄-induced liver injury in low
477 protein fed rats. *Phytother Res.* 2005;19(4):341-45.
- 478 43. Moss DW, Butterworth PJ. *Enzymology and Medicine.* Pitman Medical: London; 1974.
- 479 44. Pimple BP, et al. Protective effect of *Tamarindus indica* Linn against paracetamol
480 induced hepatotoxicity in rats. *Indian J Pharmaceut Sci.* 2007;69(6):827-31.
- 481 45. Mukherjee PK. Quality control herbal drugs. In: *An approach to evaluation of botanicals.*
482 1st ed New Delhi: Business Horizons; 2002.
- 483 46. Husain K, Somani SM. Interaction of exercise training and chronic ethanol ingestion on
484 hepatic and plasma antioxidant system in rat. *J Appl Toxicol.* 1997;17(3):189-94.

* Tel.: +91 7357000002.

E-mail address: dr.pradeepkamboj@gmail.com.

- 485 47. Hampel B, et al. Differential regulation of apoptotic cell death in senescent human cells.
486 Exp Gerontol. 2004;39(11-12):1713-21.
- 487 48. Brattin WJ, Glende Jr EA, Recknagel RO. Pathological mechanisms in carbon
488 tetrachloride hepatotoxicity. Free Radic Biol Med. 1985;1(1):27-38.
- 489 49. Souza MF, Rao VSN, Silveira ER. Inhibition of lipid peroxidation by ternatin, a
490 tetramethoxyflavone from *Egletes viscosa* L. Phytomedicine. 1997;4(1):27-31.
- 491 50. Blair IA. Endogenous glutathione adducts. Curr Drug Metabol. 2006;7:853-72.
- 492 51. Plaa GL, Charbonneau M. Detection and evaluation of chemically induced liver injury. In:
493 Hayes AW, editor. Principles and methods of toxicology. Raven Press: New York; 1989.
- 494 52. Junnela M, et al. Reduction of carbon tetrachloride induced hepatotoxic effects by oral
495 administration of betaine in male Hans-Wistar rats: A morphometric histological study.
496 Veterinary Pathology. 2000;37:231-38.
- 497 53. Gomez-Lechon MJ, et al. The use of cultured hepatocytes to investigate the metabolism
498 of drugs and mechanisms of drug hepatotoxicity. Alternatives to Laboratory Animals.
499 2001;29(3):225-31.
- 500 54. Guillouzo A. Liver cell models in *in-vitro* toxicology. Environ Health Perspect.
501 1998;106(2):511-32.
- 502 55. Knowles BB, Aden DP. Human hepatoma derived cell line, process for preparation
503 thereof, and uses therefor. US Patent 4393133; 1983.
- 504 56. Recknagel RO, et al. Mechanisms of carbon tetrachloride toxicity. Pharmacol Therapeut.
505 1989;43(43):139-54.
- 506 57. Tapas AR, Sakarkar DM, Kakde RB. Flavonoids as nutraceuticals: a review. Trop J
507 Pharmaceut Res. 2008;7(3):1089-99.