

Bacterial endophytes of the medicinal herb *Hygrophila spinosa* T. Anders and their antimicrobial activity

Arundhati Pal^{1*} and A. K. Paul²

¹Department of Botany, Serampore College,
9, William Carey Road, Serampore, Hooghly, West Bengal 712 201, India

²Microbiology Laboratory, Department of Botany, University of Calcutta,
35, Ballygunge Circular Road, Kolkata 700 019, India

ABSTRACT

Aims: The ethnobotanical herb *Hygrophila spinosa* T. Anders (Acanthaceae) is native to India and used in traditional ayurvedic medicines for its pharmacologically important phytochemicals. This study aims to isolate and characterize the culturable bacterial endophytes of *H. spinosa* and evaluate their antimicrobial properties.

Place and Duration of Study: The experiments were performed in the Department of Botany, Serampore College, Serampore as well as in the Microbiology Laboratory, Department of Botany, University of Calcutta, Kolkata during 2011 to 2012.

Methodology: Bacterial endophytes were isolated from healthy plant tissues following surface sterilization and plating on nutrient agar, glycerol asparagine agar and tryptic soy agar. They were characterized physio-biochemically following standard microbiological and biochemical methods. The endophytes were screened for production of antimicrobial compounds following cross-streak assay against test strains *Bacillus subtilis*, *B. cereus*, *Escherichia coli*, *Pseudomonas cepacia*, *Klebsiella pneumoniae* and *Staphylococcus aureus* on nutrient agar plates.

Results: Eleven phenotypically distinguishable bacterial endophytes were isolated from surface sterilized leaf, stem and root tissues and Shannon Weaver diversity index clearly revealed more diverse (0.83) types of endophytes in leaves than in stem (0.48) and root (0.41) tissues. Physio-biochemical features of the isolates clearly indicated distinct variation in their sugar fermentation profiles along with NaCl tolerance. The endophytes produced important enzymes like catalase, amylase, gelatinase, nitrate reductase and lipase. The bacterial isolates belonged to the genera *Bacillus*, *Paenibacillus*, *Pseudomonas*, *Ralstonia*, *Staphylococcus*, *Micrococcus* and *Acidomonas*. Antibiotic sensitivity profile, however, have indicated that the isolates were mostly resistant to amoxicillin and bacitracin, while they were highly susceptible to tetracycline followed by neomycin and streptomycin. Interestingly, the bacterial endophytes of *H. spinosa* give a definite stamp on their antimicrobial activity against *E. coli* and *K. pneumoniae* followed by *S. aureus*. Two isolates, *Paenibacillus* HGS 202 and *Acidomonas* HGR 302 obtained from stem and root segments respectively showed antimicrobial activity against *B. subtilis*, *B. cereus*, *E. coli*, *K. pneumoniae* and *S. aureus*.

Conclusion: This study identified eleven bacterial endophytes harboring the leaves, stem and root of *H. spinosa* which demonstrated antibacterial activity against Gram-positive as well as Gram-negative bacterial strains. Moreover these endophytic bacterial isolates could be exploited as sources of antibacterial substances.

Keywords: *Hygrophila spinosa*, Endophytic bacteria, Antibacterial activity, Antibiotic sensitivity, Enzyme profile, NaCl tolerance

21 **1. INTRODUCTION**

22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77

Medicinal plants provide valuable therapeutic agents in traditional medicines which are used on a global level for helping with a wide variety of human health issues. *Hygrophila spinosa* T. Anders, belonging to the family Acanthaceae, is a promising medicinal herb mentioned in ancient ayurvedic literature as having great economic potential. The plant is indigenous to the Indian subcontinent and is reported to contain phytosterols, fatty acids, polyphenols, proanthocyanins, alkaloids, flavonoids, terpenoids, vitamins, and glycosides as major chemical constituents. In traditional medicine, *H. spinosa* is used mainly for the treatment of hyperdipsia, vesical calculi, flatulence, diarrhea, dysentery, leukorrhea, gonorrhoea, asthma, blood diseases, gastric problems, cancer, rheumatism, etc. Many essential phytochemicals isolated from the whole plant including lupeol, stigmaterol, apigenin-7-O-glucuronide, apigenin-7-O-glucoside, betulin, 25-oxo-hentriacontanyl acetate, methyl 8-*n*-hexyltetracosanoate, oleic acid, linoleic acid, etc. have exhibited antitumor, antibacterial, antidiabetic, antiinflammatory, antipyretic, antioxidant and hepatoprotective activity [1, 2].

It has been rationalized that plants having an ethnobotanical history and exploited for human use in traditional medicine may harbor an endophytic population which may produce a plethora of microbial metabolites related closely to the plant biochemistry [3]. Endophytes, by definition, are microorganisms colonizing living internal tissues of plant either symbiotically or in mutualistic relationship. They occur ubiquitously in all plant species on earth and benefit the host plant growth by fixation of atmospheric nitrogen, production of growth promoting substances, imparting effective disease management, plant protection and stress tolerance [4]. In addition recent studies have established that secondary metabolites elaborated by these microbial endophytes could serve as prospective resources of antimicrobial substances, antioxidants, cytotoxic compounds, growth hormones and hydrolytic enzymes of biotechnological applications [5, 6].

In view of the increasing prevalence of antibiotic-resistant human and plant pathogens, there is an escalating demand for newer antimicrobials from natural sources. Bacterial and fungal endophytes residing inside the healthy plant tissues are believed to carry out a resistance mechanism to overcome pathogenic attack and have emerged as a promising source of newer antimicrobial compounds. Several antimicrobial metabolites belonging to structural classes like alkaloids, peptides, benzopyranones, flavonoids, phenolic acids, quinones, steroids, terpenoids, tetralones, xanthenes, and others have been obtained from endophytes. The occurrence of endophytic bacteria in agricultural or medicinal plants has been reported quite extensively [7-9]. A comparison of different endophytic hosts shows that nearly 35% of the endophytes possessing antimicrobial activity have been isolated from medicinal plants followed by 29% from agricultural crops [6]. The diversity and ecological distribution of fungal endophytes associated with different medicinal plants native to China, Malaysia, Australia and India have been investigated with special emphasis on their antimicrobial efficacy. A mass of bioactive natural products isolated from endophytes have been reported in recent years and majority of them have been derived from endophytic fungi [3, 7, 10, 11]. However, little information is available on the occurrence as well as on the potential significance of bacterial endophytes from medicinal plants. Although, medicinal properties of *H. spinosa* have been studied in details by many researchers [1, 2], reports on the endophytic population of this medicinal herb is lacking. Biodiversity of both culturable and unculturable endophytic microbial communities of *H. spinosa*, therefore, needs to be determined. However, culturable endophytic bacterial isolates deserve special attention for further development of microbial-based biotechnological products and formulations. In the present study, we focused on the isolation, characterization and antimicrobial evaluation of bacterial endophytes from *H. spinosa*.

78 **2. MATERIAL AND METHODS**

79

80 **2.1 Collection of plant samples**

81 Healthy plants of *Hygrophila spinosa* T. Anders (Acanthaceae) were collected from
82 Medicinal Plant Garden of Serampore College, Hooghly, West Bengal and Department
83 of Botany, University of Calcutta, Kolkata in sterile zip lock polythene bags. The collected
84 plants were brought immediately to the laboratory and stored at 4°C until used for the
85 isolation of bacterial endophytes.

86

87 **2.2 Isolation and characterization of endophytes**

88 Fresh and healthy leaf, stem and root segments were cut from the collected plants,
89 washed thoroughly under running tap water. Surface sterilization was performed in
90 sterile glass bottles by consecutive immersion in 70% ethanol (2 – 3 min), 0.5 % sodium
91 hypochlorite (5 -10 min) and again in 70% ethanol for 30 sec [7]. This was followed by
92 repeated washing of plant samples in sterile distilled water for at least three times.
93 Samples were blot dried on sterile towels and cut aseptically into small sections before
94 plating on previously prepared nutrient agar, glycerol asparagine agar and tryptic soy
95 agar plates for isolation of bacteria. The plates were incubated at 30°C for 2 – 4 days
96 and observed for growth of bacterial colonies surrounding the leaf, stem and root
97 sections. Pure cultures of bacterial endophytes were developed by dilution-streaking on
98 the same media and maintained on slopes of nutrient agar for further study. Bacterial
99 strains were characterized and identified following micromorphological and physio-
100 biochemical analysis following standard protocols [12, 13].

101

102 **2.3 Diversity of endophytes**

103 Based on the total number of samples plated and the number of samples yielding
104 isolates, colonization frequency and isolation rate were calculated. Colonization
105 frequency was calculated as the total number of plant samples infected by bacteria
106 divided by the total number of samples incubated. Isolation rate was determined as the
107 number of bacterial isolates obtained from plant samples divided by the total number of
108 samples incubated. The Shannon Weaver biodiversity index H' was calculated as:

109 $H' = -\sum P_i \times \ln P_i$, where, P_i is the proportion of individuals that species “ i ” contributes to
110 the total [7, 14].

111

112 **2.4 Antibiotic susceptibility spectrum**

113 Antibiotic sensitivity test was performed following the Kirby Bauer disc-diffusion assay
114 method [15] using antibiotic impregnated discs (6 mm diameter) from Himedia (India).
115 Based on the diameter of inhibition zone recorded to nearest mm, the organisms were
116 categorized as resistant, intermediate and sensitive following DIFCO Manual 10th edition
117 (1984). Antibiotics used include: amoxycillin (30 µg/disc), bacitracin (10 U/disc),
118 chloramphenicol (30 µg/disc), neomycin (30 µg/disc), streptomycin (30 µg/disc) and
119 tetracycline (30 µg/disc).

120

121 **2.5 Production of antimicrobial substances**

122 Bacterial endophytes were primarily screened for production of antimicrobial substances
123 following cross-streak assay method using six test organisms like *Bacillus subtilis*, *B.*
124 *cereus*, *Escherichia coli*, *Pseudomonas cepacia*, *Klebsiella pneumonia* and
125 *Staphylococcus aureus* [16]. Nutrient agar plates were inoculated with bacterial
126 endophytes as a single streak at the centre of the Petri plate and incubated for 5 days at
127 30°C. Overnight grown cultures of the test organisms were streaked at right angle to the
128 producer endophyte and observed for its growth / inhibition after 24 – 48 h of incubation
129 at 30°C. The length of inhibition zone was measured to nearest mm.

130

131

132

133 **3. RESULTS**

134
135 **3.1 Diversity of bacterial endophytes**

136 Segments of surface sterilized leaf, stem and root of *Hygrophila spinosa* (Acanthaceae)
137 incubated on nutrient agar, glycerol asparagine agar and tryptic soy agar plates showed
138 growth of morphologically distinguishable bacterial colonies surrounding the segments
139 after 48-96 h. Avoiding the repetitive strains a total of 11 phenotypically distinguishable
140 bacterial endophytes were isolated in pure form from 118 segments (39 leaf, 39 stem
141 and 40 root) of *H. spinosa*. Out of these 11 isolates, six were derived from leaf, while
142 stem and root segments yielded three and two isolates respectively (Table 1). The
143 colonization frequency was lower in leaf samples (17.9%) as compared to the stem
144 (20.5%) and root (22.5%), while the isolation rate was poor in root (0.05) but increased
145 gradually in stem (0.07) and leaf (0.15) samples. The Shannon-Weaver diversity index
146 showed that leaves (0.83) of *H. spinosa* harbor more diverse types of endophytic
147 bacteria than in its stem (0.48) and root (0.41).

148
149
150 **Table 1. Diversity of endophytic bacterial isolates in leaf, stem and root tissues**
151 **of *Hygrophila spinosa***
152

Parameters	Plant tissue			Total
	Leaf	Stem	Root	
Number of samples	39	39	40	118
Number of sample yielding isolates	07	08	09	24
Number of isolates	06	03	02	11
Colonization Frequency, % ^a	17.9	20.5	22.5	20.3
Isolation Rate ^b	0.15	0.07	0.05	0.09
Shannon-Weaver Diversity Index ^c	0.83	0.48	0.41	0.68

153 ^a Colonization frequency was calculated as the total number of plant samples infected by bacteria
154 divided by the total number of samples incubated. ^b Isolation rate was calculated as the number of
155 bacterial isolates obtained from plant samples divided by the total number of samples incubated.

156 ^c Shannon Weaver diversity index H' was calculated as: $H' = -\sum Pi \times \ln Pi$, where, Pi is the
157 proportion of individuals that species "i" contributes to the total [7, 14].
158
159

160 **Table 2. Micromorphological characteristics of bacteria isolated from leaf, stem**
161 **and root tissues of *Hygrophila spinosa***
162

Tissue	Isolate no.	Cell morphology	Gram nature	Motility	Size, µm	Endospore	Diffusible pigments
Leaf	HGL 101	cocci, in cluster	positive	non-motile	0.5 dia	absent	none
	HGL 102	cocci, single	positive	non-motile	0.4 dia	absent	yellow
	HGL 103	short rod	negative	motile	0.4 X 0.3	absent	green
	HGL 104	rod, single	positive	motile	1.1 X 0.3	present	none
	HGL 105	short rod	positive	non-motile	0.5 X 0.4	present	none
	HGL 106	short rod	negative	motile	0.5 X 0.3	absent	none
Stem	HGS 201	rod, in chain	positive	motile	1.1 X 0.5	present	none
	HGS 202	rod, single	positive	motile	0.8 X 0.4	present	none
	HGS 203	cocci, single	positive	non-motile	0.5 dia	absent	yellow
Root	HGR 301	short rod	negative	motile	0.5 X 0.4	absent	none
	HGR 302	short rod	negative	non-motile	0.5 X 0.4	absent	none

163 Colony morphology was detected in Tryptic soy agar medium after 5 days of growth at 32°C.

164 **3.2 Characterization and identification of isolates**

165 The bacterial endophytes of *H. spinosa* were characterized based on
 166 micromorphological (Table 2) and physio-biochemical characters (Table 3). Out of 11
 167 isolates seven were Gram-positive (three cocci and four rod) and four were Gram-
 168 negative (all rod). Filamentous forms were not detected in any of the plant samples. Six
 169 isolates out of 11 showed motility and only three produced yellowish to green diffusible
 170 pigments during growth on tryptic soy agar plates. All Gram-positive rods showed
 171 endospore formation.

172
 173 Enzymatic profile of endophytic bacterial isolates showed that all of them produced
 174 catalase, while about 55 and 64% of the isolates produced amylase and gelatinase
 175 respectively (Table 3). Lipolytic (55%) and nitrate reductase (36%) activities were not
 176 uncommon amongst the endophytic isolates. Production of indole by the enzyme
 177 tryptophanase was evident only in isolates HGL 103, HGL 105 and HGR 301. The
 178 isolates showed wide degree of tolerance to NaCl (2.5 – 10%) in the growth medium.
 179 The endophytes were also screened for their ability to utilize and ferment dextrose,
 180 fructose, maltose, sucrose and lactose in phenol red agar medium supplemented with
 181 1% sugar (Table 4). While dextrose was the best carbohydrate utilized by all most all the
 182 bacterial endophytes, lactose was fermented by only two isolates. The endophytic
 183 isolates were moderate in fermenting fructose, sucrose and maltose.

184
 185
 186 **Table 3. Biochemical characterization of bacterial endophytes from leaf, stem**
 187 **and root tissues of *H. spinosa***

Plant tissue	Isolate no.	Enzyme profile					Indole production	NaCl tolerance, %
		Catalase	Amylase	Gelatinase	Lipase	NO ₃ Reductase		
Leaf	HGL 101	+	+	+	+	-	-	10.0
	HGL 102	+	-	+	-	-	-	10.0
	HGL 103	+	-	+	+	-	+	3.5
	HGL 104	+	-	+	-	+	-	4.0
	HGL 105	+	-	-	+	+	+	4.0
	HGL 106	+	-	-	-	-	-	4.5
Stem	HGS 201	+	+	-	+	-	-	4.0
	HGS 202	+	+	+	-	-	-	4.0
	HGS 203	+	+	+	-	+	-	10.0
Root	HGR 301	+	+	+	+	-	+	3.0
	HGR 302	+	+	-	+	+	-	2.5

189 “+” indicate positive response, “-” indicate negative response

190
 191
 192 **Table 4. Fermentation of sugars by bacterial endophytes isolated from leaf, stem**
 193 **and root tissues of *H. spinosa***

Plant tissue	Isolate no.	Fermentation of sugars				
		Dextrose	Fructose	Lactose	Maltose	Sucrose
Leaf	HGL 101	+	+	-	+	+
	HGL 102	+	+	-	-	-
	HGL 103	+	-	-	-	-
	HGL 104	+	+	-	-	+
	HGL 105	+	+	-	+	+
	HGL 106	-	-	+	-	-
Stem	HGS 201	+	-	-	-	-
	HGS 202	+	+	-	-	+
	HGS 203	+	+	+	+	+
Root	HGR 301	+	+	-	+	-
	HGR 302	+	-	-	-	-

195 “+” indicate positive response, “-” indicate negative response

196 Fermentation of sugars was screened in phenol red agar medium supplemented with 1% sugar.

197 Based on microscopic and biochemical analysis, the bacterial isolates were tentatively identified as species of *Bacillus* (HGL 104, HGS 201),
 198 *Paenibacillus* (HGL 105, HGS 202), *Pseudomonas* (HGL 103, HGR 301), *Ralstonia* (HGL 106), *Staphylococcus* (HGL 101), *Micrococcus* (HGL
 199 102, HGS 203) and *Acidomonas* (HGR 302).
 200

201 3.3 Antibiotic sensitivity profile

202 Antibiotic sensitivity pattern of the endophytic bacterial isolates was determined by disc-diffusion method against six different antibiotics like
 203 amoxycillin, bacitracin, chloramphenicol, neomycin, streptomycin and tetracycline. Results as shown in Table 5 depict that, bacterial endophytes
 204 from leaf, stem and root tissues of *H. spinosa* were mostly resistant to amoxycillin and bacitracin, while they were mostly sensitive to tetracycline
 205 followed by neomycin and streptomycin. One leaf endophyte, *Staphylococcus* HGL 101 was highly resistant to five antibiotics and was followed
 206 by *Micrococcus* HGS 203 showing resistance to four of the six tested antibiotics. On the contrary, the isolates from leaf and stem (*Paenibacillus*
 207 HGL 105, *Bacillus* HGS 201 and *Paenibacillus* HGS 202) showed sensitive to intermediate response towards all the tested antibiotics.
 208
 209

210 **Table 5. Screening of bacterial endophytes from *Hygrophila spinosa* for their antibiotic susceptibility following disc-diffusion assay**
 211

Plant tissue	Isolate	Diameter of inhibition zone, mm					
		Antibiotics					
		Amoxycillin	Bacitracin	Chloramphenicol	Neomycin	Streptomycin	Tetracycline
Leaf	<i>Staphylococcus</i> HGL 101	08 (R)	NIL (R)	9.5 (R)	12 (R)	11 (R)	40 (S)
	<i>Micrococcus</i> HGL 102	14 (I)	12 (I)	22 (S)	20 (S)	32 (S)	10 (R)
	<i>Pseudomonas</i> HGL 103	22 (S)	14 (S)	NIL (R)	22 (S)	32 (S)	NIL (R)
	<i>Bacillus</i> HGL 104	23 (S)	NIL (R)	26 (S)	18 (S)	18 (I)	19 (S)
	<i>Paenibacillus</i> HGL 105	14 (I)	12 (I)	18 (S)	24 (S)	30 (S)	26 (S)
	<i>Ralstonia</i> HGL 106	11 (R)	12 (I)	18 (S)	28 (S)	36 (S)	44 (S)
Stem	<i>Bacillus</i> HGS 201	25 (S)	13 (S)	14 (I)	20 (S)	27 (S)	24 (S)
	<i>Paenibacillus</i> HGS 202	20 (S)	16 (S)	17 (I)	16 (I)	32 (S)	20 (S)
	<i>Micrococcus</i> HGS 203	09 (R)	NIL (R)	9.5 (R)	21 (S)	NIL (R)	19 (S)
Root	<i>Pseudomonas</i> HGR 301	7.5 (R)	NIL (R)	26 (S)	14 (I)	NIL (R)	20 (S)
	<i>Acidomonas</i> HGR 302	11 (R)	08 (R)	21 (S)	16 (I)	25 (S)	22 (S)

212
 213
 214

R=Resistant, I=Intermediate, S=Sensitive

Antibiotic susceptibility was tested on nutrient agar plates using antibiotic impregnated discs (6 mm) from HIMEDIA, India

215 **3.4 Evaluation of antimicrobial activity**

216 Antimicrobial activity of all eleven bacterial endophytes were assessed against six bacterial test organisms, *B. subtilis*, *B. cereus*, *E. coli*, *P.*
 217 *cepacia*, *K. pneumoniae* and *S. aureus* following cross-streak method on nutrient agar plates. The isolate which inhibited growth of any of the
 218 test isolate(s) was considered having antibacterial activity and the length of inhibition zone was measured (Table 6). Out of 11 endophytes
 219 screened, majority showed antibacterial activity against *E. coli* and *K. pneumoniae* followed by *S. aureus*. Isolates *Paenibacillus* HGS 202 and
 220 *Acidomonas* HGR 302 obtained from stem and root tissues respectively showed comparatively broad spectrum of antibacterial activity inhibiting
 221 both Gram-positive and Gram-negative test organisms.

222
 223
 224
 225

Table 6. Evaluation of antimicrobial activity of bacterial endophytes of *Hygrophila spinosa* following cross-streak method

Plant tissue	Isolate	Length of inhibition zone, mm					
		Test organisms					
		<i>Bacillus subtilis</i>	<i>Bacillus cereus</i>	<i>Pseudomonas cepacia</i>	<i>Escherichia coli</i>	<i>Klebsiella pneumoniae</i>	<i>Staphylococcus aureus</i>
Leaf	<i>Staphylococcus</i> HGL 101	-	-	-	20	10	-
	<i>Micrococcus</i> HGL 102	-	-	-	-	-	-
	<i>Pseudomonas</i> HGL 103	-	-	-	-	-	-
	<i>Bacillus</i> HGL 104	-	-	-	20	10	-
	<i>Paenibacillus</i> HGL 105	-	-	5	-	-	5
	<i>Ralstonia</i> HGL 106	-	-	-	-	5	-
Stem	<i>Bacillus</i> HGS 201	-	-	-	20	20	-
	<i>Paenibacillus</i> HGS 202	1	1	3	6	-	3
	<i>Micrococcus</i> HGS 203	-	-	-	20	8.5	8
Root	<i>Pseudomonas</i> HGR 301	-	-	-	20	5	-
	<i>Acidomonas</i> HGR 302	4	2	-	20	5	3

226
 227
 228

“-” means no inhibition zone produced

229 **DISCUSSION**

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

Studies on the diversity of culturable microbial endophytes in medicinal and vegetative crop plants are essential to understand their potentials and importance in different fields of biotechnology. This study is the first attempt to isolate microbial endophytes from the traditional medicinal herb *H. spinosa*. We have screened only the medicinally important plant organs like root, stem and leaf of *H. spinosa*, although endophytes could also harbor in flower, fruit and seeds. The leaves of *H. spinosa* were found to harbor more diverse types of bacterial endophytes than stem or root segments (Table 1). Such species richness in leaves may be attributed to the anatomical peculiarities of the leaves and micro-environmental conditions rich in essential nutrients which drives the selective force for survival of tissue specific endophytic taxa. Similar prevalence of endophytes in leaf tissues have been observed in *Paederia foetida* [17], *Kigelia pinnata* [18] and *Quercus ilex* [19].

Spatial distribution of endophytic genera also depends on seasonal variation, precipitation, soil parameters and location of plants, plant age and genotypes [4]. Here, we have tested only one genotype from cultivated soil of two different localities which does not reflect the true portrait of culturable endophyte diversity of *H. spinosa*. The phenotypically distinguishable bacterial endophytes harboring leaves, stem and root tissues of *H. spinosa* were characterized in details (Table 2 - 4) and tentatively identified as members belonging to bacterial genera *Bacillus*, *Paenibacillus*, *Pseudomonas*, *Ralstonia*, *Staphylococcus*, *Micrococcus* and *Acidomonas*. These isolates were mostly the fast growing endophytes and were also reported to colonize several other host plants. Occurrences of similar endophytic bacterial genera have been reported from medicinal plants like *Gynura procumbens*, *Azadirachta indica*, *Boerhaavia diffusa*, *Phyllanthus emblica*, *P. foetida* etc. [17, 20-22]. In addition, several authors have reported the presence of endophytic actinobacteria inside medicinal plants belonging to the genera *Streptomyces*, *Pseudonocardia*, *Promicromonospora*, etc. [23, 24]. However, such filamentous forms have not been recorded during the present study.

Information regarding production of enzymes by microbes of plant origin are few although endophytic bacteria isolated from leaves of maize [25], leaves and stem of *Jacaranda decurrens* [26], roots of *Chlorophytum borivilianum* [27] and leaves of mangrove plants [28] have been reported to produce hydrolytic enzymes of diverse types. All the aerobic endophytic isolates of *H. spinosa* possessed catalase responsible for the decomposition of hydrogen peroxide to less reactive oxygen and water molecules. Production of hydrolytic enzymes gelatinase, amylase and lipase (Table 3) supports earlier observations [25-28]. The presence of nitrate reductase and tryptophanase in some of the isolates appears to play a key role in the nitrogen cycle and has important agricultural, environmental and public health implications. The emergence of antibiotic resistance is not only limited to pathogenic microorganisms but also found amongst environmental isolates as a result of horizontal transfer of antibiotic resistance genes. Majority of the endophytes from *H. spinosa* showed resistance to amoxicillin and bacitracin (Table 5) similar to those encountered in bacterial endophytes of *P. foetida* [17], *Andrographis paniculata* [29] and mangrove plants [28].

In view of the ever increasing demand for novel antimicrobial substances, the endophytes have been identified as a potential source of antibiotics [6]. Several reports on the antimicrobial evaluation of endophytic fungi from medicinal plants have been presented [30-32], antimicrobial activities of endophytic bacteria are not uncommon [17, 20, 29]. Li *et al.* [30], however, have explored endophytic actinomycetes associated with pharmaceutical plants in rainforest of Yunnan, China and detected endophytic *Streptomyces* displaying antimicrobial activities against *S. aureus*, *E. coli* and *C. albicans*. In the present study, nine bacterial endophytes out of 11 from *H. spinosa* showed antibacterial activity against *B.*

282 *subtilis*, *B. cereus*, *E. coli*, *P. cepacia*, *K. pneumoniae* and *S. aureus* following cross-streak
283 assay (Table 6) and two of them showed broad spectrum antimicrobial activity indicating
284 possible biotechnological potential. However, isolation, purification and detection of active
285 compounds is in progress for their further utilization.

286

287 **4. CONCLUSION**

288

289 Endophytic bacterial isolates was found to be associated with leaves, stem and root of the
290 medicinal plant, *H. spinosa* and they differed significantly in their morphological,
291 physiological and biochemical characters. The endophytes also produced several hydrolytic
292 enzymes of commercial importance. Antimicrobial evaluation of these culturable endophytes
293 of *H. spinosa* has shown that they possess antibacterial activity against various bacterial
294 species. The endophytes of traditional medicinal plants appear to be a potential source of
295 antimicrobial metabolites as well as enzymes.

296

297 **ACKNOWLEDGEMENTS**

298

299 Financial support from University Grants Commission, New Delhi (UGC-Minor Research
300 Project PSW – 061 / 10-11 ERO) to A. Pal is duly acknowledged.

301

302 **COMPETING INTERESTS**

303

304 Authors have declared that no competing interest exists in performing this research and
305 preparation or publication of the results.

306

307

308 **AUTHORS' CONTRIBUTIONS**

309

310 This research was carried out through joint collaboration of authors AP and AKP. The
311 authors took equal responsibilities to design the study, writing protocols, literature survey,
312 performing the experiments and preparation of manuscript.

313

314 **REFERENCES**

315

- 316 1. Misra TN, Singh RS, Pandey HS, Singh BK, Pandey RP. Constituents of *Asteracantha*
317 *longifolia*. *Fitoterapia*. 2001;72(2):194–96.
- 318 2. Kshirsagar AD, Ingale KG, Vyawahare NS, Thorve VS. *Hygrophila spinosa*: A
319 comprehensive review. *Pharmacogn Rev*. 2010;4(8): 167–171.
- 320 3. Strobel G, Daisy B, Castillo U, Harper J. Natural products from endophytic
321 microorganisms. *J Nat Prod*. 2004;67(2): 257-268
- 322 4. Rosenblueth M, Martínez-Romero E. Bacterial endophytes and their Interactions with
323 hosts. *Mol Plant Microbe Interact*. 2006;19(8): 827-37.
- 324 5. Tan RX, Zou WX. Endophytes: A rich source of functional metabolites. *Nat. Prod. Rep*.
325 2001,18(4); 448-59.
- 326 6. Yu H, Zhang L, Li L, Zheng CA, Guo L, Li W. et al. Recent developments and future
327 prospects of antimicrobial metabolites produced by endophytes. *Microbiol Res*.
328 2010;165(6): 437-49.
- 329 7. Sun JQ, Guo LD, Zang W, Ping WX, Chi DF. Diversity and ecological distribution of
330 endophytic fungi associated with medicinal plants. *Sci China Ser C - Life Sci* 2008;51(8):
331 751-59
- 332 8. Chelius MK, Triplett EW. The diversity of archaea and bacteria in association with the
333 roots of *Zea mays* L. *Microbiol Ecol*. 2001;41(3): 252–63.

- 334 9. Van der Lelie D, Taghavi S, Monchy S, Schwender J, Miller L, Ferrieri R, *et al.* Poplar
335 and its bacterial endophytes: coexistence and harmony. *Crit Rev Plant Sci.* 2009;28(5):
336 346–358
- 337 10. Radu S, Kqueen CY. Preliminary screening of endophytic fungi from medicinal plants in
338 Malaysia for antimicrobial and antitumour activity. *Malaysian J Med Sci.* 2002;9(2): 23-
339 33.
- 340 11. Raviraja NS, Maria GL, Sridhar KR. Antimicrobial evaluation of endophytic fungi
341 inhabiting medicinal plants of the western ghats of India. *Engg in Life Sci.* 2006;6(5):
342 515-20.
- 343 12. Smibert RM, Krieg NR. Phenotypic characterization. In: Gerhardt P, Murray RGE, Wood
344 WA, Krieg NR, editors. *Methods for General and Molecular Bacteriology*, Washington,
345 D.C.: American Society for Microbiology; 1995.
- 346 13. Sneath PHA. *Bergey's Manual of Systematic Bacteriology.* 2nd ed. Baltimore, Williams
347 and Wilkins; 2001.
- 348 14. Pielou EC. *Ecological diversity.* New York: John Wiley and Sons Inc.; 1975.
- 349 15. Bauer AW, Kirby WMM, Sherris JC, Turck M. Antibiotic susceptibility testing by a
350 standardized single disk method. *Am J Clin Pathol.* 1996;45(4): 493–6.
- 351 16. Williston EH, Zia-Walrath P, Youmans GP. Plate methods for testing antibiotic activity of
352 actinomycetes against virulent human type tubercle bacilli. *J Bacteriol.* 1947;54(5): 563 –
353 8.
- 354 17. Pal A, Chattopadhyay A, Paul AK. Diversity and antimicrobial spectrum of endophytic
355 bacterial isolated from *Paederia foetida*. L. *Int J Curr Pharm Res.* 2012;4(3): 123-7.
- 356 18. Maheswari S, Rajagopal K. Biodiversity of endophytic fungi in *Kigelia pinnata* during two
357 different seasons. *Curr. Sci.* 2013;104(4):515-8.
- 358 19. Fisher PJ, Petrini O, Petrini LE, Sutton BC. Fungal endophytes from the leaves and
359 twigs of *Quercus ilex* L. from England, Majorca and Switzerland. *New Phytol.* 1994;
360 127(1):133–7.
- 361 20. Miller KI, Qing C, Sze DMY, Roufogalis BD, Neilan BA. Culturable endophytes of
362 medicinal plants and the genetic basis for their bioactivity. *Microbial. Ecol.*
363 2012;64(2):431-449
- 364 21. Bhoire SJ; Ravichantar N, Loh CY. Screening of endophytic bacteria isolated from leaves
365 of Sambung Nyawa [*Gynura procumbens* (Lour.) Merr.] for cytokinin-like compounds.
366 *Bioinformation* 2010;5(5): 191–7.
- 367 22. Chandrasekhara, Niranjana S; Deepak SA; Amruthesh KA; Shetty NP, Shetty HA.
368 Endophytic bacteria from different plant origin enhance growth and induce downy
369 mildew resistance in pearl millet. *Asian J. Plant Pathol.* 2007;1(1): 1-11.
- 370 23. Zhao K, Penttinen P, Guan T, Xiao J, Chen Q, Xu J, Lindstrom K, Zhang L, Zhang X,
371 Strobel GA. The diversity and anti-microbial activity of endophytic actinomycetes
372 isolated from medicinal plants in Panxi Plateau, China. *Curr. Microbiol.* 2011;62(1): 182-
373 90.
- 374 24. Li J; Zhao GZ; Huang HY; Qin S; Zhu WY; Zhao LX; *et al.* Isolation and characterization
375 of culturable endophytic actinobacteria associated with *Artemisia annua* L. *Antonie van*
376 *Leeuwenhoek* 2012;101(3):515–27.
- 377 25. Stamford T, Stamford N, Coelho L, Araujo JM. Production and characterization of a
378 thermostable glucoamylase from *Streptosporangium* sp. endophyte of maize leaves.
379 *Biores. Technol.* 2002;83(2):105–9.
- 380 26. Carrim A, Barbosa E, Vieira J. Enzymatic activity of endophytic bacterial isolates of
381 *Jacaranda decurrens* Cham. (Carobinha-do-campo). *Braz. arch. biol. technol.* 2006,
382 49(3):353-9.
- 383 27. Panchal H, Ingle SS. Isolation and characterization of endophytes from the root of
384 medicinal plant *Chlorophytum borivillianum* (Safed musli). *J. Adv. Dev. Res.* 2011;
385 2(2):205-9.

- 386 28. Gayathri S, Saravanan D, Radhakrishnan M, Balagurunathan R, Kathiresan K.
387 Bioprospecting potential of fast growing endophytic bacteria from leaves of mangrove
388 and salt-marsh plant species. *Indian J. Biotechnol.* 2010;9(4):397-402.
- 389 29. Arunachalam C, Gayathri P. Studies on bioprospecting of endophytic bacteria from the
390 medicinal plant of *Andrographis paniculata* for their antimicrobial activity and antibiotic
391 susceptibility. *Int J Curr Pharm Res.* 2010;2(4): 63-8.
- 392 30. Li H, Qing C, Zhang Y, Zhao Z. Screening of endophytic fungi with antitumour and
393 antifungal activities from Chinese medicinal plants. *W J Microbiol Biotechnol.* 2005;21(8-
394 9): 1515- 9.
- 395 31. Verma VC, Gond VC; Kumar SK, Mishra A, Kharwar RN, Gange AC. Endophytic
396 actinomycetes from *Azadirachta indica* A. Juss.: Isolation, diversity, and anti-microbial
397 activity. *Microb Ecol* 2009;57(4):749–56.
- 398 32. Sumarah MW, Kesting JR, Sorensen D, Miller JD. Antifungal metabolites from fungal
399 endophytes of *Pinus strobes*. *Phytochemistry* 2011;72(14-15), 1833-7.
- 400 33. Li J, Zhao GZ, Chen HH, Wang HB, Qin S, Zhu WY, *et al.* Antitumour and antimicrobial
401 activities of endophytic streptomycetes from pharmaceutical plants in rainforest. *Lett*
402 *Appl Microbiol.* 2008;47(6): 574–80.
403
404

405