

# Antimicrobial Resistance Evaluation of Organisms Isolated from Liquid Herbal Products Manufactured and Marketed in South Eastern Nigeria.

Ujam NT<sup>1</sup>, Oli AN<sup>1\*</sup>, Ikegbunam MN<sup>1</sup>, Adikwu MU<sup>2</sup>, Esimone CO<sup>1</sup>

(Corresponding author: Oli AN (oli\_an@yahoo.com))

<sup>1</sup>Department of Pharmaceutical Microbiology and Biotechnology, Faculty of Pharmaceutical Sciences, Nnamdi Azikiwe University, Agulu Campus

<sup>2</sup>Department of Pharmaceutics, Faculty of Pharmaceutical Sciences, University of Nigeria, Nsukka

## ABSTRACT

**Objective:** To determine the susceptibility and resistance pattern of bacteria and fungi isolates obtained from herbal anti-infective liquid preparations manufactured and marketed in South-East Nigeria to conventional antibiotics.

**Study Design:** Experimental

**Place and Duration of the study:** Pharmaceutical Microbiology and biotechnology Laboratory, Faculty of Pharmaceutical Sciences, Nnamdi Azikiwe University, Agulu Campus between October 2011 and March 2012

**Methodology:** Isolation and characterization of contaminating microorganisms were carried out using standard procedures. A total of forty-nine (49) bacteria and forty (40) fungi isolated from the herbal products were examined for susceptibility to conventional antibiotics using the disc diffusion method. The bacterial isolates were tested against ciprofloxacin, ofloxacin, amoxicillin-clavulanic acid, gentamicin, cefotaxime, ceftazidime, ceftriazone, sulphamethoxazole, tetracycline and ampicillin were employed while fungi isolates were tested against five common antifungal-griseofulvin, nystatin, ketoconazole, fluconazole and clotrimazole. The Multiple Antibiotic Resistance Index (MARI) of each of the isolated bacteria was obtained following the standard method.

**Result:** The antimicrobial susceptibility-resistance profile of the bacteria isolates revealed that most of the bacteria were sensitive to ciprofloxacin, ofloxacin, gentamicin, and ceftriaxone, On the other hand, a good number of the isolates demonstrated high level of resistance to common antibiotics like Ampicillin, amoxicillin-clavulanic acid, trimethoprim-sulphamethoxazole, and moderate level of resistance to Tetracycline, and some of the third generation cephalosporins - ceftazidime and cefotaxime. Multiple Antibiotic Resistance Index (MARI) evaluation revealed that most of the isolates were resistance to more than fifty percent (50%) of the number of antibiotics used. The fungal isolates were susceptible to nystatin, ketoconazole and clotrimazole, resistance to fluconazole and high resistance recorded against griseofulvin.

**Conclusion:** The results of this study revealed that the herbal medications can serve as a trail of spread of antibiotic-resistance genes.

**Keywords:** {Susceptibility, antibiotic resistance, herbal anti-infectives}

## 1. INTRODUCTION

The use of herbal medicine has always been part of human culture, as some plants possess important therapeutic properties, which can be used to cure human and other animal diseases [1]. Herbal medicine is becoming increasingly

\* Tel.: +234-8056306927.

E-mail address: oli\_an@yahoo.com.

popular in both developing and developed countries [2]. A World Health Organization survey indicates that about 70–80% of the world population, particularly in developing nations; rely on non-conventional medicines mainly of herbal sources in their primary health care [3]. Medicinal plant materials normally carry a large number of microbes originating from the soil. Microorganisms of various kinds are normally adhered to leaves, stems, flowers, roots and seeds. Additional contaminants may also be introduced during harvesting, handling and production of various herbal remedies since no conscious efforts are made to decontaminate the herbs other than by washing them. [4]. Herbal medicines are therefore vulnerable to attack by microorganisms and as such are disposed to spoilage. Accordingly, gross microbial contamination of herbal medicinal products commonly consumed in Nigeria has been severally demonstrated [5, 6,7]. The presence of antibiotic resistant microbial isolates in the Herbal Medicinal Products (HMPs) could lead to transfer of antibiotic resistance traits to hitherto sensitive gut or oral micro flora of consumers [8].

The emergence of multiple drug resistant bacteria (MDR) has become a major cause of failure of the treatment of infectious disease [9]. As a result, society is facing one of the most serious public health dilemmas over the emergence of infectious bacteria displaying resistance to many and in some cases, effective antibiotic [10] much like the situation in human medicine. Bacteria and fungi resistance to antimicrobial drugs has continued to grow in the last decades [11]. The increased prevalence of their resistant is due to extensive use and misuse of antimicrobials. This has rendered the current available antimicrobial agents insufficient to control microbial infections and create major public health problem.

Resistant bacteria strains may develop almost anywhere particularly in a pressurized environment containing previously non-resistant bacteria strains as contaminants. One of such environments can be created by widespread use of HMP. HMPs have been previously implicated as a pool for such contaminations [12,13]. It is of utmost importance to both monitor and ascertain the microbial purity of HMPs given the huge medical and economic implications of any such microbial contamination especially with multiple drug resistant strains. Such surveillance will both help to identify microbial contamination of herbal products and slow down and prevent the emergence of drug-resistant strains. The present study evaluated the presence of contaminating organisms and the susceptibility-resistance pattern of the isolated organisms.

## 2. EXPERIMENTAL DETAILS

### 2.1 MATERIALS: HERBAL SAMPLES

A total of twenty liquid herbal anti-infectives were purchased randomly from different shops and herbal outlets located within the five states that make up the south-east, Nigeria and were used in this study. The samples which were within their shelf lives and were kept at room temperature (as indicated by their manufacturers) were used within two weeks of collection.

### 2.2 METHODS

#### 2.2.1 Isolation and Identification of Microbial Contaminants in the Herbal

The herbal anti-infectives were serially diluted and plated on nutrient agar and sabouroud dextrose agar plates in triplicate and incubated at 37°C for 18-24 hours and 20°C- 27 °C for 72-168hours for bacteria and fungi respectively. The resultant colonies were further purified, isolated and characterized using standard methods [14].

#### 2.2.2 Characterization of Microorganisms Isolated From the Herbal Preparations

The bacteria isolates were characterized using the morphological appearance (macroscopy) of their colonies, their Gram stain reaction and confirmatory biochemical tests. The fungi isolates were also identified on the basis the morphological characteristics (macroscopy) of their colonies, microscopy, staining with ordinary stain and the appearance of their mycelia [15].

#### 2.2.3 Antibiotics Susceptibility Testing

The susceptibility tests were performed following the method M2-A6 disc diffusion method recommended by the National Committee for Clinical Laboratory Standards [16] using Mueller Hinton and Sabouraud Dextrose Agar. The bacterial isolates from the samples were reactivated by sub-culturing from agar slant onto nutrient agar plate and was incubated for 18-24 hours. The inoculum was standardized by transferring three distinct and separate colonies of the pure culture of the test organism using sterile wire loop into 3mls of sterile nutrient broth. The suspension was incubated for 3 hours at 37°C to allow for the growth of test organism till the density was equivalent to the turbidity of 0.5 McFarland. The standardized inocula were swabbed onto Mueller-Hinton agar and Sabouraud Dextrose Agar plate and the discs were placed on the inoculated plates and pressed firmly onto the agar plate for complete contact. The bacterial strains were tested against the following discs: ofloxacin (OFX, 5µg) ; ciprofloxacin (CIP,5µg) ; amoxicillin/clavulanic acid (AMC,20/10µg) ; gentamicin,(GN ,10µg) ; ceftazidime (CAZ,30µg) ; cefotaxime (CTX,30µg) ; trimethoprim-sulfamethoxazole (SXT,1.25/23.75µg); Ampicillin (AMP,10µg) ; tetracycline (TE, 30µg); ceftriaxone ( CRO, 30µg).The fungal strains were tested against the following discs: nystatin (N,20µg) ; clotrimazole (C,20µg) ; griseofulvin (G,20µg) ; ketoconazole (K,20µg) and fluconazole (F,20µg). The Plates were inverted and left on the work table for 30 minutes to allow for pre-diffusion of antibiotics into the agar. The plates were incubated at 37°C for 18-24 hours and at 25°C 24-48hours for bacteria and fungi respectively. The susceptibility of each isolate to each antibiotic was shown by a clear zone of growth inhibition and this was measured using a meter rule in millimeters and the diameter of the zones of inhibition was then interpreted using standard chart [17].

## 2.2.4 Determination of Multiple Antibiotics Resistance Index (MARI)

85 The Multiple Antibiotics Resistance Index (MARI) of ten antibiotics (ofloxacin, ciprofloxacin, gentamicin, amoxycillin-  
86 clavulanic acid, sulphamethoxazole-trimethoprim, ceftriazone, ceftazidime, cefotazime, tetracycline and ampicillin) were  
87 determined using the formula,  $MARI = a/b$

88 Where; a = the aggregate resistance of antibiotics to all isolates and b = the total number of antibiotics that was used

## 89 3. RESULTS AND DISCUSSION

### 90 3.1 RESULTS

91

92

93

Table 1: Microorganisms Isolated from the Herbal anti-infective Products.

| Samples Code | Shelf life  | Contents  | Therapeutic Claims   | Identity Of Bacteria Isolated  | Identity Of Fungi Isolated   |
|--------------|-------------|---|--|--|--|
| 1            | (48 Months) | <i>Carica papaya</i> ,<br><i>Magnifera indica</i> ,<br><i>Newbouldia leavis</i> ,<br><i>Azadricha indica</i> ,<br><i>Jaminum officionili</i> ,<br><i>Aloe barbedensis</i> ,<br><i>Ginseng</i> , Treated water 60cl. | Antibacterial, Antimalarial, Ant rheumatic, infertility, Antiviral.                              | a) <i>Staphylococcus aureus</i><br>b) <i>Proteus spp</i>   | a) <i>Microsporium spp.</i><br>b) <i>Aspergillus spp</i><br>c) <i>Nigrospora spp</i>     |
| 2            | (36 Months) | 38 African Roots, Herbs, Fruits, Barks plus ginseng, Aloe vera and Garlic.  | Antibacterial, Antirheumatic, Antifungal and Antiviral.  | a) <i>Escherichia coli</i><br>b) <i>Staphylococcus aureus</i><br>c) <i>Staphylococcus epidermidis</i><br>d) <i>Pseudomonas aeruginosa</i><br>e) <i>Bacillus spp</i><br>f) <i>Proteus spp</i> | a) <i>Candida tropicalis</i><br>b) <i>Microsporium canis</i>                             |
| 3            | (30 Months) | 60% herbs, 25% flower, 10% leaves, 5% roots.  | Antibacterial, Antirheumatic, Antifungal, Earlier Menopause, Painful and irregular menstruation. | a) <i>Staphylococcus aureus</i><br>b) <i>Bacillus subtilis</i><br>c) <i>Bacillus cereus</i><br>d) 2 <i>Salmonella spp</i>  | a) <i>Candida albicans</i><br>b) <i>Candida tropicalis</i><br>c) <i>Trichosporon spp</i> |
| 4            | (48 Months) | Aloe vera plus 31 roots and herbs ,fruits and barks   | Antibacterial, Antifungal.   | a) <i>Escherichia coli</i><br>b) <i>Staphylococcus aureus</i><br>c) <i>Streptococcus spp</i><br>d) <i>Bacillus spp</i>   | a) <i>Coccidioides immitis</i><br>b) <i>Microsporium audounii</i>                        |
| 5            | (36 Months) | Water, herbs, root and fruits.  | Antibacterials, Antimalarial, Antiparasitic, Internal heat, pile, and reduces sugar.             | a) <i>Staphylococcus epidemidis</i><br>b) <i>Streptococcus spp</i><br>c) <i>Yersinia spp</i>   | -  |
| 6            | (24 Months) | -   | Antibacterial, Treatment of all form of eye infections.  | a) <i>Escherichia.coli</i><br>b) <i>Bacillus spp</i><br>c) <i>Proteus spp</i>  | a) <i>C.topicalis</i><br>b) <i>Microsporium audounii</i><br>c) <i>Aspergillus niger</i>  |
| 7            | (12 Months) | <i>Nauclea diiderchi</i> 10%, <i>Hippocrates pallens</i> 20%, <i>Alluim sativum</i> 12.5%, <i>Cochios permum planchoni</i> 5.5%, <i>Uvaria chame</i> 5%, <i>Punica granatum</i> 47%.                                | Antibacterial, Antimalarial.   | a) <i>Escherichia coli</i><br>b) <i>Staphylococcus spp</i><br>c) <i>Pseudomonas</i><br>d) <i>Bacillus spp</i>  | a) <i>Blastomyces</i><br>b) <i>Microporium canis</i>                                     |

|    |             |  |  |   |   |
|----|-------------|--|--|---|---|
| 8  | (36 Months) | Aloe vera 40%, Olong tea 20%, Flower and roots 40%, Saracin.   | Antibacterial.   | a) <i>Staphylococcus spp</i><br>b) <i>Salmonella spp</i>                            | a) <i>Blastomyces</i>   |
| 9  | (36 Months) | Aloe Vera  | Antibacterial Anti-malarial, HBP, Cough<br>Antirheumatism, e.t.c.                      | a) <i>Staphylococcus aureus</i>   | a) <i>Blastomyces spp</i><br>b) <i>Cryptococcus neoformans</i><br>c) <i>Histoplasma</i> |
| 10 | (36 Months) | Aloe Vera, Flowers, Fruits seed barks.   | Antibacterial, Hypertension, Antiviral, fibroid, stroke.                               | a) <i>Pseudomonas aeruginosa</i>  | a) <i>Penicillum spp</i><br>b) <i>Aspergillus spp</i>                                   |
| 11 | (24 Months) | Honey (natural), Lime juice, Zingiber officillinar, Herbal seeds and roots.  | Antibacterial and Asthmatic cough.   | a) <i>Escherichia coli</i><br>b) <i>Staphylococcus</i>                              | a) <i>Candida spp</i>   |
| 12 | (31 Months) | -  | Anti-bacterial<br>Antiviral, Diabetes, Reduces cholesterol.                            | a) <i>Staphylococcus aureus</i><br>b) <i>Streptocoloccus spp</i>                    | a) <i>C.albicans</i><br>b) <i>Blastomyces</i>   |
| 13 | (24 Months) | 25 different types of roots, herbs, seeds and flowers.   | Anti-bacterial, Anti-malaria, Antirheumatic.<br>Antifungal                             | a) <i>Escherichia coli</i><br>b) <i>Corynebacteruim diphtheria</i>                  | a) <i>Aspergillus spp</i><br>b) <i>Nigrospora spp</i><br>c) <i>C.tropicalis</i>         |
| 14 | (60 Months) | Herbs, water, root and fruits.   | Antibacterial, Antiviral, Antirheumatic, Antifungal, Antiparatic, internal heat, pile. | a) <i>Escherichia coli</i><br>b) <i>Streptococcus spp</i><br>c) <i>Yersinia spp</i> | a) <i>C.albicans</i><br>b) <i>Aspergillus spp</i>                                       |
| 15 | (29 Months) | Magnifera indica, Carica papaya leaves, Psiduim guajava, Breadfruit bark, Masularia acuminata roots, Citrus lemon leaves, Zingiber Officinale roots, Cymbopogon spp. | Antibacterial, Antimalarial, Antirheumatic Antiviral.                                  | a) <i>Escherichia coli</i>  | a) <i>Microsporium spp</i><br>b) <i>Coccidioides spp</i><br>c) <i>Aspergillus spp.</i>  |
| 16 | (48 Months) | Awapa bark, white lotus, Golden seal, Mahogany, Ukor root, Aloe barbadars, Mistletoe, Osisika Aguru, Uda roots, Uvuru ilu, Lemon grass.                              | Antibacterial, Antirheumatic and Arthritis, Veneral diseases.                          | a) <i>Escherichia coli</i><br>b) <i>Pseudomonas aeruginosa</i>                      | a) <i>C.albicans</i><br>b) <i>Mucor spp</i>   |
| 17 | (15 Months) | Aloe Vera, Cadeperi salt, Lime   | Antibacterial, Treatment and prevention of toothache.                                  | a) <i>Proteus spp</i>   | a) <i>Aspergillus spp</i>   |
| 18 | (12 Months) | Lymbopogon citrates, Carica papaya leaves, Magnifera indica, bark, Treculia Africana, Citrus, Limonia, Psiduim guajava, Zingibar officinale root, Alluim sativum.    | Antibacterial, Antirheumatism reduces sugar and cholesterol.                           | a) <i>Escherichia coli</i>  | a) <i>Mucor spp</i>   |
| 19 | (24 Months) | Natural roots and barks.   | Antibacterial, Antiviral, Purifies   | a) <i>Bacillus spp</i>  | a) <i>Yeast/Blastomyces</i>   |

|    |                |  |  |  |   |
|----|----------------|--|--|--|---|
|    |                |  | blood, Detoxifies<br>toxins, Builds<br>immune system,<br>Stops dizziness,<br>weakness.                           |  | b) <i>Aspergillus spp</i><br>c) <i>Microsporium spp</i> |
| 20 | (36<br>Months) | Nuclealatifolia, Allium<br>sativum, Aloe Vera<br>bitter, Chick weed,<br>Preclina nitida,<br>Hibiscus sabdrifa,<br>Aqua, Ethanol. | Antibacterial,<br>Antiparasitic, ulcer,<br>constipation,<br>fibroid, internal<br>heat heart burn<br>and diabetes | a) <i>Staphylococcus<br/>epidermidis</i> | a) <i>Aspergillus spp</i>                               |

Table 2: Percentage of microbial isolates from the Herbal anti-infective Products

| BACTERIA ISOLATES   | % Occurrence | FUNGI ISOLATES          | % Occurrence |
|---------------------|--------------|-------------------------|--------------|
| <i>E.coli</i>       | 20.4         | <i>Aspergillus spp</i>  | 22.5         |
| <i>S.aureus</i>     | 24.5         | <i>Microsporium spp</i> | 17.5         |
| <i>P.aeruginosa</i> | 8.2          | <i>Candida spp</i>      | 22.5         |
| <i>Strep.spp</i>    | 10.2         | <i>Trichosporon spp</i> | 2.5          |
| <i>Bacillus</i>     | 16.3         | <i>Coccidiodes spp</i>  | 5.0          |
| <i>Salmonella</i>   | 6.1          | <i>Blastomyces spp</i>  | 12.5         |
| <i>Proteus spp</i>  | 8.2          | <i>Cryptococcus spp</i> | 2.5          |
| <i>Yersinia spp</i> | 4.1          | <i>Histoplasma spp</i>  | 2.5          |
| <i>C.diphtheria</i> | 2.0          | <i>Penicillium spp</i>  | 2.5          |
|                     |              | <i>Nigrospora spp</i>   | 5.0          |
|                     |              | <i>Mucor spp</i>        | 5.0          |

A total of 89 microbial strains (49 bacterial and 40 fungal strains) were isolated from the herbal preparations. The identified microbial isolates consists of nine (9) bacterial genera and eleven (11) fungal genera which include *Staphylococcus*, *E. coli*, *Bacillus*, *Streptococcus*, *Pseudomonas*, *Proteus*, *Salmonella*, *Yersinia*, *Corynebacterium diphtheria* and *Aspergillus*, *Candida*, *Microsporium*, *Trichosporon*, *Coccidiodes*, *Blastomyces*, *Cryptococcus*, *Histoplasma*, *Penicillium*, *Nigrospora*, *Mucor* respectively. The most frequently isolated bacteria and fungi specie were *Staphylococcus spp* (24.5%) and *Aspergillus spp/Candida spp* (22.5%) respectively. The least frequently isolated bacteria specie was *Corynebacterium diphtheria* (2.0%) and that of fungi were *Trichosporon spp*, *Cryptococcus spp*, *Histoplasma* and *Penicillium spp* (2.5%). See Tables 1 and 2 above.

Table 3: The Antibiotic Susceptibility-Resistance Profile of the isolated bacteria

| BACTERIA ISOLATES | Drugs and Strength (µg)   | OFX-5  | CIP-5    | SXT-1.25/23.75 | AMC-20/10 | GN-10    | CTX-30   | CAZ-30  | TE-30  | AMP-10   | CRO-30   |        |
|-------------------|---------------------------|--------|----------|----------------|-----------|----------|----------|---------|--------|----------|----------|--------|
|                   |                           | N (%)  | N (%)    | N (%)          | N (%)     | N (%)    | N (%)    | N (%)   | N (%)  | N (%)    | N (%)    |        |
| <i>E. coli</i>    | S                         | 8 (80) | 10 (100) | 0 (0)          | 0 (0)     | 9 (90)   | 0 (0)    | 3 (30)  | 1 (10) | 0 (0)    | 5 (50)   |        |
|                   | I                         | 1 (10) | 0 (0)    | 0 (0)          | 0 (0)     | 0 (0)    | 0 (0)    | 0 (0)   | 0 (0)  | 0 (0)    | 2 (20)   |        |
|                   | R                         | 1 (10) | 0 (0)    | 10 (100)       | 10 (100)  | 1 (10)   | 10 (100) | 7 (70)  | 9 (90) | 10 (100) | 3 (30)   |        |
|                   | <i>P. aeruginosa</i>      | S      | 3 (75)   | 3 (75)         | 0 (0)     | 0 (0)    | 3 (75)   | 0 (0)   | 2 (50) | 0 (0)    | 0 (0)    | 2 (50) |
|                   |                           | I      | 0 (0)    | 0 (0)          | 0 (0)     | 0 (0)    | 0 (0)    | 3 (75)  | 0 (0)  | 0 (0)    | 0 (0)    | 1 (25) |
|                   |                           | R      | 1 (25)   | 1 (25)         | 4 (100)   | 4 (100)  | 1 (25)   | 1 (25)  | 2 (50) | 4 (100)  | 4 (100)  | 1 (25) |
|                   | <i>Staphylococcus spp</i> | S      | 8 (67)   | 10 (83)        | 0 (0)     | 0 (0)    | 10 (83)  | 0 (0)   | 2 (17) | 2 (17)   | 0 (0)    | 7 (58) |
|                   |                           | I      | 4 (33)   | 2 (17)         | 0 (0)     | 0 (0)    | 0 (0)    | 2 (17)  | 1 (8)  | 2 (17)   | 0 (0)    | 4 (33) |
|                   |                           | R      | 0 (0)    | 0 (0)          | 12 (100)  | 12 (100) | 2 (17)   | 10 (83) | 9 (75) | 8 (67)   | 12 (100) | 1 (8)  |

|                          |          |            |            |         |            |            |            |            |            |            |           |
|--------------------------|----------|------------|------------|---------|------------|------------|------------|------------|------------|------------|-----------|
| <b>Salmonella spp</b>    | <b>S</b> | 1 (33)     | 1 (33)     | 0 (0)   | 0 (0)      | 3<br>(100) | 2(67)      | 2 (67)     | 1 (33)     | 0 (0)      | 1(33)     |
|                          | <b>I</b> | 1<br>(33)  | 2<br>(67)  | 0 (0)   | 0 (0)      | 0 (0)      | 0(0)       | 0 (0)      | 0 (0)      | 0 (0)      | 1(33)     |
|                          | <b>R</b> | 1 (33)     | 0 (0)      | 3 (100) | 3<br>(100) | 0 (0)      | 1 (33)     | 1 (33)     | 2 (67)     | 3<br>(100) | 1(33)     |
| <b>Streptococcus spp</b> | <b>S</b> | 3 (60)     | 4 (80)     | 0 (0)   | 0 (0)      | 4<br>(80)  | 2 (40)     | 2 (40)     | 1 (20)     | 0 (0)      | 1(20)     |
|                          | <b>I</b> | 1 (20)     | 1 (20)     | 0 (0)   | 1 (20)     | 1<br>(20)  | 0 (0)      | 1 (20)     | 1(20)      | 0 (0)      | 0(0)      |
|                          | <b>R</b> | 1 (20)     | 0 (0)      | 5 (100) | 4 (80)     | 0 (0)      | 3 (60)     | 2 (40)     | 3 (60)     | 5<br>(100) | 4<br>(80) |
| <b>Bacillus spp.</b>     | <b>S</b> | 3 (38)     | 5 (63)     | 0 (0)   | 0(0)       | 6<br>(75)  | 0 (0)      | 1 (13)     | 4 (50)     | 0 (0)      | 5(63)     |
|                          | <b>I</b> | 5<br>(63)  | 3<br>(38)  | 0 (0)   | 1(13)      | 0 (0)      | 1 (13)     | 1 (13)     | 1<br>(13)  | 1 (13)     | 1(13)     |
|                          | <b>R</b> | 0 (0)      | 0 (0)      | 8 (100) | 7(88)      | 2 (25)     | 7 (88)     | 6 (75)     | 3 (38)     | 7 (88)     | 2(25)     |
| <b>Proteus spp</b>       | <b>S</b> | 2<br>(50)  | 1<br>(25)  | 0 (0)   | 0 (0)      | 2<br>(50)  | 1 (25)     | 2<br>(50)  | 2<br>(50)  | 0 (0)      | 1<br>(25) |
|                          | <b>I</b> | 2 (50)     | 3<br>(75)  | 0 (0)   | 0 (0)      | 0 (0)      | 0 (0)      | 1 (25)     | 0 (0)      | 0 (0)      | 2<br>(50) |
|                          | <b>R</b> | 0 (0)      | 0 (0)      | 4 (100) | 4<br>(100) | 2 (50)     | 3 (75)     | 1 (25)     | 2 (50)     | 4<br>(100) | 1(25)     |
| <b>Yersinia spp</b>      | <b>S</b> | 0 (0)      | 1 (50)     | 0 (0)   | 0 (0)      | 2<br>(100) | 0 (0)      | 0 (0)      | 0 (0)      | 0 (0)      | 0 (0)     |
|                          | <b>I</b> | 2<br>(100) | 1 (50)     | 0 (0)   | 0 (0)      | 0 (0)      | 0 (0)      | 0 (0)      | 0 (0)      | 0 (0)      | 1 (50)    |
|                          | <b>R</b> | 0 (0)      | 0 (0)      | 2 (100) | 1<br>(100) | 0 (0)      | 2<br>(100) | 1 (50)     | 2<br>(100) | 2<br>(100) | 1 (50)    |
| <b>C. diphtheria</b>     | <b>S</b> | 0 (0)      | 1<br>(100) | 0 (0)   | 0 (0)      | 1<br>(100) | 0 (0)      | 1<br>(100) | 0 (0)      | 0 (0)      | 0 (0)     |
|                          | <b>I</b> | 1<br>(100) | 0 (0)      | 0 (0)   | 0 (0)      | 0 (0)      | 0 (0)      | 0 (0)      | 0 (0)      | 0 (0)      | 1(100)    |
|                          | <b>R</b> | 0 (0)      | 0 (0)      | 1 (100) | 1<br>(100) | 0 (0)      | 1<br>(100) | 0 (0)      | 1<br>(100) | 1<br>(100) | 0 (0)     |

106  
107  
108  
109  
110  
111  
112

**Key:** **S** = Sensitive, **I** = intermediate, **R** - Resistance, **N** = number of organisms, OFX= ofloxacin, CIP = ciprofloxacin, CAZ = ceftazidime, TE = tetracycline, AMP = ampicillin, SXT = trimethoprim-sulfamethoxazole, GN = gentamicin, CTX = cefotaxime, CRO = ceftriaxone, AMC = amoxicillin-clavulanic acid.

Table 3 above shows the antibiogram, of all the bacterial strains isolated from the Herbal products - a representation of the bacteria that are Susceptible, Intermediate or Resistant to the different antibiotics using the NCCLS break points [17].

Table 4: Antibiogram of Fungi Isolated from the Herbal Anti-Infectives

| Samples Code | Isolates                     | Inhibition Zone Diameter (IZD) in millimeter (mm) |          |              |              |             |
|--------------|------------------------------|---|----------|--------------|--------------|-------------|
|              |                              | Griseofulvin                                      | Nystatin | Ketoconazole | Clotrimazole | Fluconazole |
| 1            | <i>Microsporium spp.</i>     | 0   | 25       | 0            | 0            | 0           |
|              | <i>Aspergillus spp</i>       | 0   | 32       | 10           | 12           | 0           |
|              | <i>Nigrospora spp</i>        | 0   | 30       | 20           | 14           | 12          |
| 2            | <i>Candida tropicalis</i>    | 0   | 26       | 23           | 7            | 15          |
|              | <i>Microsporium canis</i>    | 0   | 25       | 0            | 0            | 0           |
| 3            | <i>Candida albicans</i>      | 0   | 22       | 12           | 10           | 7           |
|              | <i>Candida tropicalis</i>    | 0   | 23       | 11           | 10           | 8           |
|              | <i>Trichosporon spp</i>      | 7   | 30       | 32           | 18           | 8           |
| 4            | <i>Coccidioides immitis</i>  | 0   | 30       | 0            | 0            | 0           |
|              | <i>Microsporium audounii</i> | 0   | 29       | 0            | 0            | 0           |
| 6            | <i>C.tropicalis</i>          | 0   | 27       | 12           | 11           | 8           |
|              | <i>Microsporium audounii</i> | 0   | 8        | 0            | 0            | 0           |
|              | <i>Aspergillus niger</i>     | 0   | 29       | 0            | 7            | 0           |
| 7            | <i>Blastomyces</i>           | 0   | 32       | 17           | 18           | 0           |
|              | <i>Microporuim canis</i>     | 0   | 30       | 0            | 0            | 0           |

|    |                                |   |    |    |    |    |
|----|--------------------------------|---|----|----|----|----|
| 8  | <i>Blastomyces</i>             | 0 | 31 | 18 | 18 | 0  |
| 9  | <i>Blastomyces spp</i>         | 0 | 29 | 16 | 19 | 0  |
|    | <i>Cryptococcus neoformans</i> | 0 | 0  | 0  | 0  | 0  |
|    | <i>Histoplasma</i>             | 0 | 30 | 29 | 12 | 15 |
| 10 | <i>Penicillium spp</i>         | 0 | 31 | 15 | 7  | 0  |
|    | <i>Aspergillus spp</i>         | 0 | 23 | 0  | 12 | 0  |
| 11 | <i>Candida spp</i>             | 0 | 22 | 12 | 10 | 7  |
| 12 | <i>Candida albicans</i>        | 0 | 20 | 14 | 12 | 9  |
|    | <i>Blastomyces</i>             | 0 | 30 | 15 | 19 | 0  |
| 13 | <i>Aspergillus spp</i>         | 0 | 32 | 17 | 8  | 0  |
|    | <i>Nigrospora spp</i>          | 0 | 28 | 14 | 15 | 0  |
|    | <i>Candida tropicalis</i>      | 0 | 35 | 25 | 20 | 0  |
| 14 | <i>Candida albicans</i>        | 0 | 22 | 13 | 10 | 10 |
|    | <i>Aspergillus spp</i>         | 0 | 26 | 0  | 7  | 0  |
| 15 | <i>Microsporum spp</i>         | 0 | 25 | 0  | 0  | 0  |
|    | <i>Coccidioides spp</i>        | 0 | 30 | 0  | 0  | 0  |
|    | <i>Aspergillus spp.</i>        | 0 | 31 | 10 | 0  | 0  |
| 16 | <i>C.albicans</i>              | 0 | 24 | 23 | 8  | 0  |
|    | <i>Mucor spp</i>               | 0 | 26 | 0  | 0  | 0  |
| 17 | <i>Aspergillus spp</i>         | 0 | 31 | 10 | 0  | 0  |
| 18 | <i>Mucor spp</i>               | 0 | 25 | 0  | 0  | 0  |
| 19 | Yeast/ <i>Blastomyces</i>      | 0 | 28 | 18 | 16 | 0  |
|    | <i>Aspergillus spp</i>         | 0 | 29 | 0  | 9  | 0  |
|    | <i>Microsporum spp</i>         | 0 | 0  | 0  | 0  | 0  |
| 20 | <i>Aspergillus spp</i>         | 0 | 36 | 0  | 0  | 0  |

Table 5: Multiple Antibiotics Resistance Index (MARI) of the Isolated Bacteria.

| Grouping | Isolates and Samples Code   | Multiple Antibiotics Resistance Index (MARI) |
|----------|---|--|
| Group A  | Pa16  | 1  |
| Group B  | S3  | 0.8  |
| Group C  | E4,11,14,15,Sa1,P2  | 0.7  |
| Group D  | E7,15,16,13,18,Sa2,Sa5,Sa7,Sa20,St12,St4,Ba7,Ba19b,Ba3a,Ba6,Ba3b,P17,Y5 | 0.6  |
| Group E  | E2,6,Sa3,Sa4,Sa8,Sa11,Sa9,Sa2,Pa10,St5,St14,Ba9a,P6,Y14,C13             | 0.5  |
| Group F  | Sa12,Pa2,Pa7,St14,Ba2,S8,P1   | 0.4  |

Notes for Table 5: E = *E.coli*, Sa = *Staphylococcus spp*, Pa = *Pseudomonas aeruginosa*, St = *Streptococcus spp*, Ba = *Bacillus spp*, S = *Salmonella*, P = *Proteus spp*, Y = *Yesinia spp*, C = *Corynebacterium spp*. The numbers attached represent the product numbers

### 3.2 DISCUSSION

Antimicrobial susceptibility testing of the isolated microorganisms was carried out to evaluate the activity of conventional antibiotics against the isolated bacteria and fungi strains. The bacteria contaminants isolated from these herbal preparations showed wide resistance to penicillins, especially ampicillin, augmentin (amoxicillin-clavulanic acid combination) and cloxacillin, suggesting that they could be producers of penicillinases. The resistance to trimethoprim-sulphamethoxazole (co-trimoxazole) by all the isolates especially the Gram-negative isolates calls for attention. The findings of this study agree with an earlier work [12]. *Staphylococci* strains were the most frequently isolated bacteria species and it probably originated from handlers, as its habitat is human skin. *Staphylococcus* showed wide resistance to penicillins suggesting possibly that they are producers of penicillinase enzymes. Resistance to trimethoprim by *S. aureus* and *S. epidermidis* has been reported with increasing frequency [18,19, 20]. It seems probable that *S. epidermidis* serves as a reservoir for resistance, which can be transferred to *S. aureus*. Also, inter-generic transfer of resistance among different genera of Gram-positive cocci and between *Bacillus* species and *Staphylococci* and *Streptococci* has been reported [20, 21]. *Escherichia coli* were the second most frequently isolated species in these medications which is an intestinal bacterium and an indicator of faecal contaminant. Presence of *Escherichia coli* in the sample indicates poor hygiene practices and lack of adequate handling of the products. According to the European pharmacopoeia 2007 [22], no *Salmonella spp* or *Escherichia coli* strain should be present in oral medicines. The presence of *E. coli* in herbal drugs had been reported by another researcher [23]. The *Escherichia coli* isolates showed a wide resistance to ampicillin, ceftazidime, sulphamethoxazole-trimethoprim, amoxicillin-clavulanic acid, and tetracycline. *Bacillus spp.* were the third

136 most frequently found in these herbal medicaments because they are widely distributed in the soil, dust, air and because  
137 they are resistant to environmental destructive factors [20, 24]. A number of reports have described serious human  
138 infections caused by members of the genus *Bacillus* even though they have been regarded as non-pathogenic [25,26,27].  
139 All the strains of *Pseudomonas* isolated were resistant to  $\beta$ -lactam antibiotics; Inducible  $\beta$ -lactamase activity is a general  
140 property of *Pseudomonas cepacia* [28]. Gram negative rods usually have wide resistance against antimicrobial agents  
141 [20] (Esimone *et al.*, 2007a). *Streptococcus spp* showed high resistance to sulphamethoxazole-trimethoprim and  
142 ampicillin. *Salmonella spp*, were resistant to sulphamethoxazole-trimethoprim, amoxicillin clavulanic acid, and ampicillin.  
143 *Proteus spp*, *Yersinia spp* and *Corynebacterium diphtheria* showed wide resistance to sulphamethoxazole-trimethoprim,  
144 amoxicillin clavulanic acid, ceftazidime and ampicillin (Table 3). On the other hand, the bacterial isolates were susceptible  
145 to some groups of the antibiotics (ofloxacin, ciprofloxacin, gentamicin and ceftriaxone).

146 Multiple Antibiotic Resistance Index (MAR) evaluation (Table 5) revealed that species of *Escherichia coli* showed high  
147 level of multiple antibiotic resistances to the panel of antibiotics used in this study. The MAR value ranged from 0-5 -0.7,  
148 with three (30%) resistant to seven antibiotic out of the ten used, six (60%) resistant to six of the antibiotics used and two  
149 (20%) resistant to five. *Staphylococcus spp* have MAR values ranging from 0.4-0.7, with one (8.3%) resistance to seven  
150 antibiotic, four (33.3%) resistant to six antibiotics, six (50%) resistant to five antibiotics and one (8.3%) resistance to four  
151 antibiotics. The MAR result of *Bacillus spp* ranged from 0.4 – 0.6, with five (62.5%) resistant to six antibiotics, One  
152 (12.5%) being resistance to five, four and three antibiotics each. *Proteus spp* MAR value is from 0.4 - 0.7, with one (25%)  
153 each of the four isolates resistant to seven, six, five and four respectively. *Pseudomonas spp* had MAR value ranging  
154 from 0.4 - 1.0, one (25%) showed high resistance index, being resistance to ten of the antibiotics used in this study, one  
155 (25%) resistant to five antibiotics and two (50%) resistance to four antibiotics. The three species of *Streptococcus* isolated  
156 showed MAR values from 0.3-0.8, that is, one (33.3%) resistant to eight antibiotics, one (33.3%) to four antibiotics and  
157 one (33.3%) to three out of the ten antibiotics. We had two isolates of *Yersinia spp* and the MAR values are 0.5 and 0.6.  
158 Lastly, *Corynebacterium diphtheria* isolate is resistance to five antibiotics out of the ten antibiotics used in this study.  
159 **Bacteria with high MAR index originate from the environment where antibiotics are over used** [29].

160 Fungal infections are becoming an increasing cause of morbidity and mortality especially among immunocompromised  
161 patients. With the increased incidence of systemic fungal infections and the growing number of antifungal agents,  
162 laboratory methods to guide and select antifungal therapy have gained greater attention. However, determining antifungal  
163 susceptibilities of filamentous fungi is fraught with difficulties associated with slow growth of filamentous forms and  
164 subjectivity of interpreting visual endpoints [30]. In the present study, antifungal susceptibility testing of 40 fungi isolates  
165 was observed against five common antifungal agents (griseofulvin, nystatin, ketoconazole, clotrimazole, and fluconazole)  
166 using disc diffusion method, presence of inhibition zone was considered as sensitive while absence of inhibition zone was  
167 recorded as resistance. The fungi isolates were very sensitive to nystatin, ketoconazole and clotrimazole. **Most of the**  
168 **fungi isolates were resistance to fluconazole while all are resistant to Griseofulvin** (Table 5).

169 The importance of surveying resistant environmental strains is that under favourable situations, they may transfer their  
170 resistance plasmids to pathogens [31,32]. If such organisms are present in medicaments, such as herbal medicinal  
171 products they could behave as opportunist pathogens and initiate an infection, particularly in immuno-compromised  
172 patients as well as lead to transfer of antibiotic resistance traits to hitherto sensitive microorganisms co-habiting within the  
173 consumers of those products. Given the increasing rate of development of resistant bacteria strains, our challenge is to  
174 slow the rate at which resistance develops and spreads. In order to decrease the spread of resistance among antibiotics,  
175 physicians, pharmacists, researchers and consumers alike need to be more aware of the selective pressures driving  
176 these bacteria to decrease their susceptibility [33]. These selective pressures include the abuse, overuse and misuse of  
177 antimicrobials in therapy, improperly manufactured and mishandled HMPs [13, 34] as well as other numerous  
178 socioeconomic factors that govern the development of multi-drug resistant bacteria strains [35].

#### 179 4. CONCLUSION:

180 The high rate of resistance to antimicrobial agents of **microbial strains** isolated from these herbal preparations may  
181 indicate a widespread antibiotic resistance among microorganisms from different sources. It is therefore mandatory that  
182 herbal medicines should not be taken indiscriminately and that current good manufacturing practices (cGMPs) must be  
183 observed by these herbal practitioners in the production of the medicines.

#### 184 Authors' Contributions:

185 Oli Angus N - drafting of the manuscript/Corresponding author, Esimone Charles O – conceptualized and designed the  
186 work as well as revising the manuscript critically for important intellectual content, Ujam Nonye T – carried out the  
187 experiments and did analysis and interpretation of data, Adikwu Michael Umale - revised the manuscript critically for  
188 important intellectual content, Ikegbunam MN- carried out some part of the experimental work

189 **Competing Interest:** Authors have no competing interests to declare

190  
191  
192  
193



## References

1. Oyetayo, V.O. (2008). Microbial load and antimicrobial property of two Nigerian herbal remedies *Afr. J. Trad. Compl. Alt. Med.*, **5**: 74 – 76.
2. Eisenberg D, David RB, Etnner SL, Appel S, Van Rompay M, Kessler RC (1998). Trends in alternative medicine use in the United States, 1990 – 1997 *JAMA* **280**: 1569-1575.
3. WHO (1998). Regulatory Situation of Herbal Medicine. A Worldwide View. World Health Organization, Geneva.
4. Anyanwu, Chukwudi U.(2010) Fungal Contaminants of Powdered Herbal Drugs Sold in Parts of Enugu State, Southeast, Nigeria *Plant Product Research Journal* (2010), Volume 14: 46 – 50.
5. Lamikanra A, Orafidiyi L, Adediji JA (1992). A study of the microbiological quality of 'Ogun efu'. *Afr. J. Pharm. Pharm. Sci.* **22**: 222-228.
6. Esimone CO, Chah KF, Ikejide SC (2002). Microbiological quality of herbal preparations marketed in Southeast Nigeria. *J. Nat. Remedies* **2**:42-48
7. Esimone CO, Oleghe PO, Ibezim EC (2003). Effect of preservative agents on the microbial stability of some indigenous herbal preparations. *Niger. J. Pharm.* **34**: 37-42.
8. Memdie UE, Ifudu ND, Brown SA (1993). How safe are non-sterile liquid preparations? *J. W. Afr. Pharm.* **7**: 8-11.
9. Gibbons, S. (2005): Plants as source of bacterial resistance modulators and anti infective agents, *Phytochemistry Rev.* **4**:63-74.
10. Kepil, A.( 2005): The challenge of antibiotic resistance , Need to contemplate. *Indian J. med Rev.* **121**:83-91.
11. Cohen ML, Auxe RV. (1992) Drug resistant *Salmonella* in the United States: an epidemiologic perspective. *Science*; **234**: 964-70.
12. Peter AGM. (1999) Overview of herbal quality control. *Drug Information Journal*; **33**: 717–24.
13. Esimone C.O., Oleghe, P. O., Dibua, U.E and Ibezim, E.C (2007) Gross Microbial Contamination of herbal medicinal products (HMPs) Marketed in Mid-Western Nigeria. *Int.J.Mol.Med.Adv.Sci*,**3**(2):87-92.
14. Cheesbrough (2009) *District Laboratory Practice in Tropical Countries*, Second Edition Part II, Cambridge University press, London. Pg 62-70.
15. Barrow, G.I and Feltham, R.K.A (1993) *Cowan and Steel's manual for the identification of medical Bacteria*, Cambridge University Press: London, pp: 21-42.
16. National Committee for Clinical Laboratory Standards (1997). Performance Standard for Antimicrobial Disk Susceptibility Tests, 6<sup>th</sup> Edn, Approved Standards Document M2- A6. Wayne, PA: NCCLS.
17. National Committee for Clinical Laboratory Standards. (2003). NCCLS document M2-A8 volume 23, no. 1, Performance standards for antimicrobial disk; Susceptibility tests, approved standard, 8<sup>th</sup> ed. National Committee for Clinical Laboratory Standards, Villanova, PA.
18. Archer GL, Laughter JP, Johnston JL (1986). Plasmid encoded Trimethoprim resistance in Staphylococci. *Antimicrob. Agents Chemother.* **29**: 733-740.
19. Davies AJ, Stone JW (1986). Current problems of Chemotherapy of infections with coagulase – negative Staphylococci. *Eur. J. Clin. Microbiol.* **5**:277-281.
20. Esimone, C.O., Oleghe, P.O, Ibezim, E.C., Okeh, C. O. and Iroha I. R. (2007a). Susceptibility resistance profile of microorganism isolated from herbal medicine product sold in Nigeria. *African journal of biotechnology vol. 6*(24), pp-2766-2775.
21. Schaberg BR, Zerros MJ (1986). Inter-generic and interspecies gene exchange in Gram positive Cocci. *Antimicrob. Agents Chemother.* **30**: 817-822
22. European Pharmacopoeia: (2007) Directorate for the Quality of Medicines of the Council of Europe, 5<sup>th</sup> Edition. Strasbourg, France.
23. Okunlola, A., Adewoyin, A.B., and Odeku, A.O. (2007). Evaluation of pharmaceutical and microbial qualities of some herbal medicinal products in south western Nigeria. *Trop J Pharmaceut Res.* **2007**; *6*(1):661–670.
24. Martínez JL, Baquero F. (2002) Interactions among strategies associated with bacterial infection: pathogenicity, epidemicity, and antibiotic resistance. *Clinical Microbiology Reviews*; **15** (4): 647-79
25. Cotton DJ, Marshall DJ, Gress J, Thaler M, Pizzo PA (1987). Clinical features and therapeutic intervention in 17 cases of Bacillus bacteremia in an immuno-suppressed Patient Population *J. Clin. Microbiol.* **25**: 672-674.
26. Sliman R, Rehn S, Shlaces DM (1987). Serious infections caused by Bacillus species. *Medicine* **66**: 218-223.
27. Kramer JM, Gilbert RJ (1989) *Bacillus cereus* and other Bacillus species. In: Doyle MP, ed. *Foodborne Bacterial pathogens*. New York: Marcel Dekker, pp. 21-70.
28. Prince A, Wood MS, Sooug G, Xingchin N (1988). Isolation and characterization of a penicillinase from *Pseudomonas cepacia*. *Antimicrob. Agents Chemother.* **32**: 838-843.

- 251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269
29. Paul S, Bezbaruah RL, Roy MK, Ghosh AC (1997). Multiple antibiotic resistances (MAR) index and its reversion in *Pseudomonas aeruginosa*. Lett. Appl. Microb. 24: 169-171.
  30. Kumar, R., Shrivastava, S. K. and Chakraborti, A.(2010). Comparison of Broth Dilution and Disc Diffusion Method for the Antifungal Susceptibility Testing of *Aspergillus flavus* Am. J.Biomed. Sci., 2(3), 202-20
  31. O'Brien TF, Acar JF (1987). Antibiotic resistance worldwide. In; Peterson PK, Verhoef J eds. Antimicrobial Agents Annual 2nd ed. Amsterdam Elsevier, pp. 457-470.
  32. Burns JL, Hedin LA, Lieu DM (1989). Chloramphenicol resistance in *Pseudomonas cepacia* because of decreased permeability Antimicrob. Agents Chemother. 33: 136-141.
  33. Gershman K. (1997) Antibiotic resistance and judicious antibiotic use.  
[http://biology.kenyon.edu/slenc/bio38/stancikd\\_02/References.html](http://biology.kenyon.edu/slenc/bio38/stancikd_02/References.html).
  34. Lexchin J. (2000) Promoting resistance? World Health Organization Essential Drug Monitor, Geneva, Nos. 28 and 29.
  35. Toebe C. (2001) Antibiotic Resistance. City College of San Francisco.  
[http://biology.kenyon.edu/slenc/bio38/stancikd\\_02/References.html](http://biology.kenyon.edu/slenc/bio38/stancikd_02/References.html).