

AMELIORATIVE EFFECTS OF ALCOHOL ON HUMAN DIABETIC VOLUNTEERS – A PROSPECTIVE STUDY

A. Naga Vamsi Krishna¹, V. Chandra Prakash³, G. Uma Ramani⁴, N. Ch. Varadacharyulu⁵, B. Narasimha Rao², M. Pardha Saradhi², Mahendran B², L. A. Samuel⁶ and B. Venkata Raman^{2,7}

¹Department of Biochemistry, Acharya Nagarjuna University, Nagarjunanagar, Guntur district-522510, A.P., India.

²Department of Biotechnology, K L University, Vaddeswaram-522502, Guntur, A.P., India.

³Dept. of CSE, K L University, Vaddeswaram-522502, A.P., India.

⁴Dept. of Biochemistry, Katuri Medical College, Katuri Nagar, Guntur – 522019, A. P., India.

⁵Dept. of Biochemistry, Sri Krishna Devaraya University, Anantapur – 515003, A. P., India.

⁶Dept. of Biotechnology, Rajah RSRK Ranga Rao College, Bobbili, Vizianagaram, A. P., India.

⁷Dept. of Basic Sciences, Madanapalle Institute of Technology & Science (MITS), Post Box No: 14, Angallu (V), Madanapalle-517325, A. P., India.

*Authors for correspondence:

Tel. Phone: +91-8571-280295; Mobile: +918008570064

Fax: +91-8571-280433

Email: drbvraman@gmail.com

ABSTRACT

Aims: The purpose of this study is to assess and confirm the ameliorative effects of alcohol consumption on biochemical indices of blood viz. blood glucose, HbA1c, NO₂, NO₃, lipid profiles, hs-CRP (high sensitive C-Reactive protein) and membrane lipid peroxidation of diabetics.

Study Design: Pre-clinical and Biochemical experimental study.

Place and Duration of Study: Department of Biochemistry, Acharya Nagarjuna University & Dept. of Biotechnology, K L University, Guntur, A.P and Dept. of Biochemistry, Katuri Medical College, Katuri Nagar, Guntur, A.P and Dept. of Biochemistry, Sri Krishna Devaraya University, Anantapur, A.P and Dept. of Basic Sciences, Madanapalle Institute of Technology & Science (MITS), Post Box No: 14, Angallu (V), Madanapalle, A. P., India, during 2008 – 2013.

Methodology: The study is conducted on 3 groups of people of different ages ranging from 35 to 50 years at community health centers in Prakasam, Warangal, Srikakulam districts of Andhra Pradesh, India. The first group consists of type-II diabetic patients who have been consuming alcohol (arithmetic mean ranging from 14.16 to 31.61ml/day) moderately for the past 3 to 10 years. The second group consists of 110 patients who are type-II diabetics (who do not drink) taking medical treatment for minimum period of 1 year. The third group consists of 100 non-drinking, non-diabetic healthy individuals. Relationships of alcohol intake with lipid profile, hs-CRP and HBA1c are compared among the three groups.

Results: In lipid profile analysis of moderately drinking diabetic group, the HDL levels are found to be higher while the remaining factors such as total cholesterol, LDL, VLDL ($P<0.05$), triglycerides ($P<0.01$) and membrane lipid peroxidation are significantly lower. Fasting blood glucose, serum nitrites and nitrates are found to be significantly higher. These differences are not found in control group and Diabetic group who do not drink.

Conclusion: Moderate consumption of alcohol is found to have an inverse association with the risky factors like LDL cholesterol, Triglycerides, etc. that are the etiological factors for some of the sequelae of diabetes mellitus viz. coronary heart diseases, Retinopathy, etc. and has a direct association with the positive factors such as HDL and nitric oxide production. Experimental results are very significant and indicate that moderate consumption of alcohol has ameliorative effects on diabetics.

Keywords: diabetics, moderate drinkers, lipid profiles, Nitrites & Nitrates, HDL and HbA1c.

1. INTRODUCTION

Diabetes is a disorder where the body does not produce insulin or does not properly use insulin. According to recent estimates, approximately 285 million people worldwide (6.6%) are suffering from diabetes and this number is expected to rise by 438 million people (7.8%) of the adult population by 2030 [1, 2, 3, 4]. Glucose is derived from all sorts of foods that we consume. After every meal a large part of our food is converted into glucose, thereby increasing the blood glucose levels. The Insulin, a hormone secreted by pancreas carries the blood glucose to cells that need energy [5, 6]. In diabetic individuals, insulin is either not produced or not utilized properly, and hence the glucose remains in the blood causing the condition "Diabetes" [7, 8]. Today, diabetes mellitus type 2 is posing several challenges to the medical field due to its association with multiple physiological complications such as Cardiovascular complications, Microangiopathy, Neuropathy, Nephropathy, Retinopathy, Dermatopathy, etc [9]. Currently oral hypoglycaemic agents used for the treatment of Type 2 diabetes include insulin secretagogues like

sulfonylureas and metformin. Metformin acts through multiple poorly characterized mechanisms, one of which inhibits de novo glucose synthesis via indirect AMP-activated protein kinase (AMPK) activation, potentially following partial mitochondrial complex I inhibition in the liver [10]. Recently, the focus has been shifted towards the use of moderate alcohol to treat Type 2 diabetes. Alcohol consumption is increasing day by day, not only in Asian countries but also throughout the world. Alcohol is a globally abused psycho-active drug with its adverse side effects but it has also some important beneficial effects like relaxation of mental tension, vasodilatory **effect on human** health [11]. Excessive consumption of alcohol has definite adverse effect on human health. Several studies have shown that people with the habit of excessive drinking of ethanol are found to have fatty liver [12], cognitive disorders and permanent irreversible liver **damage**. On the other hand, it **is** also shown that moderate **consumption of alcohol has** beneficial health effects [13, 14, 15]. The concept of moderate consumption of ethanol (beverage alcohol) has evolved over time from considering the level of intake to be non-intoxicating and non-injurious. Moderate drinking can be defined as the level corresponding to the lowest overall rate of morbidity or mortality in a population [16].

Therefore, in our study we have evaluated the ameliorative effects of alcohol consumption on biochemical indices on human diabetic volunteers. Our results **show** that moderate alcohol consumption enhanced the levels of HDL by lowering LDL and total triglycerides pools. Moreover, enhanced levels of serum NO₂ and NO₃ **are** noticed in moderate alcohol drinking diabetic volunteers.

2. MATERIALS AND METHODS

2.1 Subjects for study

The study **is** conducted on 3 groups of people of **different** ages ranging from 35 to 50 years at community health centers in **Prakasam, Warangal, Srikakulam Districts** of Andhra Pradesh, India. The first group consists of type-II diabetic patients who have been consuming alcohol moderately for the past 3 to 10 years. This group is named as **MDD** (Moderate Drinking **Diabetics**). The second group consists of **non-drinking type-II diabetic patients who have been under** medical treatment for **a** minimum period of 1 year. This group is named as **NDD** (Non-Drinking Diabetics). The third group consists of non-drinking, non-diabetic healthy individuals. This group is **Abstainers** (Table 1). All volunteers involved in the present study **are** well informed and their consent **is** obtained. All the members of the above groups are free from Coronary Heart Diseases (CHD), Cerebro Vascular Diseases (CVD) and Cancer.

2.2 Determination of fasting blood glucose

Blood samples from every individual **are** collected into EDTA containing tubes by venipuncture. Levels of glucose in serum is estimated using monozyme diagnostic kit, which is based on the GOD-POD method [17]. In brief, glucose is oxidized by the enzyme glucose oxidase to give D-gluconic acid and hydrogen peroxide. Hydrogen peroxide in presence of enzyme peroxidase oxidizes phenol, which combines with

amino antipyrine dye to produce a red coloured quinoneimine which is measured at 505 nm against water blank.

2.3 Determination of serum triglycerides

Serum triglycerides are estimated using Qualigens diagnostic kit which is based on the method [18]. In brief, triglycerides in the sample are hydrolyzed by microbial lipase to glycerol and free fatty acids. Glycerol is further phosphorylated to glycerol 3-phosphate and is oxidized to dihydroxy acetone phosphate. Liberated hydrogen peroxide reacts with 4-amino anti pyrine and 3, 5 dichloro 2-hydroxy benzene sulphonic acid. Absorbance of quinoneimine and colour dye formed is proportional to the concentration of triglycerides.

2.4 Determination of Serum Total Cholesterol

Serum total cholesterol is estimated by the enzymatic kit method [19]. In brief 0.01ml of serum is added to 1ml of freshly reconstituted enzyme reagent, mixed well and incubated at 37°C for 5 minutes. After incubation, absorbance is measured at 505nm against blank. Simultaneously standards are run along with the test under similar conditions.

2.5 Determination of HDL and LDL -Cholesterol

Serum HDL-Cholesterol is estimated by autozyme diagnostic kit method. 0.5ml of HDL precipitant reagent (Phosphotungstic acid 2.4 mmol/L and Magnesium Chloride 40m mol/L) is added to 0.5ml of serum, mixed thoroughly, centrifuged at 4,000 rpm for 10min to obtain a clear supernatant. 1ml of working standard (enzymatic cholesterol reagent of autozyme diagnostic kit) is added to 0.05ml of supernatant, incubated for 10min at 37°C and the development of color is read at 510 nm against a blank. A standard is maintained simultaneously. LDL and VLDL cholesterol are calculated using the formula of [20].

2.6 Determination of CRP protein in serum

Cholestech LDX hs-CRP is an *in vitro* diagnostic test for the quantitative determination of hs – CRP (high sensitive C–Reactive protein) in whole blood or serum [26]. Finger stick samples are collected using a Cholestech LDX 50 µl capillary tube. The cassette is placed into the drawer of the analyzer immediately after dispensing the sample into the well. After pressing run, hs-CRP results are displayed in 6 minutes (results are displayed in 4 minutes for serum of serum sample). It is found that Hematocrit levels between 30% and 55% do not affect the results.

2.7 Determination of total blood Nitrite and Nitrate

Nitrites and Nitrates are estimated in the serum samples of the subjects [22, 23]. Serum samples are deproteinated by adding 30% ZnSO₄ followed by centrifugation at 10,000 rpm for 5 minutes. Then, 1ml of serum supernatant is mixed with 1ml Greiss reagent (1g/lit sulfanilamide, 25g/lit phosphoric acid and 0.1gm/lit N-(1-Naphthyl) ethylene diamine dihydro chloride) and incubated at room temperature for 10 minutes for color development. The absorbance is measured at 545 nm in Elico Spectrophotometer against blank.

2.8 Statistical Analysis

All the values of body weight, fasting blood sugar, and biochemical estimations were expressed as mean \pm standard deviation (S.D). Differences of mean values are assessed by paired or unpaired Student's *t* test for comparison of 2 variables and by ANOVA for comparison of multiple variables. Relationships between 2 continuous variables are assessed by a regression analysis using the Pearson correlation coefficient. Differences between Alcoholic and Non-alcoholic diabetic groups are analyzed by χ^2 test. A value of $P < 0.05$ is considered statistically significant.

3. RESULTS AND DISCUSSION

Diabetes is a complex metabolic disorder and several factors such as environmental and life style factors have shown to be responsible for the origin and development of diabetes mellitus. Although diabetes is as old as human life on earth, researchers are yet to find out a therapeutic factor with less diabetic complications. In this paper, the authors explore the possible action of alcohol intake and diabetic control by measuring several biochemical indices in the blood serum of diabetics. However, the alcohol content in different drinks viz. wine, brandy, whisky and other beverages varies considerably [24]. Therefore, a questionnaire has been prepared to know the type of drink consumed by MDD, which is shown in table 2. Based on that we calculated the arithmetic mean consumption of ethanol per day drunk by MDD, which ranges from 14.16 ml to 31.85 ml. Evaluation of the blood samples shows that moderate consumption of alcohol positively influences the indices of blood parameters of diabetics i.e., hs-CRP protein, fasting blood glucose, HbA1c, total blood Nitrite and Nitrate, total cholesterol, HDL, LDL, VLDL, Triglycerides and membrane lipid peroxidation and hence it is useful to ameliorate the deleterious effects of diabetes mellitus.

It is observed from the results that membrane lipid peroxidation is declined in moderately drinking diabetic group than diabetic and control groups (Table 3). In lipid profile analysis, only HDL levels are increased in MDD than DG while remaining factors such as total cholesterol, LDL, VLDL ($P < 0.05$), membrane lipid peroxidation and triglycerides ($P < 0.01$) are significantly reduced (Table 4). Both the study groups are compared with the control group. These results on lipid profile due to the impact of alcohol consumption are supported by several authors who conducted experiments on different animals including human beings [25]. Similar experiments are conducted on men with and without diabetes and a positive association between alcohol intake and blood pressure, triglycerides and HDL cholesterol is found [26]. Some researchers may still have a doubt whether excessive consumption of alcohol may result in obesity. But, this ambiguity is already resolved by [27] who observed that drinkers, despite their higher alcohol intake, are no more obese than nondrinkers. Their observations strongly complement the observations of present study.

The levels of serum Nitrites and Nitrates are found to be increased in MDD as compared to NDD. Earlier studies revealed that moderate alcohol consumption might have induced an increase in insulin secretion, sensitivity to insulin, increased serum nitrites and nitrates levels in MDD than NDD. Relationship between

serum Nitric Oxide (NO) production, lipid abnormalities and oxidative stress in diabetes are noticed earlier [28]. Many reports strongly support that diabetes mellitus is associated with decreased Nitric Oxide production from endothelial cells and decreased levels of serum NO₂ and NO₃ [28, 29, 30, 31]. Moderate alcohol consumption has been shown to reduce the risk of ischemic heart disease potentially through its effect on specific endothelial-derived compounds. Venkov et al., [32] have tested the hypothesis that ethanol increases the expression of endothelial Nitric Oxide Synthase (eNOS) and Nitric Oxide production in Bovine aortic endothelial cells. Luo et al., [33] and Bequette et al., [34] observed that intake of alcohol has direct influence on wound healing and ascribed this property of alcohol to increased production of NO which, as a vasodilator, helps in healing the wound. In fact, alcohol rubbed on skin dilates the blood vessels and produces a mild counter-irritant effect. In the general practice of public, whenever a small cut/injury appears on the body, people pour a few drops of alcohol on the injured part and the wound gets healed subsequently. Other reports also strongly suggest that increased production of nitric oxide in alcoholic diabetics reduces the serum glucose levels, oxidative stress, lipid and lipoprotein abnormalities [18, 35, 36, 37, 38, 39].

In the present study, it is found that moderate alcohol consumption enhances the levels of serum NO₂ and NO₃ in MDD when compared to NDD ($P < 0.05$); this observation strongly coincides with above reports. The etiological factor for most of the sequelae of diabetes mellitus of type I or II viz. Retinopathy, Nephropathy, Cardio-myopathy, Polyneuropathy, Neuritis, Erectile and Dysfunction is ischemia due to lowered levels of Nitric Oxide production. Hence, the authors opine that moderate consumption of alcohol ameliorates the severity of diabetes mellitus and its sequelae to some extent due to increased nitric oxide synthase protein expression of one or more isoforms.

The moderate consumption of alcohol causes a significant decrease in serum glucose levels ($P < 0.05$) and glycosylated hemoglobin in MDD than NDD as observed earlier through similar experiments conducted on moderately drinking type-II diabetics [40, 41]. Similar results are reported by [42] conducting experiments on rats where they demonstrated that ethanol acutely exerts substantial influences on pancreatic microcirculation by evoking a massive redistribution of pancreatic blood flow from the exocrine into the endocrine part via mechanisms mediated by nitric oxide and vagal stimuli, augmenting late-phase insulin secretion, and thereby evoking hypoglycemia. This mechanism seems to involve NO & vagal pathways and is due to the well-known hypoglycemic properties of alcohol in diabetic patients [43, 44]. A Dutch randomized trial conducted in diabetic teetotallers suggests that a glass of wine with dinner may improve glucose control, particularly in those with higher HbA1c levels to begin with. This study, while small, adds to anecdotal evidence and meta-analyses that suggest that wine may hold specific benefits for diabetics whose cardiovascular benefits have been widely touted (European Association for the study of Diabetes 2007 meeting, an unpublished report).

Consumption of white and red wines may improve coronary blood flow and improve symptoms in patients with coronary heart diseases [45]. In our experiments, it is observed that hs-CRP levels in blood serum are found to be significantly ($P < 0.05$) low in MDD when compared with that of NDD, which indicates that the probable risk of cardiovascular diseases is low in MDD (Table 3).

Glycosylated hemoglobin (Hemoglobin A1c) concentration is a hallmark of glycemic control for prognostic purpose. HbA1c levels are reported to be in correlation with, not only glycosuria but also serum glucose. Hormonal profiles and various other factors cannot influence HbA1c concentrations [34]. Our experiments on HbA1c levels in the MDD and NDD patients show that lowered levels of blood glucose exist in MDD than NDD. These results strongly support our hypothesis that moderate consumption of alcohol has an ameliorative effect on diabetes mellitus. As the results are very significant, the authors propose that moderate consumption of alcohol (ranging from 14.16 ml to 31.85 ml per day) is good for the health of the diabetics. This range is very much below the safer range i.e., 30 to 40 ml of ethanol consumption/day as advised by the UK government (International center for Alcohol Policies, USA).

4. CONCLUSION:

Moderate consumption of alcohol is found to have an inverse association with the risky factors like LDL cholesterol, Triglycerides, etc. that are the etiological factors for some of the sequelae of diabetes mellitus viz. coronary heart diseases, Retinopathy, etc. and has a direct association with the positive factors such as HDL and nitric oxide production. Experimental results are very significant and indicate that moderate consumption of alcohol has ameliorative effects on diabetics.

ACKNOWLEDGMENTS

Authors thank Prof. Dr. D. N. Rao, Dept. of Biochemistry, AIIMS, New Delhi and Prof. T. M. Radha Krishnan, Dept. of Biotechnology, AUCST, Andhra University, Visakhapatnam for their support and Dr. G. Hampamma, Professor of English, MITS, Madanapalle for English editing of this paper.

COMPETING INTERESTS

None declared

REFERENCES

1. King H, Aubert RE, Herman WH. Global burden of diabetes, 1995-2025: Prevalence, numerical estimates, and projections. *Diabetes Care*. 1998; 21:1414-1431.
2. Amos AF, McCarty DJ, Zimmet P. The rising global burden of diabetes and its complications: estimates and projections to the year 2010. *Diabet Med*. 1997; 14(5):1-85.
3. IDF Diabetes Atlas. 4th edition. International Diabetes Federation. 2009.
4. Mohan V, Pradeepa R. Epidemiology of diabetes in different regions of India. *Health Administrator*. 2009; 22:1-18.
5. Gershon, Michael D. A Groundbreaking New Understanding of Nervous Disorders of the Stomach and Intestine. *The Second Brain*. 1999; New York: HarperCollins.
6. Guyton, Arthur C., and John E. Hall. *Textbook of Medical Physiology*. 2000; 10th ed. Philadelphia: Saunders.

7. Sivitz, William I., MD."Understanding Insulin Resistance: What Are the Clinical Implications?"Postgraduate Medicine. 2004; 116:41-48.
8. Service, F. J. Hypoglycemic disorders. N Engl J Med. 1995; pp. 1144-1152.
9. King H, Aubert RE, Herman WH. Diabetes Care, 1998; 21:1414-31.
10. Shaw RJ, Lamia KA, Vasquez D, Koo SH, Bardeesy N, Depinho RA, Montminy M, Cantley LC. The kinase LKB1 mediates glucose homeostasis in liver and therapeutic effects of metformin. Science. 2005; 310:1642–1646.
11. Wang JL, Patten SB. Alcohol Consumption and Major Depression: Findings from a Follow-Up Study. Canadian Journal of Psychiatry. 2001; 46, pp 632-638.
12. Alatalo PI, Koivisto HM, Hietala JP, Puukka KS, Bloigu R, Niemelä OJ. Effect of moderate alcohol consumption on liver enzymes increases with increasing body mass index. Am J Clin Nutr. 2008; 88, 1097-1103.
13. Howard AA, Arnsten JH, Gourevitch MN. Effect of Alcohol Consumption on Diabetes MellitusA Systematic Review. Ann Intern Med. 2004; 140, 211-219.
14. Paramahansa M, Aparna S, Varadacharyulu N. Alcohol-induced alterations in blood and erythrocyte membrane in diabetics. Alcohol Alcohol. 2002; 37, 49-51.
15. Sellman D, Connor J, Robinson G, Jackson R. Alcohol cardio-protection has been talked up. N Z Med J. 2009; 122: 97-101.
16. Eckardt MJ, File SE, Gessa GL, Grant KA, Guerri C, Hoffman PL, Kalant H, Koob GF, Li TK, Tabakoff B. Effects of Moderate Alcohol Consumption on the Central Nervous System. Alcohol Clin Exp Res. 1998; 22, 998-1040.
17. Kumar K, Patel A, Shirode D, Baganal P, Rajendra S, Setty S. Influence of metronidazole on hypoglycemic activity of thiazolidinediones in normal and alloxan induced diabetic rats. Indian J. Pharm. Educ. Res. 2009; 43, 93-97.
18. Fossati P, Prencipe L. Serum triglycerides determined colorimetrically with an enzyme that produces hydrogen peroxide. Clinical chemistry.1982; 28, 2077-2080.
19. Allain CC, Poon LS, Chan CS, Richmond W, Fu PC. Enzymatic determination of total serum cholesterol. Clinical chemistry.1974; 20, 470-475.
20. Friedewald WT, Levy R I, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. Clinical chemistry. 1972; 18, 499-502.
21. Kaptoge S, Di Angelantonio E, Lowe G, Pepys MB, Thompson SG, Collins R, Danesh J. C-reactive protein concentration and risk of coronary heart disease, stroke, and mortality: an individual participant meta-analysis. Lancet. 2010; 375, 132.
22. Rajeshkumar K, Amteshwar J, Nirmal S, Bhupesh S. Ameliorative role of Atorvastatin and Pitavastatin in L-Methionine induced vascular dementia in rats. BMC Pharmacology. 2008; 8, 1-12.
23. Sastry K, Moudgal R, Mohan J, Tyagi J, Rao G. Spectrophotometric determination of serum nitrite and nitrate by copper–cadmium alloy. Anal Biochem. 2002; 306, 79-82.
24. Gual A, Martos AR, Lligoña A, Llopis JJ. Does the concept of a standard drink apply to viticultural societies? Alcohol Alcohol. 1999; 34, 153-160.
25. Estruch R, Sacanella E. Alcohol:¿ tónico o tóxico cardiovascular? Clínica e investigación en arteriosclerosis. 2005; 17, 183-195.
26. Wakabayashi I. Comparison of the relationships of alcohol intake with atherosclerotic risk factors in men with and without diabetes mellitus Alcohol Alcohol. 2011; 46, 301-307.
27. Gruchow H, Sobocinski K, Barboriak J, Scheller J. Alcohol consumption, nutrient intake and relative body weight among US adults. Am J Clin Nutr. 1985; 42, 289-295.
28. Sturgeon KM, Fenty-Stewart NM, Diaz KM, Brinkley TE, Dowling TC, Brown MD. The relationship of oxidative stress and cholesterol with dipping status before and after aerobic exercise training. Blood Press. 2009; 18, 171-179.
29. Monti LD, Barlassina C, Citterio L, Galluccio E, Berzuini C, Setola E, Valsecchi G, Lucotti P, Pozza G, Bernardinelli L. Endothelial nitric oxide synthase polymorphisms are associated with type 2 diabetes and the insulin resistance syndrome. Diabetes. 2003; 52, 1270-1275.
30. Komers R, Schutzer WE, Reed JF, Lindsley JN, Oyama TT, Buck DC, Mader S L, Anderson S. Altered endothelial nitric oxide synthase targeting and conformation and caveolin-1 expression in the diabetic kidney. Diabetes. 2006; 55, 1651-1659.

31. Kashyap SR, Roman LJ, Lamont J, Masters BSS, Bajaj M, Suraamornkul S, Belfort R, Berria R, Kellogg DL, Liu Y. Insulin resistance is associated with impaired nitric oxide synthase activity in skeletal muscle of type 2 diabetic subjects. *J Clin Endocrinol Metab.* 2005; 90, 1100-1105.
32. Venkov CD, Myers PR, Tanner MA, Su M, Vaughan DE. Ethanol increases endothelial nitric oxide production through modulation of nitric oxide synthase expression. *Thromb Haemost.* 1999; - 81, 638-642.
33. Luo JD, Wang YY, Fu WL, Wu J, Chen AF. Gene therapy of endothelial nitric oxide synthase and manganese superoxide dismutase restores delayed wound healing in type 1 diabetic mice. *Circulation.* 2004; 110, 2484-2493.
34. Bequette BW. Continuous glucose monitoring: real-time algorithms for calibration, filtering, and alarms. *J Diabetes Sci Technol.* 2010; 4, 404 – 418.
35. Alving K, Janson C, Nordvall L. Performance of a new hand-held device for exhaled nitric oxide measurement in adults and children. *Respir Res.* 2006; 7, 67.
36. Dandana A, Gammoudi I, Ferchichi S, Chahed H, Limam HB, Addad F, Miled A. Correlation of Oxidative Stress Parameters and Inflammatory Markers in Tunisian Coronary Artery Disease Patients. *Int J Biomed Sci.* 2011; 7: 6-13.
37. Hastie CE, Haw S, Pell JP. Impact of smoking cessation and lifetime exposure on C-reactive protein. *Nicotine & Tobacco Research.* 2008; 10, 637-642.
38. Szmitko PE, Wang CH, Weisel RD, de Almeida JR, Anderson TJ, Verma S. New markers of inflammation and endothelial cell activation part I. *Circulation.* 2003; 108, 1917-1923.
39. Witte M, Kiyama T, Barbul A. Nitric oxide enhances experimental wound healing in diabetes. *Br J Surg.* 2002; 89, 1594-1601.
40. Stechmiller JK, Childress B, Cowan L. Arginine supplementation and wound healing. *Nutr Clin Pract.* 2005; 20, 52-61.
41. Wotherspoon, F., Laight, D., Browne, D., Turner, C., Meeking, D., Allard, S., Munday L, Shaw K, Cummings M. Plasma homocysteine, oxidative stress and endothelial function in patients with Type 1 diabetes mellitus and microalbuminuria. *Diabet Med.* 2006; 23, 1350-1356.
42. Huang Z, Sjöholm Å. Ethanol acutely stimulates islet blood flow, amplifies insulin secretion, and induces hypoglycemia via nitric oxide and vagally mediated mechanisms. *Endocrinology.* 2008; 149, 232-236.
43. Naga Vamsi Krishna A. A prospective study of biochemical changes in membranes of chronic human alcoholic diabetic volunteers. An M.Phil Thesis submitted to Dept. of Biochemistry, Annamalai University, Chidambaram, Tamil Nadu, 2006; pp 32-36.
44. Takahashi T, Owyang C. Characterization of vagal pathways mediating gastric accommodation reflex in rats. *J Physiol.* 1997; 504, 479-488.
45. Flesch M, Schwarz A, Böhm M. Effects of red and white wine on endothelium-dependent vasorelaxation of rat aorta and human coronary arteries. *Am J Physiol.* 1998; 275, H1183-H1190.

TABLES & LEGENDS

Table -1: Profile of subject groups with and without diabetes

Variables	Alcohol consumption category		
	Moderate drinking Diabetes (MDD)	Non-drinking Diabetes (NDD)	Abstainer
Gender	Male	Male	Male
Number	1200	1200	1200
Age (Years)	52.2 ± 7.8	52.2 ± 7.8	52.2 ± 7.8
Body mass index (Kg/m ²)	25.28 ± 3.27**	28.32 ± 5.21	23.32 ± 2.39
Waist Circumference (Inches)	85.9 ± 13.2**	89.4 ± 15.2	84.3 ± 10.2
Daily consumption of moderate alcohol (>22 and ≤44 g ethanol/day)	All [†]	Nil	Nil
Smokers (%)	52.4	42.5	Nil
Systolic blood pressure (mmHg)	132.4 ± 17.2**	138.4 ± 17.8	130.3 ± 16.7
Diastolic blood pressure (mmHg)	81.4 ± 12.1**	82.4 ± 13.1	80.4 ± 11.1
Therapy for diabetes (%)	58.7**	59.3	Nil
Therapy for hypertension (%)	42.2**	35.8	8
Therapy for dyslipidemia (%)	15.9**	23.7	8.2

Mean with standard deviation or percentages of variables were compared between the non-diabetic and diabetic with drinkers and non-drinkers.

[†]Dose size is 13.0 to 40.0 % ABV (Percent Alcohol by Volume from typical beverage).

** Highly significant differences from non-drinking diabetes and abstainers (P<0.01)

Table - 2: Calculation of ethanol content in drinks consumed by **MDD**.

S. No.	Type of Drink	ABV* (%)	Daily consumption of drink ** (in ml)	Content of ethanol in the drink*** (in ml)
1	Wine	13.5	105.00	14.16
2	Brandy	40	77.65	26.76
3	Rum	37.5	80.00	30.00
4	Gin	40	71.25	28.50
5	Whisky	40	79.62	31.85
6	Cheap Liquor	40	79.02	31.61

* Typical Alcohol by Volume; ** Arithmetic mean alcohol consumption of **MDD** in a week equivalent to ethanol (i.e., 220 ml ethanol per week***)

TABLE-3: Impact of alcohol on lipid constituents of serum and erythrocyte membrane in different experimental groups

S. No.	Parameter	Alcohol consumption category		
		Moderate drinking Diabetes (MDD)*	Non-drinking Diabetes (NDD)	Abstainer
1	Fasting serum glucose (mg / dl)	130 ± 4.3	180 ± 7.0	72± 2.3
2	hs-CRP (mg/L)	2.54 ± 0.05	3.12 ± 0.03	1.3± 0.06
3	Membrane Lipid peroxidation (pmol of MDA) †	4.961 ± 1.15	8.304 ± 1.026	3.20 ± 0.15
4	HBA1c ^{††}	9.5 ± 2.3	11.4 ± 2.2	6.5± 1.0
5	Serum Nitrites (µ moles/L)	2.5 ± 0.04	2.3 ± 0.03	1.6± 1.0
6	Serum Nitrates (µ moles/L)	24.5 ± 0.4	22.7 ± 0.5	23.1 ± 8.9

†Malonaldehyde formed / mg membrane protein

††Determined using Glycated hemoglobin assay kit recommended by the American diabetes association (ADA) and is expressed as a percentage (%) of the hemoglobin

* Significant variation from non-drinking diabetes and abstainers (P<0.01 to 0.05)

TABLE-4: Variation in the lipid profiles of diabetic volunteers with and without drinking

S. No.	Parameter (mg / dl)	Alcohol consumption category*		
		Moderate drinking diabetes (MDD)**	Non-drinking Diabetes group (NDD)	Abstainers
1	Total Cholesterol	220 ± 8.4	265 ± 7.8	198±8
2	Triglycerides	170 ± 8.5	250 ± 5.3	142±29
3	HDL	82 ± 5.1	53 ± 3.7	42±1.8
4	LDL	51 ± 3.6	59 ± 4.0	60±10
5	VLDL	35 ± 3.1	48 ± 3.6	38±2.0

Mean values (n=1200) represented as mean values ± S.D.

** Significant variation from non-drinking diabetes and abstainers (P<0.01-0.05)