

# Low Vitamin D Status of Northern Italian Children in Pediatric Primary Care Setting: What to Do?

Stefano Mazzoleni<sup>1\*</sup>, Chiara Boscardin<sup>2</sup>, Daniela Toderini<sup>3</sup>

<sup>1</sup>MD, Pediatra di Libera Scelta, Polistudio Pediatrico, Piove di Sacco, Padova, Italy

<sup>2</sup>MD, Specializzanda, Clinica Pediatrica, Padova, Italy

<sup>3</sup>MD, Endocrinologa, Medico di Medicina Generale, Padova, Italy

## ABSTRACT

**Aims:** To analyze vitamin D status in a group of children living in Northeastern Italy cared by a "family pediatrician".

**Study design:** Cross-sectional study.

**Place and Duration of Study:** Pediatric primary care in a rural area near Padua (Italy, 45°N latitude), between November 2010 and September 2012.

**Methodology:** The study was conducted with 113 children (41 girls and 72 boys), aged between 1 and 15 years old. The serum level of 25-hydroxyvitamin D [25(OH)D] was measured using a chemiluminescence immunoassay methodology. Serum 25(OH)D test was included in a panel of laboratory tests ordered for different reasons. A correlation was researched between 25(OH)D level and the following variables: class of age, gender, ethnicity, skin colour, period of blood withdrawal, BMI category, results in other laboratory tests and presence of comorbidity.

**Results:** Only 26,5% of children had a normal level of 25(OH)D ( $\geq 30$  ng/ml); in 66,4% of all patients 25(OH)D level was insufficient (10-29 ng/ml) while 7,1% of children had deficiency (25(OH)D  $< 10$  ng/ml). About 40% of all children had 25(OH)D  $< 20$  ng/ml. Non-Italian ethnicity, non-white skin and blood withdrawal in January-March and April-June were significantly associated with hypovitaminosis D [25(OH)D  $< 30$  ng/ml] at univariate level. Both non-Italian ethnicity ( $P = 0.029$ ) and period of blood withdrawal ( $P = 0.0062$ ) were also significant at multivariate analysis. The combination of chronic disease or non-white skin could identify only 50% of children with 25(OH)D  $< 10$  ng/ml and 29% of children with 25(OH)D  $< 20$  ng/ml.

It is of note that rates of hypovitaminosis D were higher in children with normal or low weight than in overweight and obese children, even if this trend did not reach statistical significance.

**Conclusion:** We noted a high incidence of hypovitaminosis D in asymptomatic children without risk factors. In our region cholecalciferol supplementation should be implemented for all children between October and April. Cholecalciferol 1500 UI per day could be appropriated for northeastern Italian children while non-white skinned children need higher doses. Appropriate dose for children of Northern Italy is debated.

**Keywords:** Vitamin D, cholecalciferol, 25-hydroxycholecalciferol, hypovitaminosis D, vitamin D deficiency, vitamin D insufficiency, seasonal variation.

\* Tel./fax: +39 049 9703008. E-mail address: [stefano.mazzoleni@sanita.padova.it](mailto:stefano.mazzoleni@sanita.padova.it).

## 1. INTRODUCTION

Hypovitaminosis D is a common problem worldwide in children and adolescents [1-4], as well in adults [5-6], that places affected population at high risk for chronic diseases [7]. Although severe vitamin D deficiency is rare, there is accumulating evidence of the frequent occurrence of subclinical vitamin D deficiency or insufficiency in otherwise healthy people [8-14]. To our knowledge there are only a few studies on children living in Northeastern Italy [15-17]: they have been conducted retrospectively [15] or examining patients afferent to a Pediatric Department [16] or asthmatic [17]. On this basis an analysis of vitamin D status was prospectively conducted measuring serum 25-hydroxyvitamin D [25(OH)D] in a group of children living near Padua in a primary care setting.

## 2. MATERIALS AND METHODS

### 2.1 Patients and methods

A cross-sectional study was conducted in 113 children living in a rural area near Padua (Italy, 45°N latitude) and cared in office by a single *family paediatrician*. These patients represented 12% of about 900 children registered in his list. The 25(OH)D test was included in a panel of laboratory tests (lab tests) ordered for different reasons (suspected anemia, poor growth, fatigue, etc.) between November 2010 and September 2012. The children weren't taking vitamin D before blood collection and there was no official policy of Italian National Health Service on vitamin D prophylaxis in children > 12 months. Children aged < 12 months were excluded. In Italy they usually receive 400 IU cholecalciferol per day.

The serum level of 25(OH)D was measured using a chemiluminescence immunoassay methodology. The laboratory normal minimum was 30 ng/ml; values of 29 ng/ml or less were considered as hypovitaminosis D: laboratory defined insufficiency as 10-29 ng/ml and deficiency as < 10 ng/ml.

Levels above 100 ng/ml were considered as hypervitaminosis D. Toxicity (including hypercalcemia and hyperphosphatemia) generally occurs for serum levels of 25(OH)D greater than 150 ng/ml [18].

Data collected on patients included age, gender, self-declared ethnicity, skin colour, body mass index (BMI), results of other lab tests and presence of comorbidity. Diet was not analyzed but none of the children was vegetarian. Children were divided into the following age groups: children between 1-5 years old (preschoolers), 6-10 years old (schoolers) and 11-15 years old (adolescents). BMI was calculated and children were divided into 4 categories considering their BMI-for-age-and-gender percentile: underweight (BMI less than 5<sup>th</sup> percentile) [19], healthy weight (BMI from 5<sup>th</sup> percentile to less than 85<sup>th</sup> percentile), overweight (BMI from 85<sup>th</sup> to less than 95<sup>th</sup> percentile) and obese (95<sup>th</sup> percentile or greater) [20].

### 2.2 Statistical Analysis

All data collected were analyzed and described using descriptive statistics. Absolute and percent frequencies were used for qualitative variables; mean, standard deviation (SD), range and median were used to summarize quantitative variables. Principal data were also stratified by level of 25(OH)D (< 10 ng/ml, 10-29 ng/ml, and ≥ 30 ng/ml). Chi-squared (univariate analysis) test was used to determine the presence of a relationship between 25(OH)D stratification groups (deficiency, insufficiency, and normal level) and the following variables: month of blood withdrawal, class of age, gender, ethnicity, skin colour, BMI category, results of other lab tests (e.g. haemoglobin, serum iron, serum cholesterol) and

presence of comorbidity. The linear correlation between quantitative variables (BMI and month of blood withdrawal) and 25(OH)D level was also tested by Pearson's correlation coefficient ( $p$ ). Multinomial logistic regression was used to test factors influencing the probability of hypovitaminosis D [25(OH)D < 30 ng/ml]. The analysis of variance (ANOVA) was used to verify the different level of 25(OH)D in different periods. Tests were considered significant at  $P < 0.05$  level. SAS® software (SAS Institute Inc., Cary, North Carolina, USA) was used for statistical analysis.

### 3. RESULTS AND DISCUSSION

#### 3.1 Results

##### 3.1.1 Socio-Demographic characteristics

Between November 2010 and September 2012, 25(OH)D level was measured in 113 children. Their socio-demographic characteristics are represented in Table 1.

Table 1. Socio-Demographic characteristics of 113 children

Characteristics	N.
Females / Males	41 / 72
1-5 years	53
6-10 years	37
11-15 years	23
Italians / Non Italians	91/22
White / Non white skinned	101/12

##### 3.1.2 Prevalence of Hypovitaminosis D

Values of serum 25(OH)D varied from 4 to 70 ng/ml; the mean level was 24.2 ng/ml (SD 11,6 ng/ml; median 22.0 ng/ml; interquartile range 17-30 ng/ml). Only 30 children (26.5 %) had a normal level; in 75 children (66.4% of all patients) serum 25(OH)D was 10-29 ng/ml insufficient, and 8 children (7.1%) had 25(OH)D < 10 ng/ml vitamin D deficiency (Table 2).

Table 2. Distribution of children by level of 25(OH)D

25(OH)D level	N.	%
< 10 ng/ml	8	7.1
10-29 ng/ml	75	66.4
10-19	37	32.8
20-29	38	33.6
≥ 30 ng/ml	30	26.5
All	113	100

### 3.1.3 Gender

There were 41 females (36.3% of all patients) and 72 males (63.7%). No statistically significant difference was found between males and females regarding the levels of 25(OH)D: 3 females (7.3% of females) and 5 males (6.9% of them) had 25(OH)D < 10 ng/ml vitamin D deficiency; 28 females (68.3% of them) and 47 males (65.3% of males) had 25(OH)D 10-29 ng/ml vitamin D insufficiency. Only 10 females (24.4% of females) and 20 males (27.8% of males) had a normal level of 25(OH)D ( $\geq 30$  ng/ml) (Table 3 deleted).

### 3.1.4 Age group

53 children (46.9% of all patients) were 1-5 years old (preschoolers), 37 (32.7%) were 6-10 years old (schoolers) and 23 (20.4%) were 11 years old or older (adolescents).

Among preschoolers 11.3% had 25(OH)D < 10 ng/ml Vitamin D deficiency, 58.5% had 25(OH)D 10-29 ng/ml were insufficient and 30.2% had a normal value of Vitamin D status [25(OH)D  $\geq 30$  ng/ml]. In schoolers 25(OH)D < 10 ng/ml deficiency of Vitamin D was found only in 2.7%, 25(OH)D 10-29 ng/ml insufficiency in 75.7% and normal value in 21.6%. In the last group of age a similar distribution was found. The 69.8% of preschoolers had value of 25(OH)D lower than normal range and a similar rate (76.7%) was found in older children (> 5 y).

Children aged 5 years or less had more frequently a level of 25(OH)D < 10 ng/ml comparing with older children other groups, even if there was not statistical significance (Table 3 previously Table 4). We must however take into account that 5 among 6 of these children were withdrawn in January-April (see unit 3.6).

**Table 3. Relationship between age group and 25(OH)D level vitamin D status**

	All		25(OH)D level					
			Deficiency < 10 ng/ml		Insufficiency 10-29 ng/ml		Normal $\geq 30$ ng/ml	
	N	% col	N	% row	N	% row	N	% row
1 - 5 y	53	46.9	6	11.3	31	58.5	16	30.2
6 - 10 y	37	32.7	1	2.7	28	75.7	8	21.6
11 - 15 y	23	20.4	1	4.3	16	69.6	6	26.1
All	113	100.0	8	7.1	75	66.4	30	26.5

Chi-Square:  $P = 0.3785$

### 3.1.5 Body Mass Index

Most of the children (71.7%) had normal BMI-for-age, with similar distribution for girls (65.9%) and boys (75%). A total of 15% of the children were overweight (26.8% of females and 8.3% of males) while another 4.4% were obese (2.4% of females and 5.6% of males).

140 A relatively small percentage (8.8%) of the children was underweight for age (4.9% and  
 141 11.1% of the 72 boys and 41 girls, respectively) (Table 5 deleted).  
 142 The distribution of BMI according to different age groups was the following: among  
 143 preschoolers 15.1% were underweight, 75.5% had normal weight, 5.7% were overweight  
 144 and 3.8% were obese; among children 6 years old or more, 3.3% were underweight, 68.3%  
 145 had normal weight, 23.3% were overweight and 5% were obese (Table 6 deleted).  
 146 The prevalence of deficiency, insufficiency and normal vitamin D serum level according to  
 147 BMI groups is reported in Table 7 (deleted).  
 148 Rates of hypovitaminosis D [25(OH)D < 30 ng/ml] were higher in children with low or normal  
 149 weight (90% and 74.1% respectively) than in overweight and obese children (64.7% and  
 150 60% respectively), but this trend did not reach statistical significance ( $P = 0.4661$ ). No  
 151 correlation was found between the two variables ( $p=n.s.$ ). We must take into account that 7  
 152 among 9 of underweight children with hypovitaminosis D were withdrawn in January-April ( $n$   
 153 = 6) or in October-December ( $n = 1$ ) (see unit 3.6).

### 154 3.1.6 Ethnicity

155  
 156 The population of the study was composed by children of different ethnicity: 91 of them were  
 157 Italian (80.5%), 10 were African (8.8%), 9 were European (7.9%); Albanian (3 children),  
 158 Moldovan and Romanian (2 children each), Czech and Spanish (1 child each); 2 were  
 159 American (Central and South), and one child was Chinese. One of them was an adopted  
 160 boy.  
 161 Non-Italian children had a higher risk for 25(OH)D < 10 ng/ml vitamin D deficiency if  
 162 compared with Italian children: 22.7% versus 3.3% ( $P = 0.0056$ ) (Table 4 previously Table  
 163 8).  
 164  
 165  
 166

167 Table 4. Relationship between ethnicity and 25(OH)D level vitamin D status


	All		25(OH)D level					
			Deficiency < 10 ng/ml		Insufficiency 10-29 ng/ml		Normal ≥ 30 ng/ml	
	N	% col	N	% row	N	% row	N	% row
Italian	91	80.5	3	3.3	62	68.1	26	28.6
Non Italian	22	19.5	5	22.7*	13	59.1	4	18.2
All	113	100.0	8	7.1	75	66.4	30	26.5

169 Chi square:  $P = 0.0056$ .

### 170 3.1.7 Skin colour

171  
 172 The distribution of skin colour according to vitamin D status showed that there was a  
 173 statistically significant relation between non-white skin and 25(OH)D < 10 ng/ml vitamin D  
 174 deficiency ( $P = 0.0361$ ) (Table 9 deleted, Fig. 1).  
 175  
 176  
 177

178



179

180

181

182

183

184

185

186

187

188


189

**Fig. 1. Vitamin D status according to skin colour**

*\*Non-white children had a significantly higher rate of deficiency than white children (25.0% versus 5.0%;  $P = 0.0361$ ).*

### 3.1.8 Period of blood withdrawal

There was a seasonal effect on 25(OH)D levels that is represented in Figure 2.



190

191

192

193

**Fig. 2. 25(OH)D level according to period of blood withdrawal.**

*Bend line: Quarterly trend. Broken line: monthly level.*

*Analysis of variance:  $P = 0.0003$*

194

195

196 Children with 25(OH)D < 10 ng/ml were identified only in the first and in the second quarter  
 197 of the year. When serum samples were withdrawn between January and March only 11.8%  
 198 (4/34) of children showed a normal 25(OH)D level, while in 76.5% (26/34) it was 10-29 ng/ml  
 199 insufficient and in 11.8% (4/34) it was < 10 ng/ml deficient. In the second period (April-June),  
 200 a normal level of 25(OH)D was found in 26.7% of children (12/45); 25(OH)D was 10-29  
 201 ng/ml in 64.4% of them (29/45) were insufficient and it was < 10 ng/ml in 8.9% (4/45) were  
 202 deficient. Only samples withdrawn in the third quarter (July-September) showed mean  
 203 concentrations of serum 25(OH)D  $\geq$  30 ng/ml. The higher rate of children with normal  
 204 25(OH)D, 50.0% (8/16), was found in the third period (July-September); the remaining 50%  
 205 had 25(OH)D 10-29 ng/ml was insufficient. In the fourth quarter (October-December) 33.3%  
 206 of children had normal 25(OH)D (6/18) and the remaining 66.7% had 25(OH)D 10-29 ng/ml  
 207 was insufficient (12/18). A similar distribution was found in children whose sample was  
 208 collected between September and December. Analysis of variance showed a statistically  
 209 significant difference between periods of blood withdrawal ( $P = 0.0003$ ).

210

### 211 3.1.9 Serum 25(OH)D and other laboratory tests

212 Serum 25(OH)D test was included in a larger panel of lab tests together with haemoglobin,  
 213 white blood cells, serum iron, serum cholesterol and others. Parathyroid hormone (PTH),  
 214 alkaline phosphatase, calcium and phosphorus were done only in a few children because  
 215 this was not a structured study and its purpose wasn't to analyze biochemical consequences  
 216 of hypovitaminosis D.

217 Eighty-two percent of the children with no alteration in other parameters had 25(OH)D under  
 218 30 ng/ml compared with 66.7% of children with at least one value altered, among those  
 219 considered, this difference being not statistically significant ( $P = 0.0668$ ).

220

### 221 3.1.10 Comorbidity

222 In this study we considered the relationship between 25(OH)D stratification groups and the  
 223 presence of comorbidity (Table 5 previously 10).

224

225

226 **Table 5.** Distribution of comorbidity and vitamin D status

	25(OH)D < 10 ng/ml		25(OH)D 10-29 ng/ml		25(OH)D $\geq$ 30 ng/ml		All	
	N	% row	N	% row	N	% row	N	% row
No comorbidity	5	10.2	30	61.2	14	28.6	49	43.4
Comorbidity	3	4.7	45	70.3	16	25.0	64	56.6
All	8	7.1	75	66.4	30	26.5	113	100.0

227 **Chi-Square:  $P = 0.4335$**

228

229 Only one out of our eight children with 25(OH)D < 10 ng/ml deficiency had a chronic illness  
 230 (he was a coeliac and asthmatic boy). Among 45 children with 25(OH)D < 20 ng/ml there  
 231 were 7 patients affected by a chronic disease (coeliac disease 3, epilepsy taking specific  
 232 drugs 2, congenital hypothyroidism 1). The combination of chronic disease or non-white skin  
 233 could identify 4/8 children (50%) with 25(OH)D < 10 ng/ml deficiency and 13/45 children



(29%) with 25(OH)D < 20 ng/ml. The same combination identified 7/12 children (58%) with 25(OH)D 10 ng/ml or less that is another definition of deficiency [10].

The most common comorbidity was atopic disease, that affected 22 children; the other comorbidities were neuropsychiatric disorders (6 children), anemia (5 children), metabolic disorders (5 children), skeletal disorders (4 children), chromosomal syndromes (3 children), hypersensitivity contact dermatitis (2 children), tumors (2 children), precocious puberty (2 children), other conditions (10 children).

### **3.1.11 Multivariate analysis**

Ethnicity and period (quarter of the year) of blood withdrawal, significant at univariate level, were also tested by multivariate analysis to verify the association with hypovitaminosis D. Both non-Italian ethnicity ( $P = 0.029$ ) and period of blood withdrawal ( $P = 0.0062$ ) were significantly associated to lower level of 25(OH)D.

## **3.2 Discussion**

We studied a group of 113 children living in a rural area of Northeastern Italy (45°N latitude). All patients attended a single pediatrician office and had blood withdrawn for measurement of serum 25(OH)D, that is the best indicator of overall vitamin D status, because it reflects total vitamin D from dietary intake and sunlight exposure, as well as the conversion of vitamin D from adipose stores in the liver [21-24]. Prevalence of hypovitaminosis D in our region is similar to others of “vitamin D winter area” (latitudes above 37 degrees North and South) [8, 9, 25].

### **3.2.1 Vitamin D status in children**

As reported in the literature, patients with vitamin D deficiency are much less frequent than those with insufficiency. Our prevalence of 7.1% deficient children is similar to that of general US population reported in NHANES Study 2001-2004 that found a rate of 9% [26]. It is also remarkable that two among three of our children had an insufficient vitamin D status, that is about the same as that shown in US children in the study mentioned above [26]. Moreover, about 40% 36% of our patients had serum 25(OH)D < 20 ng/ml, that is the cut-off recently suggested to diagnose vitamin D deficiency [10, 27-31]. A similar prevalence of deficiency serum 25(OH)D < 20 ng/mL – 42% and 34.7% respectively – was reported in a sample of healthy US adolescents [11] and in a cohort of Spanish patients [12]. The importance of these data is capital because 25(OH)D is a fundamental prehormone [32]; in fact Vitamin D receptor is expressed in nearly all tissues and affects the transcription of over 900 genes in human cells [33], being critically important for the development, growth, and maintenance of a healthy body in humans.

### **3.2.2 Risk factors for hypovitaminosis D**

In our group of children factors consistently associated with low vitamin D status included blood withdrawal in January-June, non-white skin colour and non-Italian ethnicity. Mean level of 25(OH)D was < 30 ng/ml in October-December and < 20 ng/ml in January-April. In other studies a high risk of hypovitaminosis D was found only in winter season [8, 34], while in a population based study blood levels taken December-May in Great Britain had an Odd Ratio of 6.5 for vitamin D insufficiency in UK children [9]. The angle of sunlight during winter above 37 degrees latitude North is insufficient to stimulate the skin to make vitamin D, and this effect drifts in the early spring. In our group non-Italian children had a significantly higher rate of vitamin D deficiency and this was correlated with the presence of non-white skin, as reported in the literature [8, 9]. In



fact skin pigmentation blocks the absorption of ultraviolet sunlight and decreases vitamin D synthesis from photo-conversion of cutaneous 7-dehydrocholesterol. Nevertheless, it cannot be ruled out that the socioeconomic status rather than ethnicity or skin colour was responsible for this difference.

Another risk factor of hypovitaminosis D is lack of sun exposure; we can find a typical example of this condition in adopted children due to prolonged institutionalization in pre-adoptive period [35].

The relationship between hypovitaminosis D and BMI is controversial; for example it may be associated with overweight [8-10] or underweight [36], and the latter is also our experience: in fact 9 out of 10 underweight children had 25(OH)D < 30 ng/ml.

Some authors had found higher rates of hypovitaminosis D in children with greater fat mass and higher BMI [8, 34, 37, 38]. In a study the prevalence of vitamin D deficiency increased significantly with BMI: it was 21% in healthy-weight, 29% in overweight, 34% in obese, and 49% in severely obese children [39]. Because of vitamin D deposition in fatty tissue, it was suggested that the larger storage capacity in obese people may prevent 25(OH)D from circulating in the bloodstream [40]. Mechanism is probably more complex and includes other factors besides the decreased bioavailability, because hypovitaminosis D really precedes overweight [41]. Vitamin D status in underweight children with low serum 25(OH)D is probably different because these patients have both the circulating and the stored vitamin D levels poor. This depends in part on the limited quantity of vitamin D precursors in the skin due to the slim layer of fatty tissue. The fact that not always underweight children have hypovitaminosis D may be due in part to the different use of vitaminic supplements: in fact children classified as being underweight or at risk of underweight had a higher prevalence of use of vitamin D-containing supplements than did children classified as at risk of overweight or overweight [42].

Some authors reported that adolescence is significantly associated with hypovitaminosis D [8, 9], while in our experience preschool children had more frequently a level of 25(OH)D < 10 ng/ml comparing with older children, even if there was not statistical significance. The high frequency of vitamin D deficiency in preschool children This trend may depend in part on the higher rate of underweight children in this class of age (see Table 6) and in part on the fact that little children rarely go outdoors without parents surveillance.

Other factors (e.g., estimated sunlight exposure, sunscreen use, diet, etc.) were not analyzed because the purpose of our "field-study" was to examine vitamin D status of children in a primary care setting and not to determine in detail the cause of the observed levels.

### **3.2.3 To screen or not**

Dr. Holick suggested that only a few categories of patients should be screened for vitamin D deficiency/insufficiency and monitored for their 25(OH)D concentration while being treated with vitamin D, that are patients with malabsorption (inflammatory bowel disease, cystic fibrosis), and liver and kidney diseases; patients taking antiseizure medications, glucocorticoids, rifampicin, isoniazid or AIDS medications, patients with primary hyperparathyroidism or chronic granulomatous disorders [27]. On this basis, taking into account also coeliac disease, we would have 7 out of 8 children with vitamin D deficiency and 38/45 with 25(OH)D < 20 ng/ml undiagnosed. Considering the combination of chronic disease or non-white skin we would have missed 50% of children with 25(OH)D < 10 ng/ml and 71% of children with 25(OH)D < 20 ng/ml. Also lab tests other than 25(OH)D weren't predictive of hypovitaminosis D: in fact a level of 25(OH)D < 30 ng/ml was relatively more frequent in children with no alteration in other lab tests. So neither clinical/demographic factors nor lab tests other than 25(OH)D could identify all children with hypovitaminosis D. Dr. Holick suggested that it could be more cost-effective to implement a vitamin D

supplementation program for all children and adults than to measure everybody's serum 25(OH)D [27] but the question now is how much vitamin D should be given.

### **3.2.4 How much vitamin D must be given and when**

The dose to give as supplement depends both on baseline and target serum level of 25(OH)D [43]. There is no general agreement on which level is needed for an optimal health condition, even if some medical societies had stated 20 ng/ml as the minimum desirable serum 25(OH)D level during evolutive age [28-31].

A Recommended Dietary Allowance (RDA) of 600 IU per day ensures a 25(OH)D serum level of 20 ng/ml to 97-98% of population aged 1-18 years [30]. Moreover some authors suggested higher levels: 30 ng/ml [44, 45], 40 ng/ml [31, 46] or even 60 ng/ml to obtain the maximum health benefits [47, 31]. Some recommendations are derived from adults data, because little is still known about requirements of vitamin D in children. Some studies showed that 25(OH)D levels < 30 ng/ml are associated with bone demineralization [48] and an increase in PTH [49]. The target for optimal health depends on personal situations or life periods; the question is far more complex because the relationship between 25(OH)D level and effect on health can be nonlinear: emerging evidence suggests a U-shaped curve for several outcomes related to vitamin D in adults [30] and also in pediatric age [50]. In fact serum 25(OH)D levels above 50 ng/ml had potential risks for some outcomes [30].

Assuming that most of our children had less or much less than the minimum desirable [44] and considering that for every 100 IU of vitamin D taken in, there is an increase of rough 1 ng per milliliter in the serum level of 25(OH)D [51, 52], recommended doses of 600 IU per day [30, 53] probably offer no advantage when treating children with 25(OH)D < 10 ng/ml and even < 20 ng/ml. NHANES 2001-2006 data has demonstrated that 10% of children taking vitamin D supplements at doses 100-400 IU per day had a 25(OH)D level < 20 ng/ml and over 50% of children had a level < 30 ng/ml [1]. In an intervention study, 38% of peripubertal Finnish girls with vitamin D<sub>2</sub> supplementation approximately 800 IU daily from October to January had moderate hypovitaminosis D [25(OH)D 8-15 ng/ml] after treatment [54].

There is accumulating evidence to suggest that vitamin D intake should be in the range of at least 800-1,000 IU per day [50, 55]. The evidence-based recommendations by the Endocrine Society's Clinical Practice Guidelines stated 400-1,000 IU for children, 1,500-2,000 IU for adults to maintain 25(OH)D concentrations of 40-60 ng/ml for preventing and treating vitamin D deficiency [49, 56]. Dr. Holick recommends 1,000-1,500 IU/day for children 1-10 years of age, and 1,500-2,000 IU/day for teenagers [22].

While the Institute of Medicine Committee in 2011 recommended 600 IU vit D per day from 1 to 70 years of age [30] and also Italian LARN revised 2012 recommended 600 IU per day as for children aged 1-17 years as for adults aged 18-74 years [57], recent guidelines of the Italian Society for Osteoporosis, Mineral Metabolism and Bone Diseases (SIOMMMS) [58] highlighted that healthy adults with no effective sun exposure should be supplemented with 1,200 IU vitamin D per day. In our opinion children living in Northeastern Italy need about 1,500 IU cholecalciferol per day from October to April, that is about a half of tolerable (safe) upper limit for vitamin D [30]; higher doses are probably necessary in non-white skinned children.

Further studies are necessary in order to confirm vitamin D needs of Northern Italian children.

## **4. CONCLUSION**

We noted a high incidence of hypovitaminosis D in asymptomatic children living in Northeastern Italy without risk factors. In our region cholecalciferol supplementation should be implemented for all children between October and April.

392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442

**ACKNOWLEDGMENTS**

Our grateful thanks to Giovanna Magni (Statistical Doctor) for her statistical analysis and brainstorm.

**COMPETING INTERESTS**

Authors have declared that no competing interests exist.

**AUTHORS' CONTRIBUTION**

This work was carried out in collaboration between all authors. Author SM conceptualized and designed the study, did acquisition and analysis of the data, drafted the initial manuscript, and approved the final manuscript as submitted. Author CB participated in the collection of the data, in the drafting of manuscript and final approval of the manuscript. Author DT participated in the interpretation of the data, critical evaluation of the manuscript and final approval of the manuscript. All authors read and approved the final manuscript.

**CONSENT**

Not Applicable

**ETHICAL APPROVAL**

Not Applicable

**REFERENCES**

1. Mansbach M, Ginde AA, Camargo CA. Serum 25-Hydroxyvitamin D levels among US children aged 1 to 11 years: do children need more vitamin D? *Pediatrics*. 2009;124:1404-1410. PMID:19951983.
2. Rath N, Rath A. Vitamin D and child health in the 21st century. *Indian Pediatr*. 2011;48:619-625. PMID:21918267.
3. Prentice A. Vitamin D deficiency: a global perspective. *Nutr Rev*. 2008;66 Suppl:S153-164. PMID:18844843.
4. Pettifor JM. Vitamin D &/or calcium deficiency rickets in infants & children: a global perspective. *Indian J Med Res*. 2008;127:245-249. PMID:18497438.
5. Lips P. Worldwide status of vitamin D nutrition. *Journal of Steroid Biochemistry and Molecular Biology*. 2010;121:297-300. PMID:20197091
6. Looker AC, Pfeiffer CM, Lacher DA, Schleicher RL, Picciano MF, Yetley EA. Serum 25-hydroxyvitamin D status of the US population: 1988-1994 compared with 2000-2004. *Am J Clin Nutr*. 2008;88:1519-1527. PMID:19064511
7. Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *Am J Clin Nutr*. 2008;87:1080S-1086S. PMID:18400738.
8. Rovner AJ, O'Brien KO. Hypovitaminosis D Among Healthy Children in the United States. A Review of the Current Evidence. *Arch Pediatr Adolesc Med*. 2008;162:513-519. PMID:18524740.
9. Absoud M, Cummins C, Lim MJ, Wassmer E, Shaw N. Prevalence and Predictors of Vitamin D Insufficiency in Children: A Great Britain Population Based Study. *PLoS ONE*. 2011;6(7):e22179.doi:10.1371/journal.pone.0022179. PMID:21799790.

10. Saintonge S, Bang H, Gerber LM. Implications of a New Definition of Vitamin D Deficiency in a Multiracial US Adolescent Population: The National Health and Nutrition Examination Survey III. *Pediatrics*. 2009;123:797-803. PMID:19255005.
11. Gordon CM, DePeter KC, Feldman HA, Grace E, Emans SJ. Prevalence of vitamin D deficiency among healthy adolescents. *Arch Pediatr Adolesc Med*. 2004;158:531-537. PMID:15184215.
12. Gordon CM, Feldman HA, Sinclair L, Williams AL, Kleinman PK, Perez-Rossello J, Cox JE. Prevalence of vitamin D deficiency among healthy infants and toddlers. *Arch Pediatr Adolesc Med* 2008; 162: 505-512.
12. Weng FL, Shults J, Leonard MB, Stallings VA, Zemel BS. Risk factors for low serum 25-hydroxyvitamin D concentrations in otherwise healthy children and adolescents. *Am J Clin Nutr*. 2007;86:150-158. PMID:17616775.
13. El-Hajj Fuleihan GE, Nabulsi M, Choucair M, Salamoun M, Hajj Shahine C, Kizirian A, Tannous R. Hypovitaminosis D in healthy schoolchildren. *Pediatrics*. 2001;107:e53. PMID:11335774.
14. Bener A, Al-Ali M, Hoffmann GF. Vitamin D deficiency in healthy children in a sunny country: associated factors. *Int J Food Sci Nutr*. 2009;60 suppl:560-570. PMID:18946796.
15. Lippi G, Montagnana M, Targher G. Vitamin D deficiency among Italian children [Letter] *Canadian Medical Association Journal*. 2007;177:1529-1530. PMID:18056611.
16. Marrone G, Rosso I, Moretti R, Valent F, Romanello C. Is vitamin D status known among children living in Northern Italy? *Eur J Nutr*. 2011 May 4 [Epub ahead of print]. PMID:21541731.
17. Chinellato I, Piazza M, Sandri M, Peroni D, Piacentini G, Boner AL. Vitamin D serum levels and markers of asthma control in Italian children. *J Pediatr*. 2011;158:437-441. Epub 2010 Sep 26. PMID:20870246.
18. Ozkan B, Hatun S, Bereket A. Vitamin D intoxication. *Turk J Pediatr*. 2012;54:93-98. PMID:22734293.
19. World Health Organization Expert Committee on Physical Status. The Use and Interpretation of Anthropometry. Physical Status: Report of a WHO Expert Committee: WHO Technical Report Series 854, WHO, Geneva, 1996.
20. Himes, JH and Dietz, WH. Guidelines for overweight in adolescent preventive services: Recommendations from an expert committee. *American Journal of Clinical Nutrition*. 1994;59:307-316. PMID:8310979.
21. Hollis BW. Circulating 25-hydroxyvitamin D levels indicative of vitamin D sufficiency: implications for establishing a new effective dietary intake recommendation for vitamin D. *J Nutr*. 2005;135:317-322. PMID:15671234.
22. Holick MF. The D-lemma: To screen or not to screen for 25-Hydroxyvitamin D concentrations. *Clinical Chemistry*. 2010;56:729-731. PMID:20348405.
23. Adams JS, Hewison M. Update in Vitamin D. *J Clin Endocrinol Metab*. 2010;95:471-478. PMID:20133466.
23. IOM report on calcium and vitamin D. Washington, DC: Institute of Medicine, 2010. (<http://www.iom.edu/vitaminD>)
24. Heaney RP. The vitamin D requirement in health and disease. *J Steroid Biochem Mol Biol*. 2005;97:13-19. PMID:16026981.
25. Webb AR, Kline L, Holick MF. Influence of season and latitude on the cutaneous synthesis of vitamin D<sub>3</sub> exposure to winter sunlight in Boston and Edmonton will not promote vitamin D<sub>3</sub> synthesis in human skin. *J Clin Endocrinol Metab*. 1988;67:373-378. PMID:2839537.
27. Ginde AA, Liu MC, Camargo CA Jr. Demographic differences and trends of vitamin D insufficiency in the US population, 1988-2004. *Arch Intern Med*. 2009; 169: 626-632. doi: 10.1001/archinternmed.2008.604.
26. Kumar J, Muntner P, Kaskel FJ, Hailpern SM, Melamed ML. Prevalence and associations of 25-hydroxyvitamin D deficiency in US children: NHANES 2001-2004.

496 Pediatrics. 2009;124:e362-70. doi:10.1542/peds.2009-0051. Epub 2009 Aug 3.  
 497 PMID:19661054.  
 498 27. Holick MF. Vitamin D deficiency. N Engl J Med. 2007;357:266-281. PMID:17634462.  
 499 28. Wagner CL, Greer FR. American Academy of Pediatrics Section on Breastfeeding;  
 500 American Academy of Pediatrics Committee on Nutrition. Prevention of rickets and vitamin D  
 501 deficiency in infants, children, and adolescents. Pediatrics. 2008;122:1142-1152.  
 502 PMID:18977996.  
 503 29. Misra M, Pacaud D, Petryk A, Collett-Solberg PF, Kappy M. Drug and Therapeutics  
 504 Committee of the Lawson Wilkins Pediatric Endocrine Society. Vitamin D deficiency in  
 505 children and its management: review of current knowledge and recommendations.  
 506 Pediatrics. 2008;122:398-417. PMID:18676559.  
 507 30. Ross AC, Manson JAE, Abrams SA, Aloia JF, Brannon PM, Clinton SK, et al. The 2011  
 508 Report on Dietary Reference Intakes for Calcium and Vitamin D from the Institute of  
 509 Medicine: What Clinicians Need to Know. J Clin Endocrinol Metab. 2011;96:53-58.  
 510 PMID:21118827.  
 511 31. Ad hoc Committee. Prophylaxis of vitamin D deficiency. Polish Recommendations 2009,  
 512 Endokrynol Pol. 2010;61:228-232.  
 513 32. Vieth R. Why "Vitamin D" is not a hormone, and not a synonym for 1,25-dihydroxy-  
 514 vitamin D, its analogs or deltanoids. Journal of Steroid Biochemistry & Molecular Biology.  
 515 2004;89-90:571-573. PMID:15225841.  
 516 33. Norman AW. From vitamin D to hormone D: fundamentals of the vitamin D endocrine  
 517 system essential for good health. Am J Clin Nutr. 2008;88:491S-499S. PMID:18689389.  
 518 34. Alemzadeh R, Kichler J, Babar J, Calhoun M. Hypovitaminosis D in obese children and  
 519 adolescents: relationship with adiposity, insulin sensitivity, ethnicity and season. Metabolism  
 520 Clinical and Experimental. 2008;57:183-191. PMID:18191047.  
 521 35. Cataldo F, Viviano E. Health problems of internationally adopted children. Ital J Pediatr.  
 522 2007;33:92-99.  
 523 36. Bener A, Hoffmann GF. Nutritional Rickets among Children in a Sun Rich Country.  
 524 International Journal of Pediatric Endocrinology. 2010;doi:10.1155/2010/410502.  
 525 PMID:21048925.  
 526 35. el Hag AI, Karrar ZA. Nutritional vitamin D deficiency rickets in Sudanese children. Ann  
 527 Trop Paediatr 1995; 15: 69-76.  
 528 37. Reinehr T, de Sousa G, Alexy U, Kersting M, Andler W. Vitamin D status and parathyroid  
 529 hormone in obese children before and after weight loss. Eur J Endocrinol. 2007;157: 225-  
 530 232. PMID:17656603.  
 531 38. Elizondo-Montemayor L, Ugalde-Casas PA, Serrano-González M, Cuello-Garcia CA,  
 532 Borbolla-Escoboza JR. Serum 25-hydroxyvitamin D concentration, life factors and obesity in  
 533 Mexican children. Obesity. 2010;18:1805-1811. PMID:20010726.  
 534 39. Turer CB, Lin H, Flores G. Prevalence of Vitamin D Deficiency Among Overweight and  
 535 Obese US Children. Pediatrics. doi: 10.1542/peds. 2012-1711. PMID:23266927.  
 536 40. Lenders CM, Feldman HA, Von Scheven E, Merewood A, Sweeney C, Wilson DM, et al.  
 537 Relation of body fat indexes to vitamin D status and deficiency among obese adolescents.  
 538 Am J Clin Nutr. 2009;90:459-467. PMID:19640956.  
 539 41. González-Molero I, Rojo-Martínez G, Morcillo S, Gutierrez C, Rubio E, Pérez-Valero V,  
 540 at al. Hypovitaminosis D and incidence of obesity: a prospective study. European Journal of  
 541 Clinical Nutrition. 2013 | doi:10.1038/ejcn.2013.48. PMID:23422920.  
 542 42. Picciano MF, Dwyer JT, Radimer KL, Wilson DH, Fisher KD et al. Dietary supplement  
 543 use among infants, children, and adolescents in the United States, 1999-2002. Arch Pediatr  
 544 Adolesc Med. 2007;161:978-985. PMID:17909142.  
 545 43. Heaney RP. The effect of vitamin D dose on bone mineral density. Osteoporos Int.  
 546 2012;23:789-790. PMID:22113323.



44. Vieth R. Why the minimum desirable serum 25-hydroxyvitamin D level should be 75 nmol/L (30 ng/ml). *Best Pract Res Clin Endocrinol Metab.* 2011;25:681-691. PMID: 21872808.

45. Vitamin D Expert Panel. Vitamin D Expert Panel meeting, October 11-12, 2001, Atlanta, GA: final report. Accessed 10 July 2007. Available: [www.cdc.gov/nccdphp/dnpa/nutrition/pdf/Vitamin\\_D\\_Expert\\_Panel\\_Meeting.pdf](http://www.cdc.gov/nccdphp/dnpa/nutrition/pdf/Vitamin_D_Expert_Panel_Meeting.pdf).

46. Cannell JJ, Hollis BW. Use of vitamin D in clinical practice. *Alter Med Rev.* 2008;13:6-20. PMID:18377099.

47. Garland CF, Gorham ED, Mohr SB, Garland FC. Vitamin D for cancer prevention: global perspective. *Ann Epidemiol.* 2009;19:468-483. doi: 10.1016/j.annepidem.2009.03.021. PMID:19523595.

48. Priemel M, von Demarous C, Klatte TO, Kessler S, Schlie J, Meier S, et al. Bone mineralization defects and vitamin D deficiency: histomorphometric analysis of iliac crest bone biopsies and circulating 25-hydroxyvitamin D in 675 patients. *J Bone Miner Res.* 2010;25:305-312. doi: 10.1359/jbmr.090728. PMID:19594303.

49. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Guidelines for preventing and treating vitamin D deficiency and insufficiency revisited. *J Clin Endocrinol Metab.* 2012;97:1153-8. doi: 10.1210/jc.2011-2601. Epub 2012 Mar 22. PMID:22442274.

50. Mullins RJ, Clark S, Wiley V, Eyles D, Camargo CA Jr. Neonatal vitamin D status and childhood peanut allergy: a pilot study. *Ann Allergy Asthma Immunol.* 2012;109:324-328. doi: 10.1016/j.anai.2012.07.009. Epub 2012 Sep 19. PMID: 23062387.

47. Heaney RP, Davies KM, Chen TC, Holick MF, Barger-Lux MJ. Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol. *Am J Clin Nutr* 2003; 77: 204-210.

51. Maalouf J, Nabulsi M, Vieth R, Kimball S, El-Rassi R, Mahfoud Z, et al. Short- and Long-Term Safety of Weekly High-Dose Vitamin D Supplementation in School Children. *J Clin Endocrinol Metab.* 2008;93:2693-2701. Published online 2008 April 29. doi: 10.1210/jc.2007-2530. PMID:18445674.

52. Abrams SA, Hawthorne KM, and Chen Z. Supplementation with 1000 IU vitamin D/d leads to parathyroid hormone suppression, but not increased fractional calcium absorption, in 4–8-y-old children: a double-blind randomized controlled trial. *Am J Clin Nutr.* 2013;97:217-223. First published November 14, 2012, doi: 10.3945/ajcn.112.046102. PMID:23151536

53. Abrams SA. Dietary Guidelines for Calcium and Vitamin D: A New Era. *Pediatrics.* 2011;127:566-568. PMID:21339264.

54. Lehtonen-Veromaa M, Möttönen T, Nuotio I, Irjala K and Viikari J. The effect of conventional vitamin D<sub>2</sub> supplementation on serum 25(OH)D concentration is weak among peripubertal Finnish girls: a 3-y prospective study. *European Journal of Clinical Nutrition.* 2002;56:431-437. DOI: 10.1038/sj/ejcn/1601330. PMID:12001014.

55. First Nations, Inuit and Metis Health Committee, Canadian Pediatric Society. Vitamin D supplementation: Recommendations for Canadian mothers and infants. *Paediatr Child Health.* 2007;12:583-589.

56. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Endocrine Society. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* 2011;96:1911-30. doi: 10.1210/jc.2011-0385. Epub 2011 Jun 6. PMID:21646368.

57. LARN. Livelli di Assunzione di Riferimento di Nutrienti ed energia per la popolazione Italiana. Revisione 2012. Accessed 9 June 2013. Available: [www.sinu.it/pubblicazioni.asp](http://www.sinu.it/pubblicazioni.asp).

58. Adami S, Romagnoli E, Carnevale V, Scillitani A, Giusti A, Rossini M, et al. Linee guida su prevenzione e trattamento dell'ipovitaminosi D con colecalciferolo. Guidelines on prevention and treatment of vitamin D deficiency. The Italian Society for Osteoporosis,



599 Mineral Metabolism and Bone Diseases (SIOMMMS). Reumatismo. 2011;63:129-147.  
600 Italian.  
601

602