

Community-acquired CTX-M β -lactamase enteric isolates in Abeokuta, Nigeria.

ABSTRACT

Aim: Incidence of community-acquire CTX-M β -lactamase genes is fast becoming a threat to livelihood of many residents of Abeokuta, Nigeria and has also trigger multiple resistance strains to most commonly used antimicrobial agents. Therefore, the emergence and spread of CTX-M β -lactamase gene in this community is studied.

Methodology: β -lactamase and ESBL producing enteric bacteria were identified and their antimicrobial susceptibility were tested against commonly used antibiotics. Mating activity of the isolates was evaluated and their plasmid was profiled while a CTX-M chromosomal gene was assayed by PCR.

Result: Of the 27.4% β -lactamase producers obtained, only 5.1% strains were ESBL and 2.8% harbour *bla*_{CTX-M} genes. High resistant rate of 91.0% was observed to commonly used antibiotics mostly by *Eschericia coli* and *klebsiella oxytoca*.

Conclusion: There is urgent need to curb the community spread of undetected mobile genetic elements of CTX-M-encoding genes among the residents in this community and if this persists, a possible outbreak of enteric infection caused by resistant organisms is eminent.

Key words: Community residents, ESBL, *bla*_{CTX-M}.

Introduction

CTX-M β -lactamases are mutant, plasmid-mediated β -lactamases with potent hydrolytic activity against cefotaxime which is now commonly encountered among numerous enteric bacteria isolates (1, 2). In recent past, increasing prevalence of bacterial pathogens producing CTX-M β -lactamases has trigger multiple resistance strains to most commonly used 3rd Generation cephalosporin (3). However, some CTX-M-type ESBLs may actually hydrolyze ceftazidime and confer resistance to this cephalosporin. It was noted that the same organism harbouring both CTX-M-type and SHV-type ESBLs or CTX-M-type ESBLs and AmpC-type β -lactamases, may express altered antibiotic resistance phenotype (4). Thus, organisms producing CTX-M-type β -lactamases typically are now know to have high minimum inhibitory concentration (MIC) range against cefotaxime, while ceftazidime MIC is usually in the apparently susceptible range (5).

Moreover, faecal colonisation with ESBL-producing isolates is fast becoming prevalent in our society while it is a common practice in the hospital setting for implementing epidemiological measures to curtail and control this spread. Nevertheless, the increasing rate of inpatients, outpatients and healthy residents colonised with cephalosporin resistant strains are now potential risk factor for acquisition and transmission of resistant CTX-M β -lactamases in many households (6). Thus, the relationship between antibiotic consumption and occurrence of CTX-M β -lactamases has not been well studied in this locality. Although, the prevalence of resistant strains causing community-acquired diarrhoea and other

enteric diseases raises speculation that oxyimino cephalosporins available in many patent stores were taking without regulated prescription. Therefore, the emergence and spread of antibiotic resistance bacteria with CTX-M β -lactamases in this community which has become a threat to effective treatment of enteric infection were studied.

Materials and methods

Community Sample collection: Fresh faecal samples of two hundred and fifteen (215) were randomly collected from the community residents in Abeokuta, who consented to participate in the study and who have not used any antimicrobial agents in the preceding two weeks either as therapy for gastro-intestinal complication or prophylaxis between August 2010 and March 2013. Their use of antimicrobial agents, feeding habit and hygiene in the last three months was also assessed. Informed consent of all individual involved in this study was obtained while they were assured of confidentiality in view of the intricacy and sensitivity of the study.

Isolation and identification: All the faecal samples collected were cultured on Salmonella-Shigella agar (Oxoid CM 123, UK) and MacConkey agar without salt (Oxoid CM 516, UK); and incubated at 37°C for 18-24hours. Each isolates obtained was identified according to according to World Health Organisation (WHO) manual for laboratory investigation of acute enteric infections (7).

Antibiotic susceptibility tests: Standard broth micro-dilution method was used to determine the MIC of each identified isolates against the commonly used antibiotics with dilution ranges; Ampicillin (0.5–64ug/mL), Amoxicilli/Clavulanate (0.5-32ug/mL), Gentamycin (0.5–64ug/mL), Ciprofloxacin (0.5–64ug/mL), Cefuroxime (1–64ug/mL), Ceftazidime (0.25–128 ug/mL), Tetracycline (0.25–6ug/mL), Cefotaxime (0.25–64ug/mL), Azithromycin (0.5-64 ug/mL) and Imipenem (0.25–16ug/mL). The reference strains *Escherichia coli* ATCC 25922 and *Staphylococcus aureus* ATCC 25923 were used to verify the quality and accuracy of the testing procedures. MIC of each isolates was determined according to NCCLS recommended guidelines (8).

Phenotypic detection of β lactamase and ESBL: Isolates showing MIC >8ug/mL to Cefuroxime, Ceftazidime and Cefotaxime were tested for β lactamase production by a starch-iodide paper acidometric method described by Odugbemi *et al*, 1977 (9). ESBL Double Disc Synergy Test (DDST) was performed according to recommended method and criteria of NCCLS (10). 0.5 McFarland standard of bacterial suspension was evenly spread onto the surface of the Mueller–Hinton agar plate and Ceftazidime (30 μ g), Ceftazidime + clavulante (30/10 μ g) Cefotaxime + clavulante (30/10 μ g) and cefotaxime (30 μ g) discs were placed 2cm apart on the plate in a straight line. A difference of $\geq$ 5 mm between the inhibition zones of the ceftazidime (30 μ g) plus clavulanic acid(10ug) and ceftazidime (30 μ g) disc alone, was taken to indicate an ESBL producer. *Escherichia coli* ATCC 25922 was used as controls.

Mating activity: The confirmed ESBL isolates were mated with susceptible *E. coli* recipient strains, ATCC 25567 (penicillin -resistant and lactose fermentation-positive strain). Overnight cultures of donor cells of identified ESBL isolates and recipient strains (*E. coli* ATCC 25677) were grown in 0.5% sucrose enriched Luria-Bertani broth at 37°C, after mixing together at a 1:10 ratio (donor to recipient)

and incubated for 3hours and then sub-cultured on MacConkey Agar plates supplemented with 6mg/ml ceftazime (Oxoid). Transconjugants growing in the selection plates were subjected to DDST to confirm transfer of β -lactamase genes and also examine for the co-transfer of other antibiotic resistance determinants present in the donor isolates.

Plasmid DNA transformation: DNA plasmid extracted by an alkaline lysis method (10) of 20ul was added to 5ml overnight enriched Luria-Bertani broth culture of cephalosporin susceptible *Escherichia coli* ATCC 25922 isolates in the presence of 2M Calcium chloride at 37°C for 6hours according to Smith *et al*, (1981) (11). The mixture was later sub-cultured on selection plate of Muller Hinton Agar supplemented with 6mg/L ceftazidime and incubated at 37°C for 18-24hours. Transformants growing in the selection plates were subjected to DDST.

Plasmid profiling: Extra-chromosomal plasmid DNA was extracted from each isolates according to the alkaline lysis method (10), (12) using 1000 uL of the bacteria cell suspension lysed in 1% SDS, 0.2M NaOH (pH 8.0) and precipitated in 150 uL of ice-cold 3.0 M Potassium Acetate, pH 5.5. The plasmids DNA obtained were then electrophoresed on 0.8% Agarose gel, stained with 14 μ l/g Ethidium bromide. The DNA was then photographed with Polaroid camera and viewed using UV trans-illumination. The molecular weights were determined according to Meyers *et al* (12) using standard DNA molecular weight marker of 0.12-23.1kbp (Roche Diagnostic GmbH).

PCR Amplification of *bla*_{CTX-M} gene: Chromosomal DNA template was prepared by boiling pure colonies of ESBL isolates which was suspended thoroughly in 1mL DNase and RNase-free water and boiled at 55°C for 10 minutes. After centrifugation, supernatant was used as template DNA. The template DNA was quantified by Thermoscientific Nanodrop Spectrophotometer of absorbance of 260nm filter and estimated. The extracted DNA template was amplified for CTX-M gene by Polymerase Chain Reaction (PCR) using specific oligodeoxynucleotides *bla*_{CTX-M} F 5' SCS ATG TGC AGY ACC AGT AA 3' and *bla*_{CTX-M} R 5' CCG CRA TAT CRT TGG TGG TG 3'. The PCR was performed in mix of 2.5pl 10X PCR buffer, 1.5pl 25mM MgCl₂, 0.5pl of 10mM dNTPs (dATP, dGTP, dCTP, dTTP), 0.2pl *bla*_{CTX-M} F of 2500pmole, 0.2pl of *bla*_{CTX-M} R of 2500pmole, 0.2pl Taq polymerase enzymes (*Thermus aquaticus*), 17.9pl distilled water and 2.0pl of each extracted template DNA to make final reaction volume of 25.0pl in a thermal cycler block (NYX Technik Inc; Model ATC401, USA). The PCR mixture was subjected to a 5 minutes initialization at 94°C, followed by 30 cycles of 45 s at 94°C, *denaturation* at 94°C for 60 seconds, *annealing* at 60°C for 60 seconds, *elongation* at 72°C for 60 seconds. Amplified products were separated by 100V electrophoresis in a 2% agarose gel containing 0.5ug/ml ethidium bromide for 30 minutes, after which they were viewed under ultraviolet light to determine the specific *bla* gene amplicon.

Statistical analysis: Risk factors for ESBL prevalence among the studied community was assessed by using Chi square to determine the significance of ESBL isolates at p- value <0.05 at confidence interval of 95%. Significance of the high MIC to the occurrence of *bla*_{CTX-M} gene was determined by Pearson test while the correlation coefficient was calculated by the method of least squares and high level of resistance among the isolates was determined at p<0.005.

Results

Molecular characteristic of study strains: Of 215 community residents screened for enteric strains producing ESBL, 11(5.1%) strains met NCCLS screening criteria as potential ESBL producer, with very high average molecular plasmid weight of 7.91kbp. Out of the 11 ESBL producers only 7 and 8 successfully transconjugated and transformed susceptible isolates respectively. Of the 11 identified ESBL producers obtained, only 6(2.8%) harbour *bla*_{CTX-M} genes.

Prevalence of resistant enteric isolates in the studied community: The broth dilution susceptibility test to various antimicrobial agents shows a very high resistant rate of 98.8% to cefuroxime by the beta-lactamase producers while 91.0% resistant was observed among the isolates to Ampicillin, cefotaxime and ceftazidime respectively. It is worth noting that *Eschericia coli* and *klebsiella oxytoca* show a high resistant of 31.6% to most antimicrobial agents and 5.2% and 7.8% to iminepem respectively.

Table 1; Molecular characteristics of enteric isolates obtained from the community residents

Enteric isolates	Total isolates obtained n(%)	beta-lactamase n(%)	ESBL isolates n(%)	Plasmid size ($\pm$ 0.5kbp)	Conjugants (n=11)	Transformants (n=11)	<i>bla</i> _{CTX-M} n(%)
<i>Eschericial coli</i>	62(29.1)	12(5.6)	5(8.1)	8.5	2	3	2(0.9)
<i>Klebsiella oxytoca</i>	37(17.4)	9(4.2)	1(2.7)	10.2	1	1	1(0.5)
<i>Salmonella spp</i>	14(6.6)	8(3.6)	1(7.1)	7.5	1	0	1(0.5)
<i>Citrobacter freundii</i>	8(3.8)	5(2.3)	1(12.5)	6.8	0	1	0(0.0)
<i>Enterobacter cloacae</i>	22(10.3)	6(2.8)	1(4.6)	6.3	1	1	0(0.0)
<i>Pseudomonas</i>	39(18.3)	10(4.7)	1(2.4)	7.4	1	1	1(0.5)
<i>Proteus mirabilis</i>	31(14.6)	9(4.2)	1(3.2)	8.7	1	1	1(0.5)
<i>Shigella spp</i>	2(0.9)	0(0.0)	0(0.0)	0.0	0.0	0.0	0(0.0)
Total	215(100.0)	59(27.4)	11(5.1)	7.91	7	8	6(2.8)

Figure 1; *bla*_{CTX-M} genes recovered from the ESBLs positive isolates on Agarose gel




Table 2; Percentage resistance of ESBL-positive strains to commonly used drugs.

	Antimicrobial agents (MIC >8ug/ml)									
	AMP	AMC	CFX	AZT	CPX	IMP	CTX	TET	CFZ	GN
<i>Enteric isolates</i>	n(%)									
<i>Escherichia coli</i>	12(31.6)	5(13.0)	12(31.6)	3(7.8)	2(5.2)	2(5.2)	12(31.6)	2(5.2)	11(28.6)	2(5.2)
<i>Klebsiella oxytoca</i>	12(31.6)	3(7.8)	12(31.6)	8(20.8)	7(18.2)	3(7.8)	12(31.6)	6(15.6)	12(31.6)	6(15.6)
<i>Salmonella spp</i>	2(5.2)	1(2.6)	2(5.2)	0(0.0)	2(5.2)	0(0.0)	2(5.2)	2(5.2)	2(5.2)	0(0.0)
<i>Citrobacter freundii</i>	3(7.8)	1(2.6)	4(10.4)	2(5.2)	2(5.2)	0(0.0)	3(7.8)	0(0.0)	3(7.8)	0(0.0)
<i>Enterobacter cloacae</i>	1(2.6)	0(0.0)	2(5.2)	0(0.0)	1(2.6)	0(0.0)	2(5.2)	0(0.0)	2(5.2)	1(2.6)
<i>Pseudomonas aeruginosa</i>	3(7.8)	1(2.6)	3(7.8)	1(2.6)	1(2.6)	0(0.0)	2(5.2)	1(2.6)	3(7.8)	1(2.6)
<i>Proteus mirabilis</i>	2(5.2)	1(2.6)	3(7.8)	1(2.6)	2(5.2)	1(2.6)	2(5.2)	2(5.2)	2(5.2)	1(2.6)
Total	35(91.0)	12(31.6)	38(98.8)	15(39.0)	17(44.2)	6(15.6)	35(91.0)	13(33.8)	35(91.0)	11(28.6)

Key: AMP-Ampicillin, AMC-Amoxicilli/Clavulanate, CFX-Cefuroxime, AZT-Azithromycin, CPX-Ciprofloxacin, IMP-Iminepem, CTX-Cefotaxime, TET-Tetracycline, CFZ-Ceftazidime, GN-Gentamycin, n=total number of resistant isolate, %=percentage resistance

Discussion

The prevalence of CTX-M family of ESBLs is now becoming an emerging global health threat due to its unfavourable outcome in the treatment of common enteric infectious diseases in the community

and among the hospital patients. It was clearly observed from the study that out of 27.4% beta-lactamase producing enteric flora obtained from the community residents, only 5.1% were ESBL producers. This rate was lower to 11.4% recorded in Enugu, Enugu State, South-East Nigeria (13); and also 25% rate in Lagos, Nigeria (14). Although there is no report of ESBLs prevalence in any community in Nigeria, but similar study in Cambodia recorded 36.6% community-acquire ESBLs (15). Our findings suggest that there is undetected dissemination of community-acquired *bla*_{CTX-M} genes among this community residents and this could be elicited by self-medication and use of unprescribed antibiotics which is a common practice in this locality. Conjugative DNA plasmid obtained from the community resident further indicate a potential reservoir of antibiotic resistance bacteria that could contain a pool of mobile genetic elements, which could be readily disseminated to other human pathogens.

Therefore, transfers of naked resistant DNA plasmids encoding β -lactamase mostly decrease antibiotic susceptibility to commonly use antibiotics, thereby causing economic loss to the society and prolonged treatment of infection. High transformation achieved by the ESBL isolates from both the community and the hospital, suggest a very high frequency of resistant gene transfer between the same bacteria specie or other species. This transformation process could be promoted by constant release of naked DNA by dying bacteria which can be taken up by other isolates through unhygienic eating habit of most residents in poorly located food joints, restaurants and street food vendors and sachet-water supplied for drinking that usually contain this naked DNA genes (16, 17), which has a high propensity for transforming a susceptible isolates. This would definitely prone these residents to live in grave danger of increasing antibiotic resistant resulting to poor therapeutics, causing increasing morbidity.

Highly diverse antibiotics resistance rates against all cephalosporin, such as ceftazidime, cefuroxime and cefotaxime with MIC >16 μ g/mL; represent a very critical situation as compared to investigations from other regions of the world reporting resistance towards first, second and third generation of cephalosporin (18,19). The spread of antimicrobial resistance among bacterial pathogens to most antibiotics in this locality is fast becoming an important fatal challenge for the community and health institutions due to unguided use of over-the-counter drugs mostly antibiotics. Unfortunately, data regarding bacterial resistance to antimicrobial agents is limited in South-western, Nigeria.

The cause of this upsurge in resistant of community-acquired ESBL-producing organisms is not yet clear, but associations with foodstuffs, animal consumption of antibiotics, and frequent patient contact with relatives during illness may be a major factor to consider.

However, some of the antibiotic resistance genes usually associated with mobile elements were also associated with chromosomal transmission of resistant gene which maintains the spread of CTX-M extended spectrum β -lactamases among pathogenic bacteria populations, even if the subjects are not treated with extended-spectrum cephalosporins, especially in countries where antibiotics usage is unregulated. Interestingly, *bla*_{CTX-M} clone could exhibit virulence traits which could be involved in colonization, infection, and persistence in humans (20,21). These factors simultaneously explain the spread and the persistence of community spread of the CTX-M-encoding genes among enteric isolates in this community.

Conclusion: Adequate detection of new resistance mechanisms of the emerging ESBL isolates is necessary for infection control in the community and prevention of possible outbreaks of resistant organisms.

Competing interest: Neither financial nor personal relationships with other people or organisations inappropriately influence our work in any way.

CONSENT: The Authors declared that informed consent was obtained personally from all individual subjects involved in this study and any interested individuals that met the criteria of the study. They were informed of the benefit of the study and freedom to withdraw if they wish as non-invasive method of fecal collection was done.

Reference

1. Tzouvelekis LS, Tzelepi E, Tassios PT, Legakis NJ. CTX-M-type beta-lactamases: an emerging group of extended-spectrum enzymes. *Int. J. Antimicrob. Agents.* 2000;14:137-142.
2. Yu WL, Pfaller MA, Winokur PL, Jones RN. Cefepime MIC as a predictor of the extended-spectrum beta-lactamase type in *Klebsiella pneumoniae*, Taiwan. *Emerg. Infect. Dis.*,20002; 8:522-524.
3. Poirel L, Kampfer P, Nordmann P. Chromosome-encoded Ambler class A beta-lactamase of *Kluyvera georgiana*, a probable progenitor of a subgroup of CTX-M extended-spectrum beta-lactamases. *Antimicrob. Agents Chemother.* 2002;46:4038-4040.

- 219 4. Yan JJ, Ko WC, Tsai SH, Wu HM, Jin YT, Wu JJ. Dissemination of CTX-M-3 and CMY-2
220 beta-lactamases among clinical isolates of *Escherichia coli* in southern Taiwan. J. Clin.
221 Microbiol. 2000;38:4320-4325.
- 222 5. Davis J. Inactivation of antibiotics and the dissemination of resistant genes. Science.
223 1994;264: 375-382.
- 224 6. Valverde A, Grill F, Coque TM, Pintado V, Baquero F, Cantón R. High rate of intestinal
225 colonization with extended-spectrum-beta-lactamase-producing organisms in household
226 contacts of infected community patients. J. Clin Microbiol. 2008;46(8):2796-9.
- 227 7. World Health Organisation. Manual for laboratory investigation of acute enteric infections,
228 CDD/83.3 1983. WHO, 121 Geneva, 27-Switzerland.
- 229 8. National Committee for Clinical Laboratory Standards. *Performance Standards for*
230 *Antimicrobial Susceptibility Testing: Twelfth Informational Supplement M100-S12*, 2002
231 NCCLS, Villanova, PA, USA.
- 232 9. Odugbemi TO, Hafiz S, McEntegart MG. Penicillase producing *Neisseria gonorrhoea*:
233 Detection by Starch Iodide Paper Techniques. Br, Med, J. 1997;11:500.
- 234 10. Birnboim HC and Doly JA. Rapid alkaline extraction procedure for screening recombinant
235 plasmid DNA. *Nucleic Acids Res.* 1979; 7:1513-1523.
- 236 11. Smith CJ, Markowitz SM, Macrina FL, 1981. Transferable tetracycline resistance in
237 *Clostridium difficile*. *Antimicrob. Agents Chemother.* 1981;19:997–1003.
- 238 12. Meyers, J.A., Sanchez, D., Elwell, L.P. and Falkows. Simple agarose gel electrophoretic
239 method for the identification and characterization of plasmids deoxyribonuclease acid. *J.*
240 *Bacterial*, 1976;1127: 1529-1537.
- 241 13. Iroha IR, Adikwu MU, Esimone CO, Aibinu I, Amadi ES. Extended spectrum Beta –Lactamase
242 (EBSL) in *E.coli* isolated from a tertiary hospital in Enugu state, Nigeria. Parkistan Journal of
243 Medical sciences. 2009; 25:2.
- 244 14. Aibinu IE, Olaegbulam VC, Adenipakan EO, Ogunsola FT, Odugbemi JO, Mee BJ. Extended
245 spectrum beta-lactamase enzymes in clinical isolates of Enterobacteria species from Lagos.
246 Nig J Chin Microbiol. 2003; 41:2197-2200.
- 247 15. Etienne R, Sopheak H, Sovannarith L, Valerie G, Frederic A, Jean-Louis S, Didier M,
248 Guillaume A. CTX-M beta-lactamase in *Escherichia coli* from community-acquired urinary tract
249 infections, Cambodia. Emerging infectious diseases. 2009; 15,5.
- 250 16. Stobberingh, EE, Arends J, Kostanje JA, Goessens WHF, Visser MR, Buiting AGM,
251 Ossenkopp YJ, Van Ketel RJ, Vanogtrop ML, Sabbe LJ, Voorn GP, Winter HL, Van Zeijl JH.

- 252 Occurrence of extended spectrum β -lactamases in Dutch hospitals. *Infection*.2000; 27, 348-
253 354.
- 254 17. Xiong, Z, Zhu D, Zhang Y, Wang F. Extended spectrum β -lactamases in *K. pneumonia* and
255 *E. coli* isolates. *Zhonghua Yi Xue Za Zhi*;2002;82, 1476-1479.
- 256 18. Jain A, Roy I, Gupta MK, Kumar M, Agarwal SK, 2003. Prevalence of extended spectrum β -
257 lactamase producing Gram-negative bacteria in septicemia neonates in a tertiary care
258 hospital. *Journal of Medical Microbiology*. 2003;52:42-45.
- 259 19. Guillaume A. CTX-M beta-lactamase in *Escherichia coli* from community-acquired urinary tract
260 infections, Cambodia. *Emerging infectious diseases*. 2009;15,5.
- 261 20. Akinduti PA, Akinbo JA, Adenuga WF, Ejilude O, Umahoin KO, Ogunbileje JO. Occurrence of
262 Extended-Spectrum β -lactamase Producing Enterobacteriaceae (ESBLPE) among Primary
263 School Pupil in Obafemi-Owode, Nigeria. *Journal of Life Sciences*. 2011; 5:340-344.
- 264 21. Sirot D, Sirot J, Labia R, Morand A, Courvalin P, Darfeuille-Michaud A, Perroux R, Cluzel R.
265 Transferable resistance to third-generation cephalosporins in clinical isolates of *Klebsiella*
266 *pneumoniae*: identification of CTX-1, a novel beta-lactamase. *J. Antimicrob. Chemother.*
267 1987;**20**:323-334.