

Original Research Article

BIOREMEDIATION OF INDUSTRIAL SOLID WASTES.

ABSTRACT

Aims: Use of *Pleurotus ostreatus* as a veritable tool in the bioremediation of empty fruit bunches, pulp and paper wastes from respective industries.

Place and Duration of Study: Samples of pure culture *Pleurotus ostreatus* was obtained from National Roots Crops Research Institute Laboratory Umudike, Umuahia and transported to the laboratory of Starline Nigeria Group of Companies Aba, both in Abia state, Nigeria where it was used to carry out experimental work from July to September, 2013.

Methodology: The two substrates (empty fruit bunch (EFB) and paper pulp wastes) were dried in the sun for one week and further dried in Gallen Kamp hot box oven (DGH-9053A) at 50 °C for 4hours. Samples were ground, sieved and powder obtained used for proximate composition determination. Approximately 1 cm² blocks of pure culture of *Pleurotus ostreatus* mycelium previously collected and maintained on Malt extract agar (MEA) was used to inoculate sterilized wheat grain already soaked in water for 36 h, mixed with 2.5% calcium carbonate and 1% gypsum during spawn preparation. The empty fruit bunches and paper wastes were reduced to smaller sizes of 2-5cm. The respective substrates were mixed with lime and soaked in water for 4h to obtain moisture level of 70%. The substrates (2kg) was bagged in heat resistant polythene and sterilized for 3hours at 80°C. Approximately 40g individual sterilized substrate was inoculated with 2% spawn adopting multilayered technique. After inoculation, holes were made on the bags to ensure aeration, kept inside a disinfected room at 25-28 °C. When *Pleurotus* mycelium had completely colonized the substrates, bags were removed, colonized substrate transferred to sunlight illuminated room at 28°C, moisture content of 75-85% and adequate watering of the mushroom for 21days. The process was repeated in combinations of other substrates like wheat, straw and sorghum respectively with each of the substrates (EFB and Paper Waste); all prepared at 70:30 % ratio respectively.

Results: Among the different treatments, Empty fruit bunch gave maximum yield of 201.5 g/kg of substrate on 14th day and total yield of 675.1g/kg of dry substrate. The nutritional content of *Pleurotus ostreatus* grown on different substrate and substrate combination ranges from 83.5-91.4%moisture, 18.3-23.5%protein and 6.04-6.86%ash contents. The combination of EFB + sorghum best supported the bioremediation process of Empty fruit bunch.

Conclusion: *Pleurotus ostreatus* could serve as bioremediation tool for industrial solid wastes like EFB and paper. It can also alleviate poverty by providing greater income for most unemployed youth at no cost.

Keywords: [Bioremediation, Empty fruit bunch, Nutritive content, paper waste, *Pleurotus ostreatus*]

1. INTRODUCTION

Over the years pollution and management of vast quantities of wastes generated by various industrial activities have been major problems experienced by developing countries. More challenging is the unsafe disposal of these wastes into the ambient environment and biodegradation problems associated with it. These have resulted to alternative means of reclaiming the environment via bioremediation, phyto-remediation and air pollution control. Bioremediation involves the use of microorganisms to breakdown complex materials into simple end products. These micro-organisms have evolved a host of enzymes that aid in biodegradation of natural products within the ecosystem. This microbial clean-up (Bioremediation) method removes a number of other harmful pollutants and perhaps is the most environmentally safe process used today (2,7,38).

Oil palm (*Elaeis guineensis*) tree is a valuable economic crop widely grown in Nigeria. Its Empty fruit bunch (EFB) is a major solid waste in palm oil industries with 20–25% EFB being generated for every tonne of oil palm fresh fruit bunch (FFB) processed in the mill (4, 37). EFB contains lignocellulosic materials which consist of 45-50% cellulose, 25-35% of hemicellulose and lignin (12, 14). Lignin and its derivatives can form highly toxic and recalcitrant compounds. The industrial application of EFB have not been reported to our knowledge, it is rather found littered everywhere in palm oil industrial areas in Nigeria hence this research. Studies have shown that 80% EFB is used as fuel in palm oil mills thereby generating great environmental hazard to the host communities and this practice has been discouraged (27). Other ways of disposing EFB like composting incur additional operational cost and technology. However, in the south east of Nigeria EFB is used locally to prepare local delicacies like abacha (sliced boiled cassava), ugba (oil bean salad), ukwa (breadfruit) and production of local black soap because of its high potassium content (24). The brown colored filtrate from the ash of EFB is used as a sauce for preparing a local delicacy known as 'Otong' which is used in eating meat and spicing soup in south-south region of Nigeria (37). Many researchers have reported the use of lignocellulosic source of EFB as substrate in cellulase production (3,30, 37) while few reports prior to this research is known to have used EFB as a substrate for the cultivation of *Pleurotus* spp (32,31, 34).

Another source of industrial solid wastes is generated from pulp and paper manufacturing plants. Pulp and paper are manufactured from lignocellulosic raw materials from wood, recycled paper and agricultural residues. These wastes are usually disposed through landfill and incineration processes (35). These methods of disposal could be harmful to ecosystem and cause diseases to man. Many studies have been reported on the treatment of pulp and paper mill effluents using biological method such as conventional aerobic, anaerobic treatment and use of white-rot fungi in the treatment (36, 39, 11). Recently, mushrooms are regarded as the most profitable and environment-friendly method for recycling of the vast lignocellulosic waste substrates (28, 20).

Pleurotus ostreatus belong to the class of *Basidiomycetes* known as 'oyster mushroom' and mostly cultivated in the tropics using micro molecules of cellulose and starch for their growth (5). The life cycle starts from spores and germinates to form a mass of branched hyphae of mycelia on a substrate under favourable conditions (13). They are edible mushroom also used for both bioremediation and medicinal purpose. The first successful attempt to grow *Pleurotus ostreatus* for human consumption was made by (15) in Germany. *Pleurotus* species are rich source of vitamin C, B-complex (thiamin, riboflavin, folic acid and niacin pantothenic acid and biotin), minerals (Ca, P, Fe, K, Cu and Na) and protein (33,22, 25,8). Also its high content in potassium: sodium ratio makes it an ideal food for patients suffering from hypertension and heart diseases. *Pleurotus* is a proficient lignin-degrading mushroom and can grow well on different types of lignocellulosic materials which offers one of the

most realistic and economic method for the bioremediation of agro-industrial wastes (6,10). This technology can equally control air pollution associated with burning of these solid wastes and reduces environmental pollution due to unutilized agricultural wastes. The introduction of Environmental Air Quality Regulation in Nigeria is yet to be fully implemented unlike some developing countries such as Malaysia that make use of alternative disposal or management method for EFB and other solid wastes hence this research on better ways of converting these industrial solid wastes to an economically viable material which would also help to alleviate poverty in the country. Therefore, the aim of this research is to use *Pleurotus ostreatus* as a veritable tool in the bioremediation of empty fruit bunches, pulp and paper wastes from industries.

2. MATERIAL AND METHODS

2.1 Sample and Pure Cultures Collection:

The Industrial wastes (EFB and paper) used as substrates were generated from Starline palm oil mill industries and Starline corrugated packaging Owerri respectively, both in Abia State, Nigeria. The pure culture of *Pleurotus ostreatus* was obtained from National roots Crops Research Institute Laboratory Umudike, Umuahia and was transported to the laboratory of Starline Nigeria Group of Companies Aba, both in Abia state, Nigeria. The cultures were maintained on 2% malt extract agar slants at 4 °C prior to use for microbiological examination.

2.2. Preparation of Substrates for proximate analysis

The two substrates (empty fruit bunch and paper pulp wastes) used were examined, debris removed, milled and used for proximate analysis as described by (13) with slight modification. The substrates were dried in the sun for one week and further dried in Gallen Kamp hot box oven (DGH-9053A) at 50 °C for 4 hours. Samples were ground using National model kitchen electric blender and sieved through a 1mm sieve to obtain a powdered processed sample, which was used for determination of the proximate composition.

2.3.1 Spawn preparation

Clean wheat grains were commercially sourced from a Ariaria local market at Aba, Abia State, Nigeria. Two kilograms of dry wheat grain was soaked in water for 36h to absorb about 65% moisture. The grain was mixed with 2.5% calcium carbonate and 1% gypsum and packaged in different clean white glass containers of about 500 ml volume (20) modified. Each glass container was three-quarter filled with spawn substrate. The packaged spawn substrate content was then sterilized in autoclave at 121° C for 30mins. The sterilized spawn substrates were allowed to cool and aseptically inoculated with approximately 1 cm² blocks of mycelium previously maintained on MEA. Thereafter, it was incubated at a temperature of 25 ± 2°C at 150 rev/min for 48h to quicken mycelia development.

2.3.2 Preparation of Substrates for *Pleurotus ostreatus* Cultivation

The empty fruit bunches were first reduced to smaller sizes in Hammer mill then mechanically chopped into 2-5cm long bits as described (26) while the carton (paper) wastes were cut to same size manually. The respective substrates were mixed with lime and soaked in water for 4hrs; excess water was drained and maintained at optimum moisture level of 70 percent. Approximately 2kg of the substrate was transferred into a heat resistant polythene bag of 100µm thickness (60 x 35 cm size) and stoppered with a cotton plug respectively. The substrates were sterilized for 3 hours, at temperature of 80°C in an ESPEC industrial oven (PHH-302) model. The sterilized substrates were allowed to cool overnight in preparation for inoculation. 2% spawn was used to inoculate approximately 40g sterilized substrate adopting multi-layered technique respectively. After inoculation, holes were made on the bags to ensure aeration and were kept inside a disinfected room house where the

temperature and humidity were maintained at room temperature of about 25-28 °C for quick colonization of the substrate (EFB and paper wastes) respectively. After *Pleurotus* mycelium had completely colonized the substrates, the bags were removed and colonized substrate transferred to sunlight illuminated room where temperature and humidity of 28°C and 75-85% were maintained through adequate watering of the mushroom for 21days. The process was repeated with a combination of wastes from wheat straw and sorghum respectively with each of the substrates (EFB and Paper Wastes); all prepared at 70:30 % ratio respectively.

2.4 Determination of pH

The pH of substrates was determined after milling and soaking 10g of each (EFB and paper wastes) in 50ml distilled for 10 minutes using pH meter (Orion 410A+ Thermo Electron Corporation model). Preparation of adjusted pH was carried out later using Citrate-Phosphate buffer solution 5.0, 5.5, 6.0 and 6.5 as prepared by (17).

2.5 PROXIMATE ANALYSIS

2.5.1 Protein Content

The Kjeldahl method was used for determination of protein content was as described by (21, 13) and multiplying the resultant nitrogen content with factor of 6.25. About 0.2g of the respective samples was mixed with 25 ml Conc H₂SO₄ and selenium (catalyst) in a digestion flask. The mixture was heated to clear solution in a fume cupboard. The clear solution (digest) was diluted to 100ml and subsequently used for analysis. Distillation process was carried out on 10mls digest mixed with 45% (w/w) NaOH solution in Kjeldahl apparatus. The distillate was collected into 10ml of 4% boric acid solution with 3 drops of mixture indicator (Methyl red indicator and bromocressol green). A total of 50ml of distillate was collected and titrated against 0.02N H₂SO₄ solution. This process was repeated in the blank.

The N₂ content was calculated using the formula below:

$$\% N_2 = 100 \times (N \times 14 \times V_1 (T-X)).$$

$$\frac{W \times 100 V_a}{N \times 14 \times V_1 (T-X)}$$

Where W = Weight of sample

N = Normality of the titrant

V₁ = Digest volume

V_a = Volume of digest distilled

T = Titration vol of sample

X = Titration vol.Blank

$$\% \text{ Protein} = \% \text{ Nitrogen} \times 6.25$$

153

154

2.5.2 Moisture content

156

The moisture content was determined by measuring using 1g of each substrate into separate previously weighed porcelain dish. It was dried in the oven at 105°C for 3hrs, cooled in a dessicator and re-weighed. Drying, cooling and weighing were repeated at 1 hr intervals until constant weight was achieved. The weight of moisture loss was determined and expressed as percentage of the sample measured. Similar process was repeated in the case of moisture content determination of *Pleurotus ostreatus*.

163

It was calculated using this formula;

165

$$\text{Moisture content \%} = \frac{\text{Weight of fresh sample} - \text{weight dry sample}}{\text{Weight of fresh sample}} \times 100$$

$$\frac{\text{Weight of fresh sample}}{\text{Weight of fresh sample}} \times 100$$

2.5.3 Determination of Total Carbohydrate

Total carbohydrate was determined using Anthrone Method as described by (16)

2.5.4 Ash content

The ash content method was carried out as described by (25, 18) using furnace incineration. Three grammes sample (substrate/ mushroom) was ashed in a Gallenkamp furnace in previously ignited and cooled porcelain dishes of known weight at 550°C for 6hrs. After cooling in desiccators, it was re-weighed. The weight of the ash obtained was expressed as a percentage of the weight of sample analyzed.

Calculation of percentage ash was based on the formula below:-

$$\% \text{ Ash} = \frac{100 (X_2 - X_1)}{X}$$

Where X_1 = Weight of empty porcelain dish
 X_2 = Weight of porcelain dish + ashes
 X = Weight of sample

2.5.5 Crude fibre determination

Crude fibres of both substrate and mushroom samples were respectively determined according to (13) method and calculated using the formulae below:-

$$\% \text{ Crude fibre} = \frac{W_2 - W_3}{W} \times 100$$

Where W_2 = Weight of empty porcelain dish + sample after boiling, washing and drying.
 W_3 = Weight of porcelain dish + sample after ashing
 W = Weight of sample

2.5.6 Yield and Biological efficiency

Total weight of the mushroom harvested from all the three flushes were weighed as total yield of mushroom. The biological efficiency (yield of mushroom per kg substrate on dry wt. basis) was calculated by the following formula Chang (9)

$$\text{B.E.}\% = \frac{\text{Fresh weight of mushroom}}{\text{dry weight of substrate}} \times 100$$

where B.E. is Biological Efficiency.

3. RESULTS AND DISCUSSION

3.1 GROWTH SUPPORT OF *PLEUROTUS OSTREATUS* ON NUTRITIONAL CONTENT OF INDUSTRIAL SOLID WASTES

The proximate composition of unsterilized and harvested (EFB and paper wastes) substrates are presented in Tables 1 and 2 respectively. The experiment was carried out to show

possible use of both nutritional and chemical contents of 100% industrial solid used during the period of this research. Paper wastes had higher moisture content reduction of 17.64% against 7.63% in EFB (Table 2). There was no significant difference in percentage reduction of protein content (15.1%; 15%) and carbohydrate content (1.78%; 1.71%) in both EFB and paper wastes nutritional utilization respectively. The result shows that the cultivation of *Pleurotus ostreatus* on these substrates had significant effect on the ash content. This is in accordance with a report (13) on improved ash content of substrates like sawdust, cotton waste, palm oil chaff and dry plantain leaf.

225

226 **Table 1: Proximate composition of the unsterilized substrates used in the cultivation**
227 **of *Pleurotus ostreatus***

Properties	EFB	Paper wastes
Lignin	16.86 ± 2.04	18.54 ± 0.04
CHO	67.50 ± 1.09	52.60 ± 2.03
MC	23.60 ± 1.2	1.70 ± 0.08
ph	6.08 ± 0.07	7.60 ± 1.07
N	1.09 ± 0.15	0.40 ± 0.05
Dry mass	56.85 ± 0.7	21.47 ± 2.16
Protein Content	2.78 ± 1.08	0.20 ± 0.09
Ash content	4.87 ± 0.09	5.60 ± 1.3
Hemicellulose	28.50 ± 0.7	23.60 ± 1.06
cellulose	52.43 ± 1.06	35.00 ± 1.08

228 • MC: Moisture content; CHO: Carbohydrate content, N: Nitrogen (Mean ± Std. Dev)
229 of triplicate samples, All properties in percentage except pH.

230

231 **Table 2: Proximate composition of substrates harvest of *Pleurotus ostreatus***

Properties	EFB	Paper wastes
Ligin	16.56 ± 1.04	18.2 ± 1.04
CHO	66.3 ± 1.45	51.7 ± 3.06
% MC	21.8 ± 1.23	1.4 ± 0.4
ph	4.65 ± 0.07	5.09 ± 1.07
N	0.53 ± 0.15	0.32 ± 0.05
Dry mass	58.25 ± 0.72	23.17 ± 1.16
Protein Content	2.36 ± 1.25	0.17 ± 0.88
Ash content	5.07 ± 0.09	4.8 ± 1.3
hemicellulose	27.77 ± 0.70	23.32 ± 1.25
cellulose	52.03 ± 1.06	34.6 ± 0.18

232 • MC: Moisture content; CHO: Carbohydrate content, N: Nitrogen, (Mean ± Std. Dev)
233 of triplicate samples, All properties in percentage except pH.

234 **3.2 Total yield of three flushes of *Pleurotus ostreatus***

235 The yield of *P. Ostreatus* cultivated on different substrates is represented in Table 3.
236 Maximum yield of *P. ostreatus* was revealed on use of EFB + Sorghum straw (732g/kg
237 substrate) while paper wastes supported the yield of *P. ostreatus* minimally (446.1g/kg
238 substrate). The range of substrates support for the cultivation of *P. Ostreatus* should be
239 attributed to its ability to synthesize relevant hydrolytic and oxidative extracellular enzymes
240 (1).

241

242 **Table 3: Effect of substrates on average yield of *Pleurotus ostreatus***

Substrate	Yield (g/kg)	dry substrate		Total Yield(g/Kg)
type	7 th day	14 th day	21 st day	
EFB	330.3 ± 1.57	201.5 ± 0.96	143.3 ± 0.61	675.1 ± 0.87

Paper wastes		216.7 ± 0.69	117.9 ± 1.08	111.5±0.75	446.1 ±1.48
EFB+ wheat mill		375 ± 1.18	221.2 ± 1.32	99.2± 1.06	695.4 ±2.04
Paper wheat mill	+	206 ± 0.58	184.5 ± 1.86	92.0 ± 2.05	482.5 ±1.51
EFB Sorghum mill	+	375 ± 1.18	197.2 ± 1.32	159.8 ± 1.61	732 ±0.83
Paper sorghum mill	+	268.0 ± 1.58	158.0 ± 1.86	70.0± 2.15	496 ±2.58

• (Mean ± Std. Dev) of triplicate samples

3.3 Nutritional content of *Pleurotus ostreatus*

The average nutritive contents and biological efficiency of the *Pleurotus ostreatus* harvested in three flushes on different substrates is shown in Table 4. *Pleurotus ostreatus* grown on EFB + Sorghum straw showed the highest nutritive content of 91.4% moisture, 23.5% protein, carbohydrates 55.8%, crude fiber 8.04% and biological efficiency of 73.2% while minimal nutritive content of 83.5% moisture, 18.3% protein, 48.4% carbohydrate, 6.25% crude fibre and B.E 44.6% was observed on mushroom (*Pleurotus ostreatus*) harvested from paper wastes. It was observed that nutritional contents of the mushroom grown on the co-substrate sorghum was enhanced confirming previous findings (28, 23) of 90 % and 90.4% report on moisture content of mushroom grown on Empty Fruit Bunch.

Table 4: Effect of substrates in Nutritional content of *Pleurotus ostreatus*

Substrates	%age Moisture Content	% protein	%age carbohydrates	%Crude fiber	%age ash content	%BE
EFB	89.6	20.4	53.1	7.86	6.82	67.5±1.02
Paper wastes	83.5	18.3	48.4	6.25	6.04	44.6±1.05
EFB + wheat straw	90.3	22.2	53.9	6.75	7.05	69.5±2.05
Paper + wheat straw	86.7	19.6	50.7	6.55	6.85	48.2±1.85
EFB+ Sorghum straw	91.4	23.5	55.8	8.04	6.77	73.2±1.07
Paper + sorghum straw	86.3	20.1	50.2	7.05	6.43	49.6±0.07

3.4 pH variation on yield of *Pleurotus ostreatus*

Effect of pH on the yield of *Pleurotus ostreatus* on EFB and paper for the three flushes is demonstrated in Fig. 1 and 2. *P. ostreatus* yield was highest for both substrates at pH 5.0 with total yield of 793.4g/kg and 514.5g/kg on EFB and paper wastes respectively. However, pH 6.0 shows lowest support on the yield on both substrates. This is contrary to a report by (19) on optimal growth condition of *P. ostreatus* mycelium at pH=6.0 using whey permeate). It can be inferred from this result that the many factors can affect the yield of *P. ostreatus* as reported by (1).




Fig. 1. Effect of pH on Yield of *Pleurotus Ostreatus* on EFB




Fig.2. Effect of pH on Yield of *Pleurotus Ostreatus* on paper wastes

4. CONCLUSION

Air pollution associated with incineration of Empty Fruit Bunch and paper waste makes it imperative to find an alternative for the control of pollution through bioremediation process. The result from this research work reveals that these industrial wastes that pose great danger to the larger society can turn to wealth through cultivation of *Pleurotus Ostreatus* on it. Similarly this will provide an employment opportunity to the unemployed youths in the society thereby reducing crime.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Andrej G., Mirjan v and Jure P. Cultivation Techniques and Medicinal Properties of *Pleurotus* spp. Food Technol. Biotechnol. 2007: 45 (3) 238–249. ISSN 1330-9862.
2. Adriaens, P and Hickey, W. J. In: Stone, D. L. (Ed), Biotechnology for the Treatment of Hazardous Waste. – Lewis Publishers. Ann Arbor, 1993: 97-120.
3. Akamatsu, I., Kobayashi, Y., Kamishima, H., Hassa, K., Mohd Yusoff, M.N., Hushin, M *et al.* Industrial Utilization of oil Palm by-products II: Thermomechanical pulping of empty fruit bunches, cellulose chemistry and technology, 1987: 21, pp 191-197.
4. Bakar R.A, Darus S.Z, Kulaseharan S, Jamaluddin N. Effects of ten year application of empty fruit bunches in an oil palm plantation on soil chemical properties. Nutr Cycl Agroecosyst. Springer Science+Business Media B.V. Evaluation of locally available substrates for cultivation of oyster mushroom (*Pleurotus ostreatus*) in Jimma, Ethiopia 2010
5. Banjo, (1998) Cultivation of Edible Mushroom The newsletter of Industrial Microbiologist of Nigeria, 4:3
6. Bano Z, Shasirekha MN and Rajarathan S. Improvement of the bioconversion and biotransformation efficiencies of the oyster mushroom (*Pleurotus sajor-caju*) by supplementation of its rice straw with oil seed cakes. Enzymes and microbial technology. 1993: 15: 985-989.
7. Beg, M. U., Saeed, T., Al-Muzaini, S., Beg, K. R. and Al-Bahloul, M. Distribution of Petroleum hydrocarbon in Sediment from Coastal Area Receiving Industrial Effluents in Kuwait. – Ecotoxicology and Environmental Safety 2003:54: pp 47-55.
8. Caglarirmak N, The nutrients of exotic mushrooms (*Lentinula edodes* and *Pleurotus* sp.) and an estimated approach to the volatile compounds. Food chemistry, 2007;105(3), pp1188-1194.
9. Chang ST, Lau OW and Cho KY, The cultivation and nutritive value of *P. sajor-caju* . European J. Appl. Micro boil. Biotechnol. 1981: 12: 58-62.
10. Cohen R., Persky L and Hadar Y. Biotechnological applications and potential of wood degrading mushrooms of the genus *Pleurotus*. Applied Microbiology and Biotechnology, 2002:58:582-594.
11. Dalentoft E., Thulin P. The use of aerobic selectors in activated sludge systems for treatment of wastewater from the pulp and paper industry, systems for treatment of wastewater from the pulp and paper industry, Water Sci. Technol. 1997: 35:181–188.
12. Deraman M., Carbon Pellets Prepared From Fibers Of Oil Palm Empty Fruit Bunches: 1. A Quantitative X-Ray Diffraction Analysis, PORIM Bulletin Palm Oil Res. Inst. Malaysia 1993:26,
13. Dike K. S., Amuneke E. H., and Ogbulie J. N. Cultivation of *Pleurotus ostreatus*: An edible mushroom from agro base waste products, J. Microbiol. Biotech. Res., 2011: 1 (3):1-14

14. Ewulonu C., and. Igwe I. O. Properties of Oil Palm Empty Fruit Bunch Fibre Filled High Density Polyethylene. *International Journal of Engineering and Technology* 2012:3 (6), 458-471.
15. Falck, R. Über die wulkuter des Austernpilzes (*Agaricus ostreatus*) auf laubholzstubben z. Forst. Jogdwes, 1917: 49:159-165.
16. Fasidi, I.O. and M. Kadiri, Effect of sporophores maturity on chemical compoisiton of *Volvariella esculenta* (Mass) Singer, a Nigerian edible mushroom. *Die Nahrung*, 1993: 37: 269-273.
17. Gbolagade, J.S. Bacteria associated with cultures of *Psathyrella atroumbonata* (Pegler) and *Schizophyllum commune* (Fr. Ex. Fr.). *African Journal of Biomedical Research*. 2004:7: 139-143.
18. Gbolagade J., Ajayi A., Oku I. and Wankasi D. Nutritive Value of Common Wild Edible Mushrooms from Southern Nigeria. *Global Journal of Biotechnology & Biochemistry* 1 2006. :(1): 16-21, ISSN 1990-9241.
19. Ghibom B., S. Minkyung, L. Seungyong, H. Seokhwan, Response surface analysis of solid state growth of *Pleurotus ostreatus* mycelia utilizing whey permeate, *Biotechnol. Lett.*2005: 27 1537–1541.
20. Gume, B., Muleta, D. and Abate D .Evaluation of locally available substrates for cultivation of oyster mushroom (*Pleurotus ostreatus*) in Jimma, Ethiopia. *African Journal of Microbiology Research*. 2013: 7(20), pp. 2228-2237.
21. James CS. *Analytical Chemistry of Foods* , 1995:1st ed., Chapman and Hall,New York
22. Jesto M.B, Guzman G.A, Mejia EG, Diaz CLG, Martinez G and Corona EB). Composition quimica de tres cepas mexicanas de setas (*Pleurotus ostreatus*). *Archivos Latinoamericanos de Nutricion*, 1998: 48(4):359-363.
23. Kavitha, B., G. Rajannan and P. Jothimani Utilization Of Empty Fruit Bunch Of Oil Palm As Alternate Substrate For The Cultivation Of Mushroom. *International Journal of Science, Environment and Technology*, 2013: 2: (5),, 839 – 846 ISSN 2278-3687 (O)
24. Lim, B. 2000.The New Straits Times. Thursday, December 28, 2000.
25. Manzi P, Gambelli L, Marconi S, Vivanti V, Pizzoferrato L. Nutrients in edible mushrooms: An interspecies comparative study. *Food Chem.*1999: 65:477–482.
26. Patil S. S. Cultivation Of *Pleurotus Sajor-Caju* On Different Agro Wastes. *Science Research Reporter* 2012:2(3): 225-228, ISSN: 2249-2321
27. Pensamut, V., Pongrit, Intarangsi C. The Oil Palm, Department of Alternative Energy Development and Efficiency, Ministry of Energy Thailand2003,
28. Rai RD, Ahlawat OP. Edible fungi: Biotechnological approaches. *Appl. Mycol. Biotechnol.* 2002:2:87–111.
29. Rai., R. D., V. Chandrasekar and T. Arumugnathan. Post harvest technology of mushrooms. *Mush. Res.*, 2003:7:42-47.
30. Rajoka, M.I. and Malik, K. A. Cellulase production by *Cellulomonas biazotea* cultured in media containing different cellulosic substrates. *Bioresource technology*1997: 59, 1; 21-27.
31. Rizki M. and Tamai Y. Effects of different nitrogen rich substrates and their combination to the yield performance of oyster mushroom (*Pleurotus ostreatus*). *World J Microbiol Biotechnol* 2011: 27: 1695-1702.

32. Schuchardt F. *et al.* Composting of empty oil palm fruit bunch (EFB) with simultaneous evapoaration of oil mill waste water (POME). In: Proceeding of 2002.International Oil Palm Conference.2002: 1-9.
33. Sturion GL and Oetterer M. Composicao Quimica de Cogumelos comestiveis (Pleurotus sp.) originados de cultivos em diferentes substrates. Ciencia e Tecnologia de Alimentos,1995:15: (2), 189-193.
34. Sudirman L. I., Sutrisna A., Listiyowati S., Fadli L., Tarigan B. The potency of oil palm plantation wastes for mushroom production. Proceedings of the 7th International Conference on Mushroom Biology and Mushroom Products (ICMBMP7) Section: Waste conversion, substrates and casing 2011: 378-384
35. Suriyanarayanan. S., A.S. Mailappa, D. Jayakumar, K. Nanthakumar, K. Karthikeyan and S. Balasubramanian. Studies on the Characterization and Possibilities of Reutilization of Solid Wastes from a Waste Paper Based Paper Industry Global Journal of Environmental Research. 2010:4 (1): 18-22, ISSN 1990-925X.
36. Teng T.T., A.L. Ahmad, Wong S.S., Zuhairi A, Najafpour G. Treatment of pulp and paper mill wastewater by polyacrylamide (PAM) in polymer induced flocculation.2005
37. Udoetok I.A. Characterization of ash made from oil palm empty fruit bunches (OEFB) International Journal of environmental Sciences.2012:3(1). 518-524.
38. Van Hamme, J. D., Singh, A. and Ward, O. P. (2003): Recent Advances in Petroleum Microbiology. – Microbiology, Molecular Biology Review 67 (4): pp 503-549.
39. Wu V., Xiao Y.Z, Yu H.Q. Degradation of lignin in pulp mill wastewaters by white-rot fungi on biofilm, Bioresour. Technol.2005: 96, 1357–1368.