

2
3 **NOCARDIOPSIS SP. 5 ENDOPHYTIC TO TULSI**
4 **LEAVES – ISOLATION AND ANTIMICROBIAL**
5 **ACTIVITY**
6

7
8
9
10 **ABSTRACT**
11

Aims: To isolate and identify the rare endophytic *Nocardiosis* sp.5 from Tulsi (*Ocimum sanctum*) leaves of Western Ghats, India and estimates its antimicrobial activity against various human pathogens and tea pathogens.

Study design: Rare endophytic actinobacteria isolation, identification and estimation of antimicrobial activity.

Place and Duration of Study: Department of Botany and Department of Bioinformatics, Nirmala College for Women, Coimbatore – Western Ghats between December 2011 and December 2012.

Methodology: Pre-sterilization of *Ocimum sanctum* leaves and isolation of rare endophytic actinomycetes using selective media. Estimation of antimicrobial activity of the isolated rare endophytic actinobacteria by primary and secondary streaking methods. Morphological and Molecular Identification of the most bioactive isolate.

Results: Out of 11 endophytes, 4 showed antagonistic activity against tea and human pathogens. Among the four isolates, isolate no.5 showed maximum activity against all the human pathogenic microorganisms such as *Staphylococcus aureus* (21mm), *Candida albicans* (18 mm), *Enterococcus faecalis* (19mm), *Vibrio cholera* (18mm) and tea pathogens such as *Glomerella cingulata* (14mm) *Hypoxyton serpens* (17mm) and *Pestalopsis theae* (18mm). Hence, morphological and phylogenetic studies show that isolate no.5 belongs to *Nocardiosis* group.

Conclusion: Endophytic *Nocardiosis* strain 5 was shown to produce good antimicrobial activity against plant pathogens and human pathogens. Further purification, structure elucidation and characterization of the antimicrobial compound from the endophytic rare actinomycete isolate *Nocardiosis* 5 strain are recommended. There is definite scope for bioprospecting of antagonistic actinomycetes from Western Ghats once appropriate consistent studies are undertaken.

12 **Keywords:** Western Ghats, endophytic actinomycetes, *Tulsi* (*Ocimum sanctum*),
13 *Nocardiosis* 5, Phylogram analysis.
14
15
16
17

1. INTRODUCTION

Actinomycetes produce about 70% of total known antibiotics, and remaining 30% are products of filamentous fungi and non-actinomycete bacteria [1]. Hence for the past 2-3 decades, the research has been focused on the antibiotic producing actinomycetes from varied terrestrial and marine resources due to decreasing rate of discovery of novel antibiotics and the increase in the multi-drug resistant pathogens in recent years. Even, rare actinomycetes strain like *Nocardioopsis* from marine sediment exhibited antimicrobial activity against multi-resistant bacteria through the production of TP-1161 antibiotic [2] which focusses the researchers' direction towards rare microbes that proved to be the source of potent antibiotic. Soil habitats (96%) have been largely surveyed as compared to other terrestrial resources like endophytic plants (3%) and animal guts (1%) [3]. Ancient Indian literatures consider all plants as potential sources of medicinal substances. Hence, medicinal plants have been considered as an important resource of isolating endophytic actinomycetes which can induce secondary metabolites of very important value. However, the work to date is insufficient to understand the actinomycetes diversity of medicinal plants present in Western Ghats. Therefore, the present research has been targeted on exploring the rare endophytic actinomycetes from the leaves of medicinal plants of Western Ghats especially Tulsi (*Ocimum sanctum*), Tamil Nadu and examining their antimicrobial activity against tea pathogens and human pathogens. This is the first work of isolating rare endophytic actinomycetes from Tulsi leaves.

2. MATERIAL AND METHODS

2.1 Collection of sample medicinal plant

The fresh leaves of common medicinal plant Tulsi (*Ocimum sanctum*) were collected from the medicinal plant garden of Nirmala College for Women, Coimbatore (11.0183°N, 76.9725°E) belonging to Western Ghats region during December 2011 to January 2012. The voucher specimen of *Ocimum sanctum* was preserved in the Herbarium of Botany department of Nirmala College for Women. Surface sterilization of these leaves should be started within four days followed by isolation work. In the meantime, these leaves shall be preserved in refrigerator.

2.2 Isolation of endophytic actinomycetes

The collected Tulsi (*Ocimum sanctum*) leaves were washed by tap water to remove the soil particles and subjected to surface sterilization procedure as per the method of Qin *et al* [4] [5]: washing the leaves in 5% sodium hypochlorite for 4-10 minutes followed by 10-minutes wash in 2.5% sodium sulphite, a 5-minutes wash with 75% of ethanol solution, a quick rinse in sterile water followed by final rinse using 10% sodium bicarbonate solution for 10 minutes to disrupt the plant tissues and inhibit the fungal growth. The finally washed solution was inoculated onto Yeast extract Malt extract agar with nalidixic acid (50 mg/lit) and nystatin (100 mg/lit) and observed for microbial growth to check the efficiency of surface sterilization.

Then, the surface sterilized tissues were dried at 100°C for 15 minutes before inoculation.

High speed centrifugation method for selective isolation of rare actinomycetes has been carried out with slight modifications [5, 6]. Five grams of surface sterilized leaves were centrifuged with 5 ml of sterile tap water for 30 seconds. After allowing it to stand at 27°C for 60 minutes, the suspension was again centrifuged at 3000 rpm for 10 minutes in order to remove soil particles. The supernatant was then centrifuged at 10000 rpm for 10 minutes and the resulting supernatant was further centrifuged at 20000 rpm for 20 minutes. The resulting supernatant was filtered through 0.22 µm pore size membrane filter and the condensed supernatant was streaked onto Humic acid Vitamin agar supplemented with nalidixic acid (50 mg/lit) and nystatin (100 mg/lit). The media was then incubated at room temperature (28 ± 2°C) for 2 to 8 weeks for the growth of endophytic rare actinomycetes.

2.3 Identification of actinomycetes

The endophytic actinomycetes isolated from Humic acid Vitamin agar plates were purified on Yeast extract Malt extract (ISP2) media and incubated at 28 ± 2°C for 30 days. The pure colonies were identified morphologically including both microscopic and macroscopic characteristics [7, 8]. The isolates were preserved in 20% (v/v) glycerol for subsequent investigation.

2.4 Microbes used

ATCC cultures like *Escherichia coli* (25922), *Staphylococcus aureus* (25923), *Enterococcus faecalis* (29212) and *Pseudomonas aeruginosa* (27853) were obtained from Piramal Diagnostics, Mumbai. Tea plant pathogens like *Macrophoma* sp, *Hypoxyton serpens*, *Pestalopsis theae*, *Glomerella cingulate* and *Botrytiplodia theobromae* were

collected from UPASI, Valparai. Other lab isolates like *Klebsiella pneumoniae*, *Acinetobacter* sp., *Bacillus* sp., *Vibrio cholera*, *Shigella dysenteriae*, *Aspergillus niger* and *Candida albicans* were collected from PSG Hospital and Microlab, Coimbatore. All these isolates were used for this study.

2.5 Primary screening

Primary streaking was performed by cross streak method as described by Sivakumar *et al* [9]. A loop full of actinobacterial inoculum was streaked in the middle of the petri dish containing MHA agar medium. After inoculation, petri dishes were incubated at 28±2°C for 5 days for actinomycetes. Same way, a fresh colony of bacterial strain was streaked and incubated at 30°C for 24 hours to allow the isolates to secrete antibiotics into medium as described by Ahmed *et al* [10]. 24hrs old pathogenic strains were cross streaked to the growth line of antimicrobial metabolite producing actinomycetes and bacteria. Each streaking was started near the edge of the plates and streaked toward the central growth line and incubated at 37°C for 24-48 hrs. The inhibition zone produced between the bioactive strains and the pathogenic bacteria were measured.

2.6 Secondary screening

Isolates that showed broad spectrum against test pathogens in primary screening were further subjected to secondary screening by Kirby Bauer paper disc method. Bioactive isolates were inoculated into 50 ml of Yeast extract Malt extract broth and incubated at 28±2°C for 5 days at 180 rpm. The culture broth was centrifuged and the activity of the supernatant was determined against test organisms by adding 0.1 ml of the culture filtrate into 6mm sterile paper disc. The discs were then placed on fresh lawn culture of test organisms, kept at 4 °C for 30 min for the diffusion of the culture broth [11] and then incubated at their respective optimum temperature. Each test was repeated three times and the antibacterial activity was expressed as the mean of diameter of the inhibition zones (mm) produced by the secondary metabolite when compared to controls, which was recorded after 18 - 24 hours and for yeasts after 2 -3 days.

2.7 Genomic DNA Isolation

The total genomic DNA from actinomycetes was isolated by following standard methods [12]. Briefly, the strain was grown in tryptic soy broth (TSB) for five days and the mycelium was separated by centrifugation and washed thrice with distilled water. Approximately 200 mg of mycelium was resuspended in 800 µl of lysis solution (100 mM Tris HCl, pH 7.5, 20 mM EDTA, 250 mM NaCl, 2% SDS, 1 mg/ml lysozyme); to the cleared lysate, 5 µl of RNase (50 mg/ml) was added and incubated at 37°C for 3 hr. Then, 10 µl of proteinase K solution (20 mg/ml) was added and incubated at 37°C for 1 hr. The lysate was extracted with an equal volume of phenol: chloroform (24:1), then centrifuged to obtain the aqueous phase. DNA was precipitated by adding 2 volumes of 95% ice cold ethanol to the aqueous phase. After centrifugation, the DNA pellet was washed twice with 70% ethanol and resuspended in 50 µl of TE buffer (10 mM Tris HCl pH 7.4 and 1 mM EDTA pH 8). The DNA was tested for purity and quantity by spectrophotometer at 260 and 280 nm.

2.8 PCR amplification

16S rRNA gene of the actinomycetes strain was amplified between the positions 8 to 1492 using forward and reverse primers mentioned below.

Forward primer: 5'- AGAGTTTGATCMTGGCTCAG -3'
Reverse primer: 5'- TACGGYTACCTTGTTACGACTT -3'

94°C	94°C	55°C	72°C	72°C
5 min	30 sec	30 sec	1 min	5 min
Initial Denaturation	Denaturation	Annealing	Extension	Final Extension
35 cycles				

A positive control (*E.coli* genomic DNA) and a negative control in the PCR were also included to run PCR. Sequencing was performed by using Big Dye terminator cycle sequencing kit (Applied BioSystems, USA). Sequencing products were determined on an Applied Biosystems model 3500XL automated DNA sequencing system (Applied BioSystems, USA) [5, 13].

2.9 Phylogenetic identification

The 16S rRNA sequence that was retrieved from actinomycetes strain was proceeded with sequence similarity search using NCBI BLAST program. Multiple sequence alignment was carried out followed by phylogenetic tree construction using MEGA6 software [5, 14].

3. RESULTS AND DISCUSSION

The isolation procedure seems to be the significant initial step for endophytic actinomycetes studies. Selective isolation strategies need to be developed further to recover the unexploited rare endophytic actinobacteria [4]. The surface sterilization was proved effective due to absence of microbial growth by spreading the final washed solution onto Yeast extract Malt extract agar. Hence, the isolated microbes from the surface sterilized leaves when inoculated on another media will be considered as endophytes.

By inoculating the surface sterilized Tulsi leaves in Humic acid Vitamin agar, eleven actinomycetes strains (Isolate 1-11) were isolated. Of these, four actinomycetes strains (Isolate 2, 3, 5, and 9) showed antagonistic activity against tea pathogens and human pathogens. Crude extract obtained from their bioactive strains were utilized for secondary screening and the results were obtained in the Table 1.

Among the four actinobacterial strains, Isolate no. 5 showed maximum activity against all the human pathogenic microorganisms such as *Staphylococcus aureus* (21mm), *Candida albicans* (18 mm), *Enterococcus faecalis* (19mm), *Vibrio cholera* (18mm) and tea pathogens such as *Glomerella cingulata* (14mm) *Hypoxylon serpens* (17mm) and *Pestalopsis theae* (18mm) (Table 1 and Fig 1,2).

Table 1. Antimicrobial screening of the given isolates against different plant and human pathogens (Secondary screening)

S.No.	Test organisms	Inhibition zones of actinomycetes isolates (mm)			
		Isolate 2	Isolate 3	Isolate 5	Isolate 9
1	<i>Macrophoma</i> sp.	8	10	12	9
2	<i>Hypoxylon serpens</i>	-	9	17	11
3	<i>Pestalopsis theae</i>	9	10	18	8
4	<i>Glomerella cingulata</i>	11	-	14	-
5	<i>Botryplodia thebromae</i>	10	9	13	-
6	<i>Escherichia coli</i>	-	5	-	-
7	<i>Klebsiella pneumoniae</i>	9	-	14	12
8	<i>Pseudomonas aeruginosa</i>	-	10	13	-
9	<i>Acinetobacter</i> sp.	10	11	13	10
10	<i>Bacillus</i> sp.	11	12	15	7
11	<i>Vibrio cholerae</i>	-	10	18	14
12	<i>Staphylococcus aureus</i>	15	12	21	20
13	<i>Shigella dysenteriae</i>	10	10	12	11
14	<i>Enterococcus faecalis</i>	7	13	19	10
15	<i>Candida albicans</i>	11	12	18	14

Isolate no. 5 produced irregular, white powdery colonies with reddish brown color on reverse side of Yeast extract Malt extract (ISP2) agar surface (Fig. 3). Dark brown melanin pigment was produced on Tyrosine agar. By viewing under 1000X magnification, gram positive rods with rounded ends; branched hyphae with non-motile spores were found (Fig. 4). Therefore, based on actinomycetes identification key [7, 8], the isolate was assigned to the genus *Nocardopsis*.




Fig. 1. Antibacterial activity of actinobacterial isolates





Fig. 2. Antifungal activity of actinobacterial isolates

156

157



158
159 **Fig. 3. *Nocardiosis* sp. 5 growth in ISP2 agar**



161
162 **Fig. 4. Microscopic structure of *Nocardiosis* sp.5**

163
164 The amplified products were visualized on 2% ethidium bromide (10mg/ml) stained agarose gel using UV transilluminator.
165 Unincorporated PCR primers and dNTPs from PCR products were removed by using Montage PCR Clean up kit
166 (Millipore). The purified PCR products of approximately 1,400 bp were sequenced by using two primers as described
167 above [15].

168 The 16S rRNA sequence of isolate 5 (around 1088 bp) was determined and deposited in Genbank under the accession
169 number **KF909127**. Using BLAST search in the NCBI data bank, the corresponding sequences homologous to our test
170 isolate were collected and the phylogenetic tree was constructed by employing MEGA6 software (bootstrap method) [14].
171 (Fig.5). The evolutionary history was predicted by using Neighbor joining method. The percentage of replicate trees where
172 the related taxa grouped together in bootstrapping test based on thousand replicates was shown clearly next to their
173 branches [5].

174
175
176




Fig. 5. Phylogram depicting the taxonomic position of *Nocardioopsis* sp.5

The comparison of 16S rRNA sequence of *Nocardioopsis* strain A5 with corresponding homologous DNA sequences clearly revealed that the test organism form a distinct phyletic line in the *Nocardioopsis* spp. (Fig. 5) [5]. The phylogeny tree clearly depicted that the isolate A5 was related to both *Nocardioopsis synnemataformans* and *Nocardioopsis dasonvillei* by sharing a 16S rRNA gene sequence similarity of 99% each. *Nocardioopsis metallicus* was selected as outgroup. Gene cloning of 16S rRNA gene can also be considered further to obtain its nearly full length of around 1400 bp for taxonomic characterization. Still, DNA-DNA hybridizations and phenotypic comparisons need to be performed to confirm its novelty.

4. CONCLUSION

Isolate no. 5 which was shown to belong to *Nocardioopsis* genera of rare endophytic actinomycetes was shown to produce good antimicrobial activity against both plant pathogens and human pathogens. Further purification, structure elucidation and characterization of the antimicrobial compound from the endophytic rare actinomycete isolate *Nocardioopsis* 5 strain are recommended to know the quality and novelty which can be applied for the treatment of different pathogenic infections since the test isolate shows prominent antimicrobial activity. Hence, there is definite scope for bioprospecting of antagonistic actinomycetes from Western Ghats once appropriate consistent studies are undertaken.

REFERENCES

1. Akshatha M.D., Manjunatha B.K., Pooja.R., Umesh.T.M., Sreevijeth.R. Screening for Novel Antibiotic Producing Actinomycetes from Western Ghats of Karnataka State, India. Indian journal of Research. 2013; 2(2).

- 200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
2. Kerstin Engelhardt, Kristin F. Degnes, Michael Kemmler, Harald Bredholt, Espen Fjærvik, *et al.* Production of a New Thiopeptide Antibiotic, TP-1161, by a Marine *Nocardioopsis* Species. *Applied and Environmental Microbiology*. 2010; 76(15): 4969–4976.
 3. Sonashia Velho-Pereira & Nandkumar M. Kamat. Actinobacteriological research in India. *Indian Journal of Experimental Biology*. 2013; 51: 573-596.
 4. Qin, S., Li, J., Chen, H.-H., Zhao, G.-Z., Zhu, W.-Y., Jiang, C.-L., *et al.* Isolation, diversity, and antimicrobial activity of rare actinobacteria from medicinal plants of tropical rain forests in Xishuangbanna China. *Appl. Environ. Microbiol.* 2009; 75: 6176–6186.
 5. Mrinalini J Singh and Padmavathy S. Isolation, screening and characterization of endophytic PGPR actinomycetes present commonly in Neem & Tulsi leaves *-in vitro* study (Tomato). *International Journal of Recent Scientific Research* 2014; 5(3): 574-579.
 6. Kizuka M, Enokita R, Takahashi K, Okazaki T. Distribution of the actinomycetes in the Republic of South Africa investigated using a newly developed isolation method. *Actinomycetologica* 1997; 11:54–58.
 7. Shirling, E.B. and Gottlieb. D. Methods for characterization of *Nocardioopsis* species. *Int J Syst Bacteriol*. 1966; 16:313–340.
 8. Williams S.T, Sharpe M.E. and Holt J.G. *Bergey's Manual of Systematic Bacteriology*. Vol 4. Williams and Wilkins, Baltimore, 1989.
 9. Siva kumar J, Santhanam P and Masilamani Selvam M. Antimicrobial activity of actinomycetes isolated from the Western Ghats of Tamilnadu. *International Journal of Pharma and Bio Sciences* 2011; 2(1).
 10. Ahmed.N, Uzair.B, Ayaz.S, Ahmed.V. Antibacterial activity of marine bacteria from Arabian Sea of Pakistan. *The Internet Journal of Microbiology*. 2007; 4(2).
 11. Deepika Sharma, Talwinder Kaur, Chadha BS and Rajesh Kumari Manhas. Antimicrobial activity of actinomycetes against Multidrug Resistant *Staphylococcus aureus*, *E. coli* and various other pathogens. *Tropical Journal of Pharmaceutical Research* 2011; 10 (6): 801-808.
 12. Boudjella H., Bouti K, Zitouni A., Mathieu F., Lebrihi A. and Sabaou N. Taxonomy and chemical characterization of antibiotics of *Streptosporangium* Sg 10 isolated from a Saharan soil. *Microbiol. Res.* 2006; 161:288-298.
 13. Jacob H Jacob and Fawzi I. Irshaid. Biochemical and molecular taxonomy of a mild halophilic strain of *Citrobacter* isolated from hypersaline environment. *Research journal of Microbiology*. 2012; 7(4): 219-226.
 14. Tamura K, Stecher G, Peterson D, Filipski A and Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis Vesion 6.0. *Molecular Biology and Evolution* 2013; 30: 2725-2729.
 15. Kavitha JR and Devasena T. *Bacillus* species isolated from cow milk samples. *Discovery*. 2014; 20(64): 124-132