

CURRENT TAXONOMY OF RHIZOBIA

Halima BERRADA and Kawtar FIKRI BENBRAHIM

Laboratory of Microbial Biotechnology, Faculty of Sciences and Technology, Sidi Mohammed Ben Abdellah University, P. O. Box 2202, Fez, Morocco.

Abstract

Classification of rhizobia knows constant change. Recent studies have shown the existence of a great diversity among nitrogen-fixing bacteria isolated from different legumes. Currently, More than 98 species belonging to 13 genera of α - and β - proteobacteria have been described as rhizobia. The genera *Rhizobium*, *Mezorhizobium*, *Ensifer* (formerly *Sinorhizobium*), *Bradyrhizobium*, *Phyllobacterium*, *Microvirga*, *Azorhizobium*, *Ochrobacterium*, *Methylobacterium*, *Devosia*, *Shinella* (Class of α -proteobacteria), *Burkholderia*, *Cupriavidus* (formerly *Ralstonia*) (Class of β -proteobacteria) and some γ -proteobacteria, form the set of the bacteria known as legume's symbionts. There certainly is much to discover, since only 0.5% of known legumes were until now identified specifically for symbiotic relationship. The discovery of new symbionts associated with legumes is expected to create deep changes in the taxonomy of the rhizobia. A literature review of the currently recognized classification of rhizobia is presented in this paper.

Keywords: Rhizobia, Legume, Proteobacteria, Taxonomy, Classification.

1. Introduction

Rhizobia are soil bacteria able to form nodules and establish a symbiosis with the roots or the stems of leguminous plants. During the process of symbiosis, rhizobia reduce atmospheric nitrogen into a form directly assimilated by plants (ammonium).

The ability of rhizobia to fix nitrogen significantly reduced the use of chemical fertilizers in agriculture. In fact, nearly 90% of the nitrogen plant needs are well satisfied and the soil is enriched with nitrogen that will be used by subsequent crops, with no need to add chemical fertilizers. In general, this symbiosis has several advantages including improved agricultural productivity, maintenance and restoration of soil fertility, safety of expensive fertilizers currency and energy and limitation of groundwater's pollution by nitrates playing therefore a significant ecological and economical function.

The study of the rhizobial diversity is a valuable biological resource and attempts to find bacterial strains with interesting features to maximize the legumes productivity [1]. Such studies allow us not only to find new strains of rhizobia, but they also support research efforts to select efficient combinations of rhizobium-legume association.

Because of their ecological and economical importance, the diversity and taxonomy of these microorganisms have been extensively studied over the last twenty years. From one genus including

four species in 1981, the classification now includes at least 13 genus comprising more than 98 species and this number continue to increase [2].

Zakhia et al. (2004) proposed the term BNL (Bacteria Nodulating Legumes) to avoid confusion between the general term of rhizobium and genus name. Currently, all BNL described so far belong to the Proteobacteria class. The majority of them belong to genus of the α -Proteobacteria class, called *Rhizobium*, *Mesorhizobium*, *Bradyrhizobium*, *Azorhizobium*. In addition, new types were found in the α -Proteobacteria class, namely *Methylobacterium*, *Devosia*, *Microvirga*, *Ensifer* (formerly *Sinorhizobium*), *Ochrobactrum*, *Phyllobacterium* and *Shinella* [3].

The BNL were also found in the β -Proteobacteria class, such as *Burkholderia*, *Cupriavidus* (formerly *Ralstonia*) [4]. Finally, several studies reported the presence of γ -Proteobacteria class in black locust by the *Pseudomonas* sp. [5].

More than 19.700 legume species grow around the world, but only a few microsymbionts have been studied [6]. The increase in advanced study of legume's new species in different geographical regions opens new perspectives to isolate and to characterize more species of *Rhizobium*.

2. Phylogeny of rhizobia

The comparison of data obtained from bacterial strains or species with different molecular approaches allows the identification of groups of taxa. Different taxa obtained can be compared and used for tracing lines of descent commonly recognized under the name of phylogenetic tree (Figure 1). Thus, the phylogeny can be defined as the evolutionary history that reveals the existing relationships between groups of organisms with an ancestor located at a higher taxonomic level.

Nowadays, use of 16S rRNA sequence is the main tool in the study of microbial phylogeny. However, it has limitations to differentiate among close species and for this purpose several metabolic genes (housekeeping) have been proposed in several groups of bacteria. In rhizobia the two first genes analyzed were *recA* and *atpD* and currently they have been sequenced in many rhizobial species showing their usefulness in differentiation of species whose 16S rRNA genes are nearly identical. In the last years new schemes for identification and phylogenetic analysis of bacteria named MLSA (Multilocus sequence analysis) and MLST (Multilocus sequence typing) based in the analysis of several housekeeping genes have been applied phylogenetic analyses of concrete groups of rhizobia [7].

Besides the housekeeping genes also named "core" genes, some "auxiliary" or "accessory" genes involved in the legume symbiosis are commonly included in species description of rhizobia and in some MLST analysis. The symbiotic genes from rhizobia are codified in plasmids in fast and some middle growing species and in symbiotic islands in the middle and slow growing ones. From these genes, the most studied in rhizobia are *nodD*, *nodA*, *nodC* and *nifH*. Nevertheless, the symbiotic genes are not useful in taxonomy because due to their ability to be transferred in nature from plasmids to islands, from bacteria to plants and among bacteria. Therefore the analysis of symbiotic genes is

overall useful to identify non-rhizobial species able to nodulate legumes and to carry out biogeography studies of legume endosymbionts [7].

3. Polyphasic taxonomy

Originally based on phenotypic criteria, the application of polyphasic terminology has been reviewed by the revolution of different molecular techniques of which mainly the PCR or Polymerase Chain Reaction [8]. Thus, the standard principles were proposed in 1991, for the description of new genus or species on the basis of numerical analysis which is the synthesis of the results of different phenotypic and molecular tests [9]. This new definition of the polyphasic approach allowed a proper systematic study of major bacterial groups, an assessment of taxonomic resolution of different techniques of diversity study [10]. (Figure 2) and an establishment of comparable databases between different applied microbiology laboratories [11].

The most commonly used molecular techniques are: FT-IR technique (Fourier-transform infrared spectroscopy), which allows to study the diversity of bacteria at an intraspecific level [12], the technique of typing by PCR target (target PCR fingerprinting) [13] and especially the technique of DNA chips commonly known as DNA microarrays. In this last technique, the conception of the chips permits to detect simultaneously thousands of sequences or even to cover the entire genome of an organism [14]. By the same technique, Rüberg et al. (2003) have analyzed the entire genome of *Sinorhizobium meliloti* [15]. All these techniques can reveal more information about the diversity of rhizobia and can therefore provide more data available for their phylogenic study.

Other, New methods are potential alternatives to DNA-based methods for rapid and reliable characterization of bacteria. The most commonly used methods are: Matrix-assisted laser desorption ionization–time-of-flight mass spectrometry (MALDI-TOF MS) has been suggested as a fast and reliable method for bacterial identification, based on the characteristic protein profiles for each microorganism. Using this technology it has been estimated that up to 99% of strains tested are correctly identified when comparing with commercial phenotypic identification panels or *rrs* gene sequencing [16].

Thus, modern bacterial taxonomy is based on the integration of all phenotypic, genotypic and phylogenetic data for a more stable classification [17; 10]. The following paragraphs will discuss the current classification of rhizobia with this new approach.




Figure 1: phylogenetic tree of rhizobia and related species built by the neighbor-joining method from the complete sequence of 16S rRNA [18].

	Family	Genus	Species	Strain
Labeling of DNA probes				
DNA sequencing				
Phenotypic analysis (API...)				
RNA sequencing				
ADN:ARNr hybridization				
Chemotaxonomic analysis				
FAME				
ADN:ADN hybridization				
PCR/ARNr				
G+C%				
RFLP-ARDRA				
RAPD-DAF-APPCR				
AFLP				
REP-PCR				
Proteomic analysis of total cellular proteins				
Isoenzyme polymorphism				
Phage typing				
MLEE				
Serology				

Figure 2: Taxonomic resolution of the various techniques used to study the diversity of rhizobia [10;

4. Current classification of rhizobia

The first classifications were based on cross-inoculation tests between rhizobia and host plants for their identification. The host plant was not the only criteria taken into account for the classification of rhizobium which species were classified into two genus: the genus *Rhizobium* corresponding to strains with fast growth and genus *Bradyrhizobium* for slow-growing strains, based on their generation time and rate of growth on culture medium [20].

However, discordant observations between the notion of bacterial growth speed and the host range showed a lot of doubt on the validity of this classification. This makes place otherwise to comparative methods such as the serology, the coefficient of Chargaff, RNA / DNA or DNA / DNA, hybridization analysis of plasmids, etc...

This period marked the beginning of a new taxonomy based on the comparison of results from different phenotypic and biochemical analysis for the identification of symbiotic bacteria [21]. Since then, the isolation of rhizobia from an increasing number of plant species in the world and their characterization by modern polyphasic taxonomy has led to the description of other new genera and species.

The current literature describes more than 98 species grouped in 11 genus belonging to the subclass α -Proteobacteria and 2 genus belonging to the order of Burkholderiales in subclass β -Proteobacteria [17; 15; 22; 23], and finally, one genus belonging to the order Pseudomonales in the subclass γ -Proteobacteria [24].

The complete list of valid species of rhizobia is recorded in the List of Prokaryotic Names with Standing in Nomenclature that is performed and revised by Dr. Euzéby (<http://www.bacterio.cict.fr>). According to this list currently the genus *Rhizobium* contains 33 species and includes to the former genera *Agrobacterium* and *Allorhizobium*. The genus *Sinorhizobium*, currently named *Ensifer*, contains 13 species, the genus *Shinella*, contains 1 specie, the genus *Mesorhizobium*, 29 species, the genus *Bradyrhizobium*, 15 species, the genus *Phyllobacterium*, 1 specie and the genus *Azorhizobium*, 3 species, the genus *Merhylobacterium*, 1 specie, the genus *Microvirga*, 3 species.

This high number of species able to nodulate legumes (Table 1) was unexpected only two decades ago and the previsions are that it will be increased in next times.

In summary, on the basis of the DNA ribosomal 16S sequence, the legume's symbionts belong to three main distinct phylogenetic subclasses (Figure 3).

Table 1: Current classification of different species of rhizobia [28]

Genus	Species	Host plant	References
Class : Alphaproteobacteria			
Order : Rhizobiales			
Family : Rhizobiaceae			
Genus: <u>Rhizobium</u>			
<i>R. leguminosarum</i>			[25]
<i>biovar viciae</i>	<i>Pisum, viciae, Lens, Lathyrus</i>		[25; 26]
<i>biovar trifolii</i>	<i>Trifolium pratense</i>		[29; 26]
<i>biovar phaseoli</i>	<i>Phaseolus vulgaris</i>		[25; 26]
<i>R. galegae</i>	<i>Galega, Leucaena</i>		[27]
<i>biovar officinalis</i>	<i>Galega orientalis</i>		[27]
<i>biovar orientalis</i>	<i>Galega officinalis</i>		[27]
<i>R. tropici</i>	<i>Phaseolus, Medicago, Macroptilium</i>		[28]
type IIA (<i>R. leucaenae</i>)			[29]
type IIB			[28]
<i>R. endophyticum</i>	<i>Phaseolus vulgaris</i>		[30]
<i>R. phaseoli</i>	<i>Phaseolus</i>		
<i>R. fabae</i>	<i>Vicia faba</i>		[31]
<i>R. etli</i>	<i>Phaseolus,</i>		[32]
<i>biovar mimosae</i>	<i>Mimosa affinis</i>		[33]
<i>biovar phaseoli</i>	<i>Phaseolus,</i>		[24]
<i>R. undicola</i>	<i>Neptunia natans</i>		[35]
<i>R. gallicum</i>	<i>Phaseolus vulgaris</i>		[36]
<i>biovar phaseoli</i>	<i>Phaseolus vulgaris</i>		[36]
<i>biovar gallicum</i>	<i>Phaseolus vulgaris</i>		[36]
<i>R. giardinii</i>	<i>Phaseolus vulgaris</i>		[36]
<i>biovar phaseoli</i>	<i>Phaseolus,</i>		[36]
<i>biovar giardinii</i>	<i>Phaseolus vulgaris</i>		[36]
<i>R. hainanensis</i>	<i>Desmodium sinuatum,</i> <i>Centrosema, etc.</i>		[37]
<i>R. huautlense</i>	<i>Sesbania herbacea</i>		[38]
<i>R. mongolense</i>	<i>Medicago ruthenica, Phaseolus</i>		[39]
<i>R. yanglingense</i>	<i>Amphicarpaea</i>		[40]
<i>R. larrymoorei</i>	<i>Ficus benamina</i>		[41]
<i>R. indigoferae</i>	<i>Indigofera spp.</i>		[42]
<i>R. sullae</i>	<i>Hedysarum</i>		[43]
<i>R. loessense</i>	<i>Astragalus, Lespedeza</i>		[44]
<i>R. cellulolyticum</i>	<i>Populus alba</i>		[45]
<i>R. miluonense</i>	<i>Lespedeza</i>		[46]
<i>R. multihospitium</i>	Multiple legume species		[47]
<i>R. oryzae</i>	<i>Oryza alta</i>		[48]
<i>R. pisi</i>	<i>Pisum sativum</i>		[49]
<i>R. mesosinicum</i>	<i>Albizia, Kummerowia Dalbergia</i>		[50]

<i>R. alamii</i>	<i>Arabidopsis thaliana</i>	[51]
<i>R. alkalisoli</i>	<i>Caragana intermedia</i>	[52]
<i>R. tibeticum</i>	<i>Trigonella archiducis-nicolai</i>	[53]
<i>R. tubonense</i>	<i>Oxytropis glabra</i>	[54]
<i>R. halophytocola</i>	Coastal dune plant	[53]
<i>R. radiobacter</i>	*	[55]
<i>R. rhizogenes</i>	*	[55]
<i>R. rubi</i>	*	[55]
<i>R. vitis</i>	*	[55]
<i>R. nepotum</i>	*	[56]
Genus: <i>Ensifer</i>		
<i>E. meliloti</i>	<i>Medicago, Melilotus, Trigonella</i>	[57]
<i>E. fredii</i>	<i>Glycine, Vigna, Cajanus</i>	
<i>biovar fredii</i>	<i>Glycine, Vigna, Cajanus</i>	[58]
<i>biovar siensis</i>	<i>Glycine</i>	[37]
<i>E. saheli</i>	<i>Acacia, Prosopis, Neptunia, Leucaena</i>	[59]
<i>E. teranga</i>	Different host plants	[59]
<i>biovar acaciae</i>	<i>Acacia</i>	[55]
<i>biovar sesbania</i>	<i>Sesbania</i>	[60]
<i>E. medicae</i>	<i>Medicago truncatula, Melilotus</i>	[61]
<i>E. arboris</i>	<i>Acacia, Prosopis</i>	[62]
<i>E. kostiense</i>	<i>Acacia, Prosopis</i>	[62]
<i>E. xingianense</i>	<i>Glycine max</i>	[63]
<i>E. adhaerens</i>	<i>Leucaena leucocephala</i>	[64]
<i>E. kummerowiae</i>	<i>Kummerowia stipulaceae</i>	[42]
<i>E. americanum</i>	<i>Acacia</i>	[65]
<i>E. mexicanus</i>	<i>Acacia angustissima</i>	[66]
<i>E. numidicus</i>	<i>Medicago sativa</i>	[67]
Genus : <i>Shinella</i>		
<i>S. kummerowiae</i>	<i>Kummerowia stipulacea</i>	[50]
Family : <i>Phyllobacteriaceae</i>		
Genus : <i>Mesorhizobium</i>		
<i>M. loti</i>	<i>Lotus, Cicer, Anthyllis, Astragalus, etc.</i>	[68]
<i>M. huakuii</i>	<i>Astragalus sinicus</i>	[69]
<i>M. ciceri</i>	<i>Cicer arietinum</i>	[70]
<i>M. tianshanense</i>	<i>Glycyrrhiza pallidiflora</i>	[63]
<i>M. mediterraneum</i>	<i>Cicer arietinum</i>	[71]
<i>M. plurifarum</i>	<i>Acacia, Chamaecrista, Leucaena, Prosopis,</i>	[72]

<i>M. amorphae</i>	<i>Amorpha fruticosa</i>	[73]
<i>M. chacoense</i>	<i>Prosopis alba</i>	[74]
<i>M. septentrionale</i>	<i>Astragalus adsurgens</i>	[75]
<i>M. temperatum</i>	<i>Astragalus adsurgens</i>	[75]
<i>M. thiogangeticum</i>	*	[75]
<i>M. albiziae</i>	<i>Albizia kalkora</i>	[75]
<i>M. caraganae</i>	<i>Caragana spp.</i>	[64]
<i>M. gobiense</i>	Wild legumes	[47]
<i>M. tarimense</i>	Wild legumes	[47]
<i>M. australicum</i>	<i>Biserrula pelecinus L.</i>	[76]
<i>M. opportunistum</i>	<i>Biserrula pelecinus L.</i>	[76]
<i>M. metallidurans</i>	<i>Anthyllis vulneraria</i>	[77]
<i>M. alhagi</i>	<i>Alhagi</i>	[78]
<i>M. camelthorni</i>	<i>Alhagi sparsifolia.</i>	[78]
<i>M. abyssinicae</i>	Different agroforestry legume trees	[79]
<i>M. muleiense</i>	<i>Cicer arietinum L.</i>	[80]
<i>M. hawassense</i>	Different agroforestry legume trees	[79]
<i>M. qingshengii</i>	<i>Astragalus sinicus</i>	[81]
<i>M. robiniae</i>	<i>Robinia pseudoacacia</i>	[82]
<i>M. shonense</i>	Different agroforestry legume trees	[79]
<i>M. shangrilense</i>	<i>Caragana species</i>	[83]
<i>M. silamurunense</i>	<i>Astragalus species</i>	[84]
<i>M. tamadayense</i>	<i>Anagyris latifolia, Lotus berthelotii</i>	[85]
<u>Genus : Phyllobacterium</u>		
<i>P. trifolii</i>	<i>Trifolium pratense</i>	[86]
Family : <u>Methylobacteriaceae</u>		
<u>Genus : Methylobacterium</u>		
<i>M. nodulans</i>	<i>Crotalaria spp</i>	[87]
<u>Genus : Microvirga</u>		
<i>M. lupini</i>	<i>Lupinus sp.</i>	[88]
<i>M. lotononidis</i>	Different legume host	[88]
<i>M. zambiensis</i>	Different legume host	[88]
Family : <u>Brucellaceae</u>		
<u>Genus : Ochrobactrum sp.</u>		
<i>Ochrobactrum cytisi</i>	<i>Cytisus</i>	[89]
<i>Ochrobactrum lupini</i>	<i>Lupinus albus</i>	[90]
Family: <u>Hyphomicrobiaceae</u>		
<u>Genus: Azorhizobium</u>		
<i>A. caulinodans</i>	<i>Sesbania rostrata</i>	[91]
<i>A. johannense</i>	<i>Sesbania virgata</i>	[91]
<i>Azorhizobium sp</i>	<i>Sesbania rostrata</i>	[91]
<u>Genus: Devosia</u>		
<i>Devosia neptuniae</i>	<i>Neptunia natans</i>	[92]
Family : <u>Bradyrhizobiaceae</u>		

<u>Genus : Bradyrhizobium</u>		
<i>B. Japonicum</i>	<i>Glycine max, Glycine soja</i>	[26 ; 93]
<i>B. Elkanii</i>	<i>Glycine max</i>	[94]
<i>B. Liaoningensese</i>	<i>Glycine max</i>	[95]
<i>B. Yuanmingense</i>	<i>Lespedeza</i>	[96]
<i>B. Betae</i>	<i>Betae vulgaris</i>	[97]
<i>B. Canariense</i>	<i>Genisteae et Loteae</i>	[98]
<i>B. iriomotense</i>	<i>Entada koshunensis</i>	[99]
<i>B. jicamae</i>	<i>Pachyrhizus erosus</i>	[49]
<i>B. lablabi</i>	<i>Lablab purpureus</i>	[99]
<i>B. huanghuaihaiense</i>	Soybean	[100]
<i>B.cytisi</i>	<i>Cytisus villosus</i>	[101]
<i>B.daqingense</i>	Soybean	[102]
<i>B. denitrificans</i>	<i>Aeschynomene</i>	[103]
<i>B.oligotrophicum</i>		[104]
<i>B.pachyrhizi</i>	<i>Pachyrhizus erosus</i>	[105]
Class : Beta Proeobacteria		
Order : Burkholderiales		
Family: Burkholderiaceae		
<u>Genus : Burkholderia</u>		[106]
<i>B. caribensis</i>	<i>Vertisol microaggregates</i>	[107]
<i>B. cepacia</i>	<i>Alysicarpus glumaceus</i>	[4]
<i>B.tuberum</i>	<i>Aspalatus carnosa</i>	[108]
<i>B. phymatum</i>	<i>Aspalatus carnosa</i>	[108]
<i>B.nodosa</i>	<i>Mimosa bimucronata, Mimosa scabrella</i>	[109]
<i>B. sabiae</i>	<i>Mimosa caesalpiniiifolia</i>	[110]
<i>B.mimosarum</i>	<i>Mimosa spp.</i>	[111]
<i>B.rhizoxinica</i>	<i>Rhizopus microsporus</i>	[112]
<i>B.diazotrophica</i>	<i>Mimosa spp.</i>	[113]
<i>B.endofungorum</i>	<i>Rhizopus microsporus</i>	[112]
<i>B.heleia</i>	<i>Eleocharis dulcis</i>	[114]
<i>B.symbiotica</i>	<i>Mimosa spp.</i>	[115]
<u>Genus : Cupriavidus</u>	<i>Aspalatus carnosa</i>	
<i>C. taiwanensis</i>	<i>Mimosa sp.</i>	[116]
Class: Gamma-Proteobacteria		
Order: Pseudomonadales		
Family: Pseudomonaceae		
<i>Pseudomonas sp.</i>	black locust	[5]

*Species with no described nodulation ability included in traditionally considered rhizobial genera.




Figure 3: Simplified phylogenetic tree of the 16S rRNA of proteobacteria. The genus shown in bold contain rhizobia [117].

4.1. Alpha-subclass of Proteobacteria

4.1.1. Rhizobium /Ensifer/ Shinella branch

This branch is subdivided in **three** sub branches in the genus: *Rhizobium* [25], *Ensifer* (formerly *Sinorhizobium*) [37; 59], and *Shinella* [50]. These genus are grouped with:

- *Agrobacterium*, non-symbiotic bacterial genus involved in pathogenic interactions with plants;
- Bacteria ameliorative plant growth (PGPR: Plant growth-promoting rhizobacteria) as *Azospirillum*,

- Human and animal pathogens including *Brucella*, *Ochrobactrum*, *Bartonella* or soil bacteria as *Mycoplana*.

The first sub-branch corresponds to genus *Rhizobium*, define a group of thirty species, from various hosts, and includes *R. leguminosarum*, *R. tropici* [28], *R. etli* [32], *R. gallicum* [36], *R. mongolense* [39], *R.undicola* [35] and *Agrobacterium*. The members of this genus are short Gram-negative rods, of a 0.5 to 0.9µm width and a 1.2 to 3 µm length, often with an uncolored region due to the presence of polymer β-hydroxybutyrate (PHB). Rhizobia do not form endospores but are motile with polar or peritrichous flagella (4 to 6), aerobic, chemo-organotrophs. They become pleomorphic under adverse conditions [118].

There are a number of species present in these rhizobial genera that have not been observed to form nodules, and therefore do not fit the functional definition of rhizobia. These include many of the species that were formerly known as *Agrobacterium* (e.g. *R. larrymoorei*, *R. rubi*, and *R. vitis*) [55]. However, there is recent evidence that other species formerly classified as *Agrobacterium* are capable of nodulation. For example *R. radiobacter* nodulates *Phaseolus vulgaris*, *Campylotropis* spp., *Cassia* spp. [119], and *Wisteria sinensis* [120]. Both nodules and tumours were formed on *Phaseolus vulgaris* by *R. rhizogenes* strains containing a Sym plasmid [121].

The second sub-branch corresponds to genus *Ensifer* (formerly *Sinorhizobium*) and includes the species *E. fredii*, *E. meliloti*, *E. teranga* and *E. sahelense* [59; 122], *E. medicae* [61], *E. kostense* and *E. Arboris* [62]. Rods of this genus are 0.5 to 0.9 µm wide and 1.2 to 3 µm length, motile by polar flagellum or some peritrichous flagella, Gram-negative, aerobic accumulating PHB. Optimal growth of this genus is at 25-30°C (10-35°C) and at pH 6-8 (5 -10.5) and tolerates 10 g/l of NaCl [123].

Sinorhizobium was proposed as the new genus of the fast-growing soybean rhizobia. This genus is separated from *Rhizobium* and *Bradyrhizobium* based on GC content, DNA-DNA hybridization data, numerical taxonomy, serological analysis data, composition of extracellular gum, bacteriophage typing data and soluble protein pattern. However, *Rhizobium* and *Sinorhizobium* are phylogenetically fairly closely related to each other. Recently, the Judicial Commission of the International Committee on Systematics of Prokaryotes decided to transfer some species of the genus *Sinorhizobium* to the genus *Ensifer* [124].

The third sub-branch corresponds to genus *Shinella* and include one specie: *S.kummerowiae* [50], a symbiotic bacterium nodulating *Kummerowia stipulacea*. Cells of the strain are Gram-negative, strictly aerobic, non-spore-forming, motile short rods [125].

In 2001, Young et al. had revised taxonomic data and showed that the discriminatory phenotypic characteristics to distinguish between these types are not convincing, and that the phylogenetic relationships between genera deduced from the comparative analysis of 16S rDNA sequences differ depending on the chosen algorithm and more especially on the chosen sequences [55]. Therefore, despite phenotypic differences, these authors suggested to combine the three genera *Rhizobium*, *Agrobacterium* and *Allorhizobium* in a single genre: *Rhizobium*, including all species of *Agrobacterium* and *Allorhizobium* as new combinations: *R. radiobacter*, *R. rhizogenes*, *R. rubi*, *R. undicola* and *R. vitis*.

4.1.2. Mesorhizobium branch

A new genus proposed in 1997 to include the species of intermediate growth named *Mesorhizobium* [126] was included in the family *Phyllobacteriaceae* that also included the former genus *Phyllobacterium* (with one specie: *Phyllobacterium trifolii* isolated from *Trifolium pratense* nodules) [86].

In this branch *Mesorhizobium* [126] contains, *M. loti* [68], *M. huakuii* [69], *M. ciceri* [70], *M. mediterraneum* [71], *M. tianshanense* [63], *M. plurifarum* [72] and *M. amorphae* [127]. They are phylogenetically related and distinct from the large phylogenetic grouping that includes *Rhizobium*, *Agrobacterium* and *Sinorhizobium* [126]. They are characterized by a growth rate intermediate between the fast- and slowgrowing rhizobia. They are Gram-negative rods, motile by polar or subpolar flagellum, aerobic, accumulating PHB. They assimilate glucose, rhamnose, and sucrose with acid production [123].

4.1.3. Azorhizobium branch

This family regroups only one genus and only one species of symbiotic bacterium: *Azorhizobium caulinodans*. They are short rods very similar to *Rhizobium*, who present a polar and peritrichous ciliation but produce an alkaline reaction on glucose as *Bradyrhizobium*. Only this sugar is used. Strains also degrade fatty acids, organic acid and alcohols and are the only ones to use alcohols. Their growth is intermediate between the two previous genus, Generation time = 7-9 h. No strain is denitrifying. They grow up to 43 °C. They have only the dihydrolase arginine and lysine decarboxylase. They do not use mannitol, conventional substrate of the other two genera, but lactate gives good growth yields. They seem more sensitive to antibiotics [123]. In addition, it doesn't only form nodules on root parts but also on aerial parts of *Sesbania rostrata* [91]. A second species *Azorhizobium johannae* was proposed [92]. However, this species is characterized by a low level of DNA / DNA hybridization compared to the typic strain of the genus.

The analysis of the 16S rRNA sequences showed that the typic strain is mixed with species of the genus *Xanthobacter* and *Aquabacters*. The combination of these last two genres with *Azorhizobium* was investigated without being suggested because of the great dissimilarity of phenotypic traits [128; 129].

4.1.4. Bradyrhizobium branch

This genus has been defined to include all slow-growing rhizobia [26]. These are rods with a single polar or subpolar flagellum. The colonies do not exceed 1 mm diameter in agar + mannitol. These symbiotic bacteria use many sugars and organic acids but prefer pentoses, with production of polysaccharidic mucus. Some species can grow as chimolithotrophic in the presence of H₂, CO₂ and low content in O₂ due to the presence of an hydrogenase, Generation time = 10-12h. Some strains can fix N₂ in vitro. It's more resistant to antibiotics than *Rhizobium* [123].

This branch has included, for a long time, a single species: *Bradyrhizobium japonicum* that regrouped all strains nodulating the soybean (*Glycine max*). Very heterogeneous, it was then divided into three groups (I, Ia and II), based on homologies obtained by DNA/DNA hybridization [130]. A new species:

Bradyrhizobium elkanii was created in the Group II, which differs from the species *Bradyrhizobium japonicum* [94]. For other very slow growing strains, isolated from nodules of soybean *Glycine max* and *Glycine soya* in China, *Bradyrhizobium liaoningense* species has been proposed [95]. The *Bradyrhizobium* genus includes strains for which the taxonomic position is not clearly defined at the species level. The discrepancy between *Bradyrhizobium japonicum* and *Bradyrhizobium elkanii* is clear. It is more important than the divergence between *Bradyrhizobium* and non-symbiotic species as: *Afpia* (animal bacterial pathogens), *Nitrobacter* (nitrifying bacteria in the soil), *Rhodopseudomonas palustris* (photosynthetic bacteria) and *Blastobacter denitrificans* [106].

In recent years, the characterization of different strains of *Bradyrhizobium* by comparing several molecular techniques revealed the existence of 11 different genotypes, three of which correspond to known strains, while 8 genotypes are very distinct.

A new species *B. yuanmingene* was isolated from the genus *Lespedeza* [96; 131]. In the same year, a strain of *Bradyrhizobium* nodulating a wild plant of the genus *Phaseolus* was reported [132]. Other recognized species of this group are *Bradyrhizobium betae* from the roots of *Beta vulgaris* afflicted with tumor-like deformations [97] and *Bradyrhizobium canariense* from genistoid legumes from the Canary Islands [98]. Many other slow-growing rhizobia was isolated from other legume hosts and are commonly referred to as *Bradyrhizobium* sp., followed by the name of the legume host: *B. lablabi* was isolated from the genus *Lablab purpureus* [128], *B. cytisi* has been isolated from the genus *Cytisus villosus* [101] and *B. pachyrhizi* was isolated from effective nodules of *Pachyrhizus erosus* [105].

Based on ITS sequence data, the photosynthetic bradyrhizobia isolated from stem-nodules of *Aeschynomene*, form a distinct group closely related to *Blastobacter denitrificans* [106]. As a result of a comprehensive study of both groups, van Berkum et al. (2006) recently proposed to transfer *Blastobacter denitrificans* to *Bradyrhizobium* and unite the species with the isolates from *Aeschynomene indica* as the species *Bradyrhizobium denitrificans* [103].

Recently, A new species *B. Canariense* from *Genisteeae Loteae* [98], *B. Iriomotense* from *Entada koshunensis* [99], *B. jicamae* from *Pachyrhizus erosus* [85], *B. daqingense* and *B. huanghuaihaiense* from soybean was reported [102].

4.1.5. Methylobacterium branch

The *Methylobacterium* genus [133], has included a single species: *M. nodulans* that regrouped all strains nodulating the *Crotalaria*. Bacteria of this genus have the ability to grow on single-carbon compounds such as formate, formaldehyde, methylamine and methanol as only source of carbon and energy. These are chemo-organotrophic and optional methylotrophic, able to use multi-carbon compounds for growth. The cells of this genus present the rods form (0.8 - 1.0 x 1.0 to 3.0 mm), most frequently isolated and occasionally in rosette. They are often branched or pleomorphic, especially in cultures in late stationary phase. They are strictly aerobic bacteria, Gram-negative, oxydase and catalase positive and have a typical pink pigmentation due to the presence of carotenoids [134]. *Methylobacterium* strains are commonly called "pink-pigmented facultative Methylotrophs" (PPFM). However, at least one case, *M. nodulans* is not pigmented [100]. They are mesophilic and optimal

temperature is between 25 and 30°C. All these strains are motile with a single polar flagellum. Often, cells contain inclusions of poly- β -hydroxybutyrate and sometimes inclusions of polyphosphates [135].

4.2. Beta-subclass of Proteobacteria:

4.2.1. Burkholderia branch :

The *Burkholderia* [136] genus has been proposed in 1992 by Yabuuchi and al. The definition of this was based primarily on genomic considerations and on cellular lipid's composition. Regarding the main phenotypic characteristics, they are right bacilli, Gram negative, usually accumulating granules of poly-beta-hydroxybutyrate, motile with one or more polar flagella (*Burkholderia mallei* is however lacking flagella and stationary), strictly aerobic, catalase positive, oxidase variable depending on the species, can grow using as sole carbon source glucose, glycerol, inositol, galactose, sorbitol, mannose (unlike *Ralstonia sp.* that do not use this sugar) and mannitol. The different species of the genus can be differentiated by the results of auxanogrammes (137 substrates were used in the study of Gillis et al. 1995) [137].

This genus contains 39 species including rhizospheric bacteria, human pathogenic bacteria, as well as nitrogen fixing bacteria as *Burkholderia vietnamiensis* and *Burkholderia kururiensis*. Nevertheless, the genus also includes important species to the ecology of environment. In particular *Burkholderia xenovorans* (formerly known as *Pseudomonas cepacia*, then *Burkholderia cepacia* and *Burkholderia fungorum*) was studied for its faculties of deterioration of the organochlorine links in pesticides and polychlorobiphenyl (PCB).

Following the discoveries of Moulin et al. (2001) and Chen et al. (2003) [4; 111], showing the ability of bacteria to the subclass β -Proteobacteria (*Burkholderia* and *Ralstonia*) to induce nodulation in legumes, bacteria of the subclass γ -Proteobacteria (*Pseudomonas sp.*) were also isolated from legume nodules [24]. Based on these findings, it is not excluded to find other bacteria able to nodulate legumes, even outside the Proteobacteria phylum.

4.3. Other nitrogen-fixing legume symbionts

Recently, a number of isolates have been reported from legume nodules, capable of nitrogen fixation but phylogenetically located outside the traditional groups of rhizobia in the α -Proteobacteria. New lines that contain nitrogen-fixing legume symbionts include *Devosia*, *Ochrobactrum* and *Microvirga* in the α -Proteobacteria and *Cupriavidus* in the β -Proteobacteria.

In the α -Proteobacteria, *Devosia neptuniae* was proposed for strains from *Neptunia natans* from India [82]. *Ochrobactrum lupini* was described for nodule isolates from *Lupinus Albus* [90] and *Ochrobactrum cytisi* was described for nodule isolates from *Cytisus* [89]. *Microvirga lupini* was described for nodule isolates from *Lupinus sp.*, *M. lotononidis* and *M. zambiensis* were described for nodule isolates from different legume host [88].

In 2003 the nodulation of *Mimosa* by *Ralstonia taiwanensis* or *Wautersia taiwanensis* was published [86]. This species was isolated from nodules of *Mimosa pudica* and *Mimosa diplotricha*. In this case the strain was erroneously classified as *Ralstonia* and it has been later named *Cupriavidus taiwanensis* [116]. At present, *Cupriavidus* is a β - Proteobacteria, belonging to family

Burkholderiaceae of order *Burkholderiales*. Several species of this genus were isolated from soil and human clinical specimens but *C. taiwanensis* was the only species able to form effective nodules and fixes atmospheric nitrogen in legumes.

5. Conclusion

From one genus and 4 species in 1981, the rhizobia taxonomic studies have led to a total of 13 genera and 98 recognized species. This is in constant progress due to various studies in the world. Progress in taxonomy is also due to increasing numbers of effective techniques available in the characterization of bacteria, more accessible to laboratories.

So far only 23% of the total number of legumes species (between 16 500 and 19 700) were characterized for their microsymbionts. The majority (88%) of the studied legumes proved to be able of forming nodules. Specifically, tropical rhizobia are still poorly documented compared to what can be expected based on recent data, which show a great diversity in China, Brazil, Senegal, Sudan, Morocco.

In 2001, Sy et al. reported that some rhizobia isolated from *Crotalaria* sp. in Senegal were phylogenetically separated from all known rhizobia. They are a new species of the genus *Methylobacterium*, for which the name of *M. nodulans* was proposed. These finding suggests that several other new groups of symbiotic leguminous plant's bacteria (genera, species) may emerge in the future.

6. References

1. Dai J, Liu X and Wang Y. Genetic diversity and phylogeny of rhizobia isolated from *Caragana microphylla* growing in desert soil in Ningxia, China. Genet. Mol. Res. 2012; 11 (3): 2683-2693.
2. Weir BS. NZ rhizobia. Submitted by Bevan Weir on 27 April, 2013. Available: <http://www.rhizobia.co.nz/>
3. Zakhia F, Jeder H, Domergue O, Willems A, Cleyet-Marel JC, Gillis M, Dreyfus B, and de Lajudie P. Characterisation of wild legume nodulating bacteria (LNB) in the infra-arid zone of Tunisia. Syst. Appl. Microbiol. 2004; pp 380-395.
4. Moulin L, Munive A, Dreyfus B, and Boivin-Masson C. Nodulation of legumes by members of the β -sb class of Proteobacteria. Nature. N°41. 2001; 9 48-950.
5. Shiraishi A, Matsushita N, Hougetsu T. Nodulation in black locust by the Gamma- proteobacteria *Pseudomonas* sp. and the Beta-proteobacteria *Burkholderia* sp. Syst Appl. Microbiol. 2010; 33:269-74.
6. Allen EK. The Leguminosae, A Source Book of Characteristics, Uses, and Nodulation. University of Wisconsin Press. 1981.

7. Rivas R, García-Fraile P and Velázquez E. Taxonomy of Bacteria nodulating Legumes. *Microbiology Insights*. 2009; 2: 51–69.
8. Mullis KB, Faloona FA. Specific synthesis of DNA *in vitro* via a polymerase catalysed chain reaction. *Methods Enzymol*. 1987; 155: 335-350.
9. Graham PH, Sadowsky MJ, Kersters HH, Barnett YM, Bradley RS, Cooper JE, De Ley DJ, Jarvis BDW, Roslycky EB, Strijdom BW and Young JPW. Proposed minimal standards for the description of new genera and species of root-and stem-nodulating bacteria. *Int. J. Syst. Bacteriol*. 1991; 41: 582-587.
10. Vandamme P, Pot B, Gillis M, De Vos P, Kersters K and Swings J. Polyphasic taxonomy, a consensus approach to bacterial systematics. *Microbiol. Rev*. 1996; 60: 407-438.
11. Gillis MP, Vandamme P, De Vos JY, Swings Kersters K. Polyphasic taxonomy. 43-48. In *Bergey's Manual of Systematic Bacteriology*, 2nd Edition, Volume I. Boone DR, Castenholz RW, Garrity GM (eds.) 2001.
12. Oberreuter HJ, Chrzinski and S. Scherer. Intraspecific diversity of *Brevibacterium linens*, *Corynebacterium glutamium* and *Rhodococcus erythropolis* based on partial 16S rDNA sequence analysis and Fourier-transform infrared (FT-IR) spectroscopy. *Microbiology*. 2002; 148:1523-1532.
13. Perret X, and WJ Broughton. Rapid identification of *Rhizobium* strains by targeted PCR fingerprinting. *Plant & Soil*. 2002; 204: 21-34.
14. Sharon D, Smith SJ and Brown PO. A DNA microarray system for analyzing complex DNA samples using two-color fluorescent probe hybridization. *Genome Res*. 1996; 6: 639-645.
15. Rüberg S, Tian ZX, Krol E, Linke B, Meyer F, Wang Y, Pühler A, Weidner S and Becker A. Construction and validation of a *Sinorhizobium meliloti* whole genome DNA microarray: genome-wide profiling of osmoadaptive gene expression. *J. Biotechnol*. 2003, 106, 2556.
16. Ferreira L, Sánchez JF, García FP, Rivas R, Pedro FM, Martínez-Molina E, González-Buitrago JM, and Velázquez E. MALDI-TOF Mass Spectrometry Is a Fast and Reliable Platform for Identification and Ecological Studies of Species from Family *Rhizobiaceae*. *Plos One*; 2011; 6 (5) : e20223.
17. Zakhia F, De Lajudie P. Taxonomy of rhizobia. *Agronomie*. 2001; 21: 569-576.
18. Terefework Z, Nick G, Suomalainen S, Paulin L and Lindström K. Phylogeny of *Rhizobium galegae* with respect to other rhizobia and agrobacteria. *Int. J. Syst. Bacteriol*. 1998; 48: 349-356.
19. De Bruijn FJ, Mc Spadden-Gadener ME, Millcamp B, Rademaker JL, Ragatz W, Schultz D, Struffi ML and Stoltzfus J. Molecular approaches in microbial ecology to assess genomic diversity and stress induced gene expression in plant-associated diazotrophs. In *Nitrogen Fixation: fundamentals and*

applications, pp. 571-576. Edited by Elmerich C, Kondorosi A and Newton WE. Kluwer. The Netherlands. 1997.

20. Vincent JM. A Manual for the Practical Study of Root Nodule Bacteria. IBP handbook, no. 15. Blackwell Scientific Publications, Ltd., Oxford, England. 1970.

21. Saoudi M. Les bactéries nodulant les légumineuses (B.N.L) : Caractérisation des bactéries associées aux nodules de la légumineuse *Astragalus armatus*. Mémoire de Magister. Université Mentouri de Constantine. Faculté des Sciences de la Nature et de la Vie. Algérie. (2008). French.

22. Chen WM, Laevens S, Lee TM, Coenye T, De Vos P, Mergeay M, Vandamme P. *Ralstonia taiwanensis* sp. nov., isolated from root nodules of *Mimosa* species and sputum of a cystic fibrosis patient. P. Int. J. Syst. Evol. Microbiol. 2001; 51:1729-35.

23. Nogom A, Nakagawa Y, Sawada H, Tsukahara J, wakabayashi S, Uchiumi T, Nuntagij A, kotepong S, Suzuki A, Higashi S and Abe M. A novel symbiotic nitrogen-fixing member of the *Ochrobacterium* clade isolated from root nodules of *Acacia mangium*. J. gen. Appl. Microbiol. 2004; 50, 1727.

24. Benhizia Y, Goudjil H, Benguedouar A, Rosella M, Giacomini A, Squartini A. Gamma proteobacteria can nodulate legumes of the genus *Hedysarum*. System Appl Microbiol. 2004; 27: 462-468.

25. Frank B. Über die Pilzsymbiose der Leguminosen. Ber. Dtsc. Bot. Ges. 1889; 7: 332-346.

26. Jordan DC. Transfer of *rhizobium japonicum* Buchanan 1980 to *Bradyrhizobium* gen. nov., a genus of slow growing root nodule bacteria from leguminous plants, Int. J. Syst. Bacteriol. 1982; 32: 136-139.

27. Lindström K. *Rhizobium galegae*, a new species of legume root bacteria. Int. J. Syst. Bacteriol. 1989; 39: 365-367.

28. Martinez-Romero E, Segovia L, Mercante FM, Franco AA, Graham P and Pardo MA. *Rhizobium tropici*, a novel species nodulating *Phaseolus vulgaris* L. Beans and *Leucaena* sp. Trees. Int. J. Syst. Bacteriol. 1991; 41: 417-426.

29. Renan Augusto Ribeiro, Marco A. Rogel, Aline Lopez-Lopez, Ernesto Ormeno-Orrillo, Fernando Gomes Barcellos, Julio Martinez, Fabiano Lopes Thompson, Esperanza Martinez-Romero and Mariangela Hungria. Reclassification of *Rhizobium tropici* type A strains as *Rhizobium leucaenae* sp. nov. Int. J. Syst. Evol. Microbiol. 2012; 62, 1179-1184.

30. López LA, Rogel MA, Ormeño OE, Martínez-Romero J, Martínez-Romero E. Phaseolus vulgaris seed-borne endophytic community with novel bacterial species such as *Rhizobium endophyticum* sp. nov. Syst Appl Microbiol. 2010; 33: 322-7.

31. Tian CF, Wang ET, Wu LJ, Han TX, Chen WF, Gu CT, Gu JG and Chen WX. *Rhizobium fabae* sp. nov., a bacterium that nodulates *Vicia faba*. Int. J. Syst. Evol. Microbiol., 2008, **58**, 2871-2875.
32. Segovia L, Young JPW and Martinez-Romero. Reclassification of American *Rhizobium leguminosarum* biovar *phaseoli* type I strains as *Rhizobium etli* sp. nov. Int. J. Syst. Bacteriol. 1993; 43: 374-377.
33. Wang LX, Antonio Rogel M, los Santos AG, Martnez-Romero J, Cevallos MA and Martinez-Romero. *Rhizobium etli* bv. *mimosae*, a novel biovar isolated from *Mimosa affinis*. Int. J. Syst. Evol. Microbiol. 1999b; 49: 1475-1491.
34. Souza V, Eguiarte L, Avila G, Cappello R, Gallardo C, Montoya J, Piñero D. Genetic Structure of *Rhizobium etli* biovar *phaseoli* Associated with Wild and Cultivated Bean Plants (*Phaseolus vulgaris* and *Phaseolus coccineus*) in Morelos, Mexico. Appl. Envir. Microbiol. 1994; 60:1260-8.
35. de Lajudie P, Laurent-Fulele E, Willems A, Torck U, Coopman R, Collins MD, Kersters K, and Gillis M. *Allorhizobium undicola* gen. Nov. sp. nov., nitrogen-fixing bacteria that efficiently nodulate *Neptunia natans* in Senegal. Int.J. Syst.Bacteriol. 1998a; 48;1277-1290.
36. Amarger N, Macheret V and Laguerre G. *Rhizobium gallicum* sp. nov. and *Rhizobium giardinii* sp. nov. from *Phaseolus vulgaris* nodules. Int. J. Syst. Bacteriol. 1997; 47: 996-1006.
37. Chen WX, Yan GH, Li JL. Numerical taxonomic study of fast-growing soybean rhizobia and a proposal that *Rhizobium fredii* be assigned to *Sinorhizobium* gen. nov., Int. J. Syst. Bacteriol. 1988; 38: 392-397.
38. Wang ET, Van Berkum Beyene PD, Sui XH, Dorado O, Chen WX and Martinez-Romero E. *Rhizobium huautlense* sp. nov., a symbiont of *Sesbania herbacea* which has a close phylogenetic relationships with *Rhizobium galegae*. Int. J. Syst. Evol. Microbiol. 1998; 51: 1023-1026.
39. Van Berkum P, Beyene D, Bao G, Campbell TA and Eardly BD. *Rhizobium mongolense* sp. nov. in one of three rhizobial genotypes identified which nodulate and form nitrogen fixing symbioses with *Medicago ruthenica* [(L.) Ledebour]. Int. J. Syst. Evol. Microbiol. 1998; 48: 13-22.
40. Tan ZY, Kan FL, Peng GX, Wang ET, Reinhold-Hurek B and Chen WX. *Rhizobium yanglingense* sp. nov., isolated from arid and semi-arid regions in China. Int. J. Syst. Evol. Microbiol. 2001; 51, 909-914.
41. Bouzar HD. and Jones JB. *Agrobacterium Larrymoorei* sp. nov., a pathogen isolated from aerial tumors of *Ficus benjamina*. Int. J. Syst. Evol. Microbiol. 2001; 51: 1023-1026.
42. Wei GH, Wang ET, Tan ZY, Zhu ME and Chen WX. *Rhizobium indigoferae* sp. nov. and *Sinorhizobium kummerowiae* sp. nov., respectively isolated from *Indigofera spp.* and *Kummerowia stipulacea*. Int. J. Syst. Evol. Microbiol. 2002; 52: 2231-2239.
43. Squartini A, Struffi P, Döring H, Seleka-Pobell S, Tola E, Giacomini A, Vendramin E, Velásquez E, Mateos PF, Martinez-Molina E, Dazzo FB, Casella S and Nuti MP. *Rhizobium sullae* sp. nov.

- (formerly "*Rhizobium hedysari*"), the rot nodule microsymbiont of *Hedysarum coronarium* L. Int. J. Syst. Evol. Microbiol. 2002; 52: 1267-1276.
44. Wei GH, Tan ZY, Zhu ME, Wang ET, Han SZ and Chen WX. Characterization of rhizobia isolated from legume species within the genera *Astragalus* and *Lespedeza* grown in the Loess Plateau of China and description of *Rhizobium Loessense* sp. nov. Int. J. Syst. Evol. Microbiol. 2003; 53: 1575-1583.
 45. Garcia-Fraile P, Rivas R, Willems A, Peix A, Martens M, Martinez-M, Eustoquio M, Pedro F, Velazquez E. *Rhizobium cellulosilyticum* sp. nov., isolated from sawdust of *Populus alba*. Int J Syst Evol Microbiol. 2007; 57: 844-848.
 46. Gu CT, Wang ET, Tian CF, Han TX, Chen WF, Sui XH and Chen WX. *Rhizobium miluonense* sp. nov., a symbiotic bacterium isolated from *Lespedeza* root nodules. Int J Syst Evol Microbiol. 2008; 58: 1364-1368.
 47. Han TX, Han LL, Wu LJ, Chen WF, Sui XH, Gu JG, Wang ET and Chen WX. *Mesorhizobium gobiense* sp. nov. and *Mesorhizobium tarimense* sp. nov., isolated from wild legumes growing in desert soils of Xinjiang, China. Int J Syst Evol Microbiol. 2008; 58(11): 2610-2618.
 48. Peng G, Yuan Q, Li H, Zhang W, Tan Z. *Rhizobium oryzae* sp. nov., isolated from the wild rice *Oryza alta*. Int J Syst Evol Microbiol. 2008; 58: 2158-2163.
 49. Ramirez-Bahena MH, Garcia-Fraile P, Peix A, Valverde A, Rivas R, Igual Jose MM, Pedro F, Martinez-Molina E, Velazquez E. Revision of the taxonomic status of the species *Rhizobium leguminosarum* (Frank 1879) Frank 1889AL, *Rhizobium phaseoli* Dangeard 1926AL and *Rhizobium trifolii* Dangeard 1926AL. *R. trifolii* is a later synonym of *R. leguminosarum*. Reclassification of the strain *R. leguminosarum* DSM 30132 (=NCIMB 11478) as *Rhizobium pisi* sp. nov. Int J Syst Evol Microbiol. 2008; 58: 2484-2490.
 50. Lin DX, Wang ET, Tang HH, Tian XH, Yu Rong G, Su H, Chen WX. *Shinella kummerowiae* sp. nov., a symbiotic bacterium isolated from root nodules of the herbal legume *Kummerowia stipulacea*. Int J Syst Evol Microbiol. 2008; 58: 1409-1413.
 51. Berge O, Lodhi A, Brandelet G, Santaella C, Roncato MA, Christen R, Heulin T, Achouak W. *Rhizobium alamii* sp. nov., an exopolysaccharide-producing species isolated from legume and non-legume rhizospheres. Int J Syst Evol Microbiol. 2009; 59: 367-372.
 52. Lu YL, Chen W., Han LL, Wang ET, Zhang XX, Chen WX, Su HZ. *Mesorhizobium shangrilense* sp. nov., isolated from root nodules of *Caragana* spp. Int J Syst Evol Microbiol. 2009; 59: 3012-3018.
 53. Hou BC, Wang ET, Li Y, Jr J, Rui Z, Chen WF, Gao YD, Ren J, Chen WX. *Rhizobium tibeticum* sp. nov., a symbiotic bacterium isolated from *Trigonella archiducis-nicolai* (Sirj.) Vassilcz. Int. J. Syst. Evol. Microbiol. 2009; 59: 3051-3057.

54. El Akhal MR. Diversité des rhizobiums nodulant l'arachide (*Arachis hypogaea* L.) au Maroc et l'effet de la salinité sur le couple symbiotique. Thèse de doctorat de l'Université abdelmalek essaadi, Faculté des Sciences et Techniques- Tanger, Maroc; 2008. French.
55. Young JM, Kuykendall LD, Martinez-Romero E, Kerr A, Sawada H. A revision of *Rhizobium* Frank 1889, with an emended description of the genus, and the inclusion of all species of *Agrobacterium* Conn 1942 and *Allorhizobium undicola* de Lajudie et al. 1998 as new combinations: *Rhizobium radiobacter*, *R. rhizogenes*, *R. rubi*, *R. undicola* and *R. vitis*, Int. J. Syst. Evol. Microbiol. 2001; 51:89-103.
56. Pulawska J, Willems A, De Meyer SE and Sule S. *Rhizobium nepotum* sp. nov. isolated from tumors on different plant species. Syst. Appl. Microbiol., 2012, 35, 215-220.
57. Dangeard PA. Recherches sur les tubercules radicaux des légumineuses. Le botaniste, Ser.16. 270 p. Paris. 1926. French.
58. Scholla MH and Elkan GH. *Rhizobium fredii* sp. nov., a fast growing species that effectively nodulates soybean. Int. J. System. Bact. 1984; 34: 484-486.
59. de Lajudie P, Willems A, Pot B et al. Polyphasic taxonomy of rhizobia: emendation of the genus *Sinorhizobium* and description of *Sinorhizobium meliloti* comb. nov., *Sinorhizobium saheli* sp. nov., and *Sinorhizobium teranga* sp. nov. Int. j. Syst. Bacteriol. 1994; 44, 715-733.
60. Lorquin J, Lortet G, Ferro M, Mear N, Dreyfus B, Prome JC and Boivin C. Nod factors from *Sinorhizobium saheli* and *S. teranga* bv. *sesbaniae* are both arabinosylated and fucosylated, a structural feature specific to *Sesbania rostrata* symbionts. Mol. Plant-Microbe Interact. 1997b; 10: 879-890.
61. Rome S, Fernandez MP, Brunel B, Normand P and Cleyet-Marel JC. *Sinorhizobium medicae* sp. nov., isolated from annual *Medicago* spp. Int. J. Syst. Evol. Microbiol. 1996; 46: 972-980.
62. Nick G, de Lajudie P, Eardly BD, Suomalainen S, Plin L, Zhang X, Gillis M and Lindstrom K. *Sinorhizobium arboris* sp. Nov. and *Sinorhizobium kostiense* sp. Nov. isolated from leguminous trees in Sudan and Kenya. Int. J. Syst. Bacteriol. 1999; 49: 1359-1368.
63. Chen WX, Yan GH and Li JL. Numerical taxonomic study of fast-growing soybean rhizobia and a proposal that *Rhizobium fredii* be assigned to *Sinorhizobium* gen. nov. Int. J. Syst. Bacteriol., 1988, 38, 392-397.
64. Wang FQ, Wang ET, Liu J, Chen Q et al. *Mesorhizobium albiziae* sp. nov., a novel bacterium that nodulates *Albizia kalkora* in a subtropical region of China. Int. J. Syst. Evol. Microbiol. 2007; 57: 1192-1199.
65. Toledo I, Lloret L and Martinez-Romero E. *Sinorhizobium americanus* sp. nov., a new *Sinorhizobium* species nodulating native *Acacia* spp. in Mexico. Syst. Appl. Microbiol. 2003; 26: 54-64.

66. Lloret L, Ormeno-Orrillo E, Rincon R, Martinez-Romero J, Rogel-Hernandez MA, Martinez-Romero E. *Ensifer mexicanus* sp. nov. a new species nodulating *Acacia angustissima* (Mill.) Kuntze in Mexico. Syst Appl Microbiol. 2007.
67. Merabet C, Martens M, Mahdhi M, Zakhia F, Sy A, Le Roux C, Domergue O, Coopman R, Bekki A, Mars M, Willems A, de Lajudie P. Multilocus sequence analysis of root nodule isolates from *Lotus arabicus* (Senegal), *Lotus creticus*, *Argyrolobium uniflorum* and *Medicago sativa* (Tunisia) and description of *Ensifer numidicus* sp. nov. and *Ensifer garamanticus* sp. nov. Int J Syst Evol Microbiol. 2010 ; 60: 664-674 .
68. Jarvis BDW, Pankhurst CE, and Pastel JJ. *Rhizobium loti*, a new species of legume root nodule bacteria. Int. J. Syst. Bacteriol. 1982; 32: 378-380.
69. Chen H, Richardson AE, Cartner E, Diordjevic MA, Roughley RJ and Rolfe BG. Construction of an acid-tolerant *Rhizobium leguminosarum* biovar *trifolii* strain with enhanced capacity for nitrogen fixation. Appl. Environ. Microbiol. 1991; 57: 2005-2011.
70. Nour SM, Fernandez MP, Normand P and Cleyet-Marel JC. *Rhizobium ciceri* sp. nov., consisting of strains that nodulate chickpea (*Cicer arietinum* L.). Int. J. Syst. Bacteriol. 1994a; 44: 511-522.
71. Nour SM, Cleyet-Marel JC, Normand P, Fernandez MP. Genomic heterogeneity of strains nodulating chickpeas (*Cicer arietinum* L.) and description of *Rhizobium mediterraneum* sp. nov., Int. J. Syst. Bacteriol. 1995; 45: 640-648.
72. de Lajudie P, Willems A, Nick G, Moreira A, Molouba F, Hoste B, Torck U, Neyra M, Collins MD, Lindstrom K, Dreyfus B and Gillis M. Characterisation of tropical tree rhizobia and description of *Mesorhizobium plurifarum* sp. Nov. Int.J.Syst. Bacteriol. 1998b; 48: 369-382.
73. Wang ET, Van Berkum P, Sui XH, Beyene D, Chen WX and Martinez-Romero E. Diversity of rhizobia associated with *Amorpha fruticosa* isolated from Chinese soils and description of *Mesorhizobium amorpha* sp. nov. Int J Syst Evol Microbiol. 1999a; 49: 51-65.
74. Velásquez, E, Iguá JM.I, Williams Murez A, Mateos PF, Abril A, Toro N, Normand P, Cervantes E, Gillis G and Martnez-Romero E. *Mesorhizobium Chocoense* sp. nov., a novel species that nodulates *Prosopis alba* in the Chaco Arido region (Argentina). Int. J. Syst. Bacteriol. 2001; 51: 1011-1021.
75. Gao JL, Turner SL, Kan FL, Wang ET, Tan ZY, Qiu YH, Gu J, Terefework Z, Young JPW, Lindström K and Chen WX. *Mesorhizobium septentrionale* sp. nov. and *Mesorhizobium temperatum* sp. nov., isolated from *Astragalus adsurgens* growing in the northern regions of China. Int. J. Syst. Evol. Microbiol. 2004; 54: 2003-2012.
76. Nandasena, Kemanthi G, O'Hara, Graham W, Tiwari RP, Willems A, Howieson JG. *Mesorhizobium australicum* sp. nov. and *Mesorhizobium opportunistum* sp. nov., isolated from *Biserrula pelecinus* L. in Australia Int J Syst Evol Microbiol. 2009; 59: 2140-2147.

77. Vidal C, Chantreuil C, Berge OM, Lucette E, Jose BG, Brunel B, Cleyet-Marel JC. *Mesorhizobium metallidurans* sp. nov., a metal-resistant symbiont of *Anthyllis vulneraria* growing on metalicolous soil in Languedoc, France. *Int J Syst Evol Microbiol.* 2009; 59: 850-855.
78. Chen WM, Zhu WF, Bontemps C, Young J, Peter W, Wei GH. *Mesorhizobium camelthorni* sp. nov., isolated from *Alhagi sparsifolia*. *Int J Syst Evol Microbiol.* 2011 ; 61: 574-579.
79. Degefu T, Wolde-Meskel E, Liu B, Clennwerck I, Willems A and Frostegard A. *Mesorhizobium shonense* sp. nov., *Mesorhizobium hawassense* sp. nov. and *Mesorhizobium abyssinicae* sp. nov., isolated from root nodules of different agroforestry legume trees. *Int. J. Syst. Evol. Microbiol.*, 2013, 63, 1746-1753.
80. Zhang JJ, Liu TY., Chen WF, Wang ET, Sui XH, Zhang XX, Li Y, Li Y, Chen WX. *Mesorhizobium muleiense* sp. nov., nodulating with *Cicer arietinum* L. in Xinjiang, China. *Int. J. Syst. Evol. Microbiol.*, 2012 ; 62, 2737-2742.
81. Zheng WT, Li Y, Wang R, Sui XH, Zhang XX, Zhang JJ, Wang ET and Chen WX. *Mesorhizobium qingshengii* sp. nov., isolated from effective nodules of *Astragalus sinicus*. *Int. J. Syst. Evol. Microbiol.*, 2013, 63, 2002-2007.
82. Zhou PF, Chen WM and Wei GH. *Mesorhizobium robiniae* sp. nov., isolated from root nodules of *Robinia pseudoacacia*. *Int. J. Syst. Evol. Microbiol.*, 2010, 60, 2552-2556.
83. Lu YL, Chen WF, Wang ET, Han LL, Zhang XX, Chen WX and Han SZ. *Mesorhizobium shangrilense* sp. nov., isolated from root nodules of *Caragana* species. *Int. J. Syst. Evol. Microbiol.*, 2009, 59, 3012-3018.
84. Zhao CT, Wang ET, Zhang YM, Chen WF, Sui XH, Chen WX, Liu HC and Zhang XX. *Mesorhizobium silamurunense* sp. nov., isolated from root nodules of *Astragalus* species. *Int. J. Syst. Evol. Microbiol.*, 2012, 62, 2180-2186.
85. Ramirez-Bahena MH, Hernandez M, Peix A, Velaquez E and Leon-Barrios M. *Mesorhizobial* strains nodulating *Anagyris latifolia* and *Lotus berthelotii* in Tamadaya ravine (Tenerife, Canary Islands) are two symbiovars of the same species, *Mesorhizobium tamadayense* sp. nov. *Syst. Appl. Microbiol.*, 2012, 35, 334-341.
86. Valverde A, Velazquez E, Fernandez-Santos F, Vizcaino N, Rivas R, Gillis M, Mateos P F, Martinez-Molina E, Igual JM and Willems. *Phyllobacterium trifolii* sp. nov. nodulating *Trifolium* and *Lupinus* in Spanish soils. *Int. J. Syst. Evol. Microbiol.* 2005; 55: 1985-1989.
87. Jourand P, Giraud E, Bena G, Sy A, Willems A, Gillis M, Dreyfus B & de Lajudie P. *Methylobacterium nodulans* sp. nov., for a group of aerobic, facultatively methylotrophic, legume rootnodule-forming and nitrogen-fixing bacteria. *International Journal of Systematic and Evolutionary Microbiology.* 2004; 54: 2269-2273.
88. Ardley JK, Parker MA, De Meyer SE, Trengove RD, O'hara GW, Reeve WG, Yates RJ, Dilworth MJ, Willems A and Howieson JG. *Microvirga lupini* sp. nov., *Microvirga lotononidis* sp. nov. and *Microvirga zambiensis* sp. nov. are alphaproteobacterial root-nodule bacteria that specifically nodulate

and fix nitrogen with geographically and taxonomically separate legume hosts. *Int. J. Syst. Evol. Microbiol.*, 2012, 62, 2579-2588.

89. Zurdo-Piñeiro JL, Rivas R, Trujillo ME, et al. *Ochrobactrum cytisi* sp. nov., isolated from nodules of *Cytisus scoparius* in Spain. *Int J Syst Evol Microbiol.* 2007; 57:784–8.

90. Trujillo ME, Willems A, Abril A, Planchuelo AM, Rivas R, Ludena D, Mateos PF, Martinez-Molina E and Velazquez E. Nodulation of *Lupinus* by strains of *Ochrobactrum lupini* sp. nov. *Appl. Environ. Microbiol.* 2005. 71, 1318–1327.

91. Dreyfus B, Garcia JL, and Gillis M. Characterization of *Azorhizobium caulinodans* gen. Nov., sp. nov., a stem-nodulating nitrogen-fixing bacterium isolated from *Sesbania rostrata*. *Int. J. Syst. Bacteriol.* 1988; 38: 89-98.

92. Rinaudo G, Orenga S, Fernandez MP, Meugnier and Bardin R. DNA homologies among members of the genus *Azorhizobium* and other stem- and root-nodulating bacteria isolated from the tropical legume *Sesbania rostrata*. *Int. J. Syst. Bacteriol.* 1991; 41: 114-120.

93. Jordan DC. Family III. Rhizobiaceae, p.234-242. In N. R. Krieg and J. G. Holt (ed.), *Bergey's Manual of Systematic Bacteriology*, vol.1. The Williams & Wilkins Co., Baltimore. 1984.

94. Kuykendall LD, Saxena B, Devine TE and Udell SE. Genetic diversity in *bradyrhizobium japonicum* Jordan 1982 and a proposal for *bradyrhizobium elkanii* sp nov. *Can. J. Microbiol.* 1992; 38: 501-505.

95. Xu LM, Ge C, Cui Z, Li and Fan H. *Bradyrhizobium liaoningense* sp. nov. isolated from the root nodules of soybeans. *Int. J. Syst. Bacteriol.* 1995; 45: 706-711.

96. Yao ZY, Kan FL, Wang ET, Wei GH and Chen WX. Characterization of rhizobia that nodulate legume species of the genus *Lespedeza* and description of *Bradyrhizobium yuanmingense* sp. nov. *Int. J. Syst. Bacteriol.* 2002; 52: 2219-2230.

97. Rivas R, Willems A, Palomo JL, Garcia-Benavides P, Mateos PF, Martinez-Molina E, Gillis M. and Velazquez E. "*Bradyrhizobium betae* sp. nov. isolated from roots of *Beta vulgaris* affected by tumor-like deformations." *Int. J. Syst. Evol. Microbiol.* 2004; 54: 1271-1275.

98. Vinuesa P, Leon-Barrios M, Silva C, Willems A, Jarabo-Lorenzo A, Perez Galdona R, Werner D and Martinez-Romero E. *Bradyrhizobium canariense* sp. nov., an acid-tolerant endosymbiont that nodulates endemic genistoid legumes (Papilionoideae: Genisteae) from the Canary Islands, along with *Bradyrhizobium japonicum* bv. *genistearum*, *Bradyrhizobium* genospecies alpha and *Bradyrhizobium* genospecies beta. *Int. J. Syst. Evol. Microbiol.* 2005; 55: 569-575.

99. Islam MS, Kawasaki H, Muramatsu Y, Nakagawa Y and Seki T. "*Bradyrhizobium iriomotense* sp. nov., isolated from a tumor-like root of the legume *Entada koshunensis* from Iriomote Island in Japan." *Biosci. Biotechnol. Biochem.* 2008; 72:1416-1429.

100. Jourand P, Giraud E, Bena G, Sy A, Willems A, Gillis M, Dreyfus B & de Lajudie P. *Methylobacterium nodulans* sp. nov., for a group of aerobic, facultatively methylotrophic, legume rootnodule-forming and nitrogen-fixing bacteria. *International Journal of Systematic and Evolutionary Microbiology*. 2004; 54: 2269-2273.
101. Chahbourne R, Carro L, Peix A, Barrijal S, Velasquez E and Bedmar EJ. *Bradyrhizobium cytisi* sp. nov., isolated from effective nodules of *Cytisus villosus*. *Int. J. Syst. Evol. Microbiol.*, 2011, 61, 2922-2927.
102. Wang JY, Wang R, Zhang YM, Liu HC, Chen WF, Wang ET, Sui XH and Chen WX. *Bradyrhizobium daqingense* sp. nov., isolated from soybean nodules. *Int. J. Syst. Evol. Microbiol.*, 2013, 63, 616-624.
103. Van Berkum P, Leibold JM and Eardly BD. Proposal for combining *Bradyrhizobium* spp. (*Aeschynomene indica*) with *Blastobacter denitrificans* and to transfer *Blastobacter denitrificans* (Hirsh and Muller, 1985) to the genus *Bradyrhizobium* as *Bradyrhizobium denitrificans* (comb. nov.). *Syst. Appl. Microbiol.*, 2006, 29, 207-215.
104. Ramirez-Bahana MH, Chahboune R, Peix A and Velasquez E. Reclassification of *Agromonas oligotrophica* into the genus *Bradyrhizobium* as *Bradyrhizobium oligotrophicum* comb. nov. *Int. J. Syst. Evol. Microbiol.*, 2013, 63, 1013-1016.
105. Ramirez-Bahana MH, Peix A, Rivas R, Camacho M, Rodriguez-Navarro DN, Mateos PF, Martinez-Molina E, Willems A and Velasquez E. *Bradyrhizobium pachyrhizi* sp. nov. and *Bradyrhizobium jicamae* sp. nov., isolated from effective nodules of *Pachyrhizus erosus*. *Int. J. Syst. Evol. Microbiol.*, 2009, 59, 1929-1934.
106. Van Berkum P, Eardly BD. The aquatic budding bacterium *Blastobacter denitrificans* is a nitrogenfixing symbiont of *Aeschynomene indica*. *Appl and Env Microbiol.* 2002; 68 : 1132-1136.
107. Achouak W, Christen R, Barakat M, Martel MH and Heulin T. *Burkholderia caribensis* sp. nov., an exopolysaccharide-producing bacterium isolated from vertisol microaggregates in Martinique. *Int. J. Syst. Bacteriol.*, 1999, 49, 787-794.
108. Vandamme P, Goris J, Chen WM, De Vos P and Willems A. *Burkholderia tuberum* sp. nov. and *Burkholderia phymatum* sp. nov., nodulate the roots of tropical legumes. *Syst. Appl. Microbiol.*, 2002, 25, 507-512. VALIDATION LIST no. 91. *Int. J. Syst. Evol. Microbiol.*, 2003, 53, 627-628.
109. Chen WM, De Faria SM, James EK, Elliott GN, Lin KY, Chou JH, Sheu SY, Cnockaeret M, Sprent JI and Vandamme P. *Burkholderia nodosa* sp. nov., isolated from root nodules of the woody Brazilian legumes *Mimosa bimucronata* and *Mimosa scabrella*. *Int. J. Syst. Evol. Microbiol.*, 2007, 57, 1055-1059.

110. Chen WM, De Faria SM, Chou JH, James EK, Elliott GN, Sprent JI, Bontemps C, Young JPW and Vandamme P. *Burkholderia sabiae* sp. nov., isolated from root nodules of *Mimosa caesalpiniiifolia*. *Int. J. Syst. Evol. Microbiol.*, 2008, 58, 2174-2179.
111. Chen WM, Moulin L, Bontemps C, Vandamme P, Bena G, Boivin-Masson C. Legume symbiotic nitrogen fixation by beta-proteobacteria is widespread in nature. *J Bact.* 2003; 185, 7266-7272.
112. Partida-Martinez LP, Groth I, Schmitt I, Richter W, Roth M and Hertweck C. *Burkholderia rhizoxinica* sp. nov. and *Burkholderia endofungorum* sp. nov., bacterial endosymbionts of the plant-pathogenic fungus *Rhizopus microsporus*. *Int. J. Syst. Evol. Microbiol.*, 2007, 57, 2583-2590.
113. Sheu SY, Chou JH, Bontemps C, Elliot GN, Gross N, Dos Reis Junior FB, Melkonian R, Moulin L, James EK, Sprent JI, Yong JPW and Chen WM. *Burkholderia diazotrophica* sp. nov., isolated from root nodules of *Mimosa* spp. *Int. J. Syst. Evol. Microbiol.*, 2013, 63, 435-441.
114. Aizawa T, Ve NB, Nakajima M and Sunairi M. *Burkholderia heleia* sp. nov., a nitrogen-fixing bacterium isolated from an aquatic plant, *Eleocharis dulcis*, that grows in highly acidic swamps in actual acid sulfate soil areas of Vietnam. *Int. J. Syst. Evol. Microbiol.*, 2010, **60**, 1152-1157.
115. SHEU (S.Y.), Chou JH, Bontemps C, Elliott GN, GROSS (E.), JAMES (E.K.), Sprent JI, Young JPW and Chen WM. *Burkholderia symbiotica* sp. nov., isolated from root nodules of *Mimosa* spp. native to north-east Brazil. *Int. J. Syst. Evol. Microbiol.*, 2012, **62**, 2272-2278.
116. Vandamme Pand Coeyene T. Taxonomy of the genus *Cupriavidus*: a tale of lost and found. *Int. J. Syst. Evol. Microbiol.*, 2004, 54, 2285-2289.
117. Boivin MC, Bontemps C, Geoffroy G, Gris LC, Talini L. Détection et typage du gène à l'aide de biopuces à ADN : perspectives pour l'étude de la diversité et de l'écologie moléculaire des rhizobia. Les Actes du BRG. 2006 ; 6 : 97-110. French.
118. Prescott LM, Harley JP, Klein DA, Claire-Michèle BC, Dusart J. Microbiologie, chapitre 22. Les bactéries. Les protéobactérie,. Edition. De Boeck Université. p.492. 2007. French.
119. Han SZ, Wang ET, Chen WX, and Han SZ. Diverse bacteria isolated from root nodules of *Phaseolus vulgaris* and species within the genera *Campylotropis* and *Cassia* grown in China." *Systematic and Applied Microbiology*, 2005; 28(3): 265–276.
120. Liu J, Wang ET and Chen WX. Diverse rhizobia associated with woody legumes *Wisteria sinensis*, *Cercis racemosa* and *Amorpha fruticosa* grown in the temperate zone of China." *Systematic and Applied Microbiology*. 2005; 28(5): 465–477.
121. Velázquez E, Peix A, Zurdo-Piñeiro JL, Palomo JL, Mateos P, Rivas R, Muñoz Adelantado E, Toro N, García-Benavides P and Martínez-Molina E. "The coexistence of symbiosis and pathogenicity-determining genes in *Rhizobium rhizogenes* strains enables them to induce nodules and tumors or hairy roots in plants." *Molecular Plant-Microbe Interactions*. 2005; 18(12): 1325–1332.

122. Trüper HG, de Clari L. Taxonomic note: necessary correction of specific epithets formed as substantives (nouns) in "apposition", Int. J. Syst. Bacteriol. 1997; 47:908-909.
123. Garcia JL, Cayol JL et Roger P. Taxonomie des Procaryotes. *Rhizobiales*, Ordre VI des Alpha-proteobacteria, Classe 1 des Proteobactéries, Phylum BXIII du Domaine bacteria. Mis à jour en Mai 2012. http://garciajeanlouis9051.perso.neuf.fr/aaBXIII1_O6.html#Carbophilus.
124. Martens M, Dawyndt P, Coopman R, Gillis M, de Vos P, Willems A. Advantages of multilocus sequence analysis for taxonomic studies: a case study using 10 housekeeping genes in the genus *Ensifer* (including former *Sinorhizobium*). Int J Syst Evol Microbiol. 2008;58: 200–14.
125. Lin DX, Wang ET, Tang H, Han TX, He YR, Guan SH and Chen WX. *Shinella kummerowiae* sp. nov., a symbiotic bacterium isolated from root nodules of the herbal legume *Kummerowia stipulacea*. Inter J Syst Evol Microbiol. 2012; 62: 2579-2588.
126. Jarvis BDW, van Berkum P, Chen WX, Nour S, Fernandez MP, CleyetMarrel JC and Gillis M. Transfert of *Rhizobium loti*, *Rhizobium huakuii*, *Rhizobium ciceri*, *Rhizobium mediterraneum*, and *Rhizobium tianshanense* to *Mesorhizobium* gen. nov. Int. J. Syst. Bacteriol. 1997 ; 47: 895-898.
127. Wang ET, van Berkum P, Sui XH, Beyene D, Chen WX, Martinez-Romero E. Diversity of rhizobia associated with *Amorpha fruticosa* isolated from chinese soils and description of *Mesorhizobium amorphae* sp. nov., Int. J. Syst. Bacteriol. 1999; 49: 51-65.
128. Trinick MJ. Relationships amongst the fast-growing rhizobia of *Lablab purpureus*, *Leucaena leucocephala*, *Mimosa spp.*, *Acacia farnesiana* and *Sesbania grandiflora* and their affinities with other rhizobial groups. J. Appl. Bact. 1980; 49: 39-53.
129. Kuykendall LD. Genus *Azorhizobium* of the family *Hypomicrobiaceae*. Bergey's Manual of Systematic Bacteriology, 2nd Edition, 2nd Volume. 2004a.
130. Hollis AB, Kloos WE, and Elkan GH. Hybridization studies of *rhizobium japonicum* and related Rhizobiaceae. J. Gen. Microbiol. 1981; 123: 215-222.
131. Euzéby JP and Findall BJ. Status of strains that contravene rules 27(3) and 30 of the Bacteriological Code. Request for an Opinion. Int. J. Syst. Evol. Microbiol. 2004; 54:, 293-301.
132. Parker MA. Bradyrhizobia from wild *Phaseolus*, *Desmodium* and *Macroptilium* species in northern Mexico. Appl. Env. Microbiol. 2002; 68 : 2044-2048.
133. Patt TE, Cole GE, Hanson RS. *Methylobacterium*, a new genus of facultatively methylotrophic bacteria. International Journal of Systematic Bacteriology. 1976 ; 26: 226-229.
134. van Dien SJ, Marx CJ, O'Brien BN & Lidstrom ME. Genetic characterization of the carotenoid biosynthetic pathway in *Methylobacterium extorquens* AM1 and isolation of a colorless mutant. Applied and Environmental Microbiology. 2003; 69: 7563-7566.

135. Hourcade E. Dégradation du dichlorométhane et adaptation à la production intracellulaire d'acide chez *Méthylobacterium*. Thèse de Doctorat. Université Louis Pasteur, Strasbourg I. 2007. French.
136. Yabuuchi E, Kosako Y, Oyaizu H, Yano I, Hotta H, Hashimoto Y., Ezaki T, Arakawa M. Proposal of *Burkholderia* gen. nov. and transfer of seven species of the genus *Pseudomonas* homology group II to the new genus, with the type species *Burkholderia cepacia* (Palleroni and Holmes, 1981) com. nov. Microbiology and Immunology. 1992; 36, 1251-1275.
137. Gillis M, Van TV, Bardin R, Goor M, Hebbar P, Willems A, Segers P, Kersters K, Heulin T and Fernandez MP. Polyphasic taxonomy in the genus *Burkholderia* leading to an emended description of the genus and proposition of *Burkholderia vietnamiensis* sp. nov. for N₂-fixing isolates from rice in Vietnam. *Int. J. Syst. Bacteriol.* 1995; 45: 274-289.