

An improved protocol for extraction of metagenomic DNA from high humus, alkaline and saline soil of chinampa for T-RFLP fingerprinting analysis

ABSTRACT

Aims: The principle aim of this study was to obtain high quality metagenomic DNA from the high humus-containing, alkaline soils of the chinampas, an artificial sustainable agro-ecosystem.

Study design: Different protocols reported previously were tested and were modified to extract the metagenomic DNA. Quality of the DNA samples was evaluated by amplification of 16S rRNA gene with PCR and T-RFLP (Terminal Restriction Fragment Length Polymorphism) analysis.

Place and Duration of Study: This study was performed in Department of Microbiology, Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional during 2011-2012.

Methodology: Four soil samples were collected from two chinampas at the depth of 0-30 cm and 30-60 cm. A protocol started with repeated prewashing before the direct cell lysis with lysozyme followed by SDS treatments, frozen and melting cycling was developed which combined the DNA isolation and purification procedures. The 16S rRNA genes were amplified from the extracted metagenomic DNAs and were used for T-RFLP fingerprinting analysis.

Results: The 16S rRNA genes were amplified from all the DNA extracts corresponding to the four soil samples and were successfully used in the T-RFLP analysis, which generated 25 to 109 T-RFs in the four soil samples digested separately with the restriction endonucleases *HaeIII*, *HhaI* and *MspI*.

Conclusion: The protocol developed in the present study could generate high molecular weight and high quality metagenomic DNA from soils with high content of humic materials, for which the other reported protocols were not functioned. This soil harboured very diverse and unique bacterial communities belonging to at least nine phyla that might contribute to the high soil fertility.

Keywords: Metagenomics, DNA extraction, humic acids, T-RFLP fingerprinting, bacterial communities.

1. INTRODUCTION

The diversity and functions of microorganisms in different environments are two principle aspects in the study of microbial ecology. In the last decades, the application of genomic tools have dramatically improved microbial ecological studies and drastically expanded our knowledge about microbial world [1]. It has been estimated that microorganisms constitute two third of the Earth's biological diversity, and about 86% of existing species on Earth and 91% of species in the ocean still remain unrecognized [2]. Metagenomics is the genomic analysis of microorganisms based upon the DNAs extracted directly from their natural environment, through which the unexplored microbial diversity can be captured [3] and specific functional genes can be screened [4].

In the study of metagenomics, the first and basic step is the extraction of pure metagenomic DNA and/or RNA from the environmental samples, which are subsequently used for PCR analysis. For this purpose, distinct protocols of metagenomic DNA extraction have been developed referring to different environment samples [5-11], or in order to detect some special microorganisms, such as *Mycobacterium avium* [12] or fungi [13]. Some of these methods have been

compared experimentally [14] or reviewed [15-17]. Although many methods about the extraction of metagenomic DNA have been described, they cannot fit all the environmental samples, because the characteristics of this material varied dramatically, and different organisms have distinctive cellular structures and chemical compounds. Therefore, some novel or modified methods specific for different environmental samples are still emerging [18, 19], and optimized methods are screened for some specific soils [13, 20].

The chinampas refer to the artificial floating islands started about 3000 years ago in the Valley of Mexico, including Xochimilco in Mexico City, in which trunks and twigs of trees were used to construct the basis to support the soil composing of vegetable wastes and sediment of the lake. They are still used as a sustainable agro-system to produce vegetables and ornamental flowers without or with little apply of chemical fertilizers and pesticides. Since 1989 Xochimilco is a site included into World Heritage List of UNESCO. In relation to its chemical composition and floating environment, the mature soil of chinampa is rich in humic material, high pH and high salinity. Many studies have been done to evaluate the physicochemical characters of the chinampa soil [21], but little microbiological information is available about this soil [22], although the microbiota is an important factor to maintain the productivity of soil.

Aiming at investigating the bacterial diversity in soil of chinampa with metagenomic study, we realized the extraction of DNA from this soil with various methods and an improved protocol was established.

2. MATERIAL AND METHODS

2.1. Soil sampling

Bulk soils were sampled in 2012 from two chinampas far away from the touristic zone in Xochimilco, Mexico City. In each chinampa, the surface (0-30 cm depth) and subsurface (30-60 cm) were sampled. The soils samples were recollected from five points (four corners and the centre), stored in black plastic bags, and transported directly to the laboratory, where the five soil samples from each chinampa at each depth were pooled to form a compiled sample. The sampled soils showed a black colour, indicating the high content of organic matter, mainly humus with high melanisation and high molecular weight [21]. All soil samples were kept at 4°C until their utilization for soil physicochemical characterization by the routine methods [22] and for DNA extraction.

2.2 Quantification of culturable mesophilic aerobic bacteria

For each soil sample, decimal dilutions were prepared in sterilized NaCl solution (0.85%). An aliquot of 0.1 mL of the dilutions 10^{-4} , 10^{-5} and 10^{-6} was spread in duplicate in Petri dishes containing PY medium (peptone, 5 g; yeast extract, 3 g; CaCl_2 , 0.6 g; agar, 18 g; distilled water, 1 L; pH8.0). The plates were incubated at 28°C during 48 h. Colonies were counted manually and abundance of mesophilic aerobic bacteria was calculated as colony forming units (CFU) in each gram of dry soil.

2.3. Extraction of metagenomic DNA from the soil samples

In this study, several protocols [7, 9, 11, 23, 24] were employed previously, and the protocol of Ceja-Navarro et al. [23] modified by adding prewashing and substituting the flocculant agent was finally used. In starting, 5.0 g of soil were suspended with 25 ml of pyrophosphate sodium ($\text{Na}_4\text{P}_2\text{O}_7$, 0.15 M) in a 50 ml Oak Ridge tube (Nalgene™, Sigma-Aldrich®) to dissolve the humic material. The tube was vortexed for 1 min, following by 10 min of precipitation and then centrifuged 10 min at $7,700 \times g$ under room temperature. The supernatant was discarded and the washing procedure was repeated 5 times. The sediment was resuspended in 5 ml of phosphate buffer (0.15 M NaH_2PO_4 ; pH8.0). After 10 min of precipitation, the suspension was vortexed for 1 min and centrifuged 10 min at $7,700 \times g$ under 25 °C, then the supernatant was discarded and this washing procedure was repeated 4 times.

The washed sediment was suspended in 5 ml of lysis solution I (NaCl 0.15 M; EDTA 0.1M; pH =8.0; 10 mg ml^{-1} of lysozyme), vortexed and incubated at 37°C for 1 h, with briefly vortexing each 20 min. Then, 5 ml of the lysis solution II (NaCl 0.1 M; Tris-HCl 0.5 M; 12% SDS; pH 8.0) were added and the mixture was maintained at -20 °C for 20 min, and at 65°C for another 20 min. This frozen-melting cycle was repeated two times and in the first cycle, 5 ml of $\text{Al}_2(\text{SO}_4)_3$ (0.3 M) as flocculant were added to eliminate the humic material. The mixture was vortexed and the frozen-melting cycle was repeated as mentioned above. The mixture was centrifuged at $7,700 \times g$ during 10 min. The supernatant was transferred into a new sterilized Oak Ridge tube and was mixed with 2.7 ml of NaCl (5 M) and 2.1 ml of Triton X-100 (10% in 0.7 M NaCl solution). After a incubation at 65°C for 10 min, 12 ml of a chloroform:isoamyl alcohol (24:1, v/v) solution were added and the tube was mixed gently by inversion and centrifuged at $3000 \times g$ for 30 min. The aqueous phase was

transferred into a clean tube and mixed with 12 ml of Polyethyleneglycol (PEG 13%; NaCl 1.6 M) and maintained on ice overnight. The precipitated metagenomic DNA was collected by centrifugation at $12,000 \times g$ for 30 min at 4°C . The DNA pellet was resuspended in 500 μL of deionized sterile water, transferred into a sterilized Eppendorf tube, and mixed with 1 volume of absolute ethanol. After incubated at 4°C for 30 min, the tube was centrifuged at $12,000 \times g$ for 30 min at 4°C . The supernatant was discarded and the precipitated DNA was washed with 500 μL of cold ethanol (70%, v/v). The DNA pellet was dried by incubating the tube at 65°C for 10 min. Then it was dissolved in 500 μL of deionized sterile water and was stored at -20°C .

2.4. Evaluation of DNA concentration and quality

The integrity and concentration of DNA sample was estimated by electrophoresis in 1% (w/v) agarose gel with $0.5 \times \text{TBE}$ as electrode buffer as described elsewhere [25]. The DNA sample (1 μL) was loaded in the gel, and the 1 Kb ladder (Invitrogen™) was included to estimate the molecular size and concentration of extracted DNA. After the electrophoresis and stained by ethidium bromide ($0.5 \mu\text{g ml}^{-1}$), the gel was observed under UV light and was photographed with MultiImage™ Light Cabinet (Alpha Innotech Corporation). DNA concentration was measured by UV- spectrophotometry: 1 μL of DNA solution was diluted to 50 μL with sterilized deionized and the concentration of DNA was read at 260 nm directly from the spectrophotometer (Epoch-Biotek, with software Gen5 take 3).

2.5. T-RFLP analysis

For T-RFLP (Terminal Restriction Fragment Length Polymorphism) analysis, the fragments of 16S rRNA gene were amplified from the metagenomic DNA extracts by PCR as described by Sun et al. [26], with the primers 27f (5'-AGA GTT TGA TCM TGG CTC AG-3') marked with fluorescently carboxyfluorescein (6-FAM) and 1495r (5'-CTA CGG CTA CCT TGT TAC GA-3'). Then, 10 μL of each PCR product were digested with 10 U of the restriction endonucleases *MspI*, *HhaI* and *HaeIII*, respectively, at 37°C for 3 h. The digests were purified and electrophorized as reported previously [26]. The electrophoretic patterns were analyzed with the GeneMarker software (SoftGenetics®) and only the terminal restriction fragments (T-RF) between 50 and 500 pb were chosen for further analysis. The Shannon-Weaver (H') and Simpson ($1-D$) indices were calculated by using PAST 2.17c software [27] for estimation of the bacterial diversity in each soil. The T-RF patterns were used to identify the bacteria by using the data in National Center for Biotechnology Information Database (<http://trflp.limnology.wisc.edu/index.jsp>) [28]. Only the unambiguous fragments were designed into the bacterial taxa (phylum and/or class) [29].

3. RESULTS AND DISCUSSION

3.1. Soil characterization and quantification of mesophilic aerobic bacteria

The physicochemical characteristics of the chinampa soils are presented in Table 1. In general, the soils are sandy clay loam, with alkaline pH ranged between 8.0 and 8.8, with high salinity ($2054-4468 \text{ mg K kg}^{-1}$), and rich in organic materials (4.6-7.5%). The abundance of mesophilic aerobic bacteria was about 10^7 CFU g^{-1} of dry soil, implying that the major form of organic matter in the chinampa soils was humus, which is difficult for degradation and does not support the growth of bacteria.

3.2. Extraction of metagenomic DNA from soil

In our study, the DNA extracted from the chinampa soils with commonly used protocols either showed brown colour, or did not produce PCR fragments, even after the post purification procedures. The post purification procedures were 1) using of the silica columns (QUIAGEN® and MO BIO®) as recommended by the manufacturer; 2) purifying the DNA band recovered from the agarose gel (0.7%) after electrophoresis with QIAamp DNA Mini Kit (QUIAGEN) following the manufacturer's guide; and 3) flocculation of 500 μL DNA solution with 250 μL of $\text{Al}_2(\text{SO}_4)_3$ (10 mM) and 50 μL of phosphate buffer (0.15M NaH_2PO_4 ; pH8.0). These procedures could eliminate the brown colour, but the purified DNA samples were still not good for amplification by PCR. With the modified protocol of Ceja-Navarro et al. [23], the humic acids were eliminated gradually from the soil samples, because the brown colour in the supernatants was decreased time by time, until disappeared completely. The DNA extracts had high molecular size ($>12 \text{ kb}$) without degradation (Fig. 1), and the yield was 18 to 34 ng g^{-1} soil. The obtained DNA samples were good for gene amplification by PCR using BSA (Bovine Serum Albumine) as PCR additive.

Table 1. Characteristics of soil samples collected from two chinampa soils.

Soil feature	Soil samples *			
	Ch Is	Ch Id	Ch IIs	Ch IId
Organic matter (%)	7.5	4.6	7.5	6.4
Total N (%)	0.80	0.56	0.75	0.58
Total P (mg kg ⁻¹)	22.4	17.4	3.7	19.6
Total K (mg kg ⁻¹)	4468	3539	3728	2054
pH	8.6	8.2	8.7	8.0
Water Retention Capacity (%)	120.2	116.7	148.0	112.1
Humidity (%)	24	44	38	48
Humidity/ WRC (%)	19.97	37.70	25.68	42.82
CFU g ⁻¹ dry soil	5.07×10 ⁷	3.45×10 ⁷	3.45×10 ⁷	4.04×10 ⁷

* Ch Is, Ch Id, Ch IIs, and Ch IId represent soil sampled from chinampa I (0-30 cm), chinampa I (30-60 cm), chinampa II (0-30 cm), and chinampa II (30-60 cm), respectively.




Figure. 1. Metagenomic DNA extracted with the improved method developed in this study: 1, 1 Kb DNA Ladder; 2, DNA from Ch IIs; 3, DNA from Ch IId.

3.3. Analysis of T-RFLP

T-RFLP is a technique widely used for investigation of microbial diversity in environmental samples [30-32]. With the metagenomic DNAs obtained in this study, 25 to 109 T-RFs were obtained from the 16S rRNA amplicons digested by the three restriction endonucleases (Table 2). The Shannon and Simpson indices were very similar in the four samples (Table 2): both the Shannon indices greater than 3 [29] and the Simpson indices near to one [32] indicated that the bacterial communities in the chinampa soils were very diverse and composed of many species.

Table 2. T-RFs and diversity estimated from the T-RFLP analysis in the chinampa soil samples

Soil sample	<i>Hha</i> I			<i>Msp</i> I			<i>Hae</i> III			Average		
	T-RFs	H'	1-D*	T-RFs	H'	1-D	T-RFs	H'	1-D	T-RFs	H'	1-D
Ch Is	89	4.48	0.98	102	4.62	0.99	79	4.36	0.98	90.0	4.49	0.98
Ch Id	80	4.38	0.98	109	4.69	0.99	63	4.14	0.98	84.0	4.40	0.98
Ch IIs	97	4.57	0.98	53	3.97	0.98	70	4.24	0.98	73.3	4.26	0.98
Ch IId	61	4.11	0.98	81	4.39	0.98	25	3.21	0.96	55.7	3.90	0.97
Average	81.8	4.38	0.98	86.2	4.67	0.98	59.2	3.99	0.98	75.7	4.26	0.98

*. H', Shannon Index; 1-D, Simpson Diversity Index.

Comparing with the database, the T-RFs obtained in this study were defined into nine Phyla (Fig. 2). *Firmicutes* was the most abundant one in all the four samples, with 18.9% to 24.19%. The second abundant group was Phylum *Proteobacteria*, followed by Phylum *Actinobacteria*. The minor groups were Phyla *Chloroflexi*, *Cyanobacteria* and *Planctomycetes* etc. About 25% of the T-RFs could not be defined into any phylum, which might imply the existence of novel taxa or the limit of information in the database and of the T-RFLP technique.




Figure 2. Community composition of bacteria and relative abundance of different bacteria groups estimated from the T-RFLP analysis in the chinampa soils. Ch Is, Ch Id, Ch IIs, and Ch IId represent soil sampled from chinampa I (0-30 cm), chinampa I (30-60 cm), chinampa II (0-30 cm), and chinampa II (30-60 cm), respectively.

Compared with the previous results obtained from other soils with some similar characteristics, the community composition of bacteria in the chinampa soils was unique. *Proteobacteria*, *Firmicutes*, *Actinobacteria* and *Bacteroidetes* were the major bacterial groups in alkaline (pH 8.28-8.45) crop soils in China [26]. *Proteobacteria*, *Verrucomicrobia* and *Chloroflexi* were most abundant phyla in the soils of constructed wetland (pH 6.23-6.59; organic material 13.54-29.75%) in Spain [33]. While *Proteobacteria*, *Acidobacteria*, *Actinobacteria*, *Bacteroidetes* and *Firmicutes* made up more than 80% of the bacteria sequences in created wetland (pH4.2-5.8, organic material 3-6%) in Virginia, USA [34]. In addition, *Proteobacteria*, *Acidobacteria* and *Actinobacteria* were the most abundant groups in an alkaline saline soil (pH9.0, OM 1.29%) of Mexico [35]. Based upon all the previous studies mention here and the results obtained in the present study, it was clear that the *Proteobacteria* group always exist as one of the predominant group, and the other dominant groups, as well as the relative abundance of *Proteobacteria*, varied according to the soil characters, like pH, humidity, saline and maybe also the content and type of organic matter.

4. CONCLUSION

The protocol modified in the present study could generate high molecular weight and high quality metagenomic DNA from soils with high content of humic materials, for which the other reported protocols were not functioned. The key points were the prewashing of the soil to eliminate contamination of humic acids and $Al_2(SO_4)_3$ as flocculant agent. This protocol is recommendable for soils with similar characters. The DNA samples extracted with this protocol were adequate for PCR amplification and T-RFLP analysis, which revealed that the chinampa soils harboured very diverse and unique bacterial communities belonging to at least nine phyla. Further studies on the diversity are undergoing by other molecular methods.

COMPETING INTERESTS

The authors declared that we have no conflict of interest in the present publication.

ACKNOWLEDGEMENTS

This study was financially supported by the project PICS08-3 authorized by the ICyT D. F. and by the projects SIP 20120760 and SIP 20130828 authorized by IPN.

REFERENCES

1. Xu J. Microbial ecology in the age of genomics and metagenomics: concepts, tools, and recent advances. *Mol. Ecol.* 2006; 15: 1713–1731.
2. Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B. How many species are there on earth and in the ocean? *PLoS Biol.* 2011; 9(8), e1001127. doi:10.1371/journal.pbio.1001127.
3. Petrosino JF, Highlander S, Luna RA, Gibbs RA, Versalovic J. Metagenomic pyrosequencing and microbial identification. *Clin. Chem.* 2009; 55: 856–866.
4. Mitra S, Rupek P, Richter DC, Urich T, Gilbert JA, Meyer F. et al. Functional analysis of metagenomes and metatranscriptomes using SEED and KEGG. *BMC Bioinformatics* 2011; 12 Suppl 1, S 21.
5. Barzegari A, Vahed SZ, Atashpaz S, Khani S, Omid Y. Rapid and simple methodology for isolation of high quality genomic DNA from coniferous tissues (*Taxus baccata*). *Mol. Biol. Rep.* 2010; 37: 833–837.
6. Desai C, Madamwar D. Extraction of inhibitor-free metagenomic DNA from polluted sediments, compatible with molecular diversity analysis using adsorption and ion exchange treatments. *Biores. Technol.* 2007; 98: 761–768.
7. Dong D, Yan A, Liu H, Zhang X, Xu Y. Removal of humic substances from soil DNA using aluminum sulfate. *J. Microbiol. Methods* 2006; 66: 217–222.
8. Griffiths RI, Whiteley AS, O'Donnell AG, Bailey MJ. Rapid method for coextraction of DNA and RNA from natural environments for analysis of ribosomal DNA- and rRNA-based microbial community composition. *Appl. Environ. Microbiol.* 2000; 66: 5488–5491.
9. Knaebel DB, Crawford RL. Extraction and purification of microbial DNA from petroleum-contaminated soils and detection of low numbers of toluene, octane and pesticide degraders by multiplex polymerase chain reaction and Southern analysis. *Mol. Ecol.* 1995; 4: 579–591.
10. Sebastianelli A, Sen T, Bruce IJ. Extraction of DNA from soil using nanoparticles by magnetic bioseparation. *Lett. Appl. Microbiol.* 2008; 6: 488–491.
11. Zhou J, Bruns MA, Tiedje JM. DNA recovery from soils of diverse composition. *Appl. Environ. Microbiol.* 1996; 62: 316–322.
12. Cook KL, Britt JS. Optimization of methods for detecting *Mycobacterium avium* subsp. *paratuberculosis* in environmental samples using quantitative, real-time PCR. *J. Microbiol. Methods* 2007; 69: 154–160.
13. Fredricks DN, Smith C, Meier A. Comparison of six DNA extraction methods for recovery of fungal DNA as assessed by quantitative PCR. *J. Clin. Microbiol.* 2005; 43: 5122–5128.
14. Rajendhran J, Gunasekaran P. Strategies for accessing soil metagenome for desired applications. *Biotechnol. Adv.* 2008; 26: 576–590.
15. Lloyd-Jones G, Hunter DWF. Comparison of rapid DNA extraction methods applied to contrasting New Zealand soils. *Soil Biol. Biochem.* 2001; 33: 2053–2059.
16. Robe P, Nalin R, Capellano C, Vogel TM, Simonet P. Extraction of DNA from soil. *Eur. J. Soil. Biol.* 2003; 39: 183–190.
17. Schneegurt MA, Dore SY, Kulpa CF Jr. Direct extraction of DNA from soils for studies in microbial ecology. *Curr. Issues Mol. Biol.* 2003; 5: 1–8.
18. Lee S, Hallam SJ. Extraction of high molecular weight genomic DNA from soils and sediments. *J. Vis. Exp.* 2009; (33). pii: 1569. doi: 10.3791/1569.
19. Wechter P, Williamson J, Robertson A, Kluepfel D. A rapid, cost-effective procedure for the extraction of microbial DNA from soil. *World J. Microbiol. Biotech.* 2003; 19: 85–91.
20. Verma D, Satyanarayana T. An improved protocol for DNA extraction from alkaline soil and sediment samples for constructing metagenomic libraries. *Appl. Biochem. Biotechnol.* 2011; 165: 454–464.

21. Ramos-Bello R, Cajuste JL, Flores-Romá D, García NE. Metales pesados, sales y sodio en suelos de Chinampa en México. *Agrociencia* 2001; 35: 385–395.
22. Trujillo-Cabrera Y, Ponce-Mendoza A, Vásquez-Murrieta MS, Rivera-Orduña FN, Wang ET. Diverse cellulolytic bacteria isolated from the high humus, alkaline-saline chinampa soils. *Ann. Microbiol.* 2013; 63: 779–792.
23. Ceja-Navarro J, Rivera FN, Patiño-Zúñiga L, Govaerts B, Marsch R, Vila-Sanjurjo A, et al. Molecular characterization of soil bacterial communities in contrasting zero tillage systems. *Plant Soil* 2010; 329: 127–137.
24. Cullen DW, Hirsch PR. Simple and rapid method for direct extraction of microbial DNA from soil for PCR. *Soil Biol. Biochem.* 1998; 30: 983–993.
25. Green MR, Sambrook J. *Molecular Cloning: A Laboratory Manual (Fourth Edition)*. 2012; pp. 66-68. Cold Spring Harbor Laboratory Press. Cold Spring Harbor, New York.
26. Sun YM, Zhang NN, Wang ET, Yuan HL, Yang JS, Chen WX. Influence of intercropping and intercropping plus rhizobial inoculation on microbial activity and community composition in rhizosphere of alfalfa (*Medicago sativa* L.) and Siberian wild rye (*Elymus sibiricus* L.). *FEMS Microbiol. Ecol.* 2009; 70: 62–70.
27. Hammer O, Harper DAT, Ryan PD. PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Paleontologia Electronica* 2001; 4: 1-9.
28. Gil F, Iglesia RD, Mendoza L, González B, Wilkens M. Soil bacteria are differentially affected by resin of the medicinal plant *Pseudognaphalium vira* and its main component kaurenoic acid. *Microb. Ecol.* 2006; 52: 10–18.
29. De la Iglesia R, Castro D, Ginocchio R, van der Leslie D, Gonzalez B. Factors influencing the composition of bacterial communities found at abandoned copper-tailings dumps. *J. Appl. Microbiol.* 2006; 100: 537–544.
30. Liu WT, Marsh TL, Cheng H, Forney LJ. Characterization of microbial diversity by determining terminal restriction fragment length polymorphisms of genes encoding 16S rRNA. *Appl. Environ. Microbiol.* 1997; 63: 4516–4522.
31. Thies JE. Soil microbial community analysis using terminal restriction fragment length polymorphisms. *Soil Sci. Am. J.* 2007; 71: 579–591.
32. Orcutt B, Bailey B, Staudigel H, Tebo BM, Edwards KJ. An interlaboratory comparison of 16S rRNA gene-based terminal restriction fragment length polymorphism and sequencing methods for assessing microbial diversity of seafloor basalts. *Environ. Microbiol.* 2009; 11: 1728–1735.
33. Ansola G, Arroyo P, Sáenz de Miera LE. Characterization of the soil bacterial community structure and composition of natural and constructed wetlands. *Sci. Total Environ.* 2014; 473-474: 63-71.
34. Peralta RM, Ahn C, Gillet PM. Characterization of soil bacterial community structure and physicochemical properties in created and natural wetlands. *Sci. Total Environ.* 2013; 443:725-732.
35. Castro-Silva C, Ruíz-Valdiviezo VM, Valenzuela-Encinas C, Alcántara-Hernández RJ, Navarro-Noya YE, Vázquez-Núñez E, Luna-Guido M, Marsch R, Dendooven L. The bacterial community structure in an alkaline saline soil spiked with anthracene. *Electr. J. Biotechnol.* 2013; DOI: 10.2225/vol16-issue5-fulltext-14.