

Recent advances in virus-host coevolution and protective mechanisms against plant viruses

ABSTRACT

In plant pathology, the study of the interaction between the plant host and the viral pathogen has been a very active area of research in the last few decades. The infection process of a plant pathogen usually begins with the exchange of molecular signals. With particular emphasis on plant virus evolution, and focusing on quantitative and population genetics, plant virus-host interactions and coevolution allow understanding of the major factors favoring disease emergence. The exploitation of viral interaction phenomena will improve established genetic engineering strategies for viral cross-protection in plants. Also, the study of plant immunity against viral infection, both innate and adaptive (e.g. RNA silencing), has helped in the development of resistant varieties to several plant viruses based on genetic engineering. This review summarizes the recent advances in plant-virus interactions and co-evolution as well as current developments in resistant crop investigations.

Keywords: *Plant-virus interactions, virus evolution, plant immunity, antiviral resistance, cross-protection, genetic engineering*

1. INTRODUCTION

Plant infection by viruses causes physiological disorders responsible for plant diseases. This causes economic and agronomic significant losses in many crops. In 2012, the International Committee on Taxonomy of Viruses reported 92 genera of plant viruses, of which 82 were assigned to 21 different families [1]. In the last few decades, the molecular dialogue between plant hosts and their viral parasites was highlighted as a major breakthrough, providing new strategies that are directly exploitable in crop development programs. Nevertheless, the spread of crop viruses has increased dramatically in recent years. As long as there is no chemical capable of providing curative control against plant viruses, only preventive control can be considered.

Using recent knowledge in genomics and bioinformatics tools, researchers have been able to study in depth the genome of many plant viruses, the relationships between viruses and their plant hosts, and the mechanisms of virus evolution and virus-host co-evolution. Plant viruses are a convenient model system for studying viral evolution. Despite increasing efforts devoted to modeling the molecular evolution of many viral populations, such as Potyvirus [2,3,4], Carlavirus [5], Begomovirus [6,7] and Potexvirus [8] (Hasiów-Jaroszewska et al. 2011), there have been few attempts to study adaptive evolution, which could bring important information about the nature of the interaction between the host and the virus. One of the reasons for considering plant viruses as an appropriate model is that they allow for the use of infectious *in vitro*-generated viral RNAs as an inoculum for host organisms and permit controlled studies in multiple, genetically similar, intact hosts [9,10].

Another hallmark of virus evolution is the findings of viral sequences integrated into the plant genome. Some endogenous viruses could be benign components of plant genomes; others could potentially be pathogenic or, on the contrary, contribute to viral immunity [11]. Moreover, RNA viruses replicate with extremely high mutation rates due to the low fidelity of RNA-dependent RNA polymerases and reverse transcriptases. Mutation, recombination, short replicative cycles and high copy yields contribute to the generation of complex distributions of closely related mutant genomes able to adapt to fluctuating environments; these are termed quasispecies. Also, the host environment may be associated with the diversity observed in quasispecies. One biological implication of viral quasispecies, of great sanitary impact, consists in the emergence of resistant mutants after antiviral treatment. Plant virus systems are used as *in vivo* experimental model for understanding the evolution of viruses subjected to chemical mutagenesis. One of the most widely studied is *tobacco mosaic virus* (TMV) [12].

Additionally, the number of viral variants present in a single host cell, and even more the multiplicity of infection (MOI) are rarely experimentally studied in plant viruses. Yet, it is a fascinating and promising generic question in terms of the biology and evolution of viruses, since the MOI is responsible for important processes such as intracellular competition, functional complementation and recombination. Indeed, there may be a variety of control mechanisms for the entry of these infectious units in the cell, limiting them to a small number, or even exclusion mechanisms that do not allow the co-existence of two infectious units in the same cell [13].

During the process of infection, plants employ various strategies to combat viral infections. Understanding how plants defend themselves from pathogens is essential in order to protect crops and develop highly disease-resistant plant species. While cross-protection has been intensively studied for decades since the discovery of this phenomenon by McKinney in the 1920s [14], less is known about antiviral mechanisms based on genetic tools such as PAMP-triggered immunity (PTI), effector-triggered immunity (ETI) [15], or gene silencing and how they could be the source of novel plant resistance mechanisms in the field.

Most economically important crops are susceptible to viruses. Plant viruses manifest a wide variety of pathogenicity that requires individual management strategies. The use of a resistant cultivar, against either two or more types of pathogen species, called broad spectrum resistance (BSR), remains the best and cheapest control strategy in managing viral diseases. In addition, several in-depth reviews and books on the multiplication of RNA plant viruses, viral quasispecies, genetic diversity, and plant-virus interactions based on proteomic analysis have been published elsewhere [16,17,18].

With these considerations in mind, this review will focus on giving a summary of recent advances plant-virus interactions and co-evolution, as well as to indicate current developments in important strategies of combining modern molecular and physiological techniques with phytopathology for future investigations in resistant crop varieties.

2. PLANT VIRUS INFECTION

Plant viruses are obligate parasites that enter the plant cell passively through mechanical wounds caused by environmental factors or by vectors, due to lack of plant virus receptors. The next phase of virus infection proceeds in the cytoplasm after capsid disruption [19].

During the infection process, plant viruses interact with various host factors essential for their accumulation, translation, replication, cell-to-cell and long-distance movement and utilize plant proteins, normally involved in host-specific activities, for their own purposes. Considerable progress has been made in recent years toward understanding some of those virus-host interactions [20,21,22].

Most of plant viruses have an RNA genome that consists of single stranded RNA (ssRNA). A part of these genomes are positive-sense polarity in the same sense of orientation of the messenger RNAs of the cell, the other part uses ssRNAs of negative polarity. The accumulation of progeny of both DNA and RNA plant viruses involves translation and replication of viral sequences. The ability of a virus to interact with a host depends on the formation of a functional heterocomplexes between host and virus proteins [23].

Viral RNAs share characteristics with host cell mRNAs, but display a variety of structures at their 5' and 3' ends that differentiate them from cellular mRNAs. The interaction of host and viral factors at both ends 3' and 5' and sometimes along the viral RNA is mediated by RNA-RNA interactions or through the recruitment of host translation initiation factors especially the eukaryotic translation initiation factor 4G (eIF4G) [24].

In addition, virus-host interactions during DNA virus accumulation have been studied. An NAC domain protein of tomato (*Solanum lycopersicum*) interacting with a viral replication enhancer (RE) protein of Geminivirus was identified and proposed to participate in viral replication [25].

Moreover, an interaction between NAC protein of *Arabidopsis thaliana* and an RNA virus coat protein is necessary during a host resistance response [26]. This host proteins present different functions during DNA and RNA virus infection, thus making the virus-host interaction process more complex.

Cell-to-cell transport occurs through plasmodesmata (PD) connecting adjacent cells in plant tissues and involves virus proteins named movement proteins (MPs). Viruses reveal different movement strategies with specific viral and host components interacting in efficient, functional transport events [27]. A 30 kDa protein of *Tobacco mosaic virus* (TMV) was the first specific identified viral movement protein. Functions have been clarified lately and assigned to the 30 kDa-like MPs [28], including targeting of viral RNA to plasmodesmata and increasing PD

pore size (SEL, size exclusion limit) [29] to allow trafficking of ribonucleoprotein complex (RNP) formed between the TMV MP and the infectious RNA [30,31]. The PD-targeted MP transport and associated viral RNA occurs by diffusion in the endoplasmatic reticulum (ER) membranes [32].

Unlike Tobamovirus, members of Potexvirus need capsid protein for cell-to-cell transport [33]. Proteins coded by triple gene block (TGB) are also required for cell-to-cell transport of many plant virus genera [34,35].

Recently, a possible relation between virus accumulation and cell-to-cell movement was identified. Host eukaryotic translation factors eIF4E and eIF(iso)4E, which are involved in potyviruses accumulation and replication, were also shown to participate in virus cell-to-cell movement [36].

3. GENETIC EVOLUTION OF PLANT VIRUSES

3.1 Multiplicity of infection

The MOI is defined as the average number of genomes that infect and penetrate a cell. MOI is a term commonly used in cell culture, as a theoretical value, which determines the number of infectious units available in the environment per cell present in the same medium. It does not necessarily correspond to the real number of infectious units penetrating and multiplying in each cell. The number of infectious units that effectively infects each cell can be considered as the true MOI; this is often not considered, nor compared with the theoretical MOI in cell culture. The parameter of interest is precisely the real value of the MOI under conditions of “natural” infection in a multicellular host [13].

For plant viruses, it is a key parameter in the evolution of viruses, as it governs essential processes such as genetic exchange between genomes, the intensity of selection among viral genes, epistatic interactions, and the evolutionary aspect of multipartite viruses. This trait has been studied and estimated for TMV in its systemic host *Nicotiana benthamiana* [37] and *cauliflower mosaic virus* (CaMV) [13].

In a recent study, MOI levels were estimated based on the frequency of cells infected with two different TMV genotypes, discriminated using green or red fluorescent proteins (GFP and RFP). The MOI was high, but it changed during the progression of infection, reducing from an initial level of about six to a final one of one to two, with most infection cycles occurring at higher MOI levels. The decreasing MOI can be explained by mechanisms limiting superinfection and/or by genotype competition within co-infected cells, which was shown to occur in double infected tobacco protoplasts [37].

A reasonable hypothesis explaining the presence or absence of bottlenecks would be the regulation of the MOI of cells by several genomes of the viral population. Indeed, it has been shown that the viral genomes of *plum pox virus* (PPV), responsible for Sharka disease, cannot secondarily invade *Prunus* tissues that have already been infected by a closely related genome. This generates bottlenecks, thus preventing extensive mixing of genetic variants within a host [38].

3.2 Viral quasispecies

In general, RNA virus replication is characterized by high mutation rates, short replication times and high yields [39]. Also, several RNA viruses have genetically variable but closely related populations, called viral quasispecies, which act as a unit of selection and interact with the host species [12].

The quasispecies cloud size (quasispecies diversity level) may depend on adaptability and host range. In a prior study, Schneider *et al.* were limited to a single host, *Nicotiana benthamiana*, for direct comparison of quasispecies cloud size using three alpha-like RNA viruses, TMV, *cucumber mosaic virus* (CMV), and *cowpea chlorotic mottle virus* (CCMV). Yet, they indicated that evolutionarily related viruses have very different levels of diversity that correlated with their relative host range sizes. Furthermore, the quasispecies cloud size remained constant through 10 consecutives passages, i.e., all individuals had a similar average genetic distance from the consensus sequence [40].

Moreover, in a following study done by the authors cited above, it was shown that quasispecies cloud sizes are not constant; rather, they vary with different plant hosts and in changing environments. First, these differences may be due to selection pressure or bottlenecks imposed by the host-virus infection cycle. Second, this work proposed the possibility that different hosts may accelerate or decelerate the rate of viral evolution by allowing or denying high levels of diversity in viral populations [10].

Quasispecies diversity in viruses has been described previously as a mechanism to escape host resistance responses [41,42] or as a reservoir to maintain variants with selective advantages in other environments [43]. This diversity has been correlated with the ability to infect numerous hosts [40] Nevertheless, excessive diversity can generate complications if the virus is exposed to recurrent bottlenecks.

3.3 Viral bottlenecks

Most mutations are deleterious. Frequent bottlenecks can lead to the rapid loss of fitness known as Muller's ratchet [39,44]. A study has shown that the population diversity of marked CMV mutants decreased significantly and stochastically when the population moved from inoculated tobacco leaves to primary systemic leaves and reduced further as the systemic infection progressed, providing clear evidence of a genetic bottleneck [45]. Furthermore, the severity of bottlenecks varies and may depend on the structure of the minor veins and plasmodesmata of individual hosts [46].

Other studies have shown that in multicomponent RNA viruses, the function can be responsible for multiple differences in the genetic structures of different genomic segments. Populations of *tomato chlorosis virus* (ToCV) have a heterogeneous and complex genetic structure that depends on the RNA segment considered; i.e., it is more complex for RNA1 (encoding replication-associated proteins) than for RNA2 (encoding encapsidation, movement, and insect transmission) [47].

3.5 Genetic adaptation

The rate of genetic adaptation of organisms depends on the degree of mutation, the population size and the range of selection, mainly during population bottlenecks. A single nucleotide substitution may facilitate virus adaptation to the host's translation machinery due to a correlation between viral amino acid codons and those used by the host plant [48]. A study on *turnip mosaic virus* mutations genomes revealed that several nucleotide sites may be involved in adaptation to *Raphanus sativus* [49].

For pathogens that affect multiple hosts, adaptation is a key factor in determining the probability and the severity of emergent disease outbreaks and can be used as a tool for the preservation of genetic diversity both in host and pathogen species. *Tobacco etch potyvirus* (TEV) and four natural hosts were used in an experimental evolution study that revealed the strong adaptive potential of TEV to new hosts without severe evolutionary limitations. The analysis of genome consensus sequences of the evolved lineages established that all mutations shared between lineages were host-specific [50].

Recent research has provided evidence of adaptation by plant viruses to host plants. Such adaptation may alter host-plant phenotypes in ways that facilitate transmission by arthropod vectors [51].

There have been an important interest toward studying endogenous viral elements, which are knowledgeable in animals and much less well in plants. This aroused the interest of researchers who have carried out several independent studies on endogenous viral sequences in diverse plant species [11]. The sequences which have been found inserted into plant chromosomes were derived from both groups of plant DNA viruses, i.e. the single-stranded DNA geminiviruses [52] and pararetroviruses with double-stranded DNA [53,54].

Recently, researchers at National Institute of Agronomic Research (INRA) Versailles-Grignon (France) and French Agricultural Research Centre for International Development (CIRAD) described a new kind of virus belonging to caulimovirus family (*Caulimoviridae*) called "Florendovirus" whose members have colonized the genomes of a large diversity of flowering plants during evolution. The genome of the grapevine (*Vitis vinifera*) was first identified and revealed previously unknown viral sequence elements. The assembly of these endogenous sequences allowed for reconstituting the viral genome, whose genetic composition is close to that of the *Caulimoviridae* [55].

4. PLANT DEFENSE

4.1 Plant resistance mechanisms

Faced with viral invasion, plants protect themselves by a variety of complementary mechanisms in terms of defense timing, location, and targeting virus-derived molecules, i.e. the viral genome or viral proteins. There are four known defense mechanisms in plants against viruses. i) Dominant resistance relies on interactions between R gene products and viral avirulence (avr) gene products through the establishment of the so-called "gene-for-

gene” interaction; these gene products belong to the nucleotide binding site-leucine-rich repeat (NBS-LRR) class. ii) Recessive resistance, which corresponds to the absence of appropriate host factors that are required for the virus cycle; this is a type of non-inducible resistance, as it is passive and effective throughout plant colonization. It confers resistance at the infection step and requires the cellular factor of interest. iii) RNA interference (RNAi), also called gene silencing, targets viral nucleic acids. Once established after a few days, the effectiveness of this defense mechanism increases and spreads to the whole plant through a relay-amplification process. iv) Hormone-mediated resistance against viral pathogens is mediated by salicylic acid (SA) and methyl-salicylate (Me-SA) in systemic acquired resistance (SAR) [56].

4.2 Plant immunity

There are two types of immunity in plants: i) innate immunity, which relies on cellular actors already present before to the infection, and ii) adaptive immunity (e.g., gene silencing), where defense responses are acquired following an infection and are adapted to the pathogen.

For a long time, researchers believed that the frontline of defense against RNA virus infection was provided by the nonsense-mediated mRNA decay (NMD) system. The NMD system is known as a cellular quality control and regulatory system for RNA that degrades aberrant transcripts, including incorrectly fabricated and non-functional messenger RNA molecules in cells [57,58]. The viral RNA genome has some similarities with incorrect messenger RNA. This system ensures that the genome of certain RNA viruses is broken down, thereby preventing viruses from replicating in host cells [59].

Recently, researchers at ETH Zurich discovered a new form of innate immune defense during their work with human cell culture and a model virus, the *Semliki Forest virus*. In an extensive screening process, the scientists turned off individual genes inside host cells; they discovered that the cells were more susceptible to infection by the virus if the genes of the NMD system were turned off [60].

The same mechanism has been verified in plants by Garcia and colleagues in 2014. Their research was carried out using *Arabidopsis thaliana* and *potato virus X* (PVX) [61]. However, the mechanism is not 100% effective, because if this were the case, RNA viruses would not exist at all. Instead, viruses have evolved ways to avoid or actively suppress NMD. The NMD mechanism probably contributed to shaping the genomes of RNA viruses and the evolution of viruses into what we see today.

Other concepts that have been described as defense mechanisms in plant innate immunity are PAMP-triggered immunity (PTI) and effector-triggered immunity (ETI).

Pattern-triggered immunity

Plants are able to recognize conserved molecules, i.e. pathogen-associated molecular patterns (PAMPs), as a first line of defense. These elicitors are recognized by pattern recognition receptors (PRRs), including receptor-like kinases (RLKs) and receptor-like proteins, which are located on the surface of plant cells [15,62].

In a plant virus, a PAMP is identified as a non-conserved molecule, and the primary plant defense is thought to be based mainly on RNA silencing [63]. Endogenous silencing pathways generate 21-24 nt small (s)RNAs, micro (mi)RNAs and short interfering (si)RNAs that repress genes post-transcriptionally and/or transcriptionally. Four distinct Dicer-like (DCL) proteins, which normally produce endogenous miRNAs and siRNAs, are responsible for the biogenesis of viral siRNAs in infected plants. PTI-based innate responses may contribute to antiviral defense [64].

Effector-triggered immunity

To counteract PTI and establish robust infection in susceptible hosts, pathogens deploy effector proteins (virulence factors) in the plant host cell. In specific cases, plants have evolved resistance (R) genes that mediate the intracellular recognition of effector proteins, which results in ETI. Most plant antiviral R genes encode nucleotide binding site leucine-rich repeat (NBS-LRR) proteins that mediate resistance through the specific (direct or indirect) recognition of a virus avirulence (Avr) factor, and, as a result, trigger the hypersensitivity response (HR) and programmed cell death (PCD) in virus-resistant hosts [65]. In some cases, viral Avr proteins also function as silencing suppressors.

Zigzag model

The interaction between plant defense systems and its suppression by pathogens has been described as a “zigzag model” by Jones and Dangl (2006), also called an “arms race” between plant and pathogens. Counterstrategies developed by successful viruses rely on specific pathogen effector/virulence factors that support pathogen

growth by suppressing PTI, which results in effector-triggered susceptibility (ETS). In order to overcome the action of effectors, plants have evolved ETI. This evolutionary arms race between the host and the pathogen occurs with multiple rounds of ETS followed by ETI [15].

4.3 RNA silencing

RNA silencing is a simple form of adaptive immunity. It is a mechanism that regulates gene expression and chromatin and destroys invasive nucleic acids such as transgenes and viruses. This evolutionarily conserved mechanism is sequence-dependent [66]. RNA silencing is a collective term for homology dependent RNA-based mechanisms directed by small RNAs. Silencing is activated during the production of small double-stranded RNA, leading to RNA degradation of the transgene and the homologous endogenous gene [67]. Small RNAs are known to trigger silencing by initiating the assembly of protein–RNA complexes. This assembly inhibits the expression of specific target genes by downregulating their transcription levels, their mRNA stability, or the translation of their mRNAs into protein [68]. The best studied silencing trigger is double stranded RNA (dsRNA). In addition, siRNA can also be generated via *de novo* synthesis by host RNA polymerases, such as RDRP2 in *Arabidopsis*, using single-stranded RNA transcripts [69,70].

Analogous to the zigzag model, viruses have evolved diverse mechanisms to avoid silencing-mediated resistance. Silencing suppressors target RNAi pathways at different points and through diverse mechanisms, including impaired siRNA biogenesis, defective siRNA incorporation into the RNA-induced cytoplasmic silencing complex (RISC), degradation of argonaute proteins (AGOs), trapping of sRNAs, and the suppression of RNAi amplification (reviewed by Bologna and Voinnet, 2014; Li et al. 2014a) [71,72]. In turn, plants have developed specific defenses against RNA silencing suppression by pathogens, thus far providing another illustration of the never-ending molecular arms race between plant pathogens and their hosts [73,74,56].

In other words, innate (PTI and ETI) and adaptive (gene silencing) antiviral immunities act in a complementary way to fight off the pathogen. However, viruses neutralize this dual defense by effectors that suppress PTI and ETI innate responses and RNA silencing to launch successful infection [63,64].

5. CROSS-PROTECTION

It has been 50 years since researchers discovered the phenomenon of cross-protection. A plant inoculated with a mild virus, with weak infectivity, prevents the multiplication of a subsequent challenge of the same virus or closely related virus which is more virulent. This phenomenon is similar to vaccination in humans and animals and has been used on a large scale in cases where no resistant plants are available. Cross-protection may offer an alternative strategy to reducing economic losses in crops [75,76,77].

Cross-protection has been applied to control various viral diseases such as *tobacco mosaic virus* [78] (Rast, 1972), *zucchini yellow mosaic virus* [79], and *papaya ringspot virus* [80]. Tristeza disease, caused by *citrus tristeza virus* (CTV), is currently controlled by mild CTV isolates [81].

Recently, the cross-protection phenomenon was examined between pathotypes of *pepino mosaic virus* (PepMV) representing the Chilean 2 genotype. Prior inoculation with PepMV-P22, a mild Polish strain, conferred protection against two other severe strains, PepMV-P19 and PepMV-P5-IY, and reduced symptom severity [82]. Moreover, another study showed that enhanced symptom severity can occur when the mild and challenge PepMV isolates belong to different genotypes (EU or LP). This may be due to synergism between different PepMV genotypes or recombinants resulting during co-infection [83]. When a plant is co-infected by diverse viral genomes, competition may lead to decreased fitness of individual genotypes in comparison with their fitness in single infections [84].

Several hypotheses have been suggested to explain the mechanism underlying cross-protection: i) the translation of the primary virus may prevent the transcription of the incoming viral nucleic acid, ii) the production of genome length RNA could be inhibited even if the challenge virus is replicated, iii) prevention of cell-to-cell movement [85], and iv) pre-activation by the protective virus of the RISC with small interfering RNA (siRNA) [86,80].

Cross-protection can be mediated by protein or by RNA or a combination of mechanisms. For example, a recent study showed that unencapsidated viral RNA did not confer protection against challenge inoculations by a related virus. Superinfection exclusion by CTV required the production of a specific viral protein, the p33 protein, as the absence of functional p33 protein completely eliminated the ability of the virus to exclude

superinfection by the same or a closely related virus. Curiously, the protein seemed to work only in a homology-dependent manner [87].

In addition, identifying effective attenuated viruses for each virus of economic importance might be very laborious. A potential solution came recently from the development of engineered vaccines based on viral vectors carrying a genomic fragment of the virulent virus of interest [88,89,90,91].

6. ENGINEERING RESISTANCE

Plant disease resistance is crucial to the reliable production of food. Most economically important crops such as tomato, potato, and wheat are susceptible to viruses. Natural disease resistance carried by plants has not always guaranteed their durable protection against viral disease for the entire productive stage of the crop. The use of broad spectrum resistant (BSR) cultivars remains the best option in virus disease management.

6.1 Breeding and marker-assisted selection

Conventional methods of developing broad-spectrum resistance employ traditional breeding for resistance genes [92]. Employing multiplex parental material, containing three or four copies of dominant resistance genes, has been conducted with limited success due to interspecific barriers and genetic variation between crossed resistant host cultivars and resistant wild relatives. This has been overcome with somatic hybridization through the fusion of protoplasts to transfer resistance to *potato virus Y* (PVY), PVX, *potato leaf roll virus* (PLRV), and CTV from wild relatives [93,94].

Another approach that has proven to provide more durable and broad spectrum resistance is called marker-assisted selection (MAS). This approach uses gene-specific targeted transfer and pyramiding of resistance loci into an elite cultivar [95,96]. Pyramiding of Rsv resistance genes from soybean is a promising method for creating durable and broad spectrum resistance to all strains of *soybean mosaic virus* (SMV) [97]. However, gene pyramiding via conventional breeding has some limitations like losing or affecting important traits over the desired quality or the inevitable transfer of undesirable traits, including toxins and allergens. Traditional breeding for resistance is not sufficient to arm potential crops with high levels of resistance that would maximize yield and quality [98].

Plant transformation or genetic engineering has provided a way to avoid the limitations of conventional breeding. A specific gene is transferred and its expression is directed to the appropriate time or tissue, contributing to the achievement of new desired traits for virus resistance [99].

6.2 Pathogen derived resistance

The pathogen-derived resistance (PDR) approach has been adopted by many plant virus resistance researchers. It consists of introducing a sequence of the viral genome into the host plant [100,101].

- **Protein-mediated resistance:** Coat protein-mediated resistance (CPMR) has been widely used [85,102,103]. CPMR involves the interaction between the transgenic CP and the CP of the challenging virus. The efficiency of CPMR depends on the viral infection cycle [104].

Other viral transgenes including movement and replicase proteins have been used in plant resistance engineering. Replicase protein-mediated resistance (RPMR) using RNA-dependent RNA-polymerase (RdRp) to confer resistance to RNA viruses was first reported for TMV [105], then assessed for CMV and *rice yellow mottle virus* (RYMV) [106,107]. However, this type of resistance is very specific and limited to similar viral strains. For single-stranded DNA viruses, induced resistance takes advantage of viral replication (Rep) associated proteins that interact with host polymerases [108]. Expression of the *tomato yellow leaf curl Sardinia virus* (TYLCSV) C1 gene encoding Rep confers resistance to the homologous virus in *Nicotiana benthamiana* and tomato plants by repressing C1 translation and TYLCSV replication [109].

In addition, movement proteins have been utilized to engineer virus resistance in plants by producing transgenic plants with genes that modify the movement of proteins. Wild type virus encoded movement proteins and preformed dysfunctional movement proteins are in competition to bind to plasmodesmata sites, thus conferring resistance. The first description of movement protein-mediated resistance (MPMR) was shown in tobacco plants [110].

- RNA-mediated resistance: Further investigations have shown that protein-mediated resistance level is not directly related to the viral protein expression level, because high virus resistance levels could remain in transgenic lines that do not express any viral CP. These non-expressing transgenic plants are still resistant because the expressed CP mRNA triggers post-transcriptional gene silencing (PTGS) and provides RNA-mediated resistance to the virus through the siRNA pathway [111]. Based on differences in their biogenesis, two types of sRNAs have been identified, siRNA and miRNA.

The siRNAs produced are then incorporated into the RISC, which guides the cleavage of target RNAs [112]. However, this type of resistance appears to be effective only against related viral sequences. An attempt was undertaken based on the mechanism of PTGS using multiple gene inserts in a single construct. A single chimeric hairpin RNA construct containing viral 4 N gene segments, which is the main constituent of the nucleocapsid from *tomato spotted wilt virus* (TSWV), *groundnut ringspot virus* (GRSV), *tomato chlorotic spot virus* (TCSV), and *watermelon silver mottle virus* (WSMoV) was arranged as inverted repeats. Transgenic plants expressing this construct displayed broad-spectrum resistance against tospoviruses [113]. A similar strategy was used in developing multiple resistance to *Prunus* fruit viruses [114].

The second type of sRNA is miRNA; these are single-stranded RNAs about 21 nt in length, generated from the processing of longer miRNA precursors (pre-miRNA) by Dicer [115]. Recently, the miRNA pathway was shown to downregulate endogenous gene expression in plants. This pathway has been used to design artificial miRNAs (amiRNAs) whose expression in transgenic plants can confer resistance against plant viruses [116,117]. This new anti-viral approach, which has the benefit of reducing possible bio-safety risks associated with protein- and RNA-based strategies, is a first step to designing environmentally friendly virus resistance in transgenic crops [118].

As a strategy for controlling virus diseases, the expression of RNA satellites (Sat-RNAs) in transgenic plants confers resistance to some extent. Sat-RNAs are small parasitic RNAs associated with some plant viruses and are frequently able to modify the symptoms induced by their helper virus [119]. Small sat-RNAs (194 to 400 bases) do not encode proteins [119] and consequently are entirely dependent on virus association for replication and spread in plants. Symptom attenuation (resistance) mediated by sat-RNAs is widely attributed to the inhibition of helper virus replication due to competition for limited replication factors between the helper virus genome and the sat-RNA [120,121].

Induced resistance using sat-RNA has been intensively used against CMV [122,123]. However, the expression of sat-RNAs in transgenic plants may not be strong enough to protect plants from natural viral infections. Transgenic plants expressing both sat-RNA and the coat protein of CMV exhibit enhanced resistance to the virus [124]. Sat-RNA drastically reduces the replication of the viral genome, particularly in Solanaceous hosts and such plants are poor of aphid vector populations [125].

6.3 Antibody-mediated resistance

Antibody-based resistance is a novel strategy for producing transgenic resistant plants. Decades ago, it was shown that polyclonal and monoclonal antibodies can specifically bind to the surface of pathogens such as viruses, bacteria and selected fungi, and result in neutralization. This method has been improved recently by the development of recombinant antibodies (rAbs). Crop resistance can be engineered by the expression of specific antibodies, fragments of antibodies or antibody fusion proteins. This approach has allowed for the engineering of efficient rAbs targeting almost any molecule. It has been confirmed that the expression of functional pathogen-specific rAbs in plants confers effective protection against pathogens. The efficacy of the antibody-based resistance approach for plant viruses has been shown and its application to other plant pathogens is becoming widely used. However, the successful use of antibodies to generate plant pathogen resistance relies on appropriate target selection, careful antibody design, stability, efficient antibody expression, and expression the appropriate cellular compartments [126,127].

Recently, a study generated a model that simulates the complex dynamic interaction between Begomovirus genetics and adaptability to certain plants. This model indicates that the main reason for epidemic outbursts and the global spread of the disease can be found in the patterns of inter-species interactions, many of which are human-induced. In particular, the extensive use of resistant cultivars results triggers aggressive virus adaptability through accelerated mutation. It appears that the only simple option would be to develop more diverse and less concentrated cropping patterns, both in terms of crop land extent and in time [128].

7. CONCLUSIONS

Plant viruses are critical problems to several agricultural and horticultural crops around the world. These pathogens depend on the cellular machinery of their plant hosts. Several aspects of plant-virus interactions happen during the stages of virus infection process both in virus translation and/or replication and in viral cell-to-cell movement.

During the progression of plant-virus coevolution, both have evolved features to battle each other; the plant is equipped with sophisticated and rapidly mounted defense mechanisms, while the virus has developed counterstrategies to overcome those defenses. Understanding the forces that control viral quasispecies may lead to novel ways of predicting the emergence of new viral pathogens and resistance-breaking variants, and provide an overview of the structures of other organism populations. Furthermore, a detailed analysis of viral variability within hosts is needed to manage strategies of disease control, since quasispecies are reservoirs of viral variants that can potentially emerge with increased virulence or altered tropism.

As long as there are no direct methods to control viruses, as with other plant pathogens, current strategies rely on indirect measures for the management of viral diseases. Until the past few decades, classical breeding was considered an effective means to generate resistant plants, but this is costly and difficult; strong genes are rare and when known they do not always confer durable resistance. During the breeding process between plant varieties or species, not only do transferred genes confer the desired strength, but also sometimes affect undesirable genes present on neighboring loci. In addition, features such as crop quality and quantity may be compromised.

Recent advances in molecular biotechnology have made it possible to obtain and to modify genes that are useful for generating disease-resistant crops. A management strategy based on cross-protection can only be successful in areas where one genotype is dominant, otherwise co-infection may result in the emergence of virus variants with new traits after recombination between the protective and challenge isolates. In addition, identifying effective attenuated viruses for each virus of economic importance might be very arduous.

In contrast, numerous strategies, including pathogen-derived expression of sequences or anti-pathogenic agents, have been established to engineer upgraded pathogen resistance in transgenic plants. Transgenic plants expressing the RNA silencing pathway have been shown to efficiently resist viral invasion. It seems that this pathway represents the most specialized molecular strategy that plants use to combat viruses. The expression of sat-RNAs in transgenic plants confers some measure of resistance that may not be strong enough to protect plants from natural virus infections.

However, the risks of biotechnology on the environment and food security related to culture and trade in these transgenic crops are concrete, and it is necessary to define appropriate policies and strategies to protect against adverse effects. Engineering methods that aim to overcome the risks associated with transgenic plants have become more important; these approaches are based on inducing viral RNA silencing without altering the plant genome. In addition, the exploitation of changes in mRNA, cellular metabolites, and protein, once virus invasion has occurred, will help to improve our understanding of plant-virus interactions. In the near future, this will contribute to improving the efficiency of current strategies and allow the establishment of new approaches in favor of sustainable socio-economic development.

REFERENCES

1. King AMQ, Adams MJ, Carstens EB, Lefkowitz E. Virus Taxonomy: Classification and Nomenclature of Viruses: Ninth Report of the International Committee on Taxonomy of Viruses. San Diego: Elsevier Academic Press; 2012
2. Bousalem M, Douzery EJP, Fargette D. High genetic diversity, distant phylogenetic relationships and intraspecies recombination events among natural populations of *Yam mosaic virus*: a contribution to understanding potyvirus evolution. J Gen Virol. 2000;81:243–255.
3. Bateson MF, Lines RE, Revill P, Chaleeprom W, Ha CV, Gibbs AJ, Dale JL. On the evolution and molecular epidemiology of the potyvirus *Papaya ringspot virus*. J Gen Virol. 2002;83:2575–2585.

4. Moreno IM, Malpica JM, Díaz-Pendón JA, Moriones E, Fraile A, García-Arenal F. Variability and genetic structure of the population of *watermelon mosaic virus* infecting melon in Spain. *Virology*. 2004;318:451–460.
5. Ge B, He Z, Zhang Z, Wang H, Li S. Genetic variation in Potato virus M isolates infecting pepino (*Solanum muricatum*) in China. *Arch Virol*. 2014;159(12):3197-210. doi: 10.1007/s00705-014-2180-6.
6. Melgarejo TA, Kon T, Rojas MR, Paz-Carrasco L, Zerbini FM, Gilbertson RL. Characterization of a new world monopartite begomovirus causing leaf curl disease of tomato in Ecuador and Peru reveals a new direction in geminivirus evolution. *J Virol*. 2013;87(10):5397-413. doi: 10.1128/JVI.00234-13.
7. Khan AJ, Akhtar S, Singh AK, Al-Shehi AA, Al-Matrushi AM, Ammara U, **Briddon RW**. Recent evolution of a novel begomovirus causing tomato leaf curl disease in the Al-Batinah region of Oman. *Arch Virol*. 2014;159(3):445-55. doi: 10.1007/s00705-013-1853-x.
8. Hasiów-Jaroszewska B, Czerwoniec A, Pospieszny H, Elena SF. Tridimensional model structure and patterns of molecular evolution of *Pepino mosaic virus* TGBp3 protein. *Virol J*. 2011;8:318. doi: 10.1186/1743-422X-8-318.
9. Goldbach R, deHaan P. RNA viral supergroups and the evolution of RNA viruses. In *The evolutionary biology of viruses*. ed Morse S. S. (Raven Press, New York, N.Y.), 105–120;1994.
10. Schneider WL, Roossinck MJ. Genetic diversity in RNA virus quasispecies is controlled by host-virus interactions. *J Virol*. 2001;75(14):6566-6571. doi: 10.1128/JVI.75.14.6566-6571.2001.
11. Hull R, Harper G, Lockhart B. Viral sequences integrated into plant genomes. *Trends Plant Sci*. 2000;5:362–365. doi: 10.1016/S1360-1385(00)01723-4.
12. Domingo E, Martín V, Perales C, Grande-Pérez A, García-Arriaza J, Arias A. Viruses as quasispecies: biological implications. *Curr Top Microbiol Immunol*. 2006;299:51–82.
13. Gutiérrez S, Yvon M, Thébaud G, Monsion B, Michalakakis Y. Dynamics of the multiplicity of cellular infection in a plant virus. *PLoS Pathog*. 2010;6(9):e1001113. doi:10.1371/journal.ppat.1001113.
14. McKinney HH. Virus mixtures that may not be detected in young tobacco plants. *Phytopathology*. 1926;16:883.
15. Jones JDG, Dangl JL. The plant immune system. *Nature*. 2006;444:323–329. doi: 10.1038/nature05286.
16. Mandahar CL (2006) *Multiplication of RNA Plant Viruses*, Springer Netherlands, 327p, doi: 10.1007/1-4020-4725-8.
17. Domingo E, Sheldon J, Perales C. Viral quasispecies evolution. *Microbiol Mol Biol Rev*. 2012;76(2):159-216. doi: 10.1128/MMBR.05023-11.
18. Di Carli M, Benvenuto E, Donini M. Recent insights into plant-virus interactions through proteomic analysis. *J Proteome Res*. 2012;11(10):4765-4780. doi: 10.1021/pr300494e.
19. Gergerich RC, Dolja VV. Introduction to Plant Viruses, the Invisible Foe. *The Plant Health Instructor*. doi: 10.1094/PHI-I-2006-0414-01; 2006.
20. Boevink P, Oparka KJ. Virus-host interactions during movement processes. *Plant Physiol*. 2005;138:1815–1821. doi:10.1104/pp.105.066761.
21. Benítez-Alfonso Y, Faulkner C, Ritzenthaler C, Maule AJ. Plasmodesmata: gateways to local and systemic virus infection. *Mol Plant Microbe Interact*. 2010;23:1403–1412. doi: 10.1094/MPMI-05-10-0116.
22. Nagy P, Pogany J. Replication of plant RNA viruses. In C Caranta, MA Aranda, M Tepfer, JJ Lopez-Moya, eds, *Recent Advances in Plant Virology*. Caister Academic Press, Norfolk, UK, pp 19–46; 2011.
23. Thivierge K, Nicaise V, Dufresne PJ, Cotton S, Laliberté J-F, Le Gall O, Fortin MG. Plant virus RNAs: coordinated recruitment of conserved host functions by (+) ssRNA viruses during early infection events. *Plant Physiol*. 2005;138:1822–1827. doi: 10.1104/pp.105.064105.
24. Gingras AC, Raught B, Sonenberg N. eIF4 initiation factors: effectors of mRNA recruitment to ribosomes and regulators of translation. *Annu Rev Biochem*. 1999;68:913–963.
25. Selth LA, Dogra SC, Rasheed MS, Healy H, Randles JW, Rezaian MA. A NAC domain protein interacts with *Tomato leaf curl virus* replication accessory protein and enhances viral replication. *Plant Cell*. 2005;17:311–325.
26. Ren T, Qu F, Morris TJ. *HRT* gene function requires interaction between a NAC protein and viral capsid protein to confer resistance to *turnip crinkle virus*. *Plant Cell*. 2000;12:1917–1925.
27. Otulak K, Garbaczewska G. Cell-to-cell movement of three genera (+) ssRNA plant viruses. *Acta Physiol. Plant*. 2011;33:249–260. doi: 10.1007/s11738-010-0538-2.
28. Wolf S, Deom CM, Beachy RN, Lucas WJ. Movement protein of Tobacco mosaic virus modifies plasmodesmatal size exclusion limit. *Science*. 1989;246:377–379.

- 583 29. Lucas WJ. Plasmodesmata and the cell-to-cell transport of proteins and nucleoprotein
584 complexes. *J Exp Bot.* 1999;50:979–987.
- 585 30. Haywood V, Kragler F, Lucas WJ. Plasmodesmata: pathways for protein and
586 ribonucleoprotein signaling. *Plant Cell.* 2002;14(1):S303–S325.
- 587 31. Hofmann C, Niehl A, Sambade A, Steinmetz A, Heinlein M. Inhibition of tobacco mosaic virus
588 movement by expression of an actin-binding protein. *Plant Physiol.* 2009;149:1810–1823.
- 589 32. Fedorkin ON, Solovyev AG, Yelina NE, Zamyatin Jr AA, Zinovkin RA, Mäkinen K, et al. Cell-
590 to-cell movement of *potato virus X* involves distinct functions of the coat protein. *J Gen Virol.*
591 2001; 82(2):449–458.
- 592 33. Morozov SY, Solovyev AG. Triple gene block: modular design of a multifunctional machine for
593 plant virus movement. *Journal of General Virology.* 2003;84:1351–1366. doi:
594 10.1099/vir.0.18922-0.
- 595 34. Verchot-Lubicz J. A new model for cell-to-cell movement of potexviruses. *Mol Plant Microbe*
596 *Interact.* 2005;18:283–290.
- 597 35. Gao Z, Johansen E, Evers S, Thomas CL, Ellis THN, Maule AJ. The potyvirus recessive
598 resistance gene, *sbm1*, identifies a novel role for translation initiation factor eIF4E in cell-to-
599 cell trafficking. *Plant J.* 2004;40:376–385.
- 600 36. González-Jara P, Fraile A, Canto T, Garcia-Arenal F. The multiplicity of infection of a plant
601 virus varies during colonization of its eukaryotic host. *J Virol.* 2009;83:15, 7365. doi:
602 10.1128/JVI.00636-09.
- 603 37. Jridi C, Martin JF, Marie-Jeanne V, Labonne G, Blanc S. Distinct viral populations differentiate
604 and evolve independently in a single perennial host plant. *J Virol.* 2006;80:2349–2357. doi:
605 10.1128/JVI.80.5.2349–2357.2006.
- 606 38. Domingo E, Holland JJ. RNA virus mutations and fitness for survival. *Annu Rev Microbiol.*
607 1997;51:151–78.
- 608 39. Schneider WL, Roossinck MJ. Evolutionarily related sindbis-like plant viruses maintain
609 different levels of population diversity in a common host. *J Virol.* 2000;74:3130–3134.
- 610 40. Lech WJ, Wang G, Yang YL, Chee Y, Dorman K, McCrae D, et al. In vivo sequence diversity
611 of the protease of human immunodeficiency virus type 1: presence of protease inhibitor-
612 resistant variants in untreated subjects. *J Virol.* 1996;70:2038–2043.
- 613 41. Feuer R, Boone JD, Netski D, Morzunov SP, St Jeor SC. Temporal and spatial analysis of Sin
614 Nombre virus quasispecies in naturally infected rodents. *J Virol.* 1999;73:9544–9554.
- 615 42. Ruiz-Jarabo CM, Arias A, Baranowski E, Escarmí's C, Domingo E. Memory in viral
616 quasispecies. *J Virol.* 2000;74:3543–3547. doi: 10.1128/JVI.74.8.3543–3547.2000.
- 617 43. Novella IS, Quer J, Domingo E, Holland JJ. Exponential fitness gains of RNA virus populations
618 are limited by bottleneck effects. *J Virol.* 1999;73:1668–1671.
- 619 44. Li H, Roossinck MJ. Genetic bottlenecks reduce population variation in an experimental RNA
620 virus population. *Journal of Virology.* 2004;78:10582–10587. doi: 10.1128/JVI.78.19.10582-
621 10587.2004.
- 622 45. Akhtar A, and Roossinck M J. Genetic bottlenecks during systemic movement of *Cucumber*
623 *mosaic virus* vary in different host plants. *Virology.* 2010;404(2):279–283. doi:
624 10.1016/j.virol.2010.05.017.
- 625 46. Lozano G, Grande-Pérez A, Navas-Castillo J. Populations of genomic RNAs devoted to the
626 replication or spread of a bipartite virus differ in genetic structure. *J Virol.* 2009;83:12973–
627 12983. doi: 10.1128/JVI.00950-09.
- 628 47. Jończyk M, Borodynko N, Pospieszny H. Restriction analysis of genetic variability of Polish
629 isolates of *Tomato black ring virus*. *Acta Biochim Polon.* 2004;51:673–681.
- 630 48. Tan Z, Gibbs AJ, Tomitaka Y, Sánchez F, Ponz F, Ohshima K. Mutations in *Turnip mosaic*
631 *virus* genomes that have adapted to *Raphanus sativus*. *J Gen Virol.* 2005;86(2):501–10. doi:
632 10.1099/vir.0.80540-0.
- 633 49. Bedhomme S, Lafforgue G, Elena SF. Multihost experimental evolution of a plant RNA virus
634 reveals local adaptation and host specific mutations. *Mol Biol Evol.* 2011;29:1481–1492.
- 635 50. Mauck KE, De Moraes CM, Mescher MC. Evidence of local adaptation in plant virus effects on
636 host–vector interactions. *Integr Comp Biol.* 2014;54(2):193–209. doi: 10.1093/icb/icu012.
- 637 51. Bejarano ER, Khashoggi A, Witty M, Lichtenstein C. Integration of multiple repeats of
638 geminiviral DNA into the nuclear genome of tobacco during evolution. *Proc Natl Acad Sci*
639 *USA.* 1996;93:759–764.
- 640 52. Harper G, Osuji JO, Heslop-Harrison JS, Hull R. Integration of *banana streak badnavirus* into
641 the *Musa* genome: molecular and cytogenetic evidence. *Virology.* 1999;255:207–213. doi:
642 10.1006/vir.1998.9581.

- 643 53. Mao L, Wood TC, Yu Y, Budiman MA, Tomkins J, Woo S, et al. Rice transposable elements: a
644 survey of 73 000 sequence-tagged-connectors. *Genome Res.* 2000;10:982–990. doi:
645 10.1101/gr.10.7.982.
- 646 54. Geering AD, Maumus F, Copetti D, Choisine N, Zwickl DJ, Zytynicki M, et al. Endogenous
647 florendoviral elements are major components of plant genomes and hallmarks of virus
648 evolution. *Nat Commun.* 2014;10 5:5269. doi: 10.1038/ncomms6269.
- 649 55. Nicaise V. Crop immunity against viruses: outcomes and future challenges. *Front Plant Sci.*
650 2014;21(5):660. doi: 10.3389/fpls.2014.00660.
- 651 56. Maquat LE. Nonsense-mediated mRNA decay: Splicing, translation and mRNP dynamics. *Nat*
652 *Rev Mol Cell Biol.* 2004;5:89–99.
- 653 57. Kervestin S, Jacobson A. NMD: a multifaceted response to premature translational
654 termination. *Nat Rev Mol Cell Biol.* 2012;13:700–712. doi: 10.1038/nrm3454.
- 655 58. 28. Melcher U. The ‘30K’ superfamily of viral movement proteins. *J Gen Virol.* 2000;81:257–
656 266.
- 657 59. Rayson S, Arciga-Reyes L, Wootton L, De Torres Zabala M, Truman W, Graham N, et al. A
658 role for nonsense-mediated mRNA decay in plants: pathogen responses are induced in
659 *Arabidopsis thaliana* NMD mutants. *PLoS One.* 2012;7:e31917. doi:
660 10.1371/journal.pone.0031917.
- 661 60. Balistreri G, Horvath P, Schweingruber C, Zünd D, McInerney G, Merits A, Mühlemann O,
662 Azzalin C, Helenius A. The host nonsense-mediated mRNA decay pathway restricts
663 mammalian RNA virus replication. *Cell Host & Microbe.* 2014;16(3):403–411. doi:
664 10.1016/j.chom.2014.08.007.
- 665 61. Garcia D, Garcia S, Voinnet O. Nonsense-Mediated Decay Serves as a General Viral
666 Restriction Mechanism in Plants. *Cell Host & Microbe.* 2014;16(3):391–402. doi:
667 10.1016/j.chom.2014.08.001.
- 668 62. Newman M-A, Sundelin T, Nielsen JT, Erbs G. MAMP (microbe-associated molecular pattern)
669 triggered immunity in plants. *Front Plant Sci.* 2013;4:139. doi:10.3389/fpls.2013.00139.
- 670 63. Csorba T, Pantaleo V, Burgyán J. RNA silencing: an antiviral mechanism. *Adv Virus Res.*
671 2009;75:35–71. doi: 10.1016/S0065-3527(09)07502-2.
- 672 64. Zvereva AS, Pooggin MM. Silencing and innate immunity in plant defense against viral and
673 non-viral pathogens. *Viruses.* 2012;4(11):2578–2597. doi:10.3390/v4112578.
- 674 65. Coll NS, Eppe P, Dangl JL. Programmed cell death in the plant immune system. *Cell Death*
675 *Differ.* 2011;18:1247–1256. doi: 10.1038/cdd.2011.37.
- 676 66. Ding SW, Voinnet O. Antiviral immunity directed by small RNAs. *Cell.* 2007;130:413–426. doi:
677 10.1016/j.cell.2007.07.039.
- 678 67. Voinnet O. Use, tolerance and avoidance of amplified RNA silencing by plants. *Trends Plant*
679 *Sci.* 2008;13:317–328. doi: 10.1016/j.tplants.2008.05.004.
- 680 68. Tomari Y, Zamore PD. Perspective: machines for RNAi. *Genes Dev.* 2005;19:517–529.
- 681 69. Ghildiyal M, Zamore PD. Small silencing RNAs: an expanding universe. *Nat Rev Genet.*
682 2009;10, 94–108. doi: 10.1038/nrg2504.
- 683 70. Galvez LC, Banerjee J, Pinar H, Mitra A. Engineered plant virus resistance. *Plant Science.*
684 2014;228:11–25. doi: 10.1016/j.plantsci.2014.07.006.
- 685 71. Bologna NG, Voinnet O. The diversity, biogenesis, and activities of endogenous silencing
686 small RNAs in *Arabidopsis*. *Annu Rev Plant Biol.* 2014;65:473–503. doi:10.1146/annurev-
687 arplant-050213-035728.
- 688 72. Li F, Huang C, Li Z, Zhou X. Suppression of RNA silencing by a plant DNA virus satellite
689 requires a host calmodulin-like protein to repress RDR6 expression. *PLoS Pathog.*
690 2014;10(2): e1003921. doi:10.1371/journal.ppat.1003921.
- 691 73. Pumplin N, Voinnet O. RNA silencing suppression by plant pathogens: defence, counter-
692 defence and counter-counter-defence. *Nat Rev Microbiol.* 2013;11:745–760. doi:
693 10.1038/nrmicro3120.
- 694 74. Sansregret R, Dufour V, Langlois M, Daayf F, Dunoyer P, Voinnet O. Extreme resistance as a
695 host counter-counter defense against viral suppression of RNA silencing. *PLoS Pathog.*
696 2013;9:e1003435. doi:10.1371/jour- nal.ppat.1003435.
- 697 75. Ziebell H, Carr JP. Cross-protection: a century of mystery. *Adv Virus Res.* 2010;76:211–264.
698 doi: 10.1016/S0065-3527(10)76006-1.
- 699 76. Nishiguchi M, Kobayashi K. Attenuated plant viruses: preventing virus diseases and
700 understanding the molecular mechanism. *J Gen Plant Pathol.* 2011;77:221–229.
- 701 77. Zhou C, Zhou Y. Strategies for viral cross-protection in plants. *Meth Mol Biol.* 2012;894:69–81.
702 doi: 10.1007/978-1-61779-882-5.
- 703 78. Rast ATB. MII-16, an artificial symptomless mutant of *Tobacco mosaic virus* for seedling
704 inoculation of tomato crops. *Neth J Plant Pathol.* 1972;78:110–112.

79. Lecoq H, Lemaire JM. Control of *zucchini yellow mosaic virus* in squash by cross protection. *Plant Dis.* 1991;75:208–211.
80. Gal-On A, Shibolet Y. "Cross protection," in *Natural Resistance Mechanisms of Plants to Viruses*, ed. Loebeinstein G., Carr J. P., editors. (Dordrecht: Springer) 261–288;2006.
81. Lee RF, Keremane ML. Mild strain cross protection of tristeza: a review of research to protect against decline on sour orange in Florida. *Front Microbiol.* 2013;4:259. doi: 10.3389/fmicb.2013.00259.
82. Hasiów-Jaroszewska B., Minicka J, Pospieszny H. Cross-protection between different pathotypes of *Pepino mosaic virus*. *Acta Sci Pol, Hortorum Cultus.* 2014;13(5):177-185.
83. Hanssen IM, Gutierrez-Aguirre A, Paeleman A, Goen K, Wittemans L, Lievens B, et al. Cross-protection or enhanced symptom display in greenhouse tomato co-infected with different *Pepino mosaic virus* isolates. *Plant Pathol.* 2010;59:13–21. doi: 10.1111/j.1365-3059.2009.02190.x.
84. Frank SA. Multiplicity of infection and the evolution of hybrid incompatibility in segmented viruses. *Heredity.* 2001;87:522-529. doi:10.1046/j.1365-2540.2001.00911.x.
85. Beachy RN. Coat-protein-mediated resistance to *Tobacco mosaic virus*: discovery mechanisms and exploitation. *Philos Trans R Soc Lond B: Biol Sci.* 1999;354:659–664. doi: 10.1098/rstb.1999.0418.
86. Ratcliff FG, MacFarlane S, Baulcombe DD. Gene silencing without DNA: RNA-mediated cross-protection between viruses. *Plant Cell.* 1999;11:1207–1215.
87. Folimonova SY. Superinfection exclusion is an active virus-controlled function that requires a specific viral protein. *J Virol.* 2012;86(10):5554-5561. doi: 10.1128/JVI.00310-12.
88. Culver JN. Tobamovirus cross protection using a potexvirus vector. *Virology.* 1996;226:228–235. doi: 10.1006/viro.1996.0650.
89. Bazzini AA, Asurmendi S, Hopp HE, Beachy RN. *Tobacco mosaic virus* (TMV) and *Potato virus X* (PVX) coat proteins confer heterologous interference to PVX and TMV infection, respectively. *J Gen Virol.* 2006;87:1005–1012. doi: 10.1099/vir.0.81396-0.
90. Tamura A, Kato T, Taki A, Sone M, Satoh N, Yamagishi N. Preventive and curative effects of Apple latent spherical virus vectors harboring part of the target virus genome against potyvirus and cucumovirus infections. *Virology.* 2013;446:314–324. doi: 10.1016/j.virol.2013.08.019.
91. Taki A, Yamagishi N, Yoshikawa N. Development of apple latent spherical virus-based vaccines against three tospoviruses. *Virus Res.* 2013;176:251–258. doi: 10.1016/j.virusres.2013.06.015.
92. Foxe MJ. Breeding for viral resistance: conventional methods. *Netherlands J Plant Pathol.* 1992;98(2):13-20.
93. Vaikonen JPT. Natural genes and mechanisms for resistance to viruses incultivated and wild potato species (*Solanum* spp.). *Plant Breed.* 1994;112:1–16.
94. Mendes BJ, Filho FM, Farias P, Benedito V. Citrus somatic hybridization with potential for improved blight and CTV resistance. *In Vitro Cell Dev Biol Plant.* 2001;37:490–495.
95. Richardson KL, Vales MI, Kling JG, Mundt CC, Hayes PM. Pyramiding and dissecting disease resistance QTL to barley stripe rust. *Theor Appl Genet.* 2006;113:485–495.
96. Asea G, Vivek BS, Bigirwa G, Lipps PE, Pratt RC. Validation of consensus quantitative trait loci associated with resistance to multiple foliar pathogens of maize. *Phytopathology.* 2009;99:540–547. doi: 10.1094/PHYTO-99-5-0540.
97. Saghai Maroof MA, Jeong SC, Gunduz I, Tucker DM, Buss GR, Tolin SA. Pyramiding of *Soybean mosaic virus* resistance genes by marker-assisted selection. *Crop Sci.* 2008;48:517–526. doi:10.2135/cropsci2007.08.0479.
98. Rommens CM, Haring MA, Swords K, Davies HV, Belknap WR. The intragenic approach as a new extension to traditional plant breeding. *Trends Plant Sci.* 2007;12:397–403. doi: 10.1016/j.tplants.2007.08.001.
99. Binns ANTMF. Cell biology of *Agrobacterium* infection and transformation of plants. *Annu Rev Microbiol.* 1988;42:575-606.
100. Prins M, Laimer M, Noris E, Schubert J, Wassenegger M, Tepfer M. Strategies for antiviral resistance in transgenic plants. *Mol Plant Pathol.* 2008;9:73–83. doi: 10.1111/j.1364-3703.2007.00447.x.
101. Gómez P, Rodríguez-Hernández AM, Moury B, Aranda MA. Genetic resistance for the sustainable control of plant virus diseases: breeding, mechanisms and durability. *Eur J Plant Pathol.* 2009;125:1–22. doi: 10.1007/s10658-009-9468-5.
102. Mundembe R, Matibiri A, Sithole-Niang I. Transgenic plants expressing the coat protein gene of cowpea aphid-borne mosaic potyvirus predominantly convey the delayed symptom development phenotype. *Afr J Biotechnol.* 2009;8:2682–2690.

103. Panter S, Chu PG, Ludlow E, Garrett R, Kalla R, Jahufer MZ, et al. breeding of transgenic white clover (*Trifolium repens* L.) with field resistance to *Alfalfa mosaic virus* through the expression of its coat protein gene. *Transgenic Res.* 2012;21:619–632.
104. Bendahmane M, Chen I, Asurmendi S, Bazzini AA, Szecsi J, Beachy RN. Coat protein-mediated resistance to TMV infection of *Nicotiana tabacum* involves multiple modes of interference by coat protein. *Virology.* 2007;366:107–116.
105. Golemboski DB, Lomonossoff GP, Zaitlin M. Plants transformed with a *Tobacco mosaic virus* nonstructural gene sequence are resistant to the virus. *Proc Natl Acad Sci U.S.A.* 1990;87:6311–6315. doi: 10.1371/journal.pone.0015087.
106. Wintermantel WM, Zaitlin M. Transgene translatability increases effective-ness of replicase-mediated resistance to Cucumber mosaic virus. *J Gen Virol.* 2000;81:587–595.
107. Palukaitis P, Zaitlin M. Replicase-mediated resistance to plant virus disease. *Adv Virus Res.* 1997;48:349–377.
108. Shivaprasad PV, Thillaichidambaram P, Balaji V, Veluthambi K. Expression of full-length and truncated Rep genes from *mungbean yellow mosaic virus*-Vigna inhibits viral replication in transgenic tobacco. *Virus Genes.* 2006;33:365–374.
109. Brunetti A, Tavazza M, Noris E, Tavazza R, Caciagli P, Ancora G, et al. High expression of truncated viral rep protein confers resistance to *Tomato yellow leaf curl virus* in transgenic tomato plants. *Mol Plant Microbe Interact.* 1997;10:571–579.
110. Lapidot M, Gafny R, Ding B, Wolf S, Lucas WJ, Beachy RN. A dysfunctional movement protein of *Tobacco mosaic virus* that partially modifies the plasmodesmata and limits virus spread in transgenic plants. *Plant J.* 1993;4:959–970.
111. Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature.* 1998;391:806–811. doi :10.1038/35888.
112. Baulcombe DC. RNA silencing in plants. *Nature.* 2004;431:356–363. doi:10.1038/nature02874.
113. Bucher E, Lohuis D, van Poppel PM, Geerts-Dimitriadou C, Goldbach R, Prins M. Multiple virus resistance at a high frequency using a single transgene construct. *J Gen Virol.* 2006;87:3697–3701. doi :10.1099/vir.0.82276-0.
114. Liu Z, Scorza R, Hily JM, Scott SW, James D. Engineering resistance to multiple prunus fruit viruses through expression of chimeric hairpins. *J Am Soc Hortic Sci.* 2007;132:407–414.
115. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell.* 2004;116:281–297.
116. Niu QW, Lin SS, Reyes JL, Chen KC, Wu HW, Yeh SD, et al. Expression of artificial microRNAs in transgenic *Arabidopsis thaliana* confers virus resistance. *Nat Biotechnol.* 2006;24:1420–1428.
117. Qu J, Ye J, Fang R. Artificial microRNA-mediated virus resistance in plants. *J Virol.* 2007;81(12):6690-9. doi: 10.1128/JVI.02457-06.
118. Lin S S, Henriques R, Wu HW, Niu QW, Yeh SD, Chua NH. Strategies and mechanisms of plant virus resistance. *Plant Biotechnol Rep.* 2007;1:125-134.
119. Roossinck MJ, Sleat D, Palukaitis P. Satellite RNAs of plant viruses: structures and biological effects. *Microbiol Rev.* 1992 ;56(2):265–279
120. Collmer CW and SH Howell. Role of satellite RNA in the expression of symptoms caused by plant viruses. *Annu Rev Phytopathol.* 1992;30:419–442.
121. Wu G, and Kaper JM. Competition of viral and satellite RNAs of cucumber mosaic virus for replication in vitro by viral RNA-dependent RNA polymerase. *Res Virol.* 1995;146:61-67.
122. Kim SJ, Lee SJ, Kim BD, Paek KH. Satellite RNA-mediated resistance to *Cucumber mosaic virus* in transgenic plants of hot pepper (*Capsicum annuum* cv. Golden Tower). *Plant Cell Rep.* 1997;16:825–830.
123. Cillo F, Finetti-Sialer MM, Papanice MA, Gallitelli D. Analysis of mechanisms involved in the *Cucumber mosaic virus* satellite RNA-mediated transgenic resistance in tomato plants. *Mol Plant Microbe Interact.* 2004;17:98–108.
124. Yie Y, Tien P. Plant virus satellite RNAs and their role in engineering resistance to virus diseases. *Seminars in Virology.* 1993;4(6):363–368.
125. Escriu F, Perry KL, García-Arenal F. Transmissibility of *Cucumber mosaic virus* by *Aphis gossypii* correlates with viral accumulation and is affected by the presence of its satellite RNA. *Phytopathology.* 2000;90:1068–1072. doi: 10.1094/PHYTO.2000.90.10.1068.
126. Schillberg S, Zimmermann S, Zhang MY, Fischer R. Antibody-based resistance to plant pathogens. *Transgenic Res.* 2001;10(1):1-12.

- 829 127. Safarnejad MR, Jouzani GS, Tabatabaei M, Twyman RM, Schillberg S. Antibody-mediated
830 resistance against plant pathogens. *Biotechnol Adv.* 2011;29(6):961-71. doi:
831 10.1016/j.biotechadv.2011.08.011.
- 832 128. Jabłńska-Sabuka M, Kalaria R, Kauranne T. A dynamical model for epidemic outbursts by
833 begomovirus population clusters. *Ecological Modelling.* 2015;297:60–68. doi:
834 10.1016/j.ecolmodel.2014.11.008.
835