

-----Original Message-----

From: XXX YYY<xxx@yyy.zzz>

Sent: Wed, Aug 7, 2013 at 7:59 AM

To: Managing Editor

Subject: Re: Request for final decision for manuscript number 2013_CA_5783

I can not read the file "author's feedback for reviewer MA".
Could you send me again?

Comment;

First of all, PVR after TAVI procedure is an important topic. At present TAVI can be applied in many countries, so it is natural to expect that it will be dramatically increasing in case as like endovascular surgery for aortic aneurysm. I think that this article is hit the topic and should be worthy for publication.

Minor change should be required as follows,

1. in abstract, aims: The secondary aims of this study were to review the impact of moderate severe PVR....: change "moderate to severe"
2. in abstract, results: Management strategies include balloon valvuloplasty, TAVI-in-TAV,... should be spell out
3. line 22: TAVI ... should be spell out
4. line 28: "paravalvular leak" should change "paravalvular leak (PVR)"
5. line 92: what is the MCV?
6. line 95: spell out TAVI-in TAV, if you use it in first time in manuscript body
7. line 102: where did you show "patients of a similar age and Logistic EUROscores"?
8. line 127: where moderate and severe PVR... is it better to change "moderate to severe" to unity the same words? Line 143 is also; moderate/severe PVR.
9. in discussion: did you find any comments or data regarding preoperative risk factor for PVR after TAVI? I think it is better to state.

Editor's Details:

Hirofumi Kasahara

Department of Cardiovascular Surgery

National Hospital Organization Saitama Hospital, Wako, Japan

Note: Modification was done in this document ONLY to hide the email id.