

Case Study

Infantile cardiac rhabdomyoma–Pearls inside the heart

ABSTRACT

Cardiac rhabdomyoma is the most common primary pediatric tumor of the heart. We reported a 1-month old male infant who presented to our institute for routine cardiac evaluation since he was diagnosed to have a cardiac mass in the right ventricle (RV) in utero. After he was born, an echocardiogram showed two large cardiac masses occupying entire RV cavity and origin of right ventricular outflow tract (RVOT). Although our patient was asymptomatic, surgical removal of these two masses was done due to its proximity to RVOT and also because it was almost obliterating the entire RV cavity.

Keywords: Cardiac tumor; rhabdomyoma; right ventricular mass

INTRODUCTION

Cardiac tumors were first described by Boneti [1]. Cardiac rhabdomyoma usually occurs as multiple masses, and arises from ventricular free wall and septal wall. 50-80% of these patients had associated tuberous sclerosis. Although cardiac rhabdomyoma has benign prognosis and is known to undergo spontaneous regression, some may require surgical removal if the location was near or in the right ventricular outflow tract (RVOT), left ventricular outflow tract (LVOT), or when they produce intractable arrhythmias.

CASE REPORT

A 1-month-old male infant presented to our institute for routine cardiac evaluation. Fetal echo had showed intracardiac mass at 30 weeks of gestation. The baby was born to non-consanguineous parents. Tuberous sclerosis (TCS) screening was negative. The baby was asymptomatic at the time of evaluation. A routine echocardiogram (Figure 1) showed a large 3 x 1.2 cm mass occupying entire RV cavity and another 0.9 x 0.7 cm mass at the

origin of RVOT causing mild obstruction. In view of the size and location, despite being asymptomatic, the infant underwent successful surgical removal of the masses (figure 2) and was discharged uneventfully.

DISCUSSION

The prevalence of cardiac tumor is extremely rare in infants and children (0.027 to 0.17%). Cardiac rhabdomyoma is the most common type of cardiac tumor. More than 60% of antenatally diagnosed cardiac tumors are rhabdomyomas and are often associated with tuberous sclerosis [2]. Cardiac rhabdomyoma is seen in 30-80% of patients with tuberous sclerosis and usually regresses spontaneously over time [3]. In addition, 50–60% of patients with tuberous sclerosis will develop cardiac rhabdomyoma [7,8].

A meta-analysis showed that most rhabdomyomas associated with tuberous sclerosis are non-obstructive [4]. The natural history of cardiac rhabdomyoma in infants and children has been well studied [9]. The younger the age at diagnosis, the higher the chance for spontaneous regression, and complete regression being more common in the first 4 years of life. In 2D echocardiogram, they appear as round, homogenous masses [5]. Indications for surgery included life-threatening arrhythmias, and outflow tract obstructions [6]. If complete surgical resection is not possible because of the location of the tumor, a partial resection can be done and the residual tumor usually regresses. The mortality from surgery for cardiac tumor in children is about 5% [10]

A serial echocardiography may help in selecting patients who are at risk of developing ventricular outflow obstruction and who need early surgery. All cases of cardiac rhabdomyoma must be followed up closely for the future developed tuberous sclerosis.

CONCLUSION

Although watchful waiting is the general strategy for cardiac rhabdomyomas, certain clinical situations like intractable arrhythmias and locations near outflow tract causing obstruction warrant early surgical intervention. Our patient underwent surgery empirically due to its location near the RVOT and the tumor occupying almost the entire RV cavity. Because rhabdomyoma is

known to undergo spontaneous regression, general agreement was indicated for surgical treatment, which should be individualized and considered only in the presence of significant hemodynamic obstruction to the ventricular inflow or the outflow tract.

REFERENCES

1. Perlstein et al.: Sarcoma of the heart. *Am. J. M. SC.* 156: 214, 1918.
2. Isaacs H, Jr. Fetal and neonatal cardiac tumors. *PediatrCardiol* 2004;25(3):252-273.
3. Smythe JF, Dyck JD, Smallhorn JF, Freedom RM. Natural history of cardiac rhabdomyoma in infancy and childhood. *Am J Cardiol.* 1990;66:1247–1249.
4. Verhaaren H A, Vanakker O, DeWolf D, Suys B, Francois K, Matthys D. Left Ventricular outflow tract obstruction in rhabdomyoma of infancy: Meta-analysis of the literature. *J Pediatr.* 2003;143:258–263.
5. Chao AS, Chao A, Wang TH, Chang YC, Chang YL, Hsieh CC, Lien R, et al. Outcome of antenatally diagnosed cardiac rhabdomyoma: case series and a meta-analysis. *Ultrasound ObstetGynecol* 2008;31(3):289-295
6. Stiller B, Hetzer R, Meyer R, Dittrich S, Pees C, Alexi-Meskishvili V, Lange P. Primary cardiac tumours: when is surgery necessary? *Eur J of Cardio-thoracic surgery.* 2001;20:1002–06.
7. Webb DW, Thomas RD, Osborne JP .Cardiac rhabdomyomas and their association with tuberous sclerosis. *Arch Dis Child* 1993 Mar;68((3)):367-370.
8. Harding CO, Pagon RA. Incidence of tuberous sclerosis in patients with cardiac rhabdomyoma. *Am J Med Genet* 1990 Dec;37((4)):443-446.
9. Bosi G, Lintermans JP, Pellegrino PA, Svaluto-Moreolo G, Vliers A. The natural history of cardiac rhabdomyoma with and without tuberous sclerosis. *Acta Paediatr* 1996 Aug;85((8)):928-931.
10. Takach TJ, Reul GJ, Ott DA, Cooley DA. Primary cardiac tumors in infants and children: immediate and long term operative results. *Ann Thorac Surg* 1996 Aug;62((2)):559-564.

102

103


104

105 **FIGURE LEGEND**

106 Figure 1:Trans thoracic echo (TTE) showing rhabdomyoma in right ventricular
107 (RV) cavity and origin of right ventricular outflow tract (RVOT).

108

109 **FIGURE 1**




110

111

112

113 Figure 2-Gross and histopathological specimen showing rhabdomyoma
114 extracted from RV cavity



115

116

117