

Antioxidative Action of *Citrus limonum* Essential Oil on Skin

G. Bertuzzi¹, B. Tirillini², P. Angelini^{3*}, R. Venanzoni³

(*corresponding author: telephone number +39 0755856323, fax number +39 075 5856404, email paola.angelini@unipg.it)

¹Masters in Aesthetic Medicine, Dept. of Internal Medicine, University of Roma "Tor Vergata", Via Montpellier, 1 – 00133 Roma (RM) – Italy

²Dept. of Biomolecular Science, University of Urbino, Via Bramante, 28 - 61028 Urbino (PU) – Italy

³Dept. of Applied Biology, University of Perugia, Borgo XX giugno 74 - 06121 Perugia (PG) - Italy

ABSTRACT

Aims: The purpose of this study was to investigate the **action** of *Citrus limonum* essential oil to control free radical-induced lipid peroxidation and preventing tissue damage in skin.

Place and Duration of Study: Department of Internal Medicine (University of Roma "Tor Vergata") and A.R.P.A (Aging Research, Prevention and Therapy Association, www.anti-aging.it), between January 2010 and June 2011.

Methodology: The essential oil was subjected to GC-MS analysis. The superoxide anion scavenging activity of essential oil was evaluated by the enzymatic hypoxanthine/xanthine oxidase system. The same oil diluted in DMSO or grape-seed oil was spread on the face of human volunteers after UV exposition. A sample of skin lipids were collected and the presence of peroxy radicals were detected based on the measurement of light emitted (chemiluminescence) when the excited carbonyl and singlet oxygen decay to ground state.

Results: Our data demonstrate that the lemon essential oil is more active than **α -tocopherol** against O_2^- and peroxide free radical inhibition at a 1:100 dilution. A protocol for controlling free radical-induced lipid peroxidation in human skin was thus proposed.

Conclusion: The scavenging action of lemon essential oil could have a practical application for treating human skin against oxidative damage.

Keywords: anti-aging, GC-MS, grape seed oil, superoxide anion scavenging.

1. INTRODUCTION

The inhibition of lipid oxidation by essential oils such as *Origanum* spp., *Thymus* spp., *Satureja* spp., and *Rosmarinus officinalis*, have already been reported in literature (Estevez, M., Cava, R. (2006); Kulisic et al., (2005); Nakatsu et al., (2000)).

All the essential oils studied have shown a strong phenolic profile characterized by the presence of phenolic monoterpenes which are believed to be the active component of the essential oils (Teissedre, P.L., Waterhouse, A.L. (2000); Angelini, P. et al., (2006); Angelini, P. et al., (2008); Angelini, P. et al., (2009); Pagiotti, R. et al., (2011); Tirillini, B. et al.,

* Tel.: +39 075 5856423; fax: +39 075 5856404.

E-mail address: paola.angelini@unipg.it.

(2009)). *Citrus* essential oil has also been reported to have antioxidative activities against linoleic acid oxidation (Song, H.S. et al., (2001)) and to induce both Cu²⁺-induced and 2,2'-azobis(2-aminopropane)hydrochloride-induced oxidation of human low-density lipoprotein *in vitro* (Takahashi, Y. et al., (2003)). Among the compounds tested in *Citrus* essential oil, γ -terpinene had the strongest antioxidant effect (Takahashi, Y. et al., (2003)), but no clear relationship could be shown between the antioxidant activity and the essential oil composition of the extracts (Di Vaio, C. et al., (2010)). When skin is exposed to air that is irradiated by ultraviolet (UV) light consisting of UVA (320-400 nm) and UVB (290-320 nm), reactive oxygen species (ROS) including superoxide anion radical ($\text{O}_2^{\cdot-}$), hydrogen peroxide (H_2O_2), hydroxyl radical (OH^{\cdot}), singlet oxygen (O_2^1), lipid peroxides (LOOH), and their radicals (LOO $^{\cdot}$) are formed. These in turn induce skin aging, phototoxicity, inflammation and malignant tumors (Bech-Thomsen, N., Wulf, H.C. (1995); Kligman, A.M. (1969); Oikarinen, A. et al., (1985); Sakurai, H., et al., (2005); Watson, R.E.B., Griffiths, C.E.M. (2005)). Recently, consumer interest and the media have focused specifically on products that use natural ingredients, such as plant extracts. There is some evidence that these ingredients could have possible *in vitro* anti-aging activity, but the question remains whether it is possible to deliver adequate doses to the skin *in vivo*. Lemon oil, traditionally used for its aromatic properties, has recently been investigated for its effects on skin (Chiu, A., Kimbal, A.B. (2003)). The purpose of this study was to investigate the effectiveness of *Citrus limonum* Risso essential oil in controlling free radical-induced lipid peroxidation and preventing tissue damage in skin.

2. MATERIALS AND METHODS

2.1 Plant material

The *Citrus limonum* (lemon) essential oil used in this study was obtained from cold pressed oil extracted from the peel of the fruit according to the methods of Sawamura and Kuriyama (1988). The cold pressed oil was then hydrodistilled for 1h in an all-glass Clevenger apparatus.

2.2 GC and GC-MS Analysis

The GC analyses were carried out using a Varian 3300 instrument equipped with an FID and an HP-InnoWax capillary column (30 m x 0.25 mm, film thickness 0.17 μm), working from 60°C (3 min) to 210°C (15 min) at 4°C/min or an HP-5 capillary column (30 m x 0.25 mm, film thickness 0.25 μm) working from 60°C (3 min) to 300°C (15 min) at 4°C/min; The injector and detector temperature was 250°C. Helium was used as the carrier gas, with a flow rate of 1 ml/min, and the split ratio was 1 : 10.

GC-MS analyses were carried out with a Hewlett Packard 5890 GC-MS system operating in the EI mode at 70 eV, using the two above-mentioned columns. The operating conditions were analogous to those reported in the GC analyses section. The injector and transfer line temperatures were 220°C and 280°C, respectively. Helium was used as the carrier gas, with a flow rate of 1 ml/min, and the split ratio was 1 : 10.

2.3 Identification of the components

The components were identified by matching the spectra with those from mass spectral libraries; the identity of each component was confirmed by comparing the retention indices, from both columns, relative to the C6-C22 n-alkanes, with those from the literature (Adams, R.P. (2001); Davies, N.W. (1990); Heller, S.R., Milne, G.W.A. (1983); Jennings, W.G., Shibamoto, T. (1980); McLafferty, F.W., Staufner, D.B. (1989)). When reported, co-elution gas

chromatography with reference compounds was also used for an additional confirmation of the compound identity.
The percentage composition of the essential oil was obtained by the normalization method from the GC peak areas, without using correction factors.

2.4 Superoxide anion scavenging ($\text{O}_2^{\cdot-}$)

Superoxide anion was generated by a hypoxanthine-xanthine oxidase system (Arouma, O. et al., (1989)). Reaction mixtures with 100 μl EDTA (30 mmol/l), 10 μl hypoxanthine (30 mmol/l), 100 μl cytochrome c (3 mmol/l) or nitroblue tetrazolium (3 mmol/l) were added to 150 μl of lemon essential oil (solubilized in DMSO 10%) at various concentrations in a final volume of 3 ml buffered in KH_2PO_4 (50 mmol/l), pH 7.4 (Gressier, B. et al., (1993)). The reaction was started by adding 200 μl xanthine oxidase (1U/ml) and the rate of reduced cytochrome c or nitroblue tetrazolium was measured at 550 nm, and 560 nm, respectively, against a reference. The amount of $\text{O}_2^{\cdot-}$ generated was calculated using the extinction coefficient $\epsilon_{550} = 2.1 \times 10^{-2} \mu\text{mol}^{-1} \text{cm}^{-1}$ per cm and the $\text{O}_2^{\cdot-}$ inhibition was expressed as percentage values. The sample tested did not interfere with the xanthine oxidase activity (measured at 290 nm). The positive response was tested using α -tocopherol. Ten repetitions were carried out.

2.5 Randomized controlled trial

2.5.1. Subjects

Subjects were selected from among men aged 18 to 52 (mean 33 ± 11) years who were found to have no serious illness on physical checkup at A.R.P.A (Aging Research, Prevention and Therapy Association), www.anti-aging.it (Civita Castellana, VT, Italy). Eighty volunteers (average age: 33 ± 11 years) who gave their written consent to participate in the test were selected as subjects from January 2010 to June 2011.

2.5.2. Extraction of skin lipids from healthy volunteers

Skin lipids were collected with acetone-wetted cotton swabs from the forehead over a 9 cm^2 area from healthy volunteers (80 men, 18–52 years old—mean 33 ± 11) in the morning for 7 days. The sampling procedure was identical for all the subjects. The volunteers were randomly divided into four groups (A, B, C, D). In group A the forehead was treated with α -tocopherol in ethanol (20%), group B with lemon essential oil solubilized in DMSO (1:100), group C with lemon essential oil solubilized in grape-seed oil (1:100), and group D was left untreated. In accordance with the European norm EN 60335-2-27 and under medical supervision, volunteers were irradiated with UVA and UVB of $0.3 \text{ W/m}^2/\text{mm}$ from sunlamps for 7 min at each session. The participants were asked not to expose themselves to direct sunlight and to avoid the use of face creams or hair lotions for the entire duration of the experiment. Twenty-four hours after the last treatment, the skin lipids were collected.

Extracts were taken twice from the wet cotton swabs using 3 ml of chloroform/methanol (1:2.5) for two hours (10 μg heneicosanoic acid was used for the recovery test). The raw extracts were partitioned between 1% NaCl in 0.01 M HCl and chloroform. The chloroform extracts were washed with methanol/water (1:1) and dried under N_2 stream. The samples were stored at -20°C in 3 ml of chloroform/ethanol (2:1).

2.6 Lipid peroxidation analyzed by chemiluminescence

Chemiluminescence is an index of oxidative stress that quantifies lipid peroxidation and was measured according to the method of Gonzalez-Flecha, B. et al., (1991). This method is

based on the measurement of light emitted (chemiluminescence) when the excited carbonyl and singlet oxygen produced by peroxy radicals decay to ground state. This light is due to the generation of reactive oxygen species in whole lipids. Skin lipids were incubated with 3 mM t-BHP for 60 min at 37 °C. Lipid peroxidation was initiated by adding a small amount of stock solution of t-butyl hydroperoxide (80 mM) to each vial which was then maintained at 37 °C, and measured by monitoring light emission (Wright et al., 1979) with a liquid scintillation analyzer Packard 1900 TR. Chemiluminescence was measured over a 60 min period and recorded as counts per minute (cpm) every 12 min. Each reaction was terminated by adding 5 ml chloroform/methanol (2:1, v/v) containing 0.01% butylated hydroxytoluene (BHT). This also inhibited any further oxidation during the lipid extraction. The DMSO had no antioxidative action and gave a chemiluminescence curve that could be superimposed on to that of the control.

Statistical analysis

Analysis of variance, significances, correlations and other statistical analysis were performed using GraphPad Prism version 5.00, (GraphPad Software, San Diego, California, USA).

3. RESULTS AND DISCUSSION

3.1 Chemical composition of the essential oil

Citrus oils are a mixture of volatile compounds and consist mainly of monoterpene hydrocarbons (Dugo, P. et al., (1998)). Citrus essential oils, on the other hand, generally contain some amount of coumarins or furanocoumarins (Dellacassa, E. et al., (1997)), flavonoids (Miyake, Y. et al., (1997)) and tocopherols (Waters, R.D. et al., (1976)) in the non-volatile fractions of citrus oils. Coumarins and furanocoumarins may have an important role in skin photosensitization. Hydrodistillation of the cold pressed oil prevents this hazard.

Nineteen compounds were identified in the GC and GC/MS analyses. The percentage composition of *Citrus limonum* essential oil is shown in Table 1. The components are listed in the order of elution from the HP-5 column. The main component was limonene (54.6 %) followed by γ -terpinene (19.1 %) and β -pinene (14.5 %). The monoterpene hydrocarbons (87.7 %) constituted the main fraction of lemon oil. This oil's composition, as reported in the literature, is similar to other volatile fractions characterized by the high content of limonene (Espina, L. et al., (2011)).

Table 1. Percentage composition of the essential oil from *C. limonum*.

Compound	RI ^a	%
α -pinene	938	3,9
β -pinene	978	14,5
myrcene	993	1,5
α -terpinene	1019	0,3
p-cymene	1024	0,1
limonene	1028	54,6
γ -terpinene	1061	19,1

186	terpinolene	1090	0,8
187	linalool	1098	0,1
188	citronellal	1154	0,1
189	terpinen-4-ol	1176	0,1
190	α -terpineol	1189	0,3
191	citronellol	1225	0,1
192	nerol	1230	0,1
193	neral	1239	1,1
194	geraniol	1252	0,1
195	linalyl acetate	1258	0,1
196	geranial	1269	2,3
197	geranyl acetate	1383	0,8

198 ^a Retention index, relative to C₉-C₂₂ n-alkanes on the HP-5 column.

199 **3.2. *In vitro* and *in vivo* free radical scavenging activity of essential oil**

200

201 The superoxide anion scavenging activity of *Citrus limonum* essential oils was evaluated
 202 using the enzymatic hypoxanthine/xanthine oxidase system. Among the concentrations
 203 tested (Fig.1), the 1:100 dilution of lemon essential oil in DMSO had an $\cdot\text{O}_2^-$ inhibition that
 204 was comparable to that of α -tocopherol. The 1:1000 dilution inhibited $\cdot\text{O}_2^-$ less than α -
 205 tocopherol but the level of inhibition was about 76% and 65% of the α -tocopherol activity on
 206 cytochrome c and tetrazolium nitroblue, respectively.

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227




Fig.1. Percentage cytochrome c inhibition and percentage nitroblue tetrazolium reduction. Test significant from normal control ($P < 0.05$). Mean \pm S.E.M of ten experiments.

The peroxidation data as evidenced by the light emission are shown in Fig 2.




Fig. 2. Chemiluminescence over time in four groups of volunteers. α -tocopherol = group A; EO-DMSO = group B ; EO-grapeseed = group C; control = group D. Test significant from normal control ($P < 0.05$). Mean \pm S.E.M of twenty experiments

The lipids from untreated volunteers showed the highest chemiluminescence and are considered to be the normal response to the peroxy radical action. Lower emissions were recorded for the lipids from volunteers treated with antioxidative substances and the lemon essential oil was more effective than α -tocopherol as an antioxidant. The grape-seed oil showed an extra little antioxidative action that was added to the action of lemon essential oil; the chemiluminescence curve is a little lower than that of the lemon essential oil dissolved in DMSO, but the data belong to the same set according to the one-way ANOVA . These

results show that these two oils had a similar scavenging action against peroxide free radicals *in vitro* and *in vivo* (Ahn, H.S. et al., (2002)).

The exposure of human skin to UV radiation can generate ROS in both the epidermis and dermis. The depth of penetration of UV radiation, as well as its damaging potential in deeper skin cells, have been demonstrated (Katiyar, S.K. et al., (2001)). Among the scavenging substances, α -tocopherol was chosen as a reference for comparing the scavenging action of lemon essential oil. The anti-oxidant activity of oil-in-water emulsion containing α -tocopherol has been reported over a wide range of conditions and test systems (Frankel, E.N. et al., (1994)). Our data demonstrate that the lemon essential oil is more active than α -tocopherol against $\text{O}_2^{\cdot -}$ and peroxide free-radical inhibition at 1:100 dilution. Lemon essential oil is used instead of other lemon extracts, to avoid the toxic action that furanocoumarins have under UV exposure.

4. CONCLUSION

The results of this study suggest that lemon essential oil has properties that could benefit human skin as it undergoes environmental and chronological ageing.

The scavenging action of lemon essential oil solubilized in grape-seed oil could have a practical application in Aesthetic Medicine (a branch of medicine focused on satisfying the aesthetic desires and goals of patients) for treating human skin against oxidative damage. Therefore, continuous application of lemon essential oil solubilized in grape-seed oil might contribute to the prevention of lifestyle-related diseases by regulating the balance of oxidative stress.

COMPETING INTERESTS

The Authors declare that no competing interests exist.

AUTHORS' CONTRIBUTION

The work presented here was carried out with the collaboration of all the authors. GB and BT defined the research theme and designed the methods and experiments, analyzed the data, interpreted the results and wrote the paper. PA was involved in the writing process of the manuscript, RV co-designed the experiments, discussed the analyses, interpretation, and presentation of data. All authors have contributed to, seen and approved the manuscript.

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

REFERENCES

- Adams, R.P. (2001). Identification of essential oil components by gas chromatography/quadrupole mass spectroscopy. Allured Publishing Corporation.
- Ahn, H.S., Jeon, T.I., Lee, J.Y., Hwang, S.G., Lim, Y., Park, D.K. (2002). Antioxidative activity of persimmon and grape seed extract: *in vitro* and *in vivo*. Nutr. Res., 1265-1273.
- Angelini, P., Pagiotti, R., Menghini, A., Vianello, B. (2006). A antimicrobial activities of various essential oils against foodborne pathogenic or spoilage moulds. Ann. Microb., 56, 65-69.

301 Angelini, P., Granetti, B., Pagiotti, R. (2008). Effect of antimicrobial activity of *Melaleuca*
 302 *alternifolia* essential oil on antagonistic potential of *Pleurotus* species against
 303 *Trichoderma harzianum* in dual culture. World J. Microb. Biotech., 24, 197-202.
 304 Angelini, P., Pagiotti, R., Venanzoni, R., Granetti, B. (2009). Antifungal and allelopathic
 305 effects of Asafoetida against *Trichoderma harzianum* and *Pleurotus* spp. Allelopathy
 306 Journal, 23, 357-368
 307 Arouma, O., Halliwell, B., Hoey, B.M., Butler, J. (1989). The antioxidant action of N-
 308 acetylcysteine: its reaction with H_2O_2 , OH^* , OH^*_2 , $HOCl$. F.R.B.M., 6, 593-597.
 309 Bech-Thomsen, N., Wulf, H.C. (1995). Carcinogenic and melanogenic effects of a filtered
 310 metal halide UVA source and a tubular fluorescent UVA tanning source with or
 311 without additional solar-simulated UV radiation in hairless mice. Photochem.
 312 Photobiol., 62, 773-779.
 313 Chiu, A., Kimball, A.B. (2003). Topical vitamins, minerals and botanical ingredients as
 314 modulators of environmental and chronological skin damage. Br. J. Dermatol., 149,
 315 681-691.
 316 Davies, N.W. (1990). Gas chromatographic retention indices of monoterpenes and
 317 sesquiterpenes on methyl silicone and Carbowax 20M phases. Chromatogr., 503, 1-
 318 24.
 319 Dellacassa, E., Lorenzo, D., Moyna, P., Verzera, A., Mondello, L., Dugo, P. (1997). Uru-
 320 guayan essential oils. Part VI. Composition of lemon oil. Flavour Fragr. J., 12: 247-
 321 255.
 322 Di Vaio, C., Graziani, G., Gaspari, A., Scaglione, G., Nocerino, S., Ritieni, A.
 323 (2010). Essential oils content and antioxidant properties of peel ethanol extract in 18
 324 lemon cultivars. Scientia Horticulturae 126 50–55.
 325 Dugo, P., Mondello, L., Cogliandro, E., Cavazza, A., Dugo, G. (1998). On the genuineness
 326 of citrus essential oils. Part LIII. Determination of the composition of the oxygen
 327 heterocyclic fraction of lemon essential oils (*Citrus limon* (L.) Burm. f.) by normal-
 328 phase high performance liquid chromatography. Flavour Fragr. J., 13: 329-334.
 329 Espina, L., Somolinos, M., Lorán, S., Conchello, P., Pagán, D.G.R. (2011) Chemical
 330 composition of commercial citrus fruit essential oils and evaluation of their
 331 antimicrobial activity acting alone or in combined processes. Food Control 22 896-
 332 902
 333 Estevez, M., Cava, R. (2006). Effectiveness of rosemary essential oil as an inhibitor of lipid
 334 and protein oxidation: contradictory effects in different types of frankfurters. Meat
 335 Sci., 72, 348-355.
 336 Frankel, E.N., Huang, S.W., Kanner, J., German, J.B. (1994). Interfacial phenomena in the
 337 evaluation of antioxidants: bulk oils vs emulsions. J. Agric. Food Chem., 42, 1054-
 338 1059.
 339 Gonzalez-Flecha, B., Llessuy, S.F., Boveris, A. (1991.) Hydroper- oxide-initiated
 340 chemiluminescence: an assay for oxidative stress in biopsies of heart, liver and
 341 muscle. Free Rad. Biol. Med., 10, 93–100.
 342 Gressier, B., Cabanis, A., Lebegue, S., Brunet, C., Dine, T., Luyckx, M., Cazin, M., Cazin,
 343 J.C. (1993). Decrease of hypochlorous acid and hydroxyl radical generated by
 344 stimulated human neutrophils: comparison in vitro of some thiol-containing drugs.
 345 Meth. Find. Exp. Clin. Pharmacol., 15, 101-105.
 346 Heller, S.R., Milne, G.W.A. (1983). EPA/NIH mass spectral data base U. S. Government
 347 Printing Office.
 348 Jennings, W.G., Shibamoto, T. (1980). *Qualitative* analysis of flavour and fragrance volatiles
 349 by glass capillary gas chromatography. Academic Press.
 350 Katiyar, S.K., Afaq, F., Perez, A., Mukhtar, H. (2001). Green tea polyphenol (-)-
 351 epigallocatechin-3-gallate treatment of human skin inhibits ultraviolet radiation-
 352 induced oxidative stress. J. Carcinog., 22, 287-294.

353 Kligman, A.M. (1969). Early *destructive* effect of sunlight on human skin. J. Am. Med.
 354 Assoc., 210, 2377-2380.
 355 Kulisic, T., Radonic, A., Milos, M. (2005). Inhibition of lard oxidation by fractions of different
 356 essential oils. Grasas y Aceites (Sevilla, Spain), 56, 284-291.
 357 McLafferty, F.W., Stauffer, D.B. (1989). The Wiley NBS registry of mass spectral data. John
 358 Wiley and Sons.
 359 Miyake, Y., Yamamoto, K. and Osawa, T. (1997). Isolation of eriocitrin (epiodictyol 7-
 360 rutinoside) from lemon fruit (*Citrus limon* Burm. f.) and its antioxidative activity. *Food*
 361 *Sci. Technol. Int. Tokyo*, 3: 84-89.
 362 Nakatsu, T., Lupo, A.T., Chinn, J.W., Kang, R.K.L. (2000). Biological activity of essential oils
 363 and their constituents. *Stud. Nat. Prod. Chem.*, 21, 571-631.
 364 Oikarinen, A., Karvonen, J., Uitto, J., Hannuksela, M. (1985). Connective tissue alterations in
 365 skin exposed to natural and therapeutic UV-radiation. *Photoderm. J.*, 2, 15-26.
 366 Pagiotti, R., Angelini, P., Rubini, A., Tirillini, B., Granetti, B., Venanzoni, R. (2011).
 367 Identification and characterisation of human pathogenic filamentous fungi and
 368 susceptibility to *Thymus schimperi* essential oil. *Mycoses*, 54(5):e364-76.
 369 Sakurai, H., Yasui, H., Yamada, Y., Nishimura, H., Shigemoto, M. (2005). Detection of
 370 reactive oxygen species in the skin of live mice and rats exposed to UVA light: a
 371 research review on chemiluminescence and trials for UVA protection. *Photochem.*
 372 *Photobiol. Sci.*, 4, 715-720.
 373 Sawamura, M., Kuriyama, T. (1988). Quantitative determination of volatile constituents in the
 374 pummelo (*Citrus grandis* Osbeck forma Tosa-buntan). *J. Agric. Food Chem.*, 36,
 375 567-569
 376 Song, H.-S., Ukeda, H., Sawamura, M. (2001). Antioxidative activities of citrus peel essential
 377 oils and their components against linoleic acid oxidation. *Food Sci. and Tech. Res.*,
 378 7, 50-56.
 379 Takahashi, Y., Inaba, N., Kuwahara, S., Kuki, W. (2003). Antioxidative effect of citrus
 380 essential oil components on human low-density lipoprotein in vitro. *Biosc. Biotech.*
 381 *Biochem.*, 67, 195-197.
 382 Teissedre, P.L., Waterhouse, A.L. (2000). Inhibition of oxidation of human low-density
 383 lipoproteins by phenolic substances in different essential oils varieties. *J. Agric.*
 384 *Food Chem.*, 48, 3801-3805.
 385 Tirillini, B., Pagiotti, R., Angelini, P., Pintore, G., Chessa, M., Menghini L. (2009). Chemical
 386 composition and fungicidal activity of the essential oil of *Laserpitium garganicum*
 387 from Italy. *Chem. Nat. Comp.*, 45, 103-105.
 388 Waters, R. D., Kesterson, J. W., Braddock, R. J. (1976). Method for determining the α -
 389 tocopherol content of citrus essential oils. *J. Food Sci.*, 41: 370-371.
 390 Watson, R.E.B., Griffiths, C.E.M. (2005). Pathogenic aspects of cutaneous photoaging. *J.*
 391 *Cosm. Dermat.*, 4, 230-236.
 392 Wright, J.R., Rumbaugh, R.C., Colby, H.D., Miles, P.R. (1979). The relationship between
 393 chemiluminescence and lipid peroxidation in rat hepatic microsomes. *Arch.*
 394 *Biochem. Biophys.*, 192, 344-351.
 395