

2 **Kaurenoic acid isolated from the root bark of**
3 ***Annona senegalensis* induces cytotoxic and**
4 **antiproliferative effects against PANC-1 and**
5 **HeLa cells**

6
7 **Theophine C. Okoye^{1*}, Peter A. Akah¹, Chukwuemeka S. Nworu¹,**
8 **Adaobi C. Ezike¹**

9
10 ¹Department of Pharmacology and Toxicology, Faculty of Pharmaceutical Sciences,
11 University of Nigeria, Nsukka 410001, Enugu State, Nigeria.

12
13
14
15 **ABSTRACT (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)**

16
Aims: Cancer is one of the leading causes of death worldwide with an estimated 6.7 million deaths and 24.6 million people living with cancer in 2002. Presently, there is a global increase in prevalence, mortality and health burden of various malignancies. World Health Organization (WHO) report projected that cancer prevalence rates could further increase by 50% to 15 million new cases in the year 2020. The bioactivity guided isolation of the bioactive constituent, and its characterization, responsible for the anticonvulsant effects of the root bark extract of *A. senegalensis* yielded kaur-16-en-19-oic acid (KA). Therefore, the aim of this study was to evaluate the anti-proliferative activity of kaurenoic acid from *A. senegalensis* on selected cancer cell lines.
Study design: The study was designed to ascertain the antiproliferative and cytotoxic effects of kaurenoic acid, a terpenoid isolated from the root bark of Nigerian *Annona senegalensis* (Annonaceae).
Place and duration of study: Department of Pharmacology and Toxicology, Faculty of Pharmaceutical Sciences, University of Nigeria, Nsukka, Nigeria, between October 2010 and June, 2012.
Methodology: Human embryonic kidney cells expressing SV40 Large T-antigen (293 T), Pancreatic tumour (PANC-1) and Henrietta Lacks' cervical (HeLa) cell lines were used in the study using standard MTT, 3-(4, 5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazoliumbromide, assay method.
Results: Kaurenoic acid (KA) exhibited cytotoxic effects against the cells with estimated IC₅₀ values of 0.93, 0.74 and 0.52 M concentrations for 293 T, HeLa and PANC-1 cells respectively. This is an indication of the possible potentials of KA in the treatment of cervical and pancreatic cancers.
Conclusions: Kaurenoic acid (KA) a terpenoid, possesses antiproliferative effect against HeLa and PANC-1 cell lines, and could be the anticancer constituent in the root bark extract of *A. senegalensis* with potentials as a lead in the chemical synthesis of standard anti cancer agents.

17 *Tel: +234 803 6684506; E-mail: theokuba@yahoo.com

18
19 **Keywords: Kaurenoic acid, *Annona senegalensis*, antiproliferative and cytotoxic**

20
* Tel.: +xx xx 265xxxxx; fax: +xx aa 462xxxxx.
E-mail address: xyz@abc.com.

21 **1. INTRODUCTION**

22

23 Cancer is one of the leading causes of death worldwide with an estimated 6.7 million deaths
24 and 24.6 million people living with cancer in 2002 [1, 2]. The disease caused about 0.6
25 million deaths in the United States in 2011 [3]. Globally, there is an obvious increase in
26 prevalence, mortality and health burden of various malignancies. World Health Organization
27 (WHO) report [4], projected that cancer prevalence rates could further increase by 50% to 15
28 million new cases in the year 2020. In Nigeria, the burden of cancer is enormous and quite
29 on the increase and it has been projected that by 2020, cancer incidence for Nigerian males
30 and females may rise to 90.7/100,000 and 100.9/100,000, respectively, with mortality rates
31 of 72.2 and 76/100,000 for males and females respectively [4]. At the early stage diagnosis
32 and detection, cancer is usually managed by surgery and radiotherapy while advanced
33 cases of cancer could be treated with chemotherapeutic agents. Although chemotherapeutic
34 agents are effective, they are often associated with serious adverse effects and drug
35 resistance [5]. Additionally, the exorbitant costs, non-affordability and unavailability of
36 effective anticancer agents in some parts of world, have paved way for herbal therapies as
37 alternatives in the management of cancer in rural communities. Herbal therapies employing
38 plants extracts and plant bioactive compounds have long been used in the treatment of
39 cancer [6, 7]. The fact that herbs and plant-derived products lack much of the toxicity in the
40 synthetic chemicals enhances their appeal fro treating cancer and for long term preventive
41 strategies [8]. Therefore, other therapeutic options that are devoid of serious adverse effects
42 at the same time cheap and affordable are obviously desired. The use of natural sources
43 can provide an opportunity for the isolation and chemical characterization of
44 phytoconstituents that could be the desired source of lead compounds for the introduction of
45 novel chemotherapeutic agents. Hence the urgent need for novel therapeutic compounds (or
46 leads) with potent anticancer effects and minimal toxicities to normal cellular system. The
47 plant *Annona senegalensis* Pers. (Annonaceae) popularly known as African custard apple or
48 wild custard apple [9], has been reported to possess cytotoxic and anticancer effects [10, 11,
49 12]. A diterpenoid compound, kaurenoic acid, isolated from the root bark of *A. senegalensis*
50 was reported to possess antimicrobial [13] and anticonvulsant [14] effects. Also from other
51 sources, kaurenoic acid has shown to possess anti-inflammatory [15], anticonvulsant [16]
52 and antimicrobial activities [17]. Among the naturally isolated phytoconstituents used in
53 cancer chemotherapy are the diterpenoid compounds known as the taxanes (paclitaxel and
54 docetaxel), vinca alkaloids (vincristine and vinblastine) and podophyllotoxins (etoposides,
55 teniposides and etopophos) [18]. Therefore, due to the reported ethnomedicinal use of *A.*
56 *senegalensis* root bark extract cancer treatment [10, 12], we serendipitously evaluated the
57 cytotoxic effects of kaurenoic acid isolated from its root bark using some selected human
58 cancer cell lines.

59

60 **2. MATERIAL AND METHODS**

61

62 **2.1 Plant material**

63 Fresh roots of *A. senegalensis* were collected from Enugu-Ezike, Enugu State, Nigeria in the
64 month of June and authenticated by Mr. A. O. Ozioko of the International Centre for
65 Ethnomedicine and Drug Development (InterCEDD), Nsukka, Enugu State, Nigeria where a
66 voucher specimen was deposited (specimen number: BDCP/INTERCEDD 64). The root-
67 bark was peeled off, cut into pieces and allowed to dry. The dried root-bark was pulverized
68 into coarse powder. The dried powdered root-bark (2.95 kg) was extracted with a mixture of
69 methanol: methylene chloride (1:1) using Soxhlet extractor to obtain the *Annona*
70 *senegalensis* root bark extract (MME). This was evaporated using a rotary evaporator at
71 reduced pressure to obtain a yield of 375 g (12.71% w/w).

72

73

74 **2.2 Isolation and purification of A2**

75 The kaurenoic acid, A2, was isolated from the separation and activity guided fractionation of
76 MME. The detailed methods of isolation and purification of A2 from the root bark extract and
77 fractions of *A. senegalensis* have been documented [13,14].

78

79 **2.3 Identification and characterization of A2**

80 The purity of A2 was assessed by analytical HPLC using a Dionex P580 HPLC system
81 coupled to a photodiode array detector (UVD340S, Dionex Softron GmbH, Germering,
82 Germany). Detection was at 235, 254, 280 and 340 nm. The separation column (125 × 4
83 mm; length × internal diameter) was prefilled with Eurospher-10 C18 (Knauer, Germany),
84 and a linear gradient of nanopure water (adjusted to pH 2 by addition of formic acid) and
85 methanol was used as eluent. The molar mass was determined by Liquid Chromatography-
86 Electrospray Ionization Mass Spectroscopy (LC-ESI-MS) using a ThermoFinnigan LCQ-
87 Deca mass spectrometer (Germany) connected to an UV detector. Complete structural
88 characterization of the pure crystals of A2 was achieved by 1D (HNMR, 13CNMR, DEPT)
89 and 2D (HHCOSY, HMQC, HMBC) NMR spectroscopy using a Bruker ARX-500 and X-ray
90 crystallography. Spot detection was done with ultra-violet (UV) light at 254 nm and spraying
91 with vanillin sulphuric reagent. The melting point of A2 was also determined using a melting
92 point apparatus (Electrothermal®, Cat. No.: IA 6304, England).

93

94

95 **2.4 Cell lines**

96 Human embryonic kidney cells expressing SV40 Large T-antigen (293 T), Pancreatic tumour
97 (PANC-1) cell line and Henrietta Lacks' cervical (HeLa) cancer cell line were propagated in
98 D-10 medium, consisting of Dulbecco's modified Eagle's medium (DMEM) with high glucose,
99 2 mM L-glutamine and supplemented with 10% heat-inactivated fetal bovine serum (FBS),
100 100 U/ml penicillin and 100 µg/ml streptomycin. Tissue culture medium and supplements
101 were purchased from Invitrogen (Karlsruhe, Germany). The cell cultures were maintained in
102 a humidified 5% CO₂ atmosphere at 37°C.

103

104

104 **2.5 Cytotoxicity studies**

105 The cytotoxicity assay was performed in parallel to the antiviral screening using the MTT,
106 [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazoliumbromide], assay method as previously
107 described [19] on 293T, PANC-1, and HeLa cell lines. In the MTT assay, cells were seeded
108 onto a 96-well plate at a concentration of 10⁴ cells/well and a volume of 100 µl per well.
109 Different concentrations of the test extracts (31.5-1000 µg/ml) were added to culture wells in
110 triplicate. Culture medium without any drug was used as the "no-drug" control. After
111 incubation at 37°C under 5% CO₂ for 2 days, a solution of MTT (3 mg/ml, 50µl per well) was
112 added to each well and further incubated at 37°C + 5% CO₂ for 4 h to allow formazan
113 formation. Subsequently, the medium was removed and 150 µl of DMSO was used to
114 dissolve the resulting blue formazan crystals in living cells. The optical density was
115 determined at 550 nm using a multi-well microtitre plate reader (Tecan, Austria). Each single
116 value of the triplicates was expressed as percent of the mean of triplicates of the "no-drug"
117 control cultures and the mean and standard deviation of the percent values were calculated
118 for each triplicate. The concentration of 50% cellular toxicity (TC₅₀) of the test extracts was
119 calculated by non-linear regression.

120

121

121 **2.6 Statistical analysis**

122 Data were analyzed using One Way Analysis of Variance (ANOVA, SPSS Version 16) and
123 expressed as mean ± SEM and multiple comparisons was done using Dunnet test as *post*
124 *hoc*. Differences between means were regarded significant at *P* < 0.05.

125

126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177

3. RESULTS AND DISCUSSION

3.1 RESULTS

3.1.1 Identification and characterization of A2

The A2 was isolated as a white crystalline compound. The melting point range was estimated to be at 170-172 °C and in methanol it exhibited a UV maximum at 214 nm, which is typical of an unconjugated compound. It had a strong peaks at 303.2 (M + H), 650.2 (2M + 2Na) in the positive mode of LC-ESIMS and a corresponding peak at 301.6 (M - H) in the negative mode, which are consistent with the molar mass of 302. Based on this, and the analysis of ¹H and ¹³C NMR, the molecular formula of AS2 was deduced as C₂₀H₃₀O₂. The analyses of the HNMR, HHCOSY, C-13 NMR, DEPT, HMQC and HMBC (Table 1) and comparison of data with literature reports [20, 21] established the structure of A2 to be kaur-16-en-19-oic acid (Fig. 5). The absolute configuration as shown was based on the observed HNMR coupling constants, HMBC and X-ray crystallography and comparison with literature report [22, 13, 14].


3.1.2 Cytotoxicity studies

The A2 exhibited cytotoxicity effects against the three cell lines tested (Figs. 1, 2 and 3), with estimated IC₅₀ values of 281.85, 223.65, 156.35 µg/ml for 293 T, HeLa and PANC-1 cells respectively (Fig. 4). In molar concentrations, the IC₅₀ values of kaurenoic acid against the 293 T, HeLa and PANC-1 cells were estimated to be 0.93, 0.74 and 0.52 M respectively. Hence the antiproliferative activity of A2 against the cells in the order of increasing activity was; 293 T > HeLa cells > PANC-1 cells.

3.2 DISCUSSION

Kaurenoic acid, A2, exhibited cytotoxic activity against the 293 T, HeLa and PANC-1 cancer cell lines treated. It exhibited better cytotoxic and antiproliferative activity against HeLa cells than the PANC-1 cells, an indication of the anticancer effects of this diterpenoid on cervical and pancreatic cancers. On a separate reported work, kaurenoic acid has been identified to possess cytotoxic effects against HeLa cell lines [23]. However, the most potent cytotoxic effect on the PANC-1 cells showed the possible antitumor effect of A2 against cancer of the pancreas. Therefore, A2 could serve as a lead compound in the development of novel anticancer agents. Anticancer effect of kaurenoic acid on breast, leukemia and colon cancer cells has also been documented [24, 25, 26]. In addition to kaurenoic acid, other naturally isolated terpenoid compounds have shown to possess anticancer effects [27]. Similarly, betulinic acid, a triterpenoid, isolated from plants of Cactacea family has shown to possess potent anticancer effect against HeLa cells [28]. Documented studies revealed that terpenoids exhibited antitumor activities by inducing apoptosis in various cancer cells by activating various pro-apoptotic signaling cascades and by the inhibition of metastatic progression and tumor-induced angiogenesis [26]. The molecular mechanisms involved in these activities include the inhibition of various oncogenic and anti-apoptotic signaling pathways and suppression of nuclear translocation of various transcription factors including nuclear factor kappa B (NF-κB) [26]. The anti-inflammatory effects of kaurenoic acid isolated from different sources have variously been reported [29,15]. Kaurenoic acid was shown to significantly inhibit inflammatory mediators in lipopolysaccharide-induced RAW264.7 macrophages, inhibit the production of nitric oxide and reduced the secretion of


178 prostaglandin E (PGE₂), which are all potent mediators of inflammation [29]. There is the
179 possibility that the mechanism of antiproliferative activity of KA might be likely through
180 modulation of inflammatory mediators. In other reported studies, the mechanism of
181 anticancer effects of kaurenoic acid has been attributed to its inhibition of DNA
182 topoisomerases I and II [24] as well as the stimulation of p53 tumor suppressor gene [25].
183 However, the likely specific mechanism of action of kaurenoic acid against these cancer
184 cells at this stage of the work is not yet elucidated. Hence further work is encouraged **to**
185 **determine the possible** anti-cancer mechanism of kaurenoic acid.
186
187
188
189
190



191
192
193
194
195
196
197
198
199
200
201
202

Fig. 1: The effect of kaurenoic acid on the viability of 293-T cells

203
204
205
206




207
208
209
210

Fig. 2: Antiproliferative effect of kaurenoic acid on pancreatic cancer (PANC-1) cell line

211
212
213
214
215
216
217
218
219
220
221
222
223
224

225
226
227
228
229




230
231
232

Fig. 3: Antiproliferative effect of kaurenoic acid on cervical cancer (HeLa) cell line

233
234
235
236
237
238
239
240
241
242
243
244
245
246
247


248
249
250
251



252
253
254
255

Fig. 4: A comparison of the effects of kaurenoic acid on the viability of 29T cells, PANC-1 cells, and HeLA cells

256
257
258



259
260
261
262

Fig. 5: Chemical structure of Kaur-16-en-19-oic acid

263
264
265

TABLE 1: NMR spectroscopic data of A2^a

Position	δ H	δ c	HMBC
1	Ha 0.82 m Hb 1.54 m	41.48	Ha C-20 Hb C-2, C-3
2	1.42 m (2H)	19.29	1, 3, 4, 10
3	Ha 1.01 m Hb 2.17 m	37.97	1,2,4,5,19
4		43.95	
5	1.07 m	57.26	4, 6, 7, 10, 20
6	Ha 1.62 m Hb 1.84 m	22.03	4, 5, 7, 8, 10
7	Ha 1.44 m Hb 1.54 m	40.90	5, 6, 6, 8, 9, 15
8		44.05	
9	1.05 m	55.29	5,, 8, 10, 11, 14, 15, 20
10		39.85	
11	Ha 1.60 m Hb 1.88 m	18.64	9, 10, 12, 13
12	Ha 1.46 m Hb 1.62 m	33.97	11, 13, 16, 14
13	2.61 bs	43.95	C-8, C-12, C-14, C-16, C-17
14	Ha 1.16 Hb 2.02	39.9	
15	2.06	48.48	C-8, C-9, C-16, C-17
16		156.12	
17	Ha 4.72 s Hb 4.78 s	103.21	
18	1.22 s (3H)	29.18	C-2, C-3, C-4, C-5, C-19
19		184.94	
20	0.93 s (3H)	15.79	C-1, C-5, C-9, C-10

266 ^aSpectra were measured in CDCl₃ at 500 (¹H) and 150 (¹³C) MHz. Assignments were made
267 on the basis of DEPT, ¹H-¹H COSY, HMQC and HMBC experiments. The detailed NMR data
268 are available with the author for correspondence and will be readily supplied on request.
269

270
271

272 **4. CONCLUSION**

273

274 Kaurenoic acid, a diterpenoid, isolated from the root bark extract of *A. senegalensis* Pers.
275 (Annonaceae) possesses cytotoxicity and anticancer effect against PANC-1 and HeLa
276 human cell lines while the specific mechanism of anticancer activity is a point for further
277 research.

278
279

280 **5. ACKNOWLEDGEMENTS**

281

282 The authors thankfully appreciate Prof. Dr. Mark Hamann of the Department of
283 Pharmacognosy and Phytochemistry, University of Mississippi, USA, for performing the
284 proton NMR and the X-ray crystallography of A2. We appreciate Dr. FBC Okoye for doing

* Tel.: +xx xx 265xxxxx; fax: +xx aa 462xxxxx.
E-mail address: xyz@abc.com.

285 part of the NMR spectral data analysis during his stay in Germany. Part of the financial
286 support was from the Ministry of Education, Science and Technology Post Basic
287 Programme, of the Federal Government of Nigeria for the Innovators of Tomorrow (IOT)
288 Award Grant of the World Bank assisted Step-B project.

289
290
291

292 **COMPETING INTERESTS**

293
294

The authors declare that there are no conflicts of interest.

295
296

297 **AUTHORS' CONTRIBUTIONS**

298
299
300
301
302
303
304

Okoye TC and Akah PA designed, performed the experiment on the isolation of the
compound and prepare the manuscript. Nworu CS performed the cytotoxicity studies while
Ezike AC contributed in the manuscript preparation. All authors read and approve the final
manuscript.

305
306

307 **REFERENCES**

308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335

1. Parkin DM, Bray F, Ferlay J, Pisani P, Global cancer statistics, 2002, CA Cancer J Clin.
2005; 55:74-108.

2. Ji D, Ye J, Jiang Y, Qian B, Anti-tumor effect of Liqi, a traditional Chinese medicine
prescription, in tumor bearing mice, BMC Complementary and Alternative Medicine. 2009;9.
doi:10.1186/1472-6882-9-20.

3. Siegel R, Ward E, Brawley O, Jemal A, The impact of eliminating socioeconomic and
racial disparities on premature cancer deaths, CA Cancer J. Clin. 2011; 61:212-236.

4. World Health Organisation, Global cancer rates could increase by 50% to 15 million by
2020, 2003.

5. Kellof GJ, Perspective on cancer chemoprevention research and drug development, Adv.
Cancer Res. 2000; 78:199-334.

6. Hartwell JL, Plants used against cancer: a survey. Lawrence, M.A. Quarterman
Publications, 1982, pp 438-439.

7. Shoeb M, Anticancer agents from medicinal plants. Bangladesh J Pharmacol. 2000; 1:35-
41.

8. Harlev E, Nevo E, Solowey E, Bishayee A, Cancer preventive and curative attributes of
plants of the cactaceae family: a review. Planta Med. 2013; 79:713-722.

9. Okoye TC, Akah PA, Ezike AC, Okoye MO, Onyeto CA, Ndukwu F et al. Evaluation of the
acute and sub acute toxicity of *Annona senegalensis* root bark extracts. Asian Pacific J. of
Tropical Medicine. 2012; 5(4):277-282.

- 336 10. Sahpaz S, Bories CH, Loiseau PM, Cartes D, Hocquemiller R, Laurens A, Cave A,
337 Cytotoxic and antiparasitic activity from *Annona senegalensis* seeds. *Planta Medica*. 1994;
338 60:538-540.
339
- 340 11. Sahpaz S, Carmen Gonzalez M, Hocquemiller R, Zafra-Polo MC, Diego C,
341 Annosenegalin and annogalene: two cytotoxic monotetrahydrofuran acetogenins from
342 *Annona senegalensis* and *Annona cherimolia*. *Phytochemistry*. 1996; 42:103-107.
343
- 344 12. Abubakar AM, Musa AM, Ahmed A, Hussaini IM. The perception and practice of
345 traditional medicine in the treatment of cancers and inflammations by the Hausa and Fulani
346 tribes of northern Nigeria. *J Ethnopharmacol*. 2007; 111:625–629.
347
- 348 13. Okoye TC, Akah PA, Okoli CO, Ezike AC, Omeje EO, Odoh UE. Antimicrobial effects of
349 a lipophilic fraction and kaurenoic acid isolated from the root bark extracts of *Annona*
350 *senegalensis*. *Evidence-Based Comp. and Alter. Med*. 2012; doi:10.1155/2012/831327.
351
- 352 14. Okoye TC, Akah PA, Omeje EO, Okoye FBC, Nworu CS. Anticonvulsant effect of
353 kaurenoic acid isolated from the root bark of *Annona senegalensis*. *Pharmacology*
354 *Biochemistry and Behavior*. 2013; 109:38-43. Doi: 10.1016/j.pbb.2013.05.001.
355
- 356 15. Sosa-Sequera MC, Suarez O, Dalo DL. Kaurenoic acid: An *in vivo* experimental study of
357 its anti-inflammatory and antipyretic effects. *Indian J Pharmacol*. 2010; 42(5):293-296.
358
- 359 16. Dalo NL, Sosa-Sequera MC, Usubillaga A. On the anticonvulsant activity of kaurenoic
360 acid. *Invest Clin*. 2007; 48:349-358.
361
- 362 17. Davino SC, Giesbrecht AM, Roque NF. Antimicrobial activity of kaurenoic acid
363 derivatives substituted on carbon-15, *Braz J Med Biol Res*. 1989; 22:1127-9.
364
- 365 18. Alexandrova R, Alexandrov I, Velcheva M, Varadinova T. Phytoproducts and cancer,
366 *Experimental pathology and parasitology*. 2000; 4:15-26.
367
- 368 19. Romijn JC, Verkoelen CF, Schroeder FH, Application of the MTT assay to human
369 prostate cancer cell lines in vitro: Establishment of test conditions and assessment of
370 hormones-stimulated growth and drug-induced cytostatic and cytotoxic effects. *The Prostate*.
371 1988; 12(1): 99-110.
372
- 373 20. Pacheco AG, de Oliveira PM, Piló-Veloso D, Alcântara AFC, ¹³C-NMR Data of
374 Diterpenes Isolated from *Aristolochia* Species. *Molecules*. 2009; 14:1245-1262.
375
- 376 21. Lee I, Kim HJ, Youn UJ, Min BS, Jung HJ, Yoo JK et al. Absolute Configuration of a
377 Diterpene with an Acyclic 1,2-Diol Moiety and Cytotoxicity of Its Analogues from the Aerial
378 Parts of *Aralia cordata*. *Bull. Korean Chem. Soc*. 2008; 29:1839-1842.
379
- 380 22. Bruno-Colmenarez J, De Delgado GD, Peña A, Alarcon L, Usubillaga A, Delgado-
381 Méndez P. Structure of *ent*-15-hydroxy-kaur-16-en-19-oic acid. *Avances en Química*. 2011;
382 6:16-20.
383
- 384 23. Cuca LE, Coy ED, Alarcon MA, Fernandez A, Aristizabal FA. Cytotoxic effects of some
385 natural compounds isolated from Lauraceae plants and synthetic derivatives. *Biomedica*,
386 2011; 31(3):335-43.doi: 10.1590/S0120-41572011000300006.
387

- 388 24. Costa-Lotufo LV, Cunha GMA, Farias PAM, Viana GSB, Cunha KMA, Pessoa C et al.
389 The cytotoxicity and embryotoxic effects of kaurenoic acid, a diterpene isolated from
390 *Copaifera langsdorffii* Oleo-resin. *Toxicol.* 2002; 40(8): 1231-1234. doi.org/10.1016/S0041-
391 0101(02)00128-9.
392
- 393 25. Seo CS, Li G, Kim CH, Lee CS, Jahng Y, Chang HW, Son JK. Cytotoxic and DNA
394 topoisomerase I and II inhibitory constituents from the roots of *Aralia cordata*. *Arch. Pharm.*
395 *Res.* 2007; 30:1404-1409.
396
- 397 26. Peria FM, Tiezzi DG, Tirapelli DP, Neto FS, Tirapelli CR, Ambrosio S et al. *J. Clin. Oncol*
398 28(Supp: abstract e13641), 2010, ASCO Annual Meeting.
399
- 400 27. Kuttan G, Pratheeshkumar P, Manu KM, Kuttan R. Inhibition of tumor progression by
401 naturally occurring terpenoids. *Pharmaceutical Biology.* 2011; 49(10):995-1007.
402 Doi:10.3109/13880209.2011.559476.
403
- 404 28. Kinoshita K, Yang Y, Koyama K, Takahashi K, Nishino H. Inhibitory effect of some
405 triterpenes from cacti on Pi-incorporation into phospholipids of HeLa cells promoted by 12-O-
406 tetradecanoylphorbol-13-acetate. *Phytomedicine.* 1999;6:73–77.
407
- 408 29. Choi RJ, Shin EM, Jung HA, Choi JS, Kim YS. Inhibitory effects of kaurenoic acid from
409 *Aralia continentalis* on LPS-induced inflammatory response in RAW264.7 macrophages.
410 *Phytomedicine.* 2011; doi:10.1016/j.phymed.2010.11.010.
411
412
413
414
415
416
417