

1 **Selective Hematological and Biochemical Parameters in**
2 **Egyptian catfish (*Clarias lazera*) Exposed to Mercuric Chloride**
3 **with particular reference to Erythrocyte Morphological Anomalies.**
4 Author details: Wafaa Abdou Mohamed Mohamed*

5
6 Department of Clinical Pathology, Faculty of Vet. Medicine, Zagazig University,
7 El-Zeraa Street, Sharkia Province, Egypt

8
9 E-mail: waffa.clinical@yahoo.com
 waffa.clinical@yahoo.com

10
11
12
13 **ABSTRACT**

14
15 The study was undertaken to establish a sub-lethal concentrations (0.0002, 0.002,
16 0.02mg/L) toxicity of mercuric chloride in Egyptian catfish (*Clarias lazera*) after long-
17 term (14 days) exposure which achieved by investigation of some selective hematological
18 and biochemical parameters with respect to erythrocyte morphological alterations.
19 Experimental fish were set up with three groups for each concentration, plus the control
20 group. Blood samples were collected from five individuals for each concentration after
21 14 days of exposure. The hematological parameters analyzed were: total red blood cell
22 count (RBC), hemoglobin concentration (Hb), packed cell volume (PCV), mean
23 corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular
24 hemoglobin concentration (MCHC), total platelets count, total white blood cell count
25 (WBC) and differential leukocyte counts (neutrophil, eosinophil, lymphocyte, monocyte
26 and basophils). Fish exposed to 0.02 mg/L Hg for a long time showed a significant
27 decrease in RBC count, Hb, MCHC and platelets number than control groups ($P < 0.05$)
28 with moderate elevations in PCV, MCV, WBC, neutrophil, eosinophil, lymphocyte count
29 , glucose and cortisol level compared to the control groups ($P < 0.05$). However, fish
30 exposed to the other two low concentrations showed no toxic effects. Hg at the
31 concentrations studied was toxic to Egyptian catfish (*Clarias lazera*) at the higher dose
32 only (0.02 mg/L) after long term exposure.

33
34 **Keywords:** Egyptian catfish (*Clarias lazera*), Mercuric Chloride, Sublethal toxicity,
35 Erythrocyte anomalies, Cortisol, Glucose

36
37 **1. INTRODUCTION**

38
39 Mercuric chloride (HgCl_2) is one of the environmental pollutants that are currently
40 used as a catalyst or reagent in various chemical reactions, and to a lesser extent as a
41 disinfectant or pesticide. It is also a developmental toxicant in experimental animals
42 following both oral and inhalation exposure [1]. Mercury exists in nature primarily as
43 elemental mercury or as a sulfide and is found in the earth's crust at approximately 0.5
44 parts per million. Atmospheric exposures occur from outgassing from rock or through
45 volcanic activity. Human sources of atmospheric mercury include coal burning [2] and
46 mining (mercury and gold in particular). Atmospheric elemental mercury settles in the
47 water, where it is converted by microorganisms into organic (methyl or ethyl) mercury,
48 which is ingested by smaller creatures which are eventually consumed by larger fish. Fish
49 at the top of the food chain (e.g., tuna, swordfish, or shark) may concentrate considerable

mercury in their tissues [3]. Environmental pollutants induced morphological and quantitative variations in blood parameters[4,5] for this reason, assessment of cellular and plasma component of blood is very important in the detection of pathological changes in different stress conditions such as exposure to heavy metals[6]. Changes in the hematological profile of the fish exposed to mercury have been observed in *Hoplias malabaricus*[7], *Tilapia mossambica*[8], *Oreochromis aureus* [9], *Acipenser baeri* [10], *Ctenopharyngodon idella* [11], *Pleuronectes platessa* [12], *Channa punctatus* [13], *Aphanius dispar* [14]. In the present study we study the hematological and biochemical abnormalities have been found in Egyptian catfish (*Clarias lazera*) following long-term exposure to sublethal concentrations of mercuric chloride.

2. MATERIALS AND METHODS

2.1. Preparation of mercuric chloride

Mercuric chloride (HgCl_2 [99.5%] purity) was obtained from Sigma-Aldrich (St. Louis, MO, USA). A stock solution of 10 mg/L mercury was made by dissolving the salt in double-distilled water. This solution was diluted directly into aquaria in sufficient amounts to provide the following mercury concentrations in water: 0.02, 0.002 and 0.0002mg/L as LC50/10, LC50/100 and LC50/1000 [15]

2.2. Animals and Experimental design

A total of 80 Egyptian catfish (*Clarias lazera*) with average body weight (200- 220 gm) provided by the fisheries research laboratory. Fish were adapted to laboratory conditions for 7 days in a 400 L tank with dechlorinated tap water. During adaptation, all fish were fed with trading pellet twice a day and were divided into 4 groups, each containing 20 fish. PH, dissolved oxygen, temperature and conductivity were monitored during the experiment.

Group I: this group was kept as a control in which fish were conserved in a fiberglass tank without accessing toxicant.

Group II: fish in this group were exposed to mercuric chloride at a concentration of 0.0002 mg/L for 14 days according to [16]

Group III: fish in this group were exposed to mercuric chloride at a concentration of 0.002 mg/L for 14 days.

Group IV: fish in this group were exposed to mercuric chloride at a concentration of 0.02 mg/L for 14 days.

2.3. Sampling

Fish were without delay anesthetized with 200 ppm clove powder. Blood samples rapidly were taken from tail blood vessel by heparinized syringes. Blood indices were carried out on fresh blood. Another blood sample was collected without anticoagulant for serum separation and biochemical analysis.

2.4. Hematological Examination

2.4.1. Erythrogram: Total red blood cell count was carried out by the method of [17] using 1:30 dilution of diluting heparinized blood with Giemsa stain. Cells were counted using a haemocytometer Neubauer below the light microscope. Packed cell volume (PCV) was measured by putting new blood in glass capillary tubes and centrifuged in a microhematocrit centrifuge at 10000 RPM for 5 min (Hettich, Germany), then give the reading using hematocrite readerships. Hemoglobin concentration was measured by the

cyanomethemoglobin method according to [18]. Erythrocytes Indices (MCH or Mean Corpuscular Hemoglobin, MCHC or Mean Cell Hemoglobin Concentration and MCV or Mean Corpuscular Volume) were computed from RBC, PCV and Hb [18].

2.4.2. Erythrocyte anomalies determination: In all experimental groups blood smears were made, stained with May Grunwald and Giemsa solutions and fixed with Canada balsam and cover glass and examined under microscope [19].

2.4.3. Leukogram: Total leukocyte count was carried by the method of [17] using haemocytometer Neubauer. Differential leukocyte count was carried out by the method of [20] by made peripheral blood smears stained with Giemsa. In addition, the calculation of the absolute number of neutrophils lymphocytes, eosinophil and monocyte was occurred.

2.5. Biochemical Analysis:

Serum glucose was assessed according to [21]. A microtitre plate ELISA for measuring cortisol in fish according to [22].

2.6. Statistical Analysis

Results obtained were analyzed using student F-test to compare the means. Analysis was performed using computer database software from the statistical package for social sciences (version 16.0 SPSS). A P-value of < 0.05 was considered statistically significant in all clinical comparisons at 95% confidence interval.

3. RESULTS AND DISCUSSION

Heavy metals are persistent contaminants in the environment that come to the forefront of dangerous substances such as cadmium, lead, mercury, copper and zinc which causing a serious health hazard in humans and animals [23]. The toxicity data in this work showed that mercuric chloride is toxic to Egyptian cat fish. In this report, a series of hematological and biochemical parameters were examined for studying the toxicity induced by long-term exposure to sublethal concentrations of mercuric chloride in Egyptian catfish (*Clarias lazera*). Mercuric chloride at a dose of 0.02 mg/L in group (4) showed red skin erosions (Fig.1).



Fig.1. Clinical signs of 0.02 mg/L of mercuric chloride, showing red skin erosions.

Regarding to the erythrogram as in Fig.2 like RBC, Hb, PCV and other hematological indices like MCV, MCH and MCHC, we found a significant decrease in the total red blood cells count and Hb concentration following 2 weeks of exposure to 0.02 mg/L of Hg comparing with the control ($p<0.05$) which reflect an anemic situation that may be due to inhibition of RBC manufacture by the mercuric chloride or may be due to the decreased rate of production of red blood cells or an increased loss of these cells. [24] have attributed anemia to impaired erythropoiesis due to the direct effect of metal on hematopoietic centers (kidney/spleen), accelerated erythroclasia due to altered membrane permeability and/or increased mechanical fragility and defective Fe metabolism or impaired intestinal uptake of Fe due to mucosal lesions. Our results are agreed with the result of [16,25,26,27,28]




Fig.2. Changes in RBCs count (x10⁶cells/mm³), Hb concentration (gm/dl) and PCV (%) in all experimental groups.

Morphological classification of anemia was diagnosed by calculation of blood indices representative in MCV, MCH and MCHC as in Fig.3. In the present work we found a significant increase in the values of PCV and MCV following the exposure to mercuric chloride at a concentration of 0.02 mg/L which may result from an expansion of unripe RBC [29]. However, a significant decrease of MCHC was noticed with indication to macrocytic hypochromic anemia which induced by hemolysis and damaged osmoregulation which are resulted from gill injury caused by toxicants which also lead to a diminution in RBC counts [30,31]. On the other hand, no significant changes were noticed in the above mentioned parameters after 2 weeks from exposure to 0.002 and 0.0002mg/L concentrations of mercuric chloride reflecting no toxicity induced by these both doses in Egyptian catfish (*Clarias lazera*), these results are parallel to the result proved by [16]




Fig.3. Changes in MCV, MCH and MCHC values in all experimental groups.

As shown in Fig.4 regarding to the result of total platelets count, we found a moderate decrease in their number in group 4, this result agreed with [32, 33]. On the contrary, groups 2 and 3 showed no significant deviation in platelets number than the normal level and this agreed with the result of [16]




Fig.4. Changes in platelets number ($\times 10^5 \text{ cells/mm}^3$) in all experimental groups.

Erythrocyte morphological alterations and anomalies in Fig.5 showing fragmented, crenated, enlarged RBC with giant nucleus, star and semi-lunar shaped RBC with vacuolated cytoplasm, also elliptical RBC with hypochromasia in addition to micronucleus were seen in group 2 treated with mercuric chloride at a concentration of 0.02 mg/L. Our findings were parallel to the result of Gill and Pant who found erythrocyte anomalies in fish subjected to intoxication with heavy metal, these anomalies represented by erythrocyte swelling, poikilocytosis, vacuolation and deterioration of the cell membrane.




Fig.5. Blood smear for morphological anomalies.[A] showing normal RBC morphology of the control group , while [B] showing fragmented, crenated and enlarged RBC with giant nucleus , [C] showing star shape RBC,[D] showing semi-lunar shape RBC with vacuolated cytoplasm ,[E] showing elliptical RBC with hypochromasia and [F] showing presence of micronucleus indicating toxicity in fish group treated with 0.02 mg/L mercuric chloride.

Regarding to the leukogram results as in Figs.6,7 and 8 we found an increase in the total number of leukocytes ,lymphocyte , monocyte, neutrophil and eosinophil which may be attributed to the stimulation of immune system in response to tissue damage caused by Mercuric chloride [27,31,34] .In addition to the elevation in the number of lymphocytes may be due to the toxicant effects on lymphomyeloid tissue as a protection mechanism may also redound to expansion in WBC count in fish [35] or may be due to the stimulation of the immune system caused an increase in lymphocytes by an injury or tissue damage caused by heavy metals in fish [36] .Our results are agreed with [32] who found an increase in lymphocyte, monocyte, neutrophil and eosinophil cells in *Heteropneustes fossilis*, *Channa punctatus* and *Mastacmebalus puncalus* on long-term exposure to least effective concentrations of mercuric chloride. Also our results parallel to the result of [26,33,37] who observed the effect of mercuric chloride on the differential leucocyte counts and found lymphocytosis, neutrophilia, monocytosis, eosinophilia and in *Anabas testudineus*, *Tinca tinca* and *Hoplias malabricus* respectively .




Fig.6: Differences in WBCs count ($\times 10^3$ cells/mm³) in all experimental groups.





Fig.7: Variables in lymphocyte and neutrophil numbers ($\times 10^3$ cells/mm³) in all experimental groups.

232
233



234
235

Fig.8:Changes in Eosinophil and Monocyte (x10³cells/mm³) in all experimental groups.

237

238 Some plasma chemicals may be useful tools to evaluate the health and/or stress
239 condition of the fish [38, 39, 40]. Stress has been reported to elevate plasma cortisol
240 [41,42,43] and glucose levels[44,45], many researchers considered that fishes
241 undergoing stressful situations exhibit increases of plasma cortisol and glucose levels
242 [46,47,48]. Regarding to the biochemical result as seen in Fig.9, we found a significant
243 increase in blood glucose and cortisol levels in catfish exposed to 0.02mg/L Hg for 14
244 days. These parameters showed no difference in catfish exposed to 0.002 and
245 0.0002mg/L of mercuric chloride when compared with the control one, these results are
246 parallel to the result reported by [16]. Hyperglycemia in the current study might be
247 caused by gluconeogenesis to supply energy for the increased metabolic claims instituted
248 by mercury chloride stress. A significant increase in cortisol level may be due to some
249 stress which activate the hypothalamus pituitary- internal (HPI) axis, resulting in a
250 cortisol liberates that causes secondary stress answers [16]

251
252
253
254
255




Fig.9: Changes in glucose and cortisol levels in all experimental groups.

4. CONCLUSION

The LC50/100 and LC50/1000 of mercuric chloride in the present study were insufficient to cause hematological and biochemical changes in Egyptian catfish. Whereas the LC50/10 induced moderate changes in cellular and plasma constituents of blood. Therefore, further studies of Hg toxicity would be needed to be carried out at higher concentrations or using long exposure times.

COMPETING INTERESTS

Author has declared that no competing interests exist.

5. REFERENCES

- [1]US EPA. Mercury Health Effects Update: Health Issue Assessment.1984a.
- [2] Boylan H M, Cain R D , Kingston H M. "A new Method to Assess Mercury Emissions: A Study of Three Coal-Fired Electric-Generating Power Station Configurations". J Air & Waste Manage Assoc. 2003;53: 1318-25.

- [3] Bernhoft R A. Mercury toxicity and treatment: a review of the literature. Journal of Environmental and Public Health. 2012.
- [4] Juneja CJ , Mahajan CL. Hematological and haemopoietic changes in fish *Channa punctatus* due to mercury pollution in water. Indian Journal of Animal Research.1983; 17 (2): 63-71.
- [5] Ranzani-Paiva M J T, Ishikawa C M , Campos B E S, Eiras A C. Hematological characteristics associated with parasitism in mullets, *Mugil platanus* Günther, from the estuarine region of Cananéia, São Paulo. Brasil Revta Bras Zool.1997;14(2):329-39.
- [6] Nussey G ,Van Vuren J H J , du Preez H H. Effect of copper on the haematology and osmoregulation of the Mozambique tilapia, *Oreochromis mossambicus* (Cichlidae). Comp Biochem Physiol.1995; 111C (3):369-80.
- [7]Oliveira-Ribeiro C A ,Neto F F, Mela M , Silva P H, Randi M A F , Rabitto I S, Alves Costa J R M, Pelletier E. Hematological findings in neotropical fish *Hoplias malabaricus* exposed to subchronic and dietary doses of methylmercury, inorganic lead, and tributyltin chloride. Environmental Research.2006; 101: 74-80.
- [8] Menezes M R, Qasim S Z. Effects of mercury accumulation on the electrophoretic patterns of the serum , Hemoglobin and eye lens proteins of *Tilapia mossambica* (Peters). Water Research.1984; 18 (2):153-61.
- [9] Allen P. Changes in the Hematological profile of the cichlid *Oreochromis aureus* (Steindachner) during acute inorganic mercury intoxication. Comp Biochem Physiol. 1994;108C (1):117-21.
- [10] Kikrayakov VR , Lapirova TB. Influence of salts of some heavy metals on the differential blood count in Juvenile *Acipenser baeri*. J Ichthyol.1997; 37 (6): 458-62. Mercury Toxicity and Treatment: A Review of the Literature.
- [11] Shakoori A R, Iqbal M J,Mughal A L , Ali S S . Biochemical changes induced by inorganic mercury on the blood, liver and muscles of freshwater Chinese grass carp, *Ctenopharyngodon idella*. Journal of Ecotoxicology and Environmental Monitory.1994; 4 (2): 81-92.
- [12] Fletcher T C , White A. Nephrotoxic and Hematological effects of mercuric chloride in the plaice (*Pleuronectes platessa*). Aquat Toxicol.1986; 8:77-84.
- [13] Sastry KV, Sharma K. Mercury induced Hematological and biochemical anomalies in *Ophiocephalus* (*Channa*) *punctatus*. Toxicology Letters.1980; 5 (3-4): 245-9.
- [14] Hilmy A M, Shabana M B, Said M M. Hematological responses to mercury toxicity in the marine teleost, *Aphanius dispar* (RÜPP). Comp Bioch Physiol. 1980;67C: 147-58.
- [15] Ishikawa N M. Toxicidade aguda e crônica do mercúrio em tilápia "Tailandesa", *Oreochromis niloticus*. Determinação da CL50-96h e alterações hematológicas. Master's thesis. Centro de Aquicultura da Universidade Estadual Paulista, Jaboticabal, São Paulo, Brazil. 2003.

- [16] Ishikawa I NM , Ranzani-Paiva M J T , Lombardi J V , Ferreira C M. Hematological parameters in Nile Tilapia, *Oreochromis niloticus* exposed to sub-lethal concentrations of mercury. *Braz arch biol technol.*2007;50 (4).
- [17] Stevens M L. *Fundamentals of Clinical Hematology*, WB Saunders, Philadelphia, PA. 1997.
- [18] Lee R G, Foerster J, Jukens J , Paraskevas F, Greer J P, Rodgers G M. 'Wintrobe's—Clinical Hematology,' 10th ed. Lippincott Williams & Wilkins, New York, USA. 1998.
- [19] Witeska M, Kondera E , Szczypińska K. The effect of cadmium on Common Carp erythrocyte morphology. *Polish J of Environ Stud.*2010; 20(3):783-8.
- [20] Beutler E, Lichtman M A, Coller B S, Seligsohn U. 'Hematology,' sixth ed. McGraw-Hill, USA. 2001.
- [21] Trinder P. *Ann CUn Brioché.*1960; 6: 24.
- [22] Barry TP, Lapp AF, Kayes TB , Malison JA. Validation of a microtitre plate ELISA for measuring cortisol in fish and comparison of stress responses of rainbow trout (*Oncorhynchus mykiss*) and lake trout (*Salvelinus namaycush*). *Aquaculture.*1993; 117: 351–363.
- [23] Abbas H H , Zaghloul K H , Mousa M A . Effect of some heavy metal pollutants on some biochemical and histopathological changes in Blue tilapia, *Oreochromis aureus*. *Egypt J Agric Res.* 2002;80(3): 1395-411.
- [24] Gill T S , Eppele A . Stress-related changes in the Hematological profile of the American eel, *Anguilla rostrata*. *Ecotoxicol Environ Saf.*1993; 25: 227-35.
- [25] Panigrahi A K , Misra B N . Toxicological effects of mercury on a freshwater fish, *Anabas scandens*, CUV. & VAL. and their ecological implications. *Environ Pollut.*1978; 16: 31-9.
- [26] Shah S L , Altindag A . Hematological parameters of tench (*Tinca tinca* L) after acute and chronic exposure to lethal and sublethal mercury treatments. *Bull Environ Contam Toxicol.*2004a; 73: 911-8.
- [27] Saroch J D , Nisar H , Shrivastav R , Qureshi T A , Manohar S. Hematological studies of mercuric chloride affected freshwater Catfish *Clarias gariepinus* fed with *Spirulina* . *Journal of Chemical, Biological and Physical Sciences.*2012; 2(4): 1862-9.
- [28] Hedayati A , Ghaffari Z . Effect of mercuric chloride on some Hematological, biochemical parameters in Silver Carp (*Hypophthalmichthys Molitrix*). *International Journal of Veterinary Medicine: Research & Reports.*2013; 1-11.
- [29] Carvalho C S, Fernandes M N. Effect of temperature on copper toxicity and Hematological responses in the neotropical fish, *Prochilodus Scrofa* at low and high pH. *Aquaculture.*2006; 251:109-17.

- [30] Kavitha C , Malarvizhi A , Kumaran S S, Ramesh M. Toxicological effects of arsenate exposure on Hematological, biochemical and liver transaminases activity in an indian major Carp, Catla Catla. Food and Chemical Toxicology.2010; 48:2848-54.
- [31] Saravanan M , Karthika S , Malarvizhi A, Ramesh M. Ecotoxicological impacts of clofibric acid and diclofenac in Common Carp (Cyprinus Carpio) Fingerlings: Hematological, biochemical, Ionoregulatory and Enzymological Responses . Journal of Hazardous Materials.2011; 195:188-94.
- [32] Dhaneka S , Srivastava S . Studies on toxic effects of least effective concentration of mercury in fish: AHematological study.Matsya.1985; 11:75-8.
- [33] Kumar K , Priyambada P , Pandey A K . Hematological and biochemical responses of the climbing perch, Anabas testudineus (Bloch), exposed to mercury toxicity. J Ecophysio Occupl Hlth. 2004;4(1&2): 97-107.
- [34] Witeska M. The Effect of toxic chemicals of blood cell morphology in fish. Fresenius Environ Bull.2004; 13 (12):1379-84.
- [35] Ates B , Orun I , Talas Z S , Durmaz G, Yilmaz I. Effects of sodium selenite on some biochemical and Hematological parameters of Rainbow Trout (Oncorhynchus Mykiss Walbaum, 1792) Exposed to Pb²⁺ and Cu²⁺. Fish Physiology and Biochemistry.2008; 34:53-9.
- [36] Gill T S , Pant J C . Erythrocytic and leukocytic responses to cadmium poisoning in fresh water fish. Puntius conchonus Ham Environ Res.1985; 30:372-3.
- [37] Oliveira Ribeiro C A , Filipak Neto F .Hematological findings in neotropical fish Hopilas malabaricus exposed to subchronic and dietary doses of methyl mercury, inorganic lead, and tributyltin chloride. Environmental Research.2006; 101: 74-80.
- [38] Sadler J , Wells R M G, Pankhurst P M , Pankhurst N W. Blood oxygen transport, rheology and Hematological responses to confinement stress in diploid and triploid Atlantic salmon, Salmo salar Aquaculture.2000a ;184: 349-61.
- [39] Campbell T W. Blood biochemistry in lower vertebrates. In: 55th Annual Meeting of the American College of Veterinary Pathologists (ACVP) & 39th Annual Meeting of the American Society of Clinical Pathology (ASVCP), ACVP and ASVCP (ed). 2004.
- [40] Wagner T, Congleton J L. Blood chemistry correlates of nutritional condition, tissue damage, and stress in migrating juvenile chinook salmon (Oncorhynchus tshawytscha). Canadian Journal of Fisheries and Aquatic Sciences.2004; 61: 1066–74.
- [41] Wendelaar-Bonga S E. The stress response in fish. Physiological Reviews.1997; 77: 591-625.
- [42] Pottinger T G , Rand-Weaver M, Sumpter J P. Overwinter fasting and re-feeding in rainbow trout: plasma growth hormone and cortisol levels in relation to energy mobilization. Comparative Biochemistry and Physiology Part B.2003; 136: 403-17.

- [43] Haukenes A H, Barton B A , Bollig H. Cortisol responses of pallid sturgeon and yellow perch following challenge with lipopolysaccharide. *Journal of Fish Biology*.2008;72, 780–4.
- [44] Silbergeld E K. Blood glucose: A sensitive indicator of environmental stress in fish. *Bulletin of Environmental Contamination and Toxicology*.1974;11: 20-5.
- [45] David M, Shivakumar R, Mushigeri S B , Kuri R C. Blood glucose and glycogen levels as indicators of stress in the freshwater fish, *Labeo rohita* under fenvalerate intoxication. *Journal of Ecotoxicology & Environmental Monitoring*.2005;15: 1-5.
- [46] Hattingh J. Effect of tricaine methane sulphonate (MS 222) on the microhematocrit of fish blood. *Journal Fish Biology*.1977; 10: 453 – 6.
- [47] Balm P H M , Lambert J D G, Wendelaar- Bonga S E. Corticosteroid biosynthesis in the interrenal cells of the teleost fish, *Oreochromis mossambicus*. *General and Comparative Endocrinology*.1989; 76: 53-62.
- [48] Barcellos L J G, Nicolaiewsky S, de Souza S M G, Lulhier F. Plasmatic levels of cortisol in the response to acute stress in Nile tilapia, *Oreochromis niloticus* (L.), previously exposed to chronic stress. *Aquaculture Research*.1999;30: 437-444.