

Comparative Evaluation of Anti-diabetic and Hepatoprotective Effects of *Moringa oleifera* and *Bridelia ferruginea* Leaves Aqueous Extracts on Alloxan Induced Diabetic Albino Rats.

^x Aja, P.M., Nwafor, E.J., Ibiam, A. U., Orji, O. U., Ezeani, N and Nwali, B.U

Department of Biochemistry, Faculty of Biological Sciences, Ebonyi State University, Abakaliki.

^x Corresponding e-mail address: ovumte@yahoo.com

Abstract

This study was designed to evaluate comparative antidiabetic and hepatoprotective activities of aqueous extracts of *Moringa oleifera* and *Bridelia ferruginea* leaves in alloxan induced diabetic albino rats. Twenty-four (24) albino rats were grouped into A, B, C and D group. Group C and D were further subdivided into C₁, C₂, C₃, D₁, D₂ and D₃ respectively. The glucose and liver enzymes levels were determined using glucometeric and spectrophotometric methods. Diabetes was induced in all the groups, except group A (positive control). Group B (negative control) was not treated while group C and D were treated with aqueous extracts of *M. oleifera* and *B. ferruginea* leaves, which were administered orally to the animals twice daily for 1 week at varying concentrations of 200,400 and 800mg/kg body weights. The results revealed that there was significant reductions ($p < 0.05$) in glucose level in rats treated with aqueous extract of *B. ferruginea* than *M. oleifera* treated rats. There were no significant reduction ($p > 0.05$) in ALP level between the controls and the treated groups, except at 200mg/kg dose in which ALP level was high in rats treated with *M. oleifera* extract. There were significant reduction ($p < 0.05$) in ALT level in rats treated with both *M. oleifera* and *B. ferruginea* in comparison to the negative control while there was significant increase ($p < 0.05$) when compared to the positive control except at 200mg/kg dose where there was decrease in ALT level in both plant extracts. Also there was a significant decrease ($p < 0.05$) in AST level in rats treated with *M. oleifera* when compared to controls while there was significant increase ($p < 0.05$) in AST level in rats treated with *B. ferruginea* except at 200mg/kg where there was decrease when compared to the controls. These suggest that both extracts can be used therapeutically for the management of diabetes mellitus.

Key Words: Anti-diabetics, hepato-protective, *Moringa oleifera*, *Bridelia ferruginea*

Introduction

Diabetes mellitus is one of the most common endocrine diseases in domestic animals, especially man (Anaga, 2003). Diabetes mellitus is a chronic disease that occurs either when the pancreas does not produce enough insulin or when the body cannot effectively use the insulin it produces. Insulin is a hormone that regulates blood sugar (Susheela *et al.*, 2008). Diabetes is a disease condition distinguished by an inability to control blood glucose level due to the insufficient production of or heightened resistance to the hormone insulin (Ali *et al.*, 1993). It could also be described as a disorder that prevents the body using food properly (Chen *et al.*, 1995).

Diabetes mellitus is a major problem worldwide; approximately 5% of the world population suffers from diabetes (WHO, 1998). Since diabetes mellitus is a multi-factorial disease, the treatment is aimed to not only controlling blood sugar level to normal limit, but also at correcting the associated metabolic defects (Hattersley, 2002). Diabetes is characterized by chronic hyperglycemia, which causes glycation of body proteins that in turn leads to secondary complications affecting eyes, kidney, nerves and arteries. Along with hyperglycemia and abnormalities in serum lipid, diabetes is associated with micro and macro vascular complications the major causes of morbidity and death due to diabetes (Ramachandra *et al.*, 2002).

Liver is the largest internal organ. It is related by its domed upper surface to the diaphragm, which separates it from pleura, lungs, pericardium and heart (Harold, 2002). The cells of the liver, which are known as hepatocytes, house a lot of enzymes and carries out many biochemical activities. Some of these biochemical activities carried out by the hepatocytes include excretion of bile, carbohydrate metabolism, protein metabolism, synthesis of blood clotting factors, storage of iron and some vitamins, detoxification, lipid metabolism, etc (Ochei and Kolhatkar, 2007). It is therefore very obvious that any disease condition or adverse physiological conditions, which affect the hepatocytes, will cause concerted and tremendous metabolic derangement. In such conditions also, there will be increase in the serum activities of the mitochondrial-bound liver enzymes since hepatocytic damage causes their release into the serum. It is therefore very pertinent to ascertain the effect of any ingestible food or drug on the serum activities of the liver enzymes so as to ensure the hepato-protectiveness of such food or drug. This can be achieved through liver function tests, which include estimation of plasma protein, aspartate aminotransferase (AST), alkaline phosphatase (ALP), alanine aminotransferase (ALT), bilirubin, etc (Ochei and kolhatkar, 2007).

Traditionally, some plants are known to have anti-diabetic and hepato-protective properties, the folkloric use of some of these plants in the treatment of diabetes has been proven experimentally (Onukwo *et al.*, 1996). *Moringa oleifera* and *Bridelia ferruginea* are one of such plants used for the management of diabetes (Dieye *et al.*, 2008; Iwu, 1993).

Moringa oleifera belongs to the family of Moringaceae, a fast growing drought-resistant tree native of Sub-Hamalayan tracts of Northern India but now distributed worldwide in the tropics and sub tropics (Fuglie, 1999). *Moringa oleifera* has Horseradish tree, Drumstick tree and Ben oil tree, as its English names. While the local names in Nigeria include: “Gawara” (Fulani), “Zogall” (Hausa), “Ikwe oyibo” (Igbo) and “Ewe ile” (Yoruba). The Moringa tree grows mainly in semi-arid, tropical and subtropical areas. Currently it is widely grown in Africa, Central and South America, Sri-lanka, India, Mexico, Malaysia and the Philippines. It has diverse uses. The tree’s leaf and seedpod are widely consumed as

food. The bark, leaf and root have medicinal properties and have been used for centuries in folk medicine for a variety of health promoting purposes (Joseph, 2011).

Bridelia ferruginea belongs to the family Euphorbiaceae that is commonly found in savannah regions (Ekanem *et al.*, 2008). It is an indigenous medicinal plant in Nigeria. Its common names in Nigeria include “kirmi” (Hausa), “Maren” (Fulani), “Iralodan” (Yoruba) and “Ola” (Igbo). Its habitat is the savannah, especially in the moisture regions from Guinea to Zaire and Angola. *Bridelia ferruginea* has diverse uses. The leaves have been used to treat diabetes. The plant is used as a purgative and vermifuge (Cimanga *et al.*, 1999). The bark extract is being used for milk coagulation and also in limejuice for the formulation of traditional gargle “Ogunefu” (Orafidiya *et al.*, 1990).

Materials and Methods

Plants Collection

Fresh leaves of fully grown *Moringa oleifera* and *Bridelia ferruginea* were collected from Kpiripkiri in Ebonyi local government area, Ebonyi State and identified by Prof. S. S. Onyekwelu of the department of Applied Biology, Ebonyi State University, Abakaliki, Nigeria.

Preparation of Extract

The leaves of *Moringa oleifera* and *Bridelia ferruginea* were rinsed in clean water and air-dried under room temperature. The dried leaves were pulverized to fine granules using electric blender. The powder was soaked in distilled water at room temperature for 24 hours. It was then filtered using sieve cloth and the filtrate evaporated to dryness using rotator evaporator. The extracts were subsequently reconstituted in distilled water at appropriate concentrations for the experiment.

Experimental Animals

Twenty-four (24) albino rats of either sex weighing between 90 –140g (4-6-weeks old) were obtained from the animal house of the faculty of Veterinary Medicine University of Nigeria Nsukka. They were acclimatized for seven (7) days in stainless steel cages under good laboratory conditions. They were fed with commercial poultry growers mash feed (vital feed[®], Jos, Nigeria). Clean water was provided daily and access was free. The animals were weighed using triple beam weighing balance. Handling, management and use of animals for the experiment were as such that allowed minimal stress.

Experimental Design

At the end of the 7 days acclimatization period, the animals were randomly assigned into eight (8) different groups of three (3) rats each, designated as group A-D. Group A received water and feed only

and served as positive control. Diabetes were induced in group B, C and D. Group B was not treated and served as negative control while group C and D were further subdivided into C₁, C₂, C₃, D₁, D₂ and D₃ corresponding to 200,400 and 800mg/kg doses of the extracts. Group C₁ to C₃ was treated with *M. oleifera* extract while group D₁ to D₃ was treated with *B. ferruginea* twice daily for 7 days.

Induction of Diabetes:

About 2g of alloxan were dissolved in 20ml of distilled water and were administered to group B, C and D, based on the body weight of the rats and on a dosage of 100mg/kg. The administration was done intraperitoneally using diabetic syringe (Lenzen, 2008). The animals with sugar level more than 180mg/dl were considered as experimental diabetic (Stanley and Venugopal, 2001).

Collection of Blood from Animals:

After 7 days of treating the animals with the plants extract. The animals were starved for twenty-four (24) hours. The blood samples were collected from the tail vein puncture for the measurement of blood glucose. The blood glucose levels were measured in the fasting animals on 0th and 7th days. Blood samples used for liver enzyme assay was collected through ocular puncture using capillary tubes from the eyes of the animal.

Determination of Blood Glucose Level: Glucose level was determined using glucose oxidase method (Glucometer).

Determination of Liver Enzymes: Aspartate aminotransferase (AST), alanine aminotransferase (ALT), and alkaline phosphatase (ALP) were determined by method described by Reitman and Frankle, (1957), which was reviewed by Bhutia *et al.*, (2006).

Statistical Analysis: Results were expressed as mean $\pm$ standard deviation. The significance of difference between the controls and treated groups were determined, using two-way analysis of variance (ANOVA) SAS software version.




Figure 1: Effect of aqueous extracts formulated from *M. oleifera* and *B. ferruginea* leaves on the body weight of albino rats.




Figure 2: Comparison of the effects of aqueous extracts of *M. oleifera* and *B. ferruginea* leaves on serum glucose levels in diabetic albino rats.




Figure 3: Comparison of the effects of aqueous extracts of *Moringa oleifera* and *Bridelia fenuginea* leaves on the levels of Alkaline phosphatase activity in diabetic albino rats




Figure 4: Comparison of the effects of aqueous extracts of *Moringa oleifera* and *Bridelia fenuginea* leaves on the levels of Alanine aminotransferase activity in diabetic albino rats




Figure 5: Comparison of the effects of aqueous extracts of *M. oleifera* and *B. ferruginea* leaves on the levels of aspartate aminotransferase activity in diabetic albino rats.

Discussion

Figure 1 show variations on effect of aqueous extracts formulated from *M. oleifera* and *B. ferruginea* leaves on the body weight of albino rats during 7 days of administration. The treated animals and the positive control shows gradual increase in the body weight while negative control show decrease. This shows that as the glucose level decreases, the body weight improves also. This is in line with the work done by Grover *et al*, 2002.

Figure 2 shows that the glucose levels of all the animals were raised above normal after the administration of alloxan. It also demonstrated the anti-diabetics effects of the extracts on blood glucose level. The blood glucose level was seen to be lowered in the treated animals than that of the untreated. The results of this experiment validate the anti-diabetic effects of *Moringa oleifera* and *Bridelia ferruginea* as earlier asserted (Onunkwo *et al.*, 1996; Ndong *et al.*, 2007). The extracts showed significant reduction ($p < 0.05$) in the level of blood glucose in all the treated groups in dose dependent manner. The reduction of blood glucose level by *M.oleifera* leaves extract may be due to its numerous bioactive compounds such as flavonoids, alkaloid, coumarins, carotenoids, phenols, etc (Anwar *et al.*, 2007; Ferriera *et al.*, 2008). Some of these bioactive compounds may exert their hypoglycemic effects by, reducing insulin resistance, increasing release and decreasing glucagon's secretion, slowing the digestion and absorption of carbohydrates or by decreasing hepatic glucose production (Kakzung *et al.*, 2009). *B. ferruginea* also contains flavonoid, terpenoids, glycosides, alkaloids, etc (Iwu, 1993), as its bioactive compounds, which elicit their anti-diabetic effect by causing an increase in insulin output or by inhibition of the intestinal absorption of glucose or to the facilitation of metabolites in insulin dependent processes. It could also be achieved through increasing glucose transport (decreasing glycolysis and glucose oxidation in adipose tissue) (Clark *et al.*, 1985).

It can be deduced from the figure 3 that there were no significant increases ($p>0.05$) in the levels of ALP at the end of the extracts administration except at 200mg/kg of *M. oleifera*. Figure 4 shows that there were significant reductions ($p<0.05$) in the ALT levels in the treated animals when compared to the negative control. The figure 5 shows that there were significant decreases ($p<0.05$) in the AST level in animals treated with *M. oleifera* extract compared to the controls while there were increases in animals treated with *B. ferruginea* except at 400mg/kg dose where there was a decrease indicating that 400mg/kg is the therapeutic dose of *B. ferruginea*.

265

266 The results of hepatotoxicity markers showed there were no significant reduction ($p > 0.05$) in ALP level
 267 between the controls and the treated groups, except at 200mg/kg dose in which ALP level was high in rats
 268 treated with *M. oleifera* extract. There were significant reduction ($p < 0.05$) in ALT level in rats treated
 269 with both *M. oleifera* and *B. ferruginea* in comparison to the negative control while there was significant
 270 increase ($p < 0.05$) when compared to the positive control except at 200mg/kg dose in both plant extracts.
 271 There were significant reduction ($p < 0.05$) in ALT level in rats treated with both *M. oleifera* and *B.*
 272 *ferruginea* in comparison to the negative control while there was significant increase ($p < 0.05$) in ALT
 273 level when compared to the positive control except at 200mg/kg dose where there was decrease in ALT
 274 level in both plant extracts. Also there was a significant decrease in AST level in rats treated with *M.*
 275 *oleifera* compared to controls while there was significant increase ($p < 0.05$) in AST level in rats treated
 276 with *B. ferruginea* except at 200mg/kg where there was decrease when compared to the controls. The
 277 significant increase in the enzymes activities between the positive control and treated groups at the stated
 278 concentrations may be due to cholestasis in which conjugated bilirubin is equally retained (Crook, 2006).
 279 Though flavonoids and flavonoid-containing compounds also have been shown to cause hepatic disorders
 280 (Gandolfo *et al.*, 1992), and may be the result of the significant increase ($p < 0.05$) in these enzymes
 281 activities against the stated hepato-protectiveness actions of these leaves extract (Fakurazi *et al.*, 2008;
 282 Owiredu *et al.*, 2001). It could also be attributed to age variation and physical activity as most
 283 biochemical parameters are high in infancy followed by a fall towards childhood after which there is
 284 increase in level between puberty and adulthood (Owiredu *et al.*, 2011). According to Steffen and
 285 Menzel, 2009, significant increase in the serum activities of liver enzymes is an indication of hepatic
 286 injury, but since there was no significant increase in the serum activities of the enzymes except AST
 287 activity in *B. ferruginea* extract where there was reduction and elevation respectively which shows that
 288 *M. oleifera* is more safe to be used as anti-diabetic drug than *B. ferruginea* which may be as a result of the
 289 above reasons.

290

291 It therefore, means that *M. oleifera* and *B. ferruginea* leaves have hepato-protective activity and can be
 292 considered as alternative strategy to current pharmacotherapy of diabetes mellitus due to their
 293 hypoglycemic effects, less side effects of medicinal plants as compared to synthetic drugs, enormous cost
 294 of modern drug as well poor availability of the advanced therapies for many rural population in
 295 developing countries.

296 **Conclusion**

297 In conclusion, these findings suggest that *M. oleifera* and *B. ferruginea* leaf extracts have anti-diabetic
298 effect and relatively safe because of their hepato-protective properties.

299

300 **References**

301 Ali, L., Kahn, A. K. and Mosihuzzaman, M. (1993). Studies on the hypoglycemic effects of fruit pulp,
302 seed and whole plant of *Momordica cheranta* on normal and diabetic rat model. *Planta Medica*, **56**:408-
303 412.

304 Anaga, A.O. (2003). Antihyperglycemic and glucose uptake enhancing properties of *Dennettia tripetala*
305 G. baker. (Annonaceae), University of Nigeria, Nsukka, Nigeria, 203-204.
306

307 Anwar, F., Lotof, S., Ashara, M. and Gilani, A. H. (2007). *Moringa oleifera*: a food plant with multiple
308 medicinal uses. *Journal of Phytotherapy Research*, **21**: 17-25.

309

310 Chen, M. D., Lin, P. and Thouc, T. (1995). Selected metal status in patients with non-insulin dependent
311 diabetes mellitus, *Biological Trace Element Research*, **55**:119-124.

312 Cimanga, K., DeBrugne, T., Apers, S., Dietersh, T.J., Kambu, K., Tona, L., Bakana, P., Vanufford, L.Q.,
313 Benkelman.C., Labadie, R. and Veietinck, A. J. (1999). Complement-inhibiting constituent of *Bridelia*
314 *ferruginea* stem bark. *Planta Medica*, **65**(3): 213-217.

315

316 Crook, M. A. (2006). Clinical Chemistry and Metabolic Medicine, 7th edition, Edward Road Publishers,
317 London, 250-268.

318 Fakurazi, S., Wnithini, U. and Hairuszah, I. (2003). Hepatoprotective and antioxidant action *Moringa*
319 *oleifera*. *International Journal of Pharmacology*, **4**: 270-275.

320 Fuglie, L.J. (1999). The miracle tree: *Moringa oleifera* natural nutrition for the tropics. Church world
321 service, Dakar, 68-70.

322 Harold, E. (2002). Clinical Anatomy, 10th edition, Blackwell, U.K., 99-110.

323 Hattersley, A. (2002). Multiple facts of diabetes in young people. *Current Science*, **82**: 273-276

324 Iwu, M. M. (1993). Handbook of African Medicinal Plants. Chemical Rubber Company Press
325 Incorporation, Boca Raton, U.S.A., 133-135.

326 Joseph, C. (2011). *Moringa oleifera*: a review of the medicinal evidence for its nutritional,
327 therapeutic and prophylatic properties. *Trees of Life Journal*, **1**: 15-17.

328

329 Katzung, B. G., Masters, S. B. and Trevor, A. J. (2009). Basic and Clinical Pharmacology, 11th edition,
330 Large Medical Publications, Singapore, 727- 736.

331 Lenzen, S., (2008). The mechanisms of Alloxan and Streptozotocin induced diabetes. *Diabetologica*,
332 **51**:216-226.

- 333 Ochei, J. and Kolhatkar, A. (2007). Medical Laboratory Science: Theory and Practice, 6th edition, Tata
334 McGraw Hill, New Delhi, 106-158.
- 335 Onunkwo, G. C., Akah, P. A. and Udeala, O.K. (1996). Studies on *Bridelia ferruginea* leaves. Stability
336 and hypoglycemic actions of the leaf extract tablets. *Phytotherapy Research*, **10**(5): 423-420.
- 337 Ramachandra, A., Snehalata, C. and Vishwanathan, V. (2002). Burden of Type 2 diabetes and it's
338 complications-the Indian scenerio. *Current Science*, **83**:1471-77.
- 339
- 340 Reitman, S. and Frankel, S.A. (1957). A calometric method of determination of serum glutamic
341 oxaloacetic and glutamic pyruvate transminase. *American Journal of Clinical Pathology*. **28**:56-67.
- 342 Stanley, M.P. and Venugopal, M.P. (2001). Anti-oxidant action of *Tinospora cordifolia* root extract in
343 Alloxan-induced diabetic rats. *Phytotherapy Research*, **15**:213-218.
- 344 Steffen, C. and Menzel, J. (2009). Development and current status of Non-specific Enzyme therapy, 8th
345 edition, McGrew Hill companies Inc, New York, 10-14.
- 346
- 347 Susheela, T., Padma, B., Jane, T., Narender, R. and Reddy, P. (2008). Evaluation of hypoglycemic and
348 anti-diabetic effect of *Melia dubia* fruit in mice. *Current Science*, **94**(9): 1191-5.
- 349 WHO. (1998). The world health report, Life in the 21st century-A vision for all. World Health
350 Organization, Geneva.

351