

Original Research Article

Isolation of Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase from Young Leaves of *Phytolacca dioica* L.

Aims: Ribulose 1,5-bisphosphate (RuBP) carboxylase/ oxygenase (RubisCO) catalyses a key reaction by which inorganic carbon is assimilated into organic carbon found in the biosphere. Therefore, the present study aimed to isolate this enzyme from leaves of *Phytolacca dioica* L..

Study design: In this work, first crude extracts from leaves at different stages of development was assayed to isolate this enzyme, then young leaves of *P. dioica* were used considering high value of specific activity.

Methodology: Classical methods for protein isolation have been used to characterise RubisCO from *P. dioica*.

Results: RubisCO was isolated from young leaves by gel-filtration. The pure RubisCO showed two predominant bands (56- and 15-kDa) by SDS-PAGE. N-terminal sequences data on Large (56 kDa) and Small (15 kDa) subunits obtained by automatic Edman degradation show a high percentage of identity with Large and Small subunit of other RubisCo enzymes. Moreover, the N-terminal sequence of Large subunit (56 kDa) confirms the N-terminal sequence of Large subunit found by analysis from chloroplast genome of *P. dioica* (access number AFU65422).

Conclusion: The data on RubisCO from young leaves of *Phytolacca dioica* L. (RubisCO-Pd), obtained in the present work, could be used as the starting point for biological studies aimed to characterize this enzyme.

Keywords: Edman degradation, *Phytolacca dioica* L., Protein purification, RuBisCO.

1. INTRODUCTION

Ribulose 1,5-bisphosphate (RuBP) carboxylase/oxygenase (RubisCO) catalyzes the condensation of one molecule of CO₂, H₂O, and ribulose 1,5-bisphosphate (RuBP) to form two molecules of 3-phospho-D-glycerate and initiates net CO₂ fixation in autotrophs [1]. The most abundant enzyme on Earth, RubisCO, comprises around 40% of the total proteins of the plant [2]. RubisCO, however, is a bifunctional enzyme that also possesses an oxygenation activity. In this reaction, the sugar substrate is oxygenated to produce one molecule of 3-PGA and one molecule of 2-phosphoglycolate. The latter product is the substrate for photorespiration [3]. The ratio of carboxylation to oxygenation at any specified concentrations of CO₂ and O₂ is referred to as the CO₂/O₂ specificity factor [4].

RubisCO has been purified and characterized from a variety of sources. On the basis of amino acid sequences, four known forms of RubisCO, forms I, II, III and IV, were found in nature [5]. The form I has been isolated from plants, algae, proteobacteria and cyanobacteria, whereas the form II from proteobacteria, dinoflagellates. Moreover, the forms III and IV have been found in archaea or in proteobacteria, cyanobacteria, archaea and algae, respectively.

The most abundant form of RubisCO is the form I of holoenzyme, which consist of 55- and 15-kDa subunits (named L and S subunit, respectively). In higher plants the Large subunit is encoded by the chloroplast genome, whereas the Small subunit is encoded by the nuclear genome [6]. The structure of this holoenzyme is typically 550,000 in molecular weight and presumably consists of eight of each of the two types of subunits (LS)₈ [6]. The large RuBisCO subunits display a considerable sequence homology of 70-90 %. Large subunits of all three enzyme types are only for the 30-35 % identical in amino acid sequence, but highly conserved active sites. The small subunits in (LS)₈ are required for maximal catalytic rates [7].

RuBisCO is usually only active during the day as ribulose 1,5-bisphosphate. It is not regenerated in the dark. This is due to the regulation of several other enzymes in the Calvin cycle. Moreover, the RubisCO activity involves the interaction with various molecules as ions, phosphate, ATP/ADP and stromal reduction/oxidation state as well as RuBisCO activase [8,9]. In particular, RubisCO activase is the primary enzyme responsible for RubisCO's catalytic proficiency and it is heat labile. Moreover, this enzyme has even been suggested as the primary cause for reduced photosynthetic performance in response to moderate heat [10].

Phytolacca dioica L. (commonly known as "Ombú" or "Bella Sombra") is considered from our research group an important system model for the characterization of peptides, proteins and enzymes [11,12,13,14]. Therefore, in the present work, a Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase (hereafter RuBisCO-Pd) has been isolated from young leaves of *P. dioica*. The purification and partial characterization of RuBisCO-Pd from *P. dioica* are the essential steps to establish a basis for further studies.

2. MATERIAL AND METHODS

2.1 Chemicals, reagents and buffers

Young eaves of *P. dioica* were harvested at the end of March from a single female tree in the Botanical Garden of Napoli, Italy (Fig. S1).

3-phospho-D-glycerate and D-ribulose-1,5-bisphosphate were procured were from Sigma-Aldrich (Milan, Italy). The silica gel 60-coated (with specific surface area = 500 cm²/g; pore volume = 0.75 cm³/g; pore diameter = 60 Å) micro TLC plates (Merck, Darmstadt, Germany).

The solvents of highest purity analytical were purchased from Carlo Erba reagent (Milan, Italy). Coomassie Brilliant Blue R-250 was purchased from Fluka (Milan, Italy). Standard proteins for SDS-PAGE was purchased from Sigma-Aldrich. All other reagents and chemicals were of analytical grade. Reagents for automated Edman degradation were supplied by Life Technologies (Monza, Italy).

The following buffers have been used: buffer A: 25 mM Tris•Cl, 1 mM EDTA, and 1 mM dithiothreitol (DTT), pH 7.6; buffer B: 25 mM Tris•Cl containing NaCl 0.3 M, pH 8.2. buffer C: 200 mM bicine solution, pH 8.2, containing 0.4 mM EDTA and 1 mM DTT.

2.2 Fast RubisCO-Pd purification

Young leaves (10 g) were homogenized in 50 mL of buffer A by 20-s bursts at full power using a Waring Blendor (Waring Products), at 4°C. The homogenate was then filtered through Miracloth paper (Inalco, Milan, Italy) and centrifuged at 12,400g in a JA-14 rotor (Centrifuge Avanti J-25, Beckman Coulter, CA, USA) for 60 min, at 4°C. Proteins were further concentrated in Centriprep YM-10 (Millipore Corporation, Bedford, MA, USA). This concentrated fraction was first centrifuged as above and then gel-filtered on a Hi-load 26/60 Superdex 75 (GE Healthcare, Milan, Italy; separation range 100-3 kDa), equilibrated and eluted with buffer B, at a flow rate of 2.5 mL/min, using the AKTA Purifier 100 FPLC (GE Healthcare). Fractions with RubisCO activity were pooled, concentrated and stored at -20 °C.

2.3 Protein electrophoresis

Homogeneity of isolated proteins was determined by SDS-PAGE with a Mini-Protean II mini-gel apparatus (Bio-Rad; Milan, Italy), using 6% (w/v) stacking polyacrylamide gel and 15% (w/v) separation gel [15].

2.4 N-terminal amino acid sequencing by Edman degradation

Native RubisCO-Pd was separated by SDS-PAGE in reduction condition and after electrophoresis, protein was transferred to PVDF membrane (Life Technologies) by electroblotting with the mini trans-blot cell (Bio-Rad, Rome, Italy), according to the manufacturer's instructions in 10mM CAPS, pH 11.0, containing 10% methanol. PVDF membranes were then stained for 1 min with Coomassie Blue R-250, destained with the washing solution (50% methanol) [16], dried and directly analysed by Edman degradation on a Procise Model 491C sequencer (Applied Biosystem Inc. Foster City, CA, USA) [17].

2.5 Homology Studies

Amino acid sequences of other RubisCO enzymes were retrieved and aligned using the BLASTp software on the NCBI taxonomy browser. The program was available at the NCBI web site (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Alignment was performed by using the on line program ClustalW, available at <http://www.ebi.ac.uk/Tools/msa/clustalw2/>.

2.6 Enzymatic Activity

RubisCO activity was measured using a nonradioactive assay method based on thin-layer chromatographic separation of 3-phospho-D-glycerate (3-PGA) and D-ribulose-1,5-bisphosphate (RuBP) as previously reported [18].

3. RESULTS AND DISCUSSION

3.1 RubisCO-Pd Isolation

Although the presence of RubisCO in *P. dioica* has not been investigated so far, crude extracts from young leaves of *P. dioica* were assayed (data not shown). The preliminary data showed a high RubisCO activity from young leaves of *P. dioica*, thus heading towards the purification of RubisCO. Therefore, a purification procedure was undertaken in order to isolate potential novel RubisCO. The protein fraction extracted from young leaves of *P. dioica* with Tris•Cl saline buffer was submitted to gel-filtration chromatography, at pH 8.2. Each fraction showing RubisCO activity was analysed by SDS-PAGE (data not shown) and the homogeneous samples (fraction numbers 39-43) were pooled (Peak 1, Fig. 1).




Fig. 1. **A**, elution profile of the gel-filtration chromatography of the crude extract from young leaves of *P. dioica*. Fractions containing RubisCO activity are shaded (Peak 1). **B**, the chromatographic elution profile of standard proteins is also reported (BSA, bovine serum albumin; Mb, horse myoglobin; Ubi, ubiquitin).

Peak 1, showing RubisCO activity, with an elution volume with a molecular mass higher than 100 kDa was concentrated and subjected to partial enzymatic and structural characterisation (hereafter named RubisCO-Pd). Isolated RubisCO-Pd, analysed by SDS-PAGE, showed two bands of 56 and 15 kDa in the presence of β -mercaptoethanol (**Fig. 2 A**, lanes 1-2). On the other hand, when characterized by gel-filtration, a higher value than 100 kDa was obtained. These findings suggest that the native RubisCO-Pd is organised as an holoenzyme, in agreement with other forms I of RubisCO [6]. In fact, it is known that RubisCO proteins are stable as holoenzyme enzymes, supporting the prominent connection between the stability of the proteins and their enzymatic activity. Finally, in order to characterize the enzymatic activity, RubisCO-Pd_r was incubated in buffer C, at 37 °C for a fixed period in the presence of CO₂ gas, as previously reported [18]. Then, the sample was withdrawn and subjected to TLC analysis. A typical thin layer chromatogram (TLC) for the determination of RubisCO (carboxylation) activity is showed in **Fig. 2 B**. As showed in this figure, the amount of 3-PGA in the assay increases over time, vice versa the amount of RuBP decreases, confirming the carboxylation reaction of RubisCO-Pd.




Fig. 2. A, analysis by SDS-PAGE in presence of β -mercaptoethanol of RubisCO-Pd from young leaves of *P. dioica*. M, marker proteins; lanes 1-2, isolated RubisCO-Pd, (2 and 3 μ g, respectively). **B**, A thin layer chromatogram (TLC) for the determination of RubisCO-Pd (carboxylation) activity. The D-ribulose-1,5-bisphosphate (RuBP) and 3-phospho-D-glycerate (3-PGA) as standard are indicated. The assay mixture contained RuBP and purified RubisCO-Pd were incubated for the indicated time interval and analyzed by TLC.

3.2 Protein identification and homology studies

To obtain N-terminal sequence data on RubisCO-Pd, the bands of lane 2 (see **Fig. 2 A**), in a novel set of experiment, were transferred to PVDF membrane and directly subjected to automatic Edman degradation, as described in the Materials and Methods section. This analysis provided two different sequences (up to residue seventeen), 1-AGVKDYKLTY YTPQYKP-17 and 1-QVWPDLLGKK FFETLSY-17 for band 1 and band 2, respectively (**Fig. 3**). Subsequently, to obtain further sequence information on the two subunits of RubisCO-Pd, a search for sequence similarities, was performed with the BLAST program. It revealed that the both N-terminal sequences of the band 1 and band 2 of RubisCO-Pd share a high percentage of identity with large subunit and small subunit of other RubisCO enzymes. A multiple alignment of RubisCO-Pd subunits with other RubisCO sequences is reported in **Fig. 3**. Moreover, the analysis as previously reported, shows that the large subunits present highest identity with other large RubisCO subunits (100-94 % of identity), whereas, for the small subunits is lower (82 % of identity). On the other hand, the N-terminal sequence of large subunit (56 kDa band) results to be identical to that found by analysis of *P. dioica* chloroplast genome (access number AFU65422).

A, band of ~56 kDa (RubisCO large subunit)				
Plant	Accession	Sequence		Identity (%)
<i>P. dioica</i>		1-AGVKDYKLTY	YTPQYKP-17	(100)
<i>Rivina humilis</i> L.	AFI21564	1-AGVKDYKLTY	YTPQYKP-17	100
<i>Comesperma hispidulum</i> L.	CAJ81067	1-AGVKDYKLTY	YTPQYKP-17	100
<i>Phytolacca icosandra</i> L.	AFI21562	1-AGVKDYKLTY	YTPEYKP-17	94
a ***** :***				

B, band of ~15 kDa (RubisCO small subunit)				
Plant	Accession	Sequence		Identity (%)
<i>P. dioica</i>		1-QVWPDLLGKK	FFETLSY-17	(100)
<i>Ricinus communis</i> L.	XP_002521232	1-QVWP-PLGKK	KFETLSY-17	82
<i>Arachis duranensis</i>	AIF75302	1-QVWP-PLGKK	KFETLSY-17	82
<i>Litchi chinensis</i>	ADT78691	1-QVWP-PVGKK	FFETLSY-17	82
a **** .:*** *****				

Fig. 3. Alignment of the N-terminal sequences of the large and small subunits of RubisCO-Pd with those of other RubisCO. ^a, Identical residues *, conserved substitutions (:) and semiconserved substitutions (.) are reported.

4. CONCLUSION

A novel RubisCO, named RubisCO-Pd, was isolated from young leaves of *P. dioica* by gel filtration. The RubisCO-Pd showed two predominant bands by SDS-PAGE; the Large subunit, whose molecular weight was about 56 kDa and the other was the Small subunit, whose molecular was about 16 kDa, whereas a high molecular weight was obtained by gel-filtration. N-terminal sequences of both large and small subunits of RubisCO-Pd show share a high percentage of identity with those of other plants RubisCO enzymes. Further investigations on RubisCO-Pd primary structure are still in progress to extend our knowledge on the structure/function relationships of this enzyme.


Finally, it should be emphasized that primary structure data, obtained in the present work, could be used as the starting point for molecular biology studies aimed at cloning of the RubisCO-Pd gene.

REFERENCES

- 1 McFadden BA, Autotrophic CO₂ assimilation and the evolution of ribulose diphosphate carboxylase. *Bacteriol Rev* 1973;37:289-319.
- 2 Feller U, Anders I, Mae T, Rubiscolytics: fate of RubisCO after its enzymatic function in a cell is terminated. *J Exp Bot* 2008;59:1615-1624.
- 3 Bowes G, Ogren WL, Hageman RH, Phosphoglycolate production catalyzed by ribulose diphosphate carboxylase. *Biochem Biophys Res Commun* 1971;45:716-722.
- 4 Tabita FR, Hanson TE, Satagopan S, Witte BH, Kreeel NE, Phylogenetic and evolutionary relationships of RubisCO and the RubisCO-like proteins and the functional lessons provided by diverse molecular forms. *Philos Trans R Soc Lond B Biol Sci* 2008;363:2629-2640.
- 5 Andersson I, Taylor TC, Structural framework for catalysis and regulation in ribulose-1,5-bisphosphate carboxylase/oxygenase. *Arch Biochem Biophys* 2003;414:130-140.
- 6 Tabita FR, Hanson TE, Li H, Satagopan S, Singh J, Chan S, Function, structure, and evolution of the RubisCO-like proteins and their RubisCO homologs. *Microbiol Mol Biol Rev* 2007;71:576-599.
- 7 Spreitzer RJ, Role of the small subunit in ribulose-1,5-bisphosphate carboxylase/oxygenase. *Arch Biochem Biophys* 2003;414:141-149.
- 8 Parry MA, Andralojc PJ, Mitchell RA, Madgwick PJ, Keys AJ, Manipulation of RubisCO: the amount, activity, function and regulation. *J Exp Bot* 2003;54:1321-1333.
- 9 Berg JM, Tymoczko JL, Stryer L, Chapter 20: The Calvin Cycle and the Pentose Phosphate Pathway, *Biochemistry*, 5th edition, Freeman, W. H., New York, 2002.
- 10 Salvucci ME, Crafts-Brandner SJ, Inhibition of photosynthesis by heat stress: the activation state of RubisCO as a limiting factor in photosynthesis. *Physiol Plant* 2004;120:179-186.
- 11 Parente A, Berisio R, Chambery A, Di Maro A, Type 1 Ribosome-Inactivating Proteins from the Ombú Tree (*Phytolacca dioica* L.). in: L.J. M., M.R. Hartley (Eds.), *Toxic Plant Proteins*, Springer, Heidelberg, Germany, 2010, pp. 79-106.
- 12 Guida V, Criscuolo G, Tamburino R, Malorni L, Parente A, Di Maro A, Purification and enzymatic properties of a peroxidase from leaves of *Phytolacca dioica* L. (Ombu tree). *BMB Rep* 2011;44:64-69.
- 13 Di Maro A, Valbonesi P, Bolognesi A, Stirpe F, De Luca P, Siniscalco Gigliano G, Gaudio L, Delli Bovi P, Ferranti P, Malorni A, Parente A, Isolation and characterization of four type-1 ribosome-inactivating proteins, with polynucleotide:adenosine glycosidase activity, from leaves of *Phytolacca dioica* L. *Planta* 1999;208:125-131.
- 14 Di Maro A, Berisio R, Ruggiero A, Tamburino R, Severino V, Zacchia E, Parente A, Structural and enzymatic properties of an in vivo proteolytic form of PD-S2, type 1 ribosome-inactivating protein from seeds of *Phytolacca dioica* L. *Biochem Biophys Res Commun* 2012;421:514-520.
- 15 Laemmli UK, Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:680-685.
- 16 Di Maro A, Di Giovannantonio L, Delli Bovi P, De Andres Segura F, Parente A, N-terminal amino acid sequences of intact and cleaved forms of mung bean nuclease. *Planta Med* 2008;74:588-590.
- 17 Muccio C, Guida V, Di Petrillo A, Severino V, Di Maro A, A novel malate dehydrogenase from *Ceratonia siliqua* L. seeds with potential biotechnological applications. *Protein J* 2012;31:667-673.

257 18 Chakrabarti S, Bhattacharya S, Bhattacharya SK, A nonradioactive assay method for
 258 determination of enzymatic activity of D-ribulose-1,5-bisphosphate
 259 carboxylase/oxygenase (RubisCO). J Biochem Biophys Methods 2002;52:179-187.
 260
 261
 262

263 **APPENDIX**
 264
 265
 266



267 **Fig. S1** Fully expanded (A), in development (B), and young leaves from an adult, 100-year-
 268 old plant, growing in the Botanical Garden of the University of Naples "Federico II".
 269 The leaves were harvested between April and June 2013. Only young leaves were
 270 collected in March.
 271
 272
 273