

Protein fractions Functional Properties of Dried *Imbrasia* *oyemensis* Larvae Full-fat and Defatted Flours

Abstract

Imbrasia oyemensis is a caterpillar widely consumed in Côte d'Ivoire. Functional characteristics of its full-fat and defatted flours were investigated using standard methods. Defatting led to a significant ($P \leq 0.05$) reduction of emulsion capacity and stability, whereas the soluble protein fractions (such as albumin and glutelin), dispersibility, bulk density, water absorption capacity, water solubility index, oil absorption capacity, foam capacity and stability increased significantly ($p < 0.05$). The full-fat and defatted flours had high oil and water absorption capacities as desirable characteristics for use in some foods such as meats, sausages, breads and cakes. Furthermore, they showed high bulk density and good wettability, and therefore would be suitable for use as a functional ingredient in a variety of food formulations.

Keywords: Functional Properties, Protein fractions, Defatted flour, Full-fat flour, *Imbrasia oyemensis*, Larvae, Insects.

1. Introduction

Increases in world population will require the production of vast amount of foods in the latter half of the twenty-first century. However it will be difficult to increase productivity to a level that satisfies food demand, mainly because of limited availability of new farm land. This will lead to shortages of food, especially animal protein. When total food resources are insufficient, it is unwise to feed livestock with grain and other foodstuffs, which can be consumed directly by humans [1]. Therefore, it becomes necessary to look for new sources of

24 animal protein such as insects, which are rich in nutrients. Most insects are edible, although
25 there are some toxic species, and they can thrive on a diet that humans cannot consume. Some
26 insects are even scavengers, such as saprophaggers or coprophaggers. The latter can contribute
27 to recycling animal waste [2].

28 Insects represent a significant biological resource that is still not fully utilized around the
29 world. Entomophagy (eating of insects) has a long history as part of human diets and a large
30 number of insect species are consumed in many parts of the world. For approximately 2.5
31 billion people, mainly in Africa, Asia and Latin America, eating insects is part of their
32 common diets in a similar way as eating meat or fish. More than 1600 species of insects are
33 eaten worldwide [1].

34 Insects constitute high quality food for humans and animals, and there is a great potential and
35 growing global interest for utilization of insects as food resource to complement the diets of
36 continuously growing populations. Indeed, they represent an important source of proteins (20
37 to 70% of dry weight), amino acids, fat, carbohydrates, various vitamins and others micro-
38 nutrients (minerals, trace elements). Therefore insects offer an important nutritional resource
39 for humans and are worthy of development [3, 4, 5]. Recent reports in Nigeria confirm that
40 insects are indeed excellent sources of protein and other important nutrients [6]. Aside from
41 high quality protein, the study also found important supplements like minerals and vitamins,
42 even in dried form. Protein content can range as high as 29.8 percent for *Analeptes trifasciata*
43 F., and good quality vitamins A, B2 and C are found in *Apis mellifera* L. In the Philippines,
44 the nutritive value of *camaru* (mole cricket, *Gryllotalpa* sp.) was analysed and results showed
45 that a 150 g serving of the dish provided 28 and 74 percent of the daily protein and energy
46 requirement, respectively, of average Filipinos between the ages of 19 and 49 [7].

47 In Cote d'Ivoire, caterpillars are also prized and consumed. *Imbrasia oyemensis* is one of the
48 most widely eaten insects in the Western and Southern Côte d'Ivoire. This caterpillar, sold at
49 high prices on Abidjan markets, is consumed by an important population fringe of the country
50 in replacement of meat and fish [8]. The present paper focuses on the functional properties of
51 full-fat and defatted flours of dried *Imbrasia oyemensis* larvae. Indeed, functional properties
52 of foods are intrinsic physicochemical characteristics, which affect the behaviour of protein in
53 food systems during processing, manufacturing, storage and preparation [9]. This project was
54 undertaken to assess the functional properties the flour made from *Imbrasia oyemensis* larvae.
55 Such functional properties could be used to determine the suitability or otherwise of this
56 larvae as food additive.

57 **2. Materials and methods**

58 **2.1. Collection and preparation of samples**

59 Dried caterpillars *I. oyemensis* were obtained from the “Gouro” market of Adjamé (Abidjan,
60 Côte d'Ivoire). After collection, they were sorted and free from any kind of waste. Then, dried
61 caterpillars (2 kg) were ground using a porcelain mortar to get the full-fat flour. For defatted
62 flour, cold extraction method was used. Full-fat flour was placed in a conical flask and mixed
63 with hexane; the mixture was stirred by mechanical shaker for 16 hours and then filtered. The
64 residue was washed again with hexane and filtered. The resulting residue was dried in an open
65 air at room temperature. The dried flours were then stored at 0°C for further analysis.

66 **2.2. Protein Fractionation**

67 Extraction and fractionation of *Imbrasia oyemensis* proteins were carried out according to the
68 modified method of Hu and Essen [10]. Four different solvents deionized distilled water, 0.5
69 M NaCl (30 mL: 1 g), 70% Ethanol (w/v) and 0.1 M NaOH (25 mL: 1 g) were used in
70 sequence to extract virtually all of the proteins in *Imbrasia oyemensis* flours. Full-fat or

defatted flour was extracted by simply stirring, using a meal to solvent ratio of 1:10 (w/v), for 14-16 h at 4 °C. Insoluble residue was removed by centrifugation at 6000 rpm for 30 min. Extraction with each solvent was repeated three times, and all supernatants for each solvent were pooled to obtain a representative of each solubility fraction. Each fraction then was dialyzed against its own solvent followed by deionized water. Dialysates were stored in a freezer at about -20 °C. Distilled water-, NaCl, Ethanol and NaOH soluble proteins were designated “albumin”, “globulin”, “prolamine”, “glutelin”, respectively.

2.3. Protein Content

Each solubility fraction was assayed for protein content according to the method of Lowry et al. [11] using bovine serum albumin (BSA) as the protein standard. In addition, nitrogen content of unsoluble fraction (residue) was determined by micro-Kjeldahl method [12].

2.4. Dispersibility

The dispersibility of flours at selected protein concentration (2, 4, 6, 8 and 10 %) was measured according to the method of Mora-Escobedo *et al.* [13]. One gram of the flour was dispersed in distilled water in a 100 mL stoppered measuring cylinder. Then distilled water was added to reach a volume of 30 mL, the mixture was stirred vigorously and allowed to settle for three hours, the volume of settled particles was subtracted from 30 and multiplied by 100 and reported as percentage dispersibility.

2.5. Volume and bulk density measurement

The volume and bulk density were determined according to the modified method of Okezie and Bello [14], by pouring 2 g of flour into a 10 mL measuring cylinder, and then holding the cylinder on a vortex vibrator for 1 min to obtain a constant volume of the sample. The volume of the sample was recorded against the scale on the cylinder. The bulk density value was calculated as the ratio of mass of the powder and the volume occupied in the cylinder.

2.6. Determination of wettability

The method described by Onwuka [15] was adopted. One gram (1 g) of each flour sample was measured into a 10 cm³ measuring cylinder. The cylinder was inverted at 10 cm above the water contained in 600 ml beaker. The finger was used to close the cylinder disallowing the flour sample from falling. By removing the finger and giving the cylinder a gentle tap, the flour sample was discharged into the water surface. The time taken by the sample to get completely wet was recorded as the time of wettability.

2.7. Water absorption capacity and water solubility index

The water absorption capacity and solubility index of flours from *Imbrasia oyemensis* larvae were evaluated according to Phillips *et al.* [16] and Anderson *et al.* [17] methods, respectively. The full-fat and defatted flours from *Imbrasia oyemensis* larvae (2 g) were each weighed into a centrifuge tube and 50 mL distilled water added. The content of the centrifuge tube was shaken for 30 min in a KS 10 agitator. The mixture was kept in a water-bath (37°C) for 30 min and centrifuged (Ditton LAB centrifuge, UK) at 5000 rpm for 15 min. The resulting sediment (M2) was weighed and then dried at 105°C to constant weight (M1). The WAC was then calculated as follows:

$$\text{WAC (\%)} = \frac{M2 - M1}{M2} \times 100$$

While the WSI was calculated using the following equation:

$$\text{WSI (\%)} = \frac{M0 - M1}{M0} \times 100$$

M0 is the original weight of sample.

2.8. Determination of oil absorption capacity

For the oil absorption capacity, the method of Beuchat [18] was used. One gram of each flour sample was mixed with 10 mL of oil for 30 min in a mixer (Vari-whirl-mixing control set at fast speed). The sample was then allowed to stand at room temperature for 30 min. It was then centrifuged at 5000 rpm for 30 min, using a spinner (Ditton LAB centrifuge, UK) and the volume of the supernatant noted in a 10 mL graduated cylinder. The density of the oil was determined too. The volume of oil absorbed was multiplied by the density of the oil to determine the weight of oil so absorbed.

$$\text{OAC (\%)} = \frac{(V1 - V2) \times P}{W} \times 100$$

Where:

V = Initial volume of oil used 1

V = Volume remaining (not absorbed) 2

P = density of the oil used

W = Weight of sample

2.9. Foam capacity and foam stability:

The foam capacity (FC) and stability (FS) of flours from *Imbrasia oyemensis* larvae were studied according the method of Coffman and Garcia [19]. Three (3) g of each flour were transferred into clean, dry and graduated (50 mL) cylinders. The flour samples were gently levelled and the volumes noted. Distilled water (30 mL) was added to each sample; the cylinder was swirled and allowed to stand for 120 min while the change in volume was recorded every 15 min.

$$FC (\%) = \frac{V_t - V_0}{V_0} \times 100$$

$$FS (\%) = \frac{FC}{FC_0} \times 100$$

Where V_0 is the original volume of sample (mL), V_t is the total volume after different times (mL) and FC_0 is the foam capacity (FC) at 0 min.

2.10. Determination of emulsion capacity and stability

The method of Beuchat [18] was used. 2 g of each flour sample and 50 ml distilled water were blended at room temperature for 30 sec in Philips blender at 1600 rpm. After complete dispersion, vegetable oil (Gino) was added continuously in 10 mL portions from a burette. Blending continued until the emulsion breakpoint (where a separation into two layers/phases) was observed. The emulsion capacity was expressed as mL of oil emulsified per gram of sample and was expressed as %:

$$EC (\%) = \frac{V_E \times 100}{V \times W}$$

Where:

W = weight of sample

V_E = Volume of emulsion layer E

162 V = Total volume of mixture

163 For the emulsion stability, the emulsion so prepared was then allowed to stand in a 250 mL
164 graduated cylinder over time and the volume of the emulsion layer read. The stability was
165 measured in terms of the amount of oil that was retained in the emulsion layer and given by:

166

167
$$ES (\%) = \frac{V_{ET} \times 100}{V}$$

168

169 V_{ET} = Emulsion volume at Time (T)

170 V = total volume of the mixture

171

172 **2.11. Determination of swelling index**

173 The swelling capacity of the samples was determined using the method of Lin *et al.* [20], with
174 slight modification. One gram (1 g) of the flour sample was dispersed in 10 mL of cold
175 distilled water in a graduated centrifuge tube. The suspension was left at room temperature for
176 5 min to absorb water but not to swell. After 5 min the mixture was centrifuged at 2000 rpm
177 for 30 min and the volume of the sediment recorded as initial volume. Another 1 g of the
178 sample was dispersed in a centrifuge tube of known weight and the suspensions heated in
179 boiling water for 30 min. The suspension was cooled to room temperature under the tap water
180 and then centrifuged at 2000 rpm for 30 min. using a magnetic stirrer. The volume of the
181 heated sediment was recorded as final volume:

182 Swelling index = Final Vol. after heating / Initial Vol. before heating

2.12. Statistical analysis

All experiments in this study are reported as means of three replicate analyses. One-way analysis of variance (ANOVA) was carried out to compare the mean values of the two flours. Differences in the mean values were determined using Duncan's multiple range tests (SAS, 1990).

3. Results and Discussion

3.1. Soluble and insoluble fractions

Table 1 presents the Protein content of the soluble and insoluble fractions of the caterpillar flours. As regards these results, glutelin was found to be the major component in both full-fat (31.20 ± 1.47 %) and defatted (32.21 ± 0.59 %) flours. It is well known that glutelin is generally associated to elasticity and toughness properties in flours [21]. Albumin was the second component with content of 11.96 ± 0.09 % in crude flour and 18.51 ± 0.24 % in defatted flour. These results suggest that defatting of *Imbrasia oyemensis* flour significantly increase its glutelin and albumin contents. The statistical analysis showed no significant difference in globulin (4.18 ± 0.44 % in full-fat flour and 4.19 ± 0.08 % in defatted flour) and prolamin (4.44 ± 0.70 % in full-fat full-fat flour and 4.16 ± 0.14 % in defatted flour) contents.

Table 1: Protein content of the soluble and insoluble fractions *Imbrasia oyemensis* flours

Flour samples	Soluble fraction (%)				Insoluble fraction (%)
	Albumin	Globulin	Prolamin	Glutelin	
Full-fat	11.96 ± 0.09^d	4.18 ± 0.44^a	4.44 ± 0.70^a	31.20 ± 1.47^f	7.85 ± 0.30^c
Defatted	18.51 ± 0.24^e	4.19 ± 0.08^a	4.16 ± 0.14^a	32.21 ± 0.59^g	6.77 ± 0.30^b

3.2. Functional properties

The functional properties of flours play important role in the manufacturing of products. In present study the functional properties of full-fat and defatted *Imbrasia oyemensis* flours were evaluated.

3.2.1. Nitrogen solubility profile

The solubility of a protein is the thermodynamic manifestation of the equilibrium between protein–protein and protein solvent interactions [22]. Solubility of a protein is one of the critical functional attributes required for use as food ingredient, because it greatly influences other properties such as emulsification, gelation and foaming [23]. Thus, it is an important property governing the functional behavior of proteins and their potential application to food processing. The pH effect on protein solubility is shown in **Fig.1**. It reveals lower solubility in alkaline media. Higher solubility values were obtained in acid media and the isoelectric points (IEP) values are 4, and 7.6. At the isoelectric point, there is no net charge on the protein; resulting in no repulsive interactions and the protein–protein interactions disfavoring solubility [24]. The minimum solubility was found to be 0.11 ± 0.008 and 0.13 ± 0.02 mg/mL (at pH 4) and 0.0 ± 0.0 and 0.03 ± 0.0 mg/mL (at pH 7.6) for full-fat and defatted flours, respectively. On either sides of this pH, the solubility was increased up to maximum values at the extreme levels of pH. Generally, the dependency of the solubility on pH has been attributed to the change in the net charges carried by the flour protein as the pH changes [25]. The high net charge acquired at both acid and alkaline pH's caused arise in solubility due to unfolding of the flour protein with the degree of unfolding being greater at acidic than the alkaline pH [22]. The possession of 2 isoelectric points (IEP 4 and 7.6) suggests that *Imbrasia oyemensis* has more than one major protein constituent.




Figure 1: Effect of pH on protein solubility of *Imbrasia oyemensis* flours

3.2.2. Dispersibility and wettability

The dispersibility of a mixture in water indicates its reconstitutability [26]. The better temperature, ionic composition, pH and agitation degree of the solvent are major factors affecting dispersibility [23]. This property is a means of comparing the solubility of a protein in water, and this property is widely used in the flour and powder studies. The dispersibility of full-fat and defatted *I. oyemensis* flours is shown in **Table 2**. The results indicated that defatted flour had higher dispersibility than full-fat flour. Also, for both, it was higher at low concentration (2%) than high. As shown in **Table 2** the dispersibility of full-fat flour decreased significantly ($P \leq 0.05$) with increasing the level of *I. oyemensis* protein

concentration (2%, 4%, 6%, 8% and 10%). As regards defatted flour, it also decreased significantly ($P \leq 0.05$) but until concentration of 6%. These results indicated that increasing of protein concentration wasn't favourable for a better dispersibility of *I. oyemensis* flours. On the over side, defatting increased significantly this dispersibility suggesting that fats might play a negative role on this functional property. The dispersibility of a mix in water indicates its reconstitutionability. The higher the dispersibility, the better the reconstitution property [26]. Higher dispersibility enhances the emulsifying and foaming properties of proteins, which was observed during making of bread, macaroni and cookies [27].

As regards wettability, it is an important property when protein powders are dispersed to produce aqueous beverages and batters [28]. Wettability of proteins is affected by surface polarity, topography, texture, area and by the size and microstructure of the protein particles [29]. The results presented in **Table 2** shows that full-fat flour wet slowly (50.33 ± 0.58 min) while defatted sample wet very quickly (9.67 ± 0.58 min). This might be due to presence or absence of fat. So, defatting greatly influences the wettability.

Table 2: Dispersibility and wettability of *Imbrasia oyemensis* flours

Flour samples	Dispersibility (%)					Wettability (min)
	2%	4%	6%	8%	10%	
Full-fat	88.73 ± 2.63^c	80.90 ± 2.32^a	74.64 ± 0.63^f	71.61 ± 0.67^e	59.71 ± 1.65^d	50.33 ± 0.58^b
Defatted	94.61 ± 2.26^g	91.12 ± 2.31^c	84.89 ± 1.86^b	83.51 ± 0.31^{ab}	81.63 ± 0.33^a	9.67 ± 0.58^a

3.2.3. Bulk density

Bulk density is a measure of heaviness of a flour sample. It is important for determining packaging requirements, material handling and application in wet processing in the food

industry [28]. The bulk density of full-fat and defatted *I. oyemensis* flour was found to be 1.00 ± 0.0005 and 1.04 ± 0.02 g/mL (**Table 3**). It worth to note that the bulk density reported in the present study for the two samples is higher than the previously reported values for *I. belina* (0.67–0.71 g/mL) [30]. Higher bulk density is desirable for the greater ease of dispersibility and reduction of paste thickness which is an important factor in convalescent and child feeding [31]. Thus, the full-fat and defatted *I. oyemensis* flour could possibly be used in food formulation as supplement for child food. The high bulk density of *I. oyemensis* flours indicates that they would serve as good thickeners in food products.

Table 3: Bulk density of *Imbrasia oyemensis* flours

Flour samples	Bulk Density (g/mL)
Full-fat	1.00 ± 0.0005^a
Defatted	1.04 ± 0.02^b

3.2.4. Water and oil absorption capacity

Hydration or rehydration is the first and perhaps most critical step in imparting desirable functional properties to proteins in a food system. Interactions of water and oil with flours are very important in food transformation because of their effects on the flavour and texture of foods. Intrinsic factors affecting water binding properties of food flours with relatively high protein contents include amino acid composition, protein conformation and surface polarity/hydrophobicity [32]. Water absorption capacity describes flour – water association ability under limited water supply. The results of water and oil absorption capacity of the two samples are shown in **Table 4**. The defatted *I. oyemensis* flour (65.44 ± 5.81 to 86.89 ± 1.38 %) had higher water absorption capacity (WAC) than the full-fat flour (56.90 ± 1.39 to 66.07 ± 0.55 %). These values are higher than those reported for the Jackfruit seed flour (25 %) [33]

and for African yam bean (17 %) [34]. The results obtained are however lower than those of *Cirina forda* (300.00 ± 0.15 %) reported by Omotoso [35] and than those of mushroom flours reported by Aremu et al. [36]. This shows that *I. oyemensis* is slowly hydrophilic. The difference in protein structure and the presence of different hydrophilic carbohydrates might be responsible for variation in the WAC of the flours. Indeed, flours with high WAC have more hydrophilic constituents such as polysaccharides [36].

The highest oil absorption capacity (OAC) was found in defatted flour at 2 % of sample (84.08 ± 0.43 %) and the lowest in full-fat flour at 10 % of sample (56.71 ± 0.74 %). Oil absorption capacity is important since oil acts as flavour retainer and increases the palatability of foods [23]. The result obtained shows that *I. oyemensis* flours are high flavour retainer and may therefore find useful application in food systems such as ground meat formulations.

289

290 **Table 4:** Water and oil absorption capacity of *Imbrasia oyemensis* flours

Flour samples	Water Absorption Capacity (WAC) (%)					Oil Absorption Capacity (OAC) (%)				
	2%	4%	6%	8%	10%	2%	4%	6%	8%	10%
Full-fat	56,90 ± 1,39 ^d	61,20 ± 0,89 ^b	66,07 ± 0,55 ^a	64,61 ± 0,43 ^{ab}	63,40 ± 0,48 ^{ab}	78.12 ± 0.45 ^b	67.09 ± 0.65 ^g	64.93± 0.80 ^f	60.23± 1.47 ^d	56.71± 0.74 ^c
Defatted	65,44 ± 5,81 ^a	79,06 ± 0,07 ^c	86,89 ± 1,38 ^e	79,15 ± 0,00 ^c	78,58 ± 0,55 ^c	84.08 ± 0.43 ^h	79.30 ± 0.44 ^b	69.92± 1.15 ^a	70.26± 0.44 ^a	62.68 ± 0.57 ^e

291

3.2.5. Emulsifying properties

The emulsifying properties are usually attributed to the flexibility of solutes and exposure of hydrophobic domains. Food emulsions are thermodynamically unstable mixtures of immiscible liquids. The formation and stability of emulsion is very important in food systems such as salad dressing [37, 38]. As shown in **Table 5** the emulsion capacity was higher for defatted flour than full-fat flour at 2 % proteins concentration and, also for both, it was higher at low concentration (2%) than high. The emulsion capacity of full-fat as defatted flour decreased significantly ($P \leq 0.05$) with increasing the level of *I. oyemensis* proteins concentration. These values were higher than those reported for defatted groundnut cake flour (28.10 %) [37] and for roasted peanut [38]. The emulsifying activity results were 80.85 ± 0.98 and 86.46 ± 1.38 % respectively for full-fat and defatted flour, whereas the emulsion stability was 84.76 ± 0.34 and 87.21 ± 0.31 % for the two samples, respectively. The results obtained were much higher than those reported by Fekria et al. [37] for defatted groundnut cake flour and Omotoso [35] for *Cirina forda* larva protein. The capacity of proteins to enhance the formation and stabilization of emulsion is important for many applications in cakes, coffee whiteners, and frozen desserts. In these products varying emulsifying and stabilizing capacities are required because of different compositions and stresses to which these products are subjected [39].

310

311 **Table 5:** Emulsifying properties of *Imbrasia oyemensis* flours

Flour samples	Emulsion Capacity (EC) (%)					Emulsion Stability (ES) (%)				
	2%	4%	6%	8%	10%	2%	4%	6%	8%	10%
Full-fat	80,85 ± 0,98 ^g	46,66 ± 1,20 ^f	25,90 ± 0,60 ^b	21,65 ± 0,58 ^a	16,84 ± 0,23 ^d	84,76 ± 0,34 ^a	91,22 ± 0,73 ^h	79,72 ± 0,25 ^f	88,14 ± 0,35 ^c	84,58 ± 0,61 ^a
Defatted	86,46 ± 1,38 ^h	43,80 ± 1,63 ^e	24,43 ± 0,72 ^b	21,61 ± 0,64 ^a	14,93 ± 0,11 ^c	87,21 ± 0,31 ^{bc}	86,40 ± 1,02 ^b	71,70 ± 1,42 ^d	82,78 ± 0,93 ^g	74,22 ± 0,63 ^e

312

313

3.2.6. Foaming capacity and foam stability

The foaming capacity (FC) of a protein refers to the amount of interfacial area that can be created by the protein and foam stability (FS) refers to the ability of protein to stabilize against gravitational and mechanical stresses [40]. Foam formation and foam stability are a function of the type of protein, pH, processing methods, viscosity and surface tension. The full-fat *I. oyemensis* flour showed no foam whereas the defatted flour had foaming capacity of 18.006 ± 2.37 % and foaming stability of 27.19 ± 0.98 % (Table 6). These values are greater than some insect flours as reported by some workers: *I. belina*, 11.8 ± 1.15 % (Ekpo et al., 2008) and *Rhyncophorus phoenicis*, 10 % [41]. Akubor and Chukwu [42] reported that foams are used to improve the texture, consistency and appearance of foods. Graham and Phillips [43] linked good foamability with flexible protein molecules, which reduces surface tension. Low foamability on the other hand can be related to highly ordered globular proteins, which resists surface denaturation. The basic requirements of proteins as good foaming agents are the ability to (i) adsorb rapidly at air water interface during bubbling, (ii) undergo rapid conformational change and rearrangement at the interface, and (iii) form a cohesive viscoelastic film via intermolecular interactions. The first two factors are essential for better foamability whereas the third is important for the stability of the foam [40].

Table 6: Foaming properties and bulk density of *Imbrasia oyemensis* flours

Flour samples	Foaming Capacity (FC) (%)	Foaming Stability (FS) (%)	Bulk Density (g/mL)
Full-fat	0.00 ^a	0.00 ^a	1.00 ± 0.0005^a
Defatted	18.006 ± 2.37^b	27.19 ± 0.98^b	1.04 ± 0.02^b

4. Conclusion

The data obtained in this study showed that defatting have significant effect on functional properties of dried *I. oyemensis* flours. The high protein content of both flours indicated that they could be a valuable protein. Moreover, high solubility of both flours at acid pH makes them suitable in beverages. Furthermore, full-fat and defatted flours had high oil and water absorption capacities as desirable characteristics for use in some foods such as meats, sausages, breads and cakes. Both the full-fat and defatted flours showed high bulk density and would be suitable for use in weaning food. The good wettability of the two flours makes them suitable for use in textured meats and baked products.

6. References cited

- [1] FAO 2004. *Contribution des insectes de la forêt à la sécurité alimentaire: L'exemple des chenilles d'Afrique Centrale*. Fao, Rome, Programme des Produits Forestiers non Ligneux. 112 pp.
- [2] Raubenheimer D, Rothman JM (2012). Nutritional Ecology of Entomophagy in Humans and Other Primates. *Annu. Rev. Entomol.* 58:141–60.
- [3] Ramos-Elorduy, J. & Pino, M.J.M. 1989. *Los insectos comestibles en el Mexico antiguo*. 108 pp.
- [4] DeFoliart, G. 1995. Edible insects as minilivestock. *Biodiversity and Conservation*, 4: 306-321.
- [5] Chen Xiaoming & Feng Ying. 1999. *The edible insects of China*. Beijing, Science and Technology Publishing House. 180 pp.

- 356 [6] Banjo AD, Lawal OA, Songonuga EA (2006). The nutritional value of fourteen species of
357 edible insects in Southwestern Nigeria. *Afr. J. Biotechnol.* 5(3): 298- 301.
- 358 [7] Barrion-Dupo, A.L.A., Sheree, A. & Lit, I.L., Jr. 2008. Nutritive value of camaro.
359 *Transactions of the National Academy of Science and Technology Philippines*, 30(1): 199.
- 360 [8] Akpossan RA, Ahipo DE, Kouadio JPEN et Kouamé LP (2009). Valeur nutritionnelle et
361 caractérisation physicochimique de la matière grasse de la chenille (*Imbrasia oyemensis*)
362 séchée et vendue au marché d'Adjamé (Abidjan, Côte d'Ivoire). *Journal of Animal & Plant*
363 *Sciences*, 3 (3): 243 - 250.
- 364 [9] Onimawo, I. A. and Akubor, P. I. (2005). Functional properties of food. In: *Food*
365 *Chemistry, integrated approach with biochemical background*. Ambik Press Ltd, Benin City.
366 Pp 208–221.
- 367 [10] Hu, B.; Esen, A. Heterogeneity of soybean seed proteins: one dimensional electrophoretic
368 profiles of six different solubility fractions. *J. Agric. Food Chem.* **1981**, 29, 497-501.
- 369 [11] Lowry O.H., Rosebrough N.J., Farr A.L. and Randall R.J. (1951). Protein measurement
370 with the folin reagent. *Journal of Biological Chemistry*, 193, 265.
- 371 [12] AOAC. (1995). *Official Methods of Analysis of the Association of Official Analytical*
372 *Chemists*, 16th ed. Virginia. U.S.A.
- 373 [13] Mora-Escobedo, R.; Paredes-Lopes, O. & Gutierrez-Lopes, G. F. (1991), Effect of
374 germination on the rheological and functional properties of amaranth seeds. *Lebensmittel -*
375 *Wissenschaft und Technol.*, 24, 241-244.
- 376 [14] Okezie, B.O. and Bello, A.B. (1988). Physicochemical and functional properties of
377 winged bean flour and isolate compared with soy isolate, *J. Food Sci.* 53:450-454.

- 378 [15] Onwuka, G.I. (2005). Food Analysis and Instrumentation. Theory and Practice. Naphtali
379 Prints, Lagos, Nigeria .pp 133-137.
- 380 [16] Phillips RD, Chinnan MS, Branch AL, Miller J, Mcwatters KH (1988) Effects of pre-
381 treatment on functional and nutritional properties of cowpea meal. J Food Sci 53(3):805–809.
- 382 [17] Anderson RA, Conway HF, Pfeifer VF, Griffin EL (1969) Roll and extrusion-cooking of
383 grain sorghum grits. Cereal Sci Today 14:372–380.
- 384 [18] Beuchat, L.R., 1977. Functional and electrophoretic characteristics of succinylated peanut
385 flour protein. *Journal of Agricultural and Food Chemistry*, 25:258-261.
- 386 [19] Coffman, C.W. and V.V. Gracia, 1977. Functional properties of amino acid content of a
387 protein isolate from mung bean flour. J. Food Technol., 12: 473-484.
- 388 [20] Lin, M. J. V., Humbert, E. S. and Sosulski, F.W. (1974). Certain functional properties of
389 sunflower meal products. J. Food Sci, 39, 368- 370.
- 390 [21] Tatham, A.S., Shewry, P.R., 1985. The conformation of wheat gluten proteins. The secondary
391 structures and thermal stabilities of a-, b-, g and o-gliadins. J. Cereal Sci. 3, 104–113.
- 392 [22] Damodaran, S. (1996). Amino acids peptides and proteins. In: Owen, R., Fennema (ed)
393 food chemistry, 3rd edn. Marcel Dekker, Inc., 270 Madison Avenue. New York, 10016, USA
394 pp. 327-425.
- 395 [23] J. E. Kinsella, 1976. Functional Properties of Proteins in Foods, a Survey,” *Critical*
396 *Review of Food Science and Nutrition*, 7: 219-222. [doi:10.1080/10408397609527208](https://doi.org/10.1080/10408397609527208)
- 397 [24] Singh N., Kaur M. and Sandhu, K.S. (2005). Physicochemical and functional properties
398 of freeze-dried and oven dried corn gluten meals. Drying Technology, 23, 1–14.

- 399 [25] Fagbemi TN, Oshodi AA & Ipinmoroti KO (2006). Effects of processing in the
400 functional properties of full fat and defatted fluted pumpkin seed flours. *J Food Tech* 4: 70 –
401 79.
- 402 [26] K. D. Kulkarni, D. N. Kulkarni and U. M. Ingle. 1991. Sorghum Malt Based Weaning
403 Food Formulations: Preparation, Functional Properties and Nutritive Value. *Food and*
404 *Nutrition Bulletin*, 13: 4322-4327.
- 405 [27] J. E. Kinsella. 1979. Functional Properties of Soy Proteins. *Journal of American Oil*
406 *Chemists Society*, 56: 242-249. [doi:10.1007/BF02671468](https://doi.org/10.1007/BF02671468)
- 407 [28] Zayas, J.F. 1997. Water holding capacity of proteins. In; *Functionality of proteins in*
408 *Food*. (1st Ed), Springer.
- 409 [29] Hagerdal, B. and Lofqvist, B. 1978. Wettability and surface pressure of myoglobin
410 treated with acetone. *Journal of Food Science* 43: 27-30.
- 411 [30] Ekpo, K.E., Azeke, M.A. And Ugbenyen, A.M 2008. Functional properties of dried
412 *Imbrasia belina* larvae flour as Affected by mesh size and pH. *Nigerian annals of natural*
413 *sciences*, 8(1): 6-9.
- 414 [31] Padmashree, T.S., Vijayalakshmi, L. and Puttaraj, S. 1987. Effect of traditional
415 processing on the functional properties of cowpea (*Vigna Catjang*) flour. *Journal of Food*
416 *Science and Technology* 24: 221-224.
- 417 [32] Barbut, S. (1999). Determining water and fat holding. In G. M. Hall (Ed.), *Methods of*
418 *testing protein functionality*, pp. 186–225. New York: Blackie Academic and Professional.

419 [33] Ocloo F.C.K., Bansa D., Boatın R., Adom T., Agbemavor W.S. 2010. Physico-chemical,
 420 functional and pasting characteristics of flour produced from Jackfruits (*Artocarpus*
 421 *heterophyllus*) seeds. Agriculture And Biology Journal Of North America. 15: 903-908.

422 [34] Eke OS, Akobundu ENT (1993). Functional properties of African yam bean
 423 (*Sphenostylis stenocarpa*) seed flour as affected by processing. Food Chem. 48: 337-340.

424 [35] Omotoso O.T. 2006. Nutritional quality, functional properties and anti-nutrient
 425 compositions of the larva of *Cirina forda* (Westwood) (Lepidoptera: Saturniidae). *J Zhejiang*
 426 *Univ SCIENCE B* 7(1):51-55.

427 [36] Aremu MO, Basu SK, Gyar SD, Goyal A, Bhowmik PK & Datta Banik S 2009.
 428 Proximate Composition and Functional Properties of Mushroom Flours from *Ganoderma*
 429 *spp.*, *Omphalotus olearius* (DC.) Sing. and *Hebeloma mesophaeum* (Pers.) Quél. Used in
 430 Nasarawa State, Nigeria. *Mal J Nutr* 15(2): 233 – 241.

431 [37] Fekria, A. M., Isam, A. M. A., Suha, O. A. and Elfadil, E. B. 2012. Nutritional and
 432 functional characterization of defatted seed cake flour of two Sudanese groundnut (*Arachis*
 433 *hypogaea*) cultivars Kinsella, JE (1979). Functional properties of soy proteins. *J Amer Oil*
 434 *Chemists' Soc* 56: 242 – 258.

435 [38] Yu, J., Ahmedna, M. and Goktepe, I. 2007. Peanut protein concentrates: production and
 436 functional properties as affected by processing. *Food Chemistry* 103:121–129.

437 [39] Elkhailifa, A. E. O. and Bernhardt, R. 2010. Influence of grain germination on functional
 438 properties of sorghum flour. *Food Chemistry*, 12: 387–392.

439 [40] Fennema, R.O. (1996). Food chemistry, (3rd edition). Marcel Dekker, Inc. New York,
 440 Basel. Hongkong. pp. 365-39.

441 [41] Adeyeye E I and Aye P A (2013). The Chemical Composition and Functional Properties
442 of the Flesh and Viscera of Raffia Palm Tree Grub (*Rhyncophoros Phoenicis*). Open Journal
443 of Analytical Chemistry Research, 1(2): 18-25. DOI: 10.12966/ojacr.08.01.2013.

444 [42] Akubor, P.I., Chukwu, J.K., 1999. Proximate composition and selected functional
445 properties of fermented and unfermented African oil bean (*Pentaclethra macrophylla*) seed
446 flour. *Plant Foods for Human Nutrition*, 54:227-238. [doi:10.1023/A:1008100930856]

447 [43] Graham DE, Phillips MC (1976). Foams. Academic Press, London. p. 237.

448