

1 Original Research Article
2 **GC-MS Analysis of Leaf, Fruits and Latex of**
3 ***Croton bonplandianum* Bail.**
4
5
6
7
8
9

10 **ABSTRACT**
11

Aims: To characterize the phytochemicals of different parts of *Croton bonplandianum* using GC-MS.

Study Design: GC-MS analysis of bioactive constituents of *C. bonplandianum*.

Place and Duration of Study: Post Graduate and Research Department of Biochemistry at Government Arts College (Autonomous), Kumbakonam and Food Safety and Quality Testing Laboratory, Indian Institute of Crop Processing Technology, Thanjavur, Tamilnadu, India, between May, 2011 to June, 2012.

Methodology: The *C. bonplandianum* leaf and fruit (25 grams) powder was soaked in 60 ml of ethanol each and kept at room temperature for 12 hours and the fresh latex of 10 ml was mixed with 90 ml of ethanol and kept at shaker for 3 hours. The samples were filtered and concentrated. Each sample was subjected to phytochemical analysis using GC-MS.

Results: The GC-MS analysis showed peaks of twenty one phytochemicals from different parts of *C. bonplandianum*. Out of which five compounds were found in leaf, one compound in latex and fifteen compounds in fruits. The highest peak area of 88.69% for 16-Hexadecanoyl hydrazide and the lowest peak area of 1.39% for Phytol were obtained in leaf extract. The latex of *C. bonplandianum* showed that the presence of Myo-Inositol, 2-C-methyl with the peak area of 30.8%. The fruits of *C. bonplandianum* showed that the presence of 9,12,15-Octadecatrienoic acid, methyl ester (z,z,z)- with the highest peak area of 41.81% and 2-Hexen-1-ol, 2-ethyl with the lowest peak area of 0.69%.

Conclusions: The phytochemical constituents of *C. bonplandianum* have been screened and the isolation of individual bioactive compounds from *C. bonplandianum* will be helpful to find new drugs.

12
13 **Keywords:** GC-MS analysis; *Croton bonplandianum*; bioactive compounds; leaf; fruit; latex.
14
15

16 **1. INTRODUCTION**
17

18 Medicinal plants are the richest bio-resources of folk medicines, traditional systems of
19 medicine, food supplements, nutraceuticals, pharmaceutical industries and chemical
20 entities for synthetic drugs [1]. Modern medicine has evolved from folk medicine and
21 traditional system only after through chemical and pharmaceutical screening [2]. India is
22 the birth place of renewed system of indigenous medicine such as Siddha, Ayurvedha
23 and Unani [3]. Traditional systems of medicines are prepared from a single plant or
24 combinations of a number of plants. The efficacy depends on the use of proper plant part
25 and its biological potency which in turn depends upon the presence of required quantity and
26 nature of secondary metabolite in a raw drug [4,5]. There is growing awareness in
27 correlating the phytochemical constituents of a medicinal plant with its pharmacological

activity [6]. Screening of active compounds from plants has lead to the invention of new medicinal drugs and they have an efficient protection against various diseases including cancer [7] and Alzheimer's diseases [8].

Phytochemicals are responsible for medicinal activity of plants [9]. These are non-nutritive chemicals that have protected human from various diseases [3]. Phytochemicals are basically divided into two groups that is primary and secondary metabolites based on the function in plant metabolism. Primary metabolites are comprise common carbohydrates, amino acids, proteins and chlorophylls while secondary metabolites consist of alkaloids, saponins, steroids, flavonoids, tannins and phenolic compounds [10,11]. Phytochemical constituents are the basic source for the establishment of several pharmaceutical industries. The phytochemical constituents are playing a significant role in the identification of crude drugs [8].

In recent years, the interest for the study of organic compounds from the plants and their activity has increased [12]. A knowledge of the chemical constituents of plant is desirable not only for the discovery of therapeutic agents, but also helps in disclosing new sources of economic phytochemicals for the synthesis of complex chemical substances and for discovering the actual significance of folkloric remedies [13]. There is an increasing interest on the phytochemical compounds, which could be relevant to their nutritional incidence and their role in health and disease [14]. The combination of an ideal separation technique - Gas Chromatography (GC) with the best identification technique - Mass Spectrum (MS) made GC-MS, which is an ideal technique for qualitative and quantitative analysis of volatile and semi volatile compounds. This technique has proved to be a valuable method for the analysis of non polar components and volatile oils, fatty acids, lipids and alkaloids [15]. *Croton bonplandianum* Bail belongs to the family of Euphorbiaceae and it is distributed in Southern Bolivia, Paraguay, South Western Brazil, Northern Argentina and India. *C. bonplandianum* was reported to have many medicinal uses including the repellent property against the insects [16] and showed antibacterial [17], antifungal [18], antioxidant [19], analgesic [20], nematocide [21], anticoronary [22,23,24], hepatoprotective [25] and wound healing activities [26,27]. Local people in the remote area of West Bengal in India are using its root against snake bite and leaf extract against high fever [28]. The methanol extract of *C. bonplandianum* has been found to exhibit antitumour properties in plants caused by *Agrobacterium tumefaciens* [29] and larvicidal activities [16]. Although the plant is used in Ayurvedic medicine for the treatment of various ailments but there are no detailed reports on the analysis of phytoconstituents of different parts of *C. bonplandianum*. With this background, the present study was aimed to identify the phytoconstituents present in leaf, fruit and latex of *C. bonplandianum* using GC-MS analysis.

2. MATERIAL AND METHODS

2.1 Collection of Plant Material

C. bonplandianum was collected from the waste lands in and around Mayiladuthurai, Tamilnadu, India, where it was found naturally. The plant was identified by Rev.Dr.John Britto, The Director of the Rabinat herbarium and Centre for Molecular Systematics, St.Joseph's College, Tiruchirappalli, Tamilnadu, India. The leaves and fruits were separated and washed thoroughly in running tap water to remove soil particles and other adhered debris and then finally washed with sterile distilled water. The whole plants were shade dried and ground well into fine powder. The powdered materials were stored in air tight container until the time of use [30].

The latex of *C. bonplandianum* was obtained as exudates by and plucking of fresh leaves from actively growing plants. The latex was collected into sterile, plastic containers by pressing and squeezing in between fingers, the apex of the leaves to release as much as possible latex into the containers. After collection, the latex was centrifuged with 5000 rpm for 15 minutes to remove any solid particles present in it. The latex was decanted off into containers and plugged with cotton and stored at 4 °C until required for use. Collections were made in the mornings during the days of each analysis [31].

2.2 Identification of Phytocompounds by GC-MS

2.2.1 Sample Preparation

25 grams of *C. bonplandianum* leaf and fruit powder were soaked in 60 ml of ethanol separately and kept at room temperature for 12 hours and 10 ml of fresh latex was mixed with 90 ml of ethanol and kept at shaker for 3 hours [30,32]. The samples were filtered and concentrated separately through nitrogen flushing to 1 ml. From this 2 µl of each prepared sample was injected into the GC-MS instrument for phytochemical analysis.

2.2.2 Equipment

The GC-MS analysis was carried out using a Perkin Elmer Clarus 500 for the analysis of phytocompounds of *C. bonplandianum* [33]. The data were obtained on a Capillary Column Elite-5MS (5% phenyl 95% dimethyl poly siloxane). Helium was used as the carrier gas with a flow rate of 1ml/min in the split mode (10:1). An aliquot of 2 µl of ethanol solution of the sample was injected into the column with the injector temperature at 250 °C. GC oven temperature started at 110 °C and holding for 2 min and it was raised to 200 °C at the rate of 10 °C /min without holding. Holding was allowed at 280 °C for 9 min with program rate of 5 °C /min. The injector and detector temperatures were set at 250 °C and 280 °C respectively. GC interface and Ion source temperature was maintained at 200 °C. The mass spectrum of compounds in the samples was obtained by electron ionization at 70 eV and the detector was operated in scan mode from 40-450 amu (atomic mass units). A scan interval of 0.5 second and fragments from 40 to 450 Da were maintained. The total running time was 36 minutes.

2.2.3 Interpretation of Mass Spectrum (MS)

In the MS Programme, NIST Version 2.0 library database of National Institute of Standard and Technology (NIST) having more than 2,00,000 patterns was used for identifying the phytochemical components of *C. bonplandianum*. The spectrum of the unknown component was compared with the spectrum of the known components stored in the NIST library. The name, molecular weight and structure of the components of the test materials were ascertained.

3. RESULTS

The name of identified phytocompounds of ethanolic extract of different parts like leaf, fruit and latex of *C. bonplandianum* with their retention time (RT), peak area percentage, molecular weight and their biological activities are given in Table 1-3. The phytocompound prediction is based on NIST Version 2.0 library database of National Institute of Standard and Technology (NIST). The GC-MS results showed that the presence of twenty one major compounds in different parts of *C. bonplandianum*. Among these five phytocompounds such as 16-Hexadecanoyl hydrazide (88.69%), 1,2-Benzenedicarboxylic acid, diisooctyl ester (5.56%), 2-Piperidinone, N-[4-bromo-n-butyl] (2.56%), Phthalic acid, bis (7-methyloctyl) ester

133 (1.80%) and Phytol (1.39%) were found in leaf. The latex of *C. bonplandianum* showed the
 134 presence only one phytocompound Myo-Inositol, 2-C-methyl (30.8%). The fruits of *C.*
 135 *bonplandianum* showed the presence of fifteen major phytocompounds including 9,12,15-
 136 Octadecatrienoic acid, methyl ester (z,z,z)- (41.81%), Diazoprogerone (19.03%),
 137 Decanoic acid, ethyl ester (4.86%), 1-Propene, 2- nitro-3-(1-cyclooctenyl) (4.58%) and
 138 6,9,12- Octadecatrienoic acid, 13-Tetradec-11-yn-1-ol (3.47%).
 139
 140
 141




Fig. 1. GC-MS chromatogram of ethanolic leaf extract of *C. bonplandianum*





Fig. 2. GC-MS chromatogram of ethanolic latex extract of *C. bonplandianum*

156
157
158



159
160
161
162
163

Fig. 3. GC-MS chromatogram of ethanolic fruit extract of *C. bonplandianum*

Table 1. Screening of the phytocompounds of ethanolic leaf extract of *C. bonplandianum* by GC-MS

Sl.No.	RT	Name of the compound	Molecular Formula	MW	Peak Area %	Nature of compound	**Activity
1.	14.97	Phytol	C ₂₀ H ₄₀ O	296	1.39	Diterpene compound	Antimicrobial Antiinflammatory Anticancer Diuretic
2.	20.89	1,2-Benzenedicarboxylic acid, diisooctyl ester	C ₂₄ H ₃₈ O ₄	390	5.56	Plasticizer compound	Antimicrobial Antifouling
3.	24.42	2-Piperidinone, N-[4-bromo-n-butyl]-	C ₉ H ₁₆ BrNO	233	2.56	Alkaloid compound	Antimicrobial Anti-inflammatory
4.	24.79	Phthalic acid, bis(7-methyloctyl) ester	C ₂₆ H ₄₂ O ₄	418	1.8	Plasticizer compound	Antimicrobial Antifouling
5.	30.33	16-Hexadecanoyl hydrazide	C ₁₆ H ₃₄ N ₂ O	270	88.69	Nitrogen compound	Antimicrobial

RT – Retention Time, MW – Molecular Weight, **Dr.Duke`s phytochemical and Ethnobotanical Databases

Table 2. Screening of the phytocompounds of ethanolic latex extract of *C. bonplandianum* by GC-MS

Sl.No.	RT	Name of the compound	Molecular Formula	MW	Peak Area %	Nature of compound	**Activity
1.	10.48	Myo-Inositol, 2-C-methyl	C ₇ H ₁₇ O ₆	194	30.8	Inositol compound	Antimicrobial

RT – Retention Time, MW – Molecular Weight, **Dr.Duke`s phytochemical and Ethnobotanical Databases

Table 3. Screening of the phytocompounds of ethanolic fruit extract of *C. bonplandianum* by GC-MS

Sl.No.	RT	Name of the compound	Molecular Formula	MW	Peak Area %	Nature of compound	**Activity
1.	13.42	Decanoic acid, ethyl ester	C ₁₂ H ₂₄ O ₂	200	4.86	Fatty acid ester	Insecticide compound
2.	15.68	9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)-	C ₁₉ H ₃₂ O ₂	292	41.81	Linolenic acid ester	Antiinflammatory Hypocholesterolemic Cancer preventive Hepatoprotective Nematicide Insectifuge Antihistaminic Antieczemic Antiacne 5-Alpha reductase inhibitor Antiandrogenic Antiarthritic Anticoronary
3.	18.46	1,2:4,5:9,10-Trieoxydecane	C ₁₀ H ₁₆ O ₃	184	2.92	Epoxy compound	No activity reported
4.	19.35	1-Octadecyne	C ₁₈ H ₃₄	250	1.25	Alkene compound	No activity reported
5.	20.79	1-Tridecyne	C ₁₃ H ₂₄	180	1.25	Alkene compound	No activity reported
6.	23.09	2-Decen-1-ol	C ₁₀ H ₂₀ O	156	8.33	Alcoholic compound	No activity reported
7.	24.63	6,11-Dimethyl-2,6,10-dodecatrien-1-ol	C ₁₄ H ₂₄ O	208	2.22	Alcoholic compound	No activity reported
8.	25.8	2-Hexen-1-ol, 2-ethyl-	C ₈ H ₁₆ O	128	0.69	Alcoholic compound	Antimicrobial

9.	27.07	Pseudoephedrine, (+)-	C ₁₀ H ₁₅ NO	165	2.64	Alkaloid compound	Antimicrobial Antiinflammatory Analgesic compound No activity reported
11.	27.76	Cyclohexane, (2-ethyl-1-methylbutylidene)-	C ₁₃ H ₂₄	180	1.94	Alkene compound	No activity reported
12.	30.5	13-Tetradecene-11-yn-1-ol	C ₁₄ H ₂₄ O	208	3.47	Alcoholic compound	No activity reported
13.	30.99	1-Propene, 2-nitro-3-(1-cyclooctenyl)	C ₁₁ H ₁₇ NO ₂	195	4.58	Nitrogen compound	Antimicrobial
14.	32.25	Diazoprogestone	C ₂₁ H ₃₀ N ₄	338	19.03	Steroid compound	Antimicrobial Antiinflammatory Anticancer Antiarthritic Asthma Diuretic
15.	33.74	6,9,12-Octadecatrienoic acid, phenyl methyl ester, (Z,Z,Z)-	C ₂₅ H ₃₆ O ₂	368	2.22	Linolenic acid ester compound	Antiinflammatory Hypocholesterolemic Cancer preventive Hepatoprotective Nematicide Insectifuge Antihistaminic Antieczemic Antiacne 5-Alpha reductase inhibitor Antiandrogenic Antiarthritic Anticoronary

RT – Retention Time, MW – Molecular Weight, **Dr.Duke`s phytochemical and Ethnobotanical Databases

4.DISCUSSION

In the present study twenty one major phytochemicals were identified in different parts like leaf, fruit and latex of *C. bonplandianum*. The biological activities of identified phytochemicals in *C. bonplandianum* were predicted based on Dr. Duke's phytochemical and Ethnobotanical Databases. In terms of peak area percentage of 16-Hexadecanoyl hydrazide was predominant in ethanolic leaf extract where as 9,12,15-Octadecatrienoic acid, methyl ester (z,z,z)- and Diazoprogestosterone were predominant in fruits of *C. bonplandianum*. The compound like Myo-Inositol, 2-C-methyl was identified in latex of *C. bonplandianum*. The leaf of *C. bonplandianum* extract showed five major compounds, among these phytol has antimicrobial, anti-inflammatory, anticancer and diuretic activities, 1,2-Benzenedicarboxylic acid, diisooctyl ester and Phthalic acid, bis (7-methyloctyl) ester compounds have antimicrobial and antifouling activities, 2-Piperidinone, N-[4-bromo-n-butyl] has antimicrobial and anti-inflammatory activities and 16-Hexadecanoyl hydrazide has antimicrobial activity (Table 1). Latex showed that the phytochemical of Myo-Inositol, 2-C-methyl and it has antimicrobial activity (Table 2). The *C. bonplandianum* fruit extract exhibits the phytochemicals like Decanoic acid, ethyl ester has insecticide activity, where as 9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z) and 6,9,12-Octadecatrienoic acid, phenylmethyl ester, (Z,Z,Z) compounds have anti-inflammatory, hypocholesterolemic, cancer preventive, hepatoprotective, nematocidal, insecticide, antihistaminic, antieczemic, antiacne, 5-alpha reductase inhibitor, antiandrogenic, anticoronary and antiarthritic activities, 2-Hexen-1-ol, 2-ethyl and 1-Propene, 2-nitro-3-(1-cyclooctenyl) compounds have antimicrobial activity, Pseudoephedrine has analgesic, antimicrobial and anti-inflammatory activities and Diazoprogestosterone has antimicrobial, anti-inflammatory, anticancer, antiarthritic, antiasthma and diuretic activities (Table 3).

There is growing awareness in correlating the phytochemical components and their biological activities [35,36,37]. Figure-1 showed the highest peak area 88.69% for the phytochemical of 16-Hexadecanoyl hydrazide and lowest peak area 1.39% for the compound of phytol in leaf extract. Phytol is one among the five compounds of the leaf extract were observed in the present study. Similarly, the researchers reported that the presence of phytol in the leaf of other plants like *Lantana camara*, *Mimosa pudica* and *Erythralium scandens* [38-41]. The antibacterial activity of phytol against *Staphylococcus aureus* was reported and it cause damage the cell membrane as a result there is a leakage of potassium ions from bacterial cells [42]. Phytol constitute a promising novel class of pharmaceuticals for the treatment of rheumatoid arthritis and possibly other chronic inflammatory diseases [43]. Phytol is an acyclic diterpene alcohol and it is also a precursor for vitamins E and K1. Different solvent extracts of leaf of *C. bonplandianum* showed effective inhibition activity against bacteria and fungi [44,45]. *In vitro* anti-inflammatory activity of ethanolic leaf extract of *C. bonplandianum* was reported [46]. It is also used as antiseptic and antidote [22,23,24]. Tumour formation by *Agrobacterium* in potato plants was distinctly inhibited by the methanolic leaf extract of *C. bonplandianum* [29].

As shown in figure-2, the peak area 30.8% for Myo-Inositol, 2-C-methyl was observed in the latex extract of *C. bonplandianum* and it possesses antimicrobial activity was recorded in Dr.Duke's phytochemical and ethnobotanical database. Similarly, the presence of Myo-Inositol, 2-C-methyl was reported in *Prosopis spicigera* [47]. The antifungal activity of latex of *C. bonplandianum* against *Microsporium gypsum* and *Trichophyton mentagrophyt* was reported that [18]. The medicinal plants have healing effect on wounds and cuts [25,26] and it may be due to their antimicrobial activity which seems to be responsible for wound contraction and increased rate of epithelialisation [48].

Figure-3 showed that the highest peak area 41.81% for the 9,12,15-Octadecatrienoic acid, methyl ester (z,z,z)- in fruit extract which is an Omega-3-fatty acid and this compound have the anti-inflammatory and antiarthritic properties [49,50]. Similarly 9, 12, 15-Octadecatrienoic acid methyl ester (z,z,z)- identified in the ethanol leaf extract of *Aloe vera* and *Vitex negundo* [51,52]. Omega-3-fatty acids have been found to be essential for normal growth and development. They play an important role in the prevention and treatment of coronary artery disease, hypertension, diabetes, arthritis, inflammation, autoimmune disorders and cancer [53]. The hepatoprotective compounds like 9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)- and 6,9,12-Octadecatrienoic acid, phenyl methyl ester, (Z,Z,Z)- were observed in the fruits of *C. bonplandianum*. *C. bonplandianum* is used for the treatment of jaundice [20] and liver diseases [45]. The plant has hypotensive effect [54] and also used in the treatment of high blood pressure [22,23,24]. The extracts of this plant showed antimicrobial and antitumour activities [55]. *C. bonplandianum* is also considered as chologogue and purgative [56]. The fresh juice of *C. bonplandianum* is used in the treatment of headache by ethnic groups [21]. *C. bonplandianum* has hepatoprotective and antihelmenthic properties [21,22,23] and also posses repellent property against the insects [21,57]. The lowest peak area 0.69% was observed for 2-Hexen-1-ol, 2-ethyl in the fruit extract of *C. bonplandianum*. It is used as Ayurvedic medicine, however there are no reports on phytochemical analysis in fruits of *C. bonplandianum* using GC-MS. The researchers were reported eleven phytochemicals in chloroform leaf extract and sixteen phytochemicals in ethanolic leaf extract of *C. bonplandianum* [58,59]. But in the present study, we found only five compounds in the ethanolic leaf extract, among these the compound phytol is commonly reported in both studies remaining four compounds are newly reported. Fifteen compounds in fruits extract and one compound in the latex extract were newly reported in this study. In the present study, we screened totally twenty one phytochemicals from different parts like leaf, fruit and latex of *C. bonplandianum* by GC-MS analysis. Up to our knowledge, this study may be a first report on the analysis of phytochemicals in different parts especially fruits and latex of *C. bonplandianum* using GC-MS analysis. So, the present study is the first step to find out the nature of active principles in different parts of *C. bonplandianum* and this study will also be helpful for further research.

5. CONCLUSION

The medicinal plant *C. bonplandianum* could be a potential source for useful drugs like anti-inflammatory, anticancer, antimicrobial, insectifuge, nematicide, anticorony, wound healing, hepatoprotective activities. So, further research is needed to isolate, identify, characterize and elucidate the structure of these bioactive compounds responsible for medicinal values of *C. bonplandianum*.

REFERENCES

1. Ncube NS, Afolayan AJ, Okoh AI. Assessment techniques of antimicrobial properties of natural compounds of plant origin: Current methods and future trends. *Afr J Biotechnol.* 2008;7:1797-1806.
2. Boopathi AC, Sivakumar R. Phytochemical screening studies on the leaves and stem of *Andrographis neesiana* wight - An endemic medicinal plant from India. *World App Sci J.* 2011;12(3):307-311.
3. Kalimuthu K, Prabakaran R. Preliminary phytochemical screening and GC-MS analysis of methanol extract of *Ceropegia pusilla*. *International journal of Research in Applied, Natural and Social Sciences.* 2013;1(3),49-58.

- 104 4. Vinoth S, Rajesh Kanna P, Gurusaravanan P, Jayabalan N. Evaluation of
105 phytochemical, antimicrobial and GC-MS analysis of extracts of *Indigofera trita* L. Int J
106 Agric Res. 2011;6(4):358-367.
- 107 5. Savithramma N, Venkateswarlu P, Suhrulatha D, Basha SKM, Venkataramanadevi CH.
108 Studies of *Boswellia ovalifoliolata* Bal. and *Henry* - An endemic and endangered
109 medicinal plant. *The Bioscan*. 2010;5:359-362.
- 110 6. Turker AU, Usta C. Biological screening of some Turkish medicinal plants for
111 antimicrobial and toxicity studies. Nat Prod. 2008;22:136-146.
- 112 7. Sheeja K, Kuttan G. Activation of cytotoxic T lymphocyte responses and attenuation of
113 tumour growth *in vivo* by *Andrographis paniculata* extract and andrographolide.
114 Immunopharmacol Immunotoxicol. 2007;29:81-93.
- 115 8. Mukherjee PK, Kumar V, Houghton PJ. Screening of Indian medicinal plants for acetyl
116 cholinesterase inhibitory activity. Phytother Res. 2007;21:1142-1145.
- 117 9. Savithramma N, Linga Rao, Beenaprabha M. Phytochemical studies of *Dysophylla*
118 *myosuroides* (Roth.) Benth. In. wall. and *Talinum cuneifolium* (Vahl.) Willd. Res J Phyto.
119 2011;5(3):163-169.
- 120 10. Parekh Jigna, Chanda V Sumitra. *In vitro* antimicrobial activity and phytochemical
121 analysis of some Indian medicinal plants. Turk J Biol. 2007;31:53-58.
- 122 11. Kumar A, Ilavarasan R, Jayachandran T, Decaraman M, Aravindhan P,
123 Padmanaban N, Krishna MRV. Phytochemical investigation on a tropical plant. Pak J
124 Nutri. 2009;8:83-85.
- 125 12. Neelam Singh, Mukesh Kumar Meena, Vidya Patni. Phytochemical profiling and GC-MS
126 analysis of bioactive constituents callus of *Naringi crenulata*. Int J Pharm Sci Rev Res.
127 2014;6:29-34.
- 128 13. Milne A. Inhalational and local anesthetics reduce tactile and thermal responses in
129 *Mimosa pudica* Linn. Masui. 1993;1190-1193.
- 130 14. Steinmetz KA, Potter JD. Vegetables, fruits, cancer prevention: A review. J Am Diet
131 Assoc. 1996;1027-1039.
- 132 15. Nishaa S, Vishnupriya M, Sasikumar JM, Gobalakrishnan VK. Phytochemical screening
133 and GC-MS analysis of ethanolic extract of rhizomes of *Maranta arundinacea* L. *Res J*
134 *Pharm BiolChemSci*. 2013;2:52-59.
- 135 16. Jeeshna MV, Mallikadevi T, Paulsamy S. Screening of the weed plant species, *Croton*
136 *bonplandianum* Bail. for larvicidal activity of *Aedes aegypti*. J Biopesticides.
137 2010;3(1):192 – 194.
- 138 17. Vadlapudi V. *In vitro* antimicrobial activity of methanolic extract of selected Indian
139 medicinal plants. Pharmacophore. 2010;1(3):214-219.
- 140 18. Asthana A, Mall HV, Dixit K, Gupta S. Fungitoxic properties of latex of plants with
141 special reference to that of *Croton bonplandianum* Bail. Pharm Biol. 1989;27(1):25-
142 28.
- 143 19. Divya S, Naveen Krishna K, Ramachandran S, Dhanaraju MD. Wound healing and *In*
144 *vitro* antioxidant activities of *Croton bonplandianum* leaf extract in rats. Global J
145 Pharmacol. 2011;5(3):159-163.
- 146 20. Saggoo MIS, Walia S, Kaur R. Evaluation of genotoxic and antimicrobial potential of
147 *Croton bonplandianum* Bail. Arch Appl Sci Res. 2010;2(2):211-216.

- 148 21. Maria CMT, Joao CA, Gilvandete MPS, Manoel AN, Edilberto RS, Leticia VCL, Daniel
149 PB, José DBMF, Francisco AV, Otilia DLP. Larvicidal and nematocidal activities of the leaf
150 essential oil of *Croton regelianus*. J Chem Biodiv. 2008;5(12):2724-2728.
- 151 22. Bhakat RK, Sen UK. Ethno medicinal plant conservation through sacred groves. Tribes
152 and Tribals. 2008;2:55-58.
- 153 23. Chaudhuri AB. Endangered medicinal plants. Daya publishing House, Delhi. 2007;226.
- 154 24. Nishanta R, Harris CS, Towers GHN. Antimicrobial activity of plants collected from
155 serpentine outcrops in Sri Lanka. Pharm Biol. 2002;40(3):235-244.
- 156 25. Das AJ, Dutta BK, Sharma GD. Medicinal plants used by different tribes of Cachar
157 district, Assam. Indian J Trad Knowledge. 2008;7(3):446-454.
- 158 26. Chandel KPS, Shukla G, Sharma N. Biodiversity in medicinal and aromatic plants in
159 India. National Bureau of Plant Genetic Resources, New Delhi, 1996.
- 160 27. Reddy KR. Folk medicines from Chittor District Andra Pradesh, used in treatment of
161 jaundice. Pharmaceutical Biol. 1995;26:137-140.
- 162 28. Pranab Ghosh, Amitava Mandal, Mohammad Golam Rasul. A new bioactive ursane-type
163 triterpenoid from *Croton bonplandianum* Bail. J Chem Sci. 2013;125(2):359-364.
- 164 29. Islam MS, Rahman MM, Rahman MA, Qayum MA, Alam MF. *In vitro* evaluation of *Croton*
165 *bonplandianum* Bail. as potential antitumor properties using *Agrobacterium*
166 *tumefaciens*. J Agri Tech. 2010;6(1):79-86.
- 167 30. Selvamangai G, Anusha Bhaskar. GC-MS analysis of phytocomponents in the
168 methanolic extract of *Eupatorium triplinerve*. Asian Pac J Trop Biomed. 2012;S1329-
169 S1332.
- 170 31. Oyi R, Onaolapo JA, Haruna AK, Morah CO. Antimicrobial screening and stability studies
171 of the crude extract of *Jatropha curcas* (L) latex. Nig J Pharmac Sci. 2007;6(2):14-20.
- 172 32. Arekemase MO, Kayode RMO, Ajiboye AE. Antimicrobial activity and phytochemical
173 analysis of *Jatropha curcas* plant against some selected microorganisms. Int J Biol.
174 2011;3(3):52-59.
- 175 33. Janakiraman N, Johnson M, Sahaya Sathish S. GC-MS analysis of bioactive constituents
176 of *Peristrophe bicalyculata* (Retz.) Nees. (Acanthaceae). Asian Pac J Trop
177 Biomed. 2012;S46-S49.
- 178 34. Gopalakrishnan S. Vadivel E. GC-MS Analysis of some bioactive constituents of
179 *Mussaenda frondosa* Linn. Int J Pharm Bio Sci. 2011;2(1):313-320.
- 180 35. Fernie AR, Trethewey RN, Krotzky AJ and Willmitzer L. Metabolite profiling: from
181 diagnostics to systems biology. Nat Rev Mol Cell Biol. 2004;5:763-769.
- 182 36. Summer LW, Mendes PJ, Dixon RA. Plant Metabolomics: Large-scale phytochemistry in
183 the functional genomics. Phytochemistry. 2003;62(6):817-836.
- 184 37. Robertson DG. Metabolomics in toxicology: A review. J Toxicol Sci. 2005;85:809-822.
- 185 38. Maria Jancy Rani P, Kannan PSM, Kumaravel S. GC-MS analysis of *Lantana*
186 *camara* L. leaves. Journal of pharmaceutical Research and Development. 2011; 2(11):
187 63-66.
- 188 39. Sridharan S, Meenaa V, Kavitha V, Agnel Arul John Nayagam. GC-MS study and
189 phytochemical profiling of *Mimosa pudica* Linn. J Pharm Res. 2011;4(3):741-742.
- 190 40. Sathish kumar M, Manimegalai S. Evaluation of larvicidal effect of *Lantana camara*
191 Linn against mosquito species *Aedes aegypti* and *Culex quinquefasciatus*. Advan.
192 Biol. Res. 2008;2(3-4):39-43.

193 41. Sutha S, Kalpana Devi V, Mohan VR. GC-MS determination of bioactive components of
194 *Erythralum scandens* Bl., Bijdr. J App Pharm Sci. 2011; 01(9):170-173

195 42. Inoue Y, Hada TA, Shiraishi K, Hirore, Hamashima H, Kobayashi S. Biphasic effects
196 of Geranylgeraniol, Terpenone and Phytol on the growth of *Staphylococcus*
197 *aureus*. Antimicrob Agents and Chemother. 2005;49(5):1770-1774.

198 43. Ogunlesi M, Okiei W, Ofor E, Osibole AE. Analysis of the essential oil from the
199 dried leaves of *Euphorbia hirta* Linn (Euphorbiaceae), a potential medication for
200 asthma. African J Biotech. 2009; 8: 7042-7050.

201 44. Jeeshna MV, Paulsamy S, Mallikadevi T . Phytochemical constituents and antimicrobial
202 studies of the exotic plant species, *Croton bonplandianum* Bail. J Life Sci. 2011;3(1):23-
203 27.

204 45. Bapuji WJ, Ratnam VS. Ethnobotany. Leaflets. 2009;13:388-389.

205 46. Thenmozhi M, Vasuki K, Dhanalakshmi M, Devi K, Manjula. Evaluation of anti
206 inflammatory potential of *Croton bonplandianum* Bail leaf. Int J Pharm Bio Sci.
207 2013;7(1):9.

208 47. Siddabathuni Aneela, Akalanka Dey, Somnath De. GC-MS analysis of methanolic extract
209 of *Prosopis spicigera*. International Journal of Phytopharmacology. 2014; 5(3): 168-171.

210 48. Dash GK, Murthy PN. Evaluation of *Argemone Mexicana* Linn leaves for wound healing
211 activity. J Nat Prod Plant Resour. 2011;1(1):46-56.

212 49. Lalitharani S, Mohan VR, Regini GS, Kalidass C. GC-MS analysis of ethanolic
213 extract of *Pothos scandens* L. leaf. J Herb Medi Toxicology. 2009;3:159-160.

214 50. Maruthupandian A, Mohan VR. GC-MS analysis of ethanol extract of *Wattakaka*
215 *volubilis* (L.f) Stapf. leaf. Int J Phytomedicine. 2011;3:59-62.

216 51. Arunkumar S, Muthuselvam M. Analysis of phytochemical constituents and
217 antimicrobial activities of *Aloe vera* L. against clinical pathogens. World J
218 Agricultural Sci. 2009;5(5):572-576.

219 52. Praveen kumar P, Kumaravel S, Lalitha C. Screening of antioxidant activity, total
220 phenolics and GC-MS study of *Vitex negundo*. Afr J Biochemistry Res. 2010;4 (7):191-
221 195.

222 53. Simopoulos AP. Omega-3 fatty acids and antioxidants in edible wild plants. Biol Res.
223 2004; 37: 263–277.

224 54. Asolkar LV, Kakkar KK, Chakra OJ. Second supplement to glossary of Indian medicinal
225 plants with active principles. Part-1 (A-K). Publication and Information Division (CSIR),
226 New Delhi, 1992.

227 55. Ramesh VM, Hilda A, Manjula VK. Fungi toxic effect of leaf extract of *Croton sparciflorus*
228 on phytopathic fungi. Acta Botanica Indica. 1995;23:63-66.

229 56. Dutt N, Kumar D, Singh S, Mahajan R. Taxonomy, ethnobotany and antimicrobial activity
230 of *Croton bonplandianum*, *Euphorbia hirta* and *Phyllanthus fraternus*. J Adv Dev Res.
231 2011;2(1):21-29

232 57. Krebs HC, Miarantsoa H. Clerodane diterpenes of *Croton hovarum*. Phytochemistry.
233 1997;40: 931.

234 58. Keerthana G, kalaivani MK, sumathy A. *In-vitro* alpha amylase inhibitory and anti-oxidant
235 activities of ethanolic leaf extract of *Croton bonplandianum*. Asian J Pharm Clin Res.
236 2013;6(4):32-36.

237 59. Kalaivani M, Bhavana J, Sumathy A. GC-MS analysis of chloroform extract of *Croton*
238 *bonplandianum*. Int J Pharm Bio Sci. 2013;4(4):613-617.