

Mini-review:

The influence of respiratory and pH imbalance in cancer development ~~Respiratory and pH imbalance at the early precancerous stage~~

Abstract

Carcinogenesis is a process by which normal cells are transformed into cancer cells. Cancer is a multifactorial disease with contributions from environmental, genetic, and lifestyle factors. Cancer prevention is a global priority, yet the proximate causes of most cancers are still little understood ~~even though this is what we want to learn how to avoid~~. In this review, we bring further evidence in support of the hypothesis on the so-called respiratory and pH imbalance (RpHI) as a cause of long standing hypoxia within the whole organism, characteristic to the ~~preneoplastic preneoplastic precancerous~~ stage. Since hypoxia depends on the intensity and duration of action of the stress agent, human and animal organisms may compensate hypoxia only if the causal agent stop acting continuously. In brief Moreover, under hypoxic conditions, the overbusy cells, getting less oxygen than needed, turn into anaerobic adenosine triphosphate (ATP) production, excessive multiplication and, finally, tumor development. Cancer prevention might be associated with an increase in oxygen delivery to overstressed cells, carbon dioxide removing, as well as pH and glucose concentration balancing. The role of RpHI in chemical carcinogenesis and ~~cancer incidence most cancer diseases~~ is also discussed. Evidence from the literature data is brought to show also suggest that cancers develop at the physiological level, while the molecular changes in cancer cells are the consequence and not the main cause of malign processes. A link between the Warburg effect and the Macovschi's biostructural theory was suggested.

Formatted: Justified

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Not Bold

Formatted: Not Highlight

Keywords: Preneoplastic stage~~Precancerous stage~~; Cancer prevention; Hypoxia; Respiratory and pH imbalance-RpHI; Cachexia; The Warburg effect~~;~~ Warburg's theory.

Formatted: Space Before: 0 pt, After: 0 pt, Line spacing: 1.5 lines

Introduction

Ancient Asian medicine was mostly a preventive medicine, while current western medicine is mainly a curative one. However, the query still remains: should the medical act balance the body or repair the already occurred damage? Imagine that we want to prevent cancer by administering the anti-cancer drug, cyclophosphamide. In a few years we will have become ill from cancer! Presently, the gap between the two medical approaches seems to be even more evident.

Cancer prevention and etiology are medical priorities, but history study suggests that the journey towards achieving this goal is difficult and full of detours and roadblocks [1].⁺ The proximate causes of most cancers are still less understood even though this is what we want to learn how to avoid [2].² For instance, it appears difficult to understand why does it have to be such a long time interval between the exposure of a certain cell to a mutagen and the expression of the resulting mutations. It is hard to imagine how the numerous genetic changes found in cancer cells could have been produced in any cell as the result of a single exposure to a DNA-damaging agent, or why months or years have to elapse before the effect of these changes are observed. Malignant tumors exhibit increased dependence on glycolysis, resulting in abundant export of lactic acid, a hypothesized key step in carcinogenesis [3,4],~~3,4~~ It is also well-known that the development of cancer has been associated with microbial infection, injury, and inflammation and tissue repair [5-7],~~5-7~~ Chronic inflammation conditions, caused by exposure to environmental factors can increase the risk of cancer [8-10],~~8-10~~ Most cancer cases are mainly attributable to long-term exposure to environmental factors, via a multi-year, multi-step, and multi-path process of tumorigenesis [11,12],~~11,12~~ They may involve cumulative genetic and epigenetic alterations in the chronic carcinogenesis of body cells from a non-cancerous stage to precancerous and cancerous stages [13],~~13~~ Chemopreventive effects of dietary

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

59 phytochemicals used at the precancerous stage can hardly be explained in light of current
60 theories [14].⁴⁴ Several dietary phytochemicals such as β -carotene, curcumin, resveratrol,
61 gallic acid, chlorogenic acid, caffeic acid, carnosol, capsaicin, 6-shogaol, 6-gingerol, and
62 their related derivatives have been largely investigated against cancer invasion and
63 metastasis [15,16].^{45,16} Since carcinogenesis is a multistep process, these bioactive
64 compounds might act on a variety of stages of the precancerous process to prevent
65 inflammation, enzyme activities, or lactic production [17,18].^{47,18} Evidence for cancer-
66 related anti-inflammatory activity of natural and synthetic phenols was provided by
67 Aspirin, which inhibits colorectal carcinogenesis [19].⁴⁹

68 ———The role of metabolic reprogramming in carcinogenesis is being increasingly
69 recognized [Schulze & Harris, 2012]. The identification of metabolic processes of cancer
70 cells has been used to create strategies for treating cancer. Warburg stated that there are
71 fundamental differences in the central metabolic pathways operating in tumorigenesis [21-
72 25].^{Wu W, Zhao, 2013, 20-22} In recent years, the metabolism of malignant tissue has once again
73 become one of the most intense areas of research in cancer biology [26-29].^{Upadhyay et al, 2013;}
74^{23 Gogvadze et al, 2010; Ferreira, 2010;} Considering all these issues, new ideas are needed to clarify
75 what happens during carcinogenesis. In spite of the fact that Otto Warburg studied aerobic
76 glycolysis long ago, the molecular mechanisms how increased glycolysis is regulated by
77 oncogenic and/or tumor suppressive signaling pathways are unclear [Hitosugi & Chen,
78 2014³⁰].

79 Starting with the observation that the cellular structure is partially broken down
80 during hypoxia, a so-called respiratory and pH imbalance (RpHI) was supposed to be
81 involved in cancer etiology [31-33].²⁴⁻²⁶

82 Therefore, this work aims at presenting evidence for the role of RpHI during the
83 precancerous stage, as well as some concluding remarks that could be useful to design new
84 preventive strategies.

85 **RpHI hypothesis**

86 **RpHI, a definition**

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Formatted: Space Before: 0 pt, After: 0 pt

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Font: Bold

88 Normally, various stress agents, such as physical and chemical stressors, viruses,
89 other infectious agents, hormones in excess, but also long lasting anxiety, emotions,
90 conflict states, etc., are able to affect the body as a whole [34-37]. In order to preserve the
91 identity and relative independence of living organisms confronted with the external
92 environment, some co-ordinate physiological processes are involved in keeping their
93 internal states of equilibrium [38-40]. The living bodies react against aggressions using
94 metabolic energy obtained by the oxidation of organic substances, including organic acids
95 in the mitochondria [41,42]. Under normal conditions, when the supply of oxygen is
96 sufficient, cells can carry out aerobic respiration [43,44]. Hence, when busy, cells produce
97 the energy they need from the food stores, including the organic acids they possess. During
98 the Krebs cycle, carbon dioxide, of a weaker acid type, is released, normally out [45,46].
99 Since stronger organic acids such as succinic, malic, 2-ketoglutaric and oxalylacetic acids
100 are replaced by carbon dioxide, the cell milieu tends to be more alkaline, just blood and
101 urine [47]. The tendency of blood alkalization entails retaining of carbon dioxide, so that
102 the pH of blood should alter as little as possible. The retaining of carbon dioxide in blood
103 does not allow the oxygen shift at a normal rate in lung [48,49]. As a consequence, oxygen
104 pressure in blood decreases albeit it stays normal all around. However, oxygen may reach
105 quite a low level without affecting cell breathing in any drastic way. Unfortunately, the
106 busiest cells in the body need more oxygen. Oxygen partial pressure getting lower, these
107 cells receive less oxygen than they need, and fermentative processes develop alongside
108 with breathing.

109 In principle, the ~~RpHIrespiratory and pH imbalance~~, which can be regarded as a
110 physiological reaction against any stress agents, may cause an oxygen crisis within the
111 body, and, consequently, the busiest cells divide faster and faster producing first
112 preneoplastic cells, and, in time, malignant tumors~~faster producing benign or malignant~~
113 ~~tumors~~. The tumors thus acquired, as well as their fermentation products, may disturb the
114 normal functions of most cells and tissues in the body [50].²⁷

115 ~~RpHI Respiratory and pH imbalance~~ was discovered by putting together the main
116 experimental data in cancer with the most important findings in physiology, biochemistry,
117 cybernetics, etc., and treating them as a whole; it represents the mechanism that could

Formatted: Font: Bold

create a long standing intracellular hypoxia state, which, over time, generates the conditions for malignant transformation [51,52].^{28,29} Nevertheless, it is hard to believe that a tumor is generated starting from a physiological imbalance, prior to the first real malign cell, while the modern biology is looking for precise molecular mechanisms involved in carcinogenesis. However, the impairment of cellular breathing, altered intracellular and extracellular pH, as well as hypoxic conditions might mostly be related to a precancerous pathology. Besides, cancer-associated impairment of respiration and changes in pH are less explored than the genetic mutations.

—Therefore, this work aims at presenting evidence for the role of RpH during the precancerous stage, as well as some concluding remarks that could be useful to design new preventive strategies.

CA hypothesis on cancer-related physiological changes

Since the oncogene revolution pushed tumor metabolism to the margins of cancer research [23,53],^{20,30} interest in the fundamental differences of central metabolic pathways operating in malignant tissue has only been renewed in the recent years. The potential causes of deranged metabolism in cancer should be discussed with emphasis on changes in energy metabolism of glucose, fat and protein [53].³⁰

In brief, various stress agents, such as physical and chemical stressors, viruses, other infectious agents, hormones in excess, but also long-lasting anxiety, emotions, conflict states, etc., are able to affect the body as a whole.³¹⁻³⁴ In order to preserve the identity and relative independence of living organisms confronted with the external environment, some co-ordinate physiological processes are involved in keeping their internal states of equilibrium.³⁵⁻³⁷ The living bodies react against aggressions using metabolic energy obtained by the oxidation of organic substances, including organic acids in the mitochondria.^{38,39} Under normal conditions, when the supply of oxygen is sufficient, cells can carry out aerobic respiration.^{40,41} Hence, when busy, cells produce the energy they need from the food stores, including the organic acids they possess. During the Krebs cycle, carbon dioxide, of a weaker acid type, is released, normally out.^{42,43} Since stronger organic acids such as succinic, malic, 2-ketoglutaric and oxalylacetic acids are replaced by carbon

dioxide, the cell milieu tends to be more alkaline, just blood and urine.⁴⁴ The tendency of blood alkalization entails retaining of carbon dioxide, so that the pH of blood should alter as little as possible. The retaining of carbon dioxide in blood does not allow the oxygen shift at a normal rate in lungs.^{45,46} As a consequence, oxygen pressure in blood decreases albeit it stays normal all around. However, oxygen may reach quite a low level without affecting cell breathing in any drastic way. Unfortunately, the busiest cells in the body need more oxygen. Oxygen partial pressure getting lower, these cells receive less oxygen than they need, and fermentative processes develop alongside with breathing.

However, hypoxia is widely considered as the natural environment in which life appears and proliferates and DNA auto-replication and transcription takes place *in vivo* in all eukaryotes [54-57].⁴⁷⁻⁵⁰ Nuclear division unfolds anaerobically using the energy produced by glycolysis [58-60].⁵¹⁻⁵³ Consequently, the cells forced to manifest themselves in rather anaerobic conditions will divide more intensely [61].⁵⁴ Aerobic breathing, which provides cells with a great amount of energy, creates the necessary conditions for the existence of fine structures of the cells, the specific functions running unimpaired [62].⁵⁵ Lack of oxygen, even partial, causes rupture of these structures, hence the gradual disappearance of specific functions of cells, as well as contact inhibition, while it entails cell-division to a greater extent than necessary for the tissue in question [63].⁵⁶ Then comes the first stage of a so-called respiratory and pH imbalance the RpHI, which affects the cell respiration and division. Nevertheless, the organism possesses buffer systems, lung ventilation and the kidney mechanisms to control the concentration of hydrogen ions within the cellular milieu (Figure 1) [64].⁵⁷ Alveolar ventilation is responsible for carbon dioxide elimination [65]. A slight respiratory acidosis results primarily when alveolar ventilation is decreased or if carbon dioxide production is increased [65]. Lung ventilation is only able to control the blood pH, and not the oxygen pressure.⁵⁸ However, the organism has several compensatory systems to minimize a decrease in pH. For example, deoxygenated hemoglobin readily accepts hydrogen ions to prevent substantial changes in pH. Normally, carbon dioxide, a respiratory stimulant, induces an increase in minute ventilation to normalize the pH by eliminating increased quantities of CO₂. Unfortunately, this effect is attenuated if the carbon dioxide concentrations remain elevated for more than several hours. The kidneys are

Formatted: Highlight

also capable to control both the blood pH and some other blood parameters; however, this process is slow and lasts for several hours or days. In fact, renal compensation for begins in 6-12 hours, but maximal compensation occurs in 3-5 days. The kidneys increase excretion of hydrogen ions predominantly in the form of ammonium. If the stress agents act continuously, the blood will become slightly more alkaline than usual, and the blood oxygen concentration will be lower than normal. The overstrained cells, getting less oxygen than needed, will lead to anaerobic fermentation and also to a lower content of $\text{NADH}+\text{H}^+$ and $\text{NADPH}+\text{H}^+$, and a higher one of NAD^+ and NADP^+ . Therefore, a decrease of the oxidation-reduction potential will occur as well. The quantity of sulphhydryl groups in the blood and tissues also decreases. Besides, marked decrease in succinic dehydrogenase and slight increase in cytochrom oxidase levels could be found, suggesting the alteration of Krebs cycle. For this reason, the cell with an excessive fermentation will not reach an upper energetic state, if the neighbor cells and blood cannot interfere with its metabolism.

The second stage of the ~~RpHI respiratory and pH imbalance~~ is reached when lactic acid is produced due to hypoxic conditions. This time, the CO_2 concentration may decrease; however, part of the produced carbon dioxide is not removed, because the cell content may remain slightly alkaline in spite of the lactate production. Again, the kidneys should control the hydrogen concentration in blood, releasing acidic species, such as ammonium ions or phosphates NaH_2PO_4 , into urine. The process is complicated by the existence of lactic acid in blood which decreases blood pH, while intracellular pH of overstrained cells is increased.




Figure 1. Suggestive scheme for the role of the Respiratory and pH Imbalance (RpHI) in cancer development. Relative RpHI means both the relative intensity and duration of respiratory and pH imbalance within the body, whereas local/cellular stress may have different intensities or is variably long standing. Schematic presentation of the role of the Respiratory and pH Imbalance (RpHI) in the precancerous stage.

However, if the stress agents act continuously, the neighbor cells involved in curing or rebalancing the overbusy cells will work hard, also being overstrained and deprived of oxygen. There follows a third stage of the RpHIrespiratory and pH imbalance in which a real state of illness (infections, viruses) occurs. A very special balance between the two types of cell (attacked cells and neighbor ones) -is established. As a result, the pH value of the blood is little altered. If the stress agent is very strong, it can be lethal to the organism, and this one does no longer reach the cancer stage. A stronger action of the stress agents could kill the body at this stage. It is the case of microorganism- or virus-induced diseases, which, if untreated, have a bad prognostic. Contrary, a long-standing, still mild action of the stress agents may cause a slow shift of this imbalance, resulting in a stepwise decrease of blood pH value. Long standing infectious pathologies may thus result in a RpHI, which create the conditions for either the transformation of normal cells into preneoplastic cells followed by preneoplastic to neoplastic pathway or for the multiplication of malignant cells. It is well-known that only the bodies with a significant amount of morbidity may become cancerous. Most cells become glycolytic while fewer remain normal, but overstressed. This is the pathway to reach the fourth stage of the RpHIrespiratory and pH imbalance, which is that of tumor formation.

Formatted: Highlight

238 The ~~is so-called RpHI~~respiratory and pH imbalance causes an oxygen crisis and the
239 busiest cells divide faster and faster under anaerobic conditions. The preneoplastic cells
240 thus appeared may become neoplastic, which, in their turn, will generate tumors, if the
241 imbalance persists; these ones ~~thus acquired~~ and their fermentation products can kill the
242 body, if they are not surgically or by other means removed. The ~~RpHI~~respiratory and pH
243 ~~imbalance~~ cannot be easily measured, since the affected individuals seem to be normal and
244 look like they are in good health. Otherwise, if stress agents are more powerful, patients
245 might die during the first stages of body interaction with the environment stressors, without
246 reaching the cancerous stage ~~(IV)~~. At this stage neoplastic cells and tumors may be found in
247 the body. To reach cancer, the RpHI should be less strong, yet long-standing in order to get
248 a chronic lack of oxygen in the blood and cells. Cancer could be considered an impairment
249 of the whole body, while malignant cells are quite normal cells that, ~~because although are~~
250 overstrained, get less oxygen than needed and turn into anaerobic fermentation, being
251 divided to a greater extent than what the body requires.

252 The local/cellular stress means the reaction of the individual cell against the external
253 environmental agents. There is a reciprocal relationship between the ~~RpHI~~respiratory and
254 ~~pH imbalance~~, which affects the entire body and the local solicitation, which created a local
255 (in a target cell or tissue) RpHI. Thus, an imbalanced body, found in apparently good
256 health, will die when confronted to a less important local solicitation (~~Cancer~~ ~~Imbalanced~~
257 ~~body, suitable environment for malign transformation~~). On the other hand, as in the case of
258 experimental chemically produced tumors, a long-standing local stress induced by the
259 carcinogen may cause a general imbalance, which could also promote cancer (Figure 1;
260 ~~Cancer~~ ~~H~~). There are many intermediary states between these extreme situations, giving
261 cancer its characteristic complexity. The ~~RpHI~~respiratory and pH imbalance caused by the
262 difficulty of the lung-kidney system to remove the excess of CO₂ and alkaline ions
263 produced during the stress reaction means/determines a general weakness of the entire
264 body. At the same time, such degree of weakness may be associated either with cancer or
265 other diseases.

266 The living body can be considered a relatively independent entity, which possesses
267 the physiological mechanisms to preserve its specific structure in order to adapt to the

environmental conditions. The [R_pH_irespiratory and pH imbalance](#) is the expression of exceeding the adaptive capability of the body. Generally, the body fights against the alien disturbing agents through a well-known phenomenon, involving the humoral immunity and the capacity of immune response mediated by cells. At the same time, the organism possesses metabolizing and transformation systems, which transform alien and complex substances into simpler ones to be released.

The body must also release its own excessive cells because they are no longer necessary for the organism structure and functions. This process is more clearly understood if one considers wound healing. Thus, the injury puts the inner cells and the external environmental agents together. To react against the changes of environmental conditions, these cells use energy, alkalize the surrounding cell environment, get less oxygen and pass to an excessive fermentative (glycolysis) state. Indeed, because of the fundamental role of oxygen in metabolism, a drop in tissue oxygen level to the point where oxygen demand exceeds supply leads rapidly to metabolic crisis [\[43\]](#).⁴⁰

They divide excessively even after the healing of the wound, but not for too long. Then several cells, which have multiplied in excess for that tissue and are being no longer necessary, are released; thus, the body's own structure is restored. This process takes place in balanced bodies only. Otherwise, the useless cells divide continuously, becoming cancerous. Their neighbors cannot destroy them or help them integrate into the body.

In order for maintaining its specific structure, the body tends to remove the existing malignant cells, too, even if they are its own cells. This process seems to be dependent on the size of the [R_pH_irespiratory and pH imbalance](#). The surgical removing off the tumor cells causes a decrease in this imbalance. As a result, both the immunity and the capacity of restoration of the body's own structure, will increase because the tumors are strong stress agents, as well.

Cancer cells can thus be considered as normal; however, they behave abnormally under hypoxic conditions. Consequently, carcinogenesis only develops in exhausted organisms, hence where a [R_pH_irespiratory and pH imbalance](#) occurs. A carcinogenic promoter acts in a similar way as those agents that provoke a wound, but lasts longer; a local [R_pH_irespiratory and pH imbalance](#) causing cellular hypoxia develops.

Chemical carcinogenicity and the abnormal metabolism of carcinogens

The carcinogenic substances may cause a RpHI, which respiratory and pH imbalance that causes in its turn a slight intracellular hypoxia, which this one provides the normal specific conditions for expressing the ancestral gene existing in every cell. It follows that a carcinogen entering a living tissue must be released at any price, because the affected living cell will become exhausted in its effort to destroy and release that compound and its metabolites. Also, this effort is subsequently assumed by the neighbor cells and a RpHI respiratory and pH imbalance at the level of the whole organism takes place. The RpHI will cause in its turn the cellular hypoxia that represents the specific medium of for malignant transformation.

The degradation pathway of carcinogens is based on oxidation reactions, particularly those classified as microsomal monooxygenation reactions. These reactions occur with oxygen as the oxidizing agent, generating a long standing hypoxia if carcinogen degradation is a multi-step process. Many carcinogens are polyaromatic hydrocarbons that required activation to electrophiles to form covalent adducts with cellular macromolecules. Conney and the Millers identified microsomal enzymes (P450s) that activate many drugs and chemical carcinogens [66].

Initiation involves a brief and irreversible interaction between a carcinogen and the target tissue, and implies a local cellular stress/solicitation. Gradually, the stress caused by carcinogen removal may promote an overstraining of the neighbor cells, and then of the whole body, causing a general imbalance. The two stages of cancer occurrence provoked by carcinogenic substances can easily be explained by the existence of the RpHI respiratory and pH imbalance. In the two-stage skin carcinogenesis, the initiation phase is carried out by the application of a sub-carcinogenic dose of a carcinogen [Abel et al. Nat Protoc. 2009 67].

Suppose that a small amount of carcinogen enters a cell. Although all cells are able to degrade chemicals, this capacity differs from one tissue to another, and depends on age, sex, health status, etc. Cell will degrade the foreign compound through a series of eight

Formatted: English (United States)

Formatted: English (United States)

consecutive chemical reactions within the microsomal mixed function oxidase system [68].⁵⁹ Cytochromes P450, existing in each cell, provide the principal, initial source of biotransformation of xenobiotics that cell comes in contact with. The role of intestinal P450 in detoxifying specific carcinogens like 3-methylcholanthrene has clearly been demonstrated [Kaminsky LS, Fasco MJ. Small intestinal cytochromes P450. Crit Rev Toxicol. 1992;21(6):407-22. DOI:10.3109/1040844920908988169]. Moreover, elevated intestinal P450 levels have been indirectly linked to gastrointestinal cancer. Intestinal metabolism of 2,2,2-trifluoroethanol, which is not an carcinogen, produces intestinal lesions with consequent systemic bacterial infection. The activity of these enzymes requires nicotinamide adenine dinucleotide phosphate (NADPH) and molecular oxygen; in a typical reaction, one molecule of oxygen is consumed (reduced) per substrate molecule, with one oxygen atom appearing in the product and the other in the form of water. However, noxious reactive oxygen species (ROS) and H₂O₂ may appear. Some other degradation reactions may also be possible, depending on the type of cell. To form NADH from NAD⁺, glucose, fatty acids, amino acids, citric acid, malic acid, etc., are consumed. During this process, cell content will become more alkaline, and, if oxygen supply is not enough for removing the carcinogen, fermentation will develop alongside with respiration. Glucose from blood will help the fermentative processes, because it forms acetyl-CoA, both in the presence and absence of oxygen. Acetyl-CoA cannot be used in the absence of oxygen; therefore, glucose will give only pyruvate, which, in turn, is reduced by NADH to lactate.

Whenever a cell is damaged, its neighbors undergo inflammation by which this cell can be destroyed. The interesting dependence between chemical carcinogenesis and inflammation was reviewed by Loeb and Harris [200866]. In this process, the need of oxygen of the neighbor cells is increased. They undergo an inflammatory process, during which they become slightly alkaline, and retain carbon dioxide. A hypoxia state may occur in the surrounding tissue. Paradoxically, blood will provide a cell with less oxygen if it acts against a carcinogen than if it is in the neighborhood. If the carcinogen does not react directly with the cell chemical components, so as to be degraded and removed, the cell will continue to act against it. The degradation process, in which the carcinogen is transformed into metabolites may last long enough. Whereas a long standing hypoxia state is created,

the fine cellular structure is damaged by ROS, and the lactic acid is released from the contaminated cell, being then reconverted into glucose with ATP consumption as energy source. The neighbor cells support the carcinogen-contaminated cell, undergoing an inflammatory reaction. Both the degradation and the inflammatory reaction may damage the involved tissue. Omega (omega)-3 polyunsaturated fatty acids have been recognized as having anticancer activity [69],⁶⁹ probably as a consequence of their anti-inflammatory effect.

Logically, both the carcinogen and the poisoned cell should be removed. ~~Then the tissue will recover.~~ If not, more oxygen is needed to help cells ~~fight-destroying against that~~ carcinogen.

Chromosomal instability during precancerous stage

Epidemiology and clinical evidence clearly indicate that specific genetic, environmental and behavioral factors are associated with an increased risk for cancer development [1,70].⁺ ~~However, our findings are concordant to the observation that intracellular alkalinizing and extracellular acidification play a key role in the growth and metastasis of tumor cells.~~⁶¹ The stepwise progression from preneoplastic lesions to cancer seems to be associated gradually increases in genomic instability [71] ~~from an early dysplastic lesion to full blown metastasis malignancy seems to be associated with increases in genomic instability.~~⁶² ~~Still~~ Nevertheless, at each step of carcinogenesis, there are natural mechanisms protecting the body against development of cancer [72].⁶³ ~~Besides, carcinogens can also be removed from the body.~~ If DNA damage occurs, it is repaired in most of the cases.

Changes in DNA methylation consisting of either DNA hypomethylation or hypermethylation are commonly seen in cancer cells [73].⁶⁴ ~~The significance of these alterations in the genesis of cancer is, however, unclear.~~⁶⁴ Nevertheless, hypermethylation of tumor-suppressor genes in cancer could merely be a secondary event to help maintain gene inactivity which has been induced by some other mechanism. DNA methylation-DNA methylation changes have been shown to be associated with gene expression, sequence, and copy number changes [Weisenberger, 201474]. Cancer cells usually exhibit important changes in DNA methylation and histone modification [Jones, 201475]. Hence, targeting

Formatted: Font color: Auto, Superscript

epigenome-modifying enzymes could be an attractive therapeutic strategy. The cancers with poor prognosis demonstrate evidence of a metabolic shift, involving downregulation of genes involved in the TCA cycle, increased glycolysis, decreased AMPK and PTEN protein levels, glutamine-dependent lipogenesis, fumarate hydratase, upregulation of the pentose phosphate pathway and the glutamine transporter genes, increased acetyl-CoA carboxylase protein, and altered promoter methylation of miR-21 [76](Nature, 2013). Nevertheless, epigenetic changes such as DNA methylation act to regulate gene expression in normal mammalian development [Baylin 200577]. In fact, DNA methylation helps to maintain transcriptional silence in noncoding or nonexpressed regions of the genome. Such findings suggest that gene silencing by hypermethylation is relatively common both in human cancer and many other human diseases [Christensen et al. 200978]; interindividual epigenetic variations in normal tissues due to aging, environmental factors, or innate susceptibility are also characterized. Besides, the variability of gene expression is related to epigenomic states that vary across individuals. For instance, methylation profile classes are significantly associated with age, being significant predictors of tissue origin as well [Christensen et al. 200978].

Nevertheless, renal carcinogenesis is associated with neither chronic inflammation nor persistent viral infection, and any histological change is hardly evident [79].⁶⁵ However, regional DNA hypermethylation on C-type CpG islands has already accumulated in such non-cancerous renal tissues, suggesting that, from the viewpoint of altered DNA methylation, the presence of precancerous conditions can be recognized even in the kidney. Since DNA methylation of genes is age-dependent, being related to the tissue origin, it cannot be a hallmark of cancer as some authors claim [Teschendorff et al. 201080], but a phenomenon associated with excessive cell multiplication in a specific tissue.

Malign process is preceded by the genetic deficiencies, yet, only 0.3 per cent of the recessive-lethal mutations give neoplasm in *Drosophila melanogaster* [81,82].^{66,67} Moreover, tumor cells of this species become easily reversible because of the fact that these mutations do not seem to be the cause of cancer. In addition, even if cancer occurrence is

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Normal (Web), Indent: First line: 0"

directly related to the recessive-lethal mutations, the resulted abnormal proteins are normally hydrolyzed to amino acids.

The accumulation of unfolded proteins in the endoplasmic reticulum (ER) represents a cellular stress induced by multiple stimuli and pathological conditions. These include hypoxia, oxidative injury, high-fat diet, hypoglycemia, protein inclusion bodies and viral infection [83].⁶⁸

~~Eugen Macovschi stated that the biostructure is partially broken down during hypoxia [84].^{Macovschi, 1969} Since the biostructured matter is found under an altered, abnormal state in cancer cell [85,86].^{69, Macovschi, 1981} the RpHI hypothesis was advanced to explain first the transformation of a normal cell into a preneoplastic one, and second, the transition from preneoplastic cells to malignant ones [31,32,50].^{Drochioiu, 2006} According to Eugen Macovschi,⁶⁹ the structure in the cancer cell is found under an altered, abnormal state; the investigation of this state is essential for the knowledge of carcinogenesis.~~ The RpHI hypothesis is concordant with Macovschi's statement, because it considers the role of hypoxia in the malignant transformation of cell biostructure. Genomic instability may also be associated with hypoxia, which might be the causal phenomenon. However, on using metabolic inhibitors like dinitrophenol and sodium azide, Macovschi demonstrated partial and reversible biostructure breakdown [84].^{Macovschi, 1969} He concluded that these compounds inhibit oxidative phosphorylation and, consequently, the ATP content of cells decreases. Since ATP is needed to support the biostructure, then any carcinogenic agent alter the state of living matter by decreasing the ATP pool. Cancerogenesis is therefore accompanied by the partial breakdown of the biostructure, leading to alteration of cell behavior. Nevertheless, further research showed that cellular ATP concentrations only slightly decrease by switching from aerobic production of ATP to glycolysis. Under such circumstances, RpHI hypothesis admits that carcinogens are chemicals difficult to be degraded and eliminated from the body, and not very reactive compounds. Metabolic processes of carcinogens occur in cells in organisms [Manahan SE. Toxicological chemistry and biochemistry. 3rd Ed., CRC Press LLC, Boca Raton London New York Washington, D.C., 2003, pp. 162-16687]. The most important degradation pathways are oxidation reactions, particularly those classified as microsomal monooxygenation reactions.

Formatted: Tab stops: 0.58", Left + Not at 1.79"

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Monooxidations occur with oxygen as the oxidizing agent, generating a long standing hypoxia if carcinogen degradation is a multi-step process, this one is not too toxic to kill the cell, and its degradation products are also difficult to be eliminated. Dinitrophenols also induce hypoxia but by inhibiting respiration, without affecting too much the ATP concentration in poisoned cell. Under hypoxic conditions, ATP production is stepwise shifted from respiration to glycolysis, resulting in extracellular pH decrease, enhanced proteolysis [33] etc. In fact, these biochemical processes as well as some others like the hydrolysis of proteins involved in oxidative phosphorylation, cytochrome c or K⁺ leaking, affect the biostructure integrity. Hence, RpHI hypothesis may bring has thus a contribution to rationale explanation understanding for of biostructure breakdown in cancer, since it refers to the cause for long standing hypoxia, which creates the conditions for biostructure breaking down.

Formatted: Font: Not Bold

Increased glucose metabolism of cancers

In solid regions of tumors the most consistent finding is: a higher glucose utilization rate, a higher lactate production, and a higher-lower pH than in the surrounding tissues [88].⁷⁰ The acidic extracellular pH that is lower than pH in surrounding tissues is a key feature of solid malignant tumors. Tumor ATP was found to be slightly higher and glucose slightly lower than in the neighbor tissues. Most measurements of regional glucose and oxygen consumption by positron emission tomography have confirmed that oxygen extraction is lower than in normal tissue, and that the oxygen/glucose uptake ratio corresponds to a glycolytic rate of more than 65% [89].⁷¹ Since ATP production by glycolysis is less than 6% of that of oxidative phosphorylation, restriction of glucose delivery to the tumor by the blood was supposed to be more important than acidosis for inducing energy failure and subsequent tissue necrosis [88].⁷⁰

Formatted: Justified, Line spacing: 1.5 lines

Formatted: Font: 12 pt

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

It was suggested that glycolysis is elevated because it produces acid, which provides an evolutionary advantage to cancer cells vis-à-vis normal parenchyma which they invade [90].⁷² The RpHI hypothesis states that the slight hypoxia in the precancerous stage will

induce an increased consumption of glucose ~~in order as required by cells~~ to produce the same amount of ATP as that obtained by normal respiration.

However, hypoxia occurs in physiological situations such as during embryonic development, as well as in pathological conditions such as ischemia, wound healing, and cancer [35].³²

The link between hypoxia and the regulation of angiogenesis is an area of intense research [91]. ~~The molecular details of this connection are still being elaborated.~~⁷³ Moreover, molecular insights into these processes are rapidly being generated [Carmeliet, 200392]. A primary effector of the adaptive response to hypoxia in mammals is the hypoxia-inducible family factor of transcription regulators. These proteins activate the expression of a broad range of genes that mediate many of the responses to the decreased oxygen concentration, including enhanced glucose uptake, increased red blood cell production, and the formation of new blood vessels via angiogenesis [91].⁷³

The limiting role of glucose - and not of oxygen - for maintenance of energy state is also reflected by the relationship between ATP and glucose content under conditions of critical blood perfusion. Diabetics treated with an insulin-lowering drug, known as metformin, have less chances of cancer than those who receive insulin as treatment [93].⁷⁴ Indeed, metformin may lower cancer risk in diabetics and improve outcomes of many cancers [94]. Metformin may act by impacting mitochondrial respiration leading to the activation of the AMP-activated protein kinase, which controls energy homeostasis in cells. This drug also seems to inhibit cancer cell growth in vitro and delays the onset of carcinogen-induced cancer in mice. However, the undertaken studies are incapable of establishing a causal relationship. According to RpHI hypothesis, A relationship between that this drug and may lower the glucose concentration may be suspected in cells, reducing thus oxygen consumption and glycolysis.

The tumors were shown to be slightly more alkaline than normal brain tissue, although lactate was consistently increased [95].⁷⁵ Thus, in animals with pronounced peritumoral brain edema, the latter is distinctly more alkaline than the healthy tissue. ~~Therefore, increased lactate content from tumors is usually not associated with an acid shift of pH.~~⁷⁰ Moreover, high CO₂ levels decrease O₂ consumption and ATP production, and

impair cell proliferation independently of acidosis and hypoxia [96].⁷⁶ Quantification of tissue pH in solid regions of glioma reveals an average value of 7.26. This value is higher than in the cortex of the opposite hemisphere and lower than that in adjacent edematous cortex.

In tumors, ATP content generally decreases when glucose begins to fall, whereas in the normal brain, ATP declines before glucose is reduced [97].⁷⁷ Although these considerations and the observed lactate increase are in support of a preferential glycolytic pathway for ATP production, tumor pH was slightly more alkaline than in normal brain tissue [98-100].⁷⁸⁻⁸⁰ Nevertheless, at an extracellular pH of 7.25, the oxygen consumption rate of cells increases by nearly a factor of 2 as the glucose concentration decreases from 5.5 mM to 0.4 mM [101].⁸¹ This effect of glucose concentration on the oxygen consumption rate, however, was slight at an extracellular pH of 6.95 and disappeared completely at an extracellular pH of 6.60. Hence, low glucose concentrations, associated with a normal extracellular pH will increase oxygen utilization, slowdown glycolysis, increase respiration and aerobic ATP production. An ill animal will consume less food and will avoid effort. This behavior could be a lesson for human beings to reduce blood glucose concentration according to physiological requirements.

~~The new approach against cancer, as represented by i~~Inhibition of proton pumps in cancer and, changes in pH, ~~and are associated with the~~ mechanism ~~used by which~~ tumor cells ~~to avoid intracellular accumulation of toxic substances~~ ~~was already reviewed~~ [102].⁸²

Respiration and glycolysis

It is well known that respiration and fermentation are two independent biochemical processes, which may occur simultaneously in the living cell. Given an oxygen concentration of 10 per cent or more in the surrounding atmosphere, respiration occurs in the living cell. Given a concentration below 3% of oxygen, fermentation (glycolysis) will occur. Both processes occur in the range from 3% to 10% oxygen (Figure 2). The growth and lifespan of human diploid cell strains at oxygen concentrations of less than 20% is enhanced, and an extension of about 25% of the lifespan of both cell types was carried out by long term cultivation below 10% oxygen [103].⁸³

Whenever a cell within the metabolic system (muscle, liver, kidney, lung etc.) uses oxygen at a faster rate than it can be supplied by the circulatory system, the cell begins functioning anaerobically, reducing the pyruvate to lactate instead of further oxidizing it, as it would occur if oxygen supplies were adequate. Lactate thus accumulates in that cell, diffuses out through the bloodstream, and eventually finds its way to the liver, where it is reoxidized to pyruvate and converted back to glucose by the gluconeogenic pathway operative there.




Figure 2. Schematic presentation of the relative intensities of cellular respiration and glycolysis as a function of oxygen partial pressure.

Mitochondria play an important role in the functional responses of eukaryotic cells to changes in O_2 concentration, being involved in the adaptation to hypoxia. Previous investigations showed that when eukaryotic cells are exposed to hypoxia, glycolysis is stimulated and it can compensate the low ATP production within the oxidative phosphorylation process. Moreover, Otto Warburg showed that normal cells would become irreversibly cancerous if the environment they rested in had their oxygen levels lowered by 35% for two days. Besides, long-term cell cultures may become malignant without any other influences.

The glucose consumption rate of cancer cells increases much when the oxygen concentration is reduced [101];⁸⁴ however, severe conditions are required to stop cell growth (8.2 μ M oxygen and an extracellular pH of 6.6).

Hypoxia inducible factor 1, a marker of the Warburg effect and altered tumor metabolism, is associated with worse overall and recurrence free survival in peripheral tumor regions [104]. Meanwhile, central tumor expression of HIF-1 α has no significant correlation with outcome. These findings highlight the importance of assessing metabolic markers in the context of tumor microenvironment and oxygen tension.

Indeed, hypoxia and the hypoxia-inducible factors (HIFs) step in the metabolic crosstalk between cancer cells and their microenvironment [105]. Nevertheless, tumor cells are able to maintain a high glycolytic rate even if enough oxygen is supplied. HIFs were suspected to control the response of malignant cells to this specific microenvironment by shifting energy production from oxidative phosphorylation towards glycolysis. According to RpHI hypothesis, although enough oxygen may be present in the microenvironment, the cells cannot exchange it with carbon dioxide, which is kept by these cells. Furthermore, the mechanisms by which hypoxia and HIFs regulate tumor metabolism should be reconsidered. Possibly, RpHI may induce hypoxia, which in its turn, creates the necessary conditions for HIF production, a marker of hypoxic state.

Hypoxia prevents normal ATP production by oxidative phosphorylation [106].⁹⁷ Hypoxic cells swell, whereas the swollen mitochondria produce less energy. Chronic inflammation begins following continuous exposure to noxious exogenous influences [106].⁹⁷ Most cancers are mainly attributable to long-term exposure to environmental factors, via a multi-year, multi-step, and multi-path process of tumorigenesis [11].⁴⁴ Despite ATP synthesis rate is reduced when oxygen availability decreases, the exposure of cells to low oxygen level do not significantly affect the total ATP content, which is quite the same level of normoxia [107].⁹⁸ These data suggest that cells respond to lowered oxygen tension by increasing the activity of glycolytic enzymes and glucose transporters via the hypoxia-inducible factor-1 (HIF) [107].⁹⁸ HIF up regulates pyruvate dehydrogenase kinase 1, that inhibits the pyruvate dehydrogenase decreasing TCA cycle and consequently oxidative phosphorylation in

Formatted: Indent: First line: 0.5"

mitochondria. HIF, which mediates the transcriptional response to hypoxia, is a strong promoter of tumor growth and invasion [108,109].^{99,100}

Formatted: English (United States)

Formatted: Indent: First line: 0.5"

Formatted: Romanian (Romania)

Stressed cells and neighbor cells

As the metabolic control mechanism which regulates the transition from aerobic to anaerobic metabolism seems to be of key importance in the development of pathological changes, several studies were initiated to correlate the effects of anoxia on normal and aged mitochondria to the various biochemical and functional parameters of these organelles.

In isolated mitochondria inhibition of respiration and energy production have been demonstrated to be important factors in cell injury especially when nitrogen-induced anaerobiosis was investigated [110].⁸⁴ It was observed that aging, especially under nitrogen, leads to a decrease of the affinity constants of all mitochondria. The effect of anaerobiosis is increased by the combination with decreased pH; it was demonstrated that decreased pH exerts an effect on the redox state of cytochrome *a* in anaerobic, ATP-treated intact mitochondria [111].⁸⁵

Cytochrome *c* oxidase (EC 1.9.3.1.) acts as a final electron acceptor in the mitochondrial respiratory chain and catalyzes the molecular oxygen reduction to water. The fact that the radical intermediates of oxygen reduction have a carcinogenic potential calls to a special consideration of the cytochrome *c* oxidase pathway [112].⁸⁶

The cytochrome *c* oxidase activity is the highest in all forms of host liver mitochondrion; aging of mitochondria was associated with a decrease in enzyme activity, which was more intense in host liver mitochondria than in the tumor ones. As depicted in Figure 3, ageing and anaerobiosis provoke a partial breakdown of the mitochondrial biostructure; the cytochrome *c* oxidase activity mirrors this process [103].⁸³ The biostructure of ascitic mitochondria is also broken-down; anaerobiosis and carcinogenic factors induce a similar breaking down of mitochondria. Synthesis of mitochondrial-encoded subunits of cytochrome *c* oxidase in *Saccharomyces cerevisiae* is also suppressed in response to oxygen deprivation both *in vivo* and *in vitro*. [113].⁸⁷ Cytochrome *c* oxidase

activity among other respiratory enzymes monitored an imbalance, which characterizes both a precancerous stage and cancer. While cancer cell is hypoxic, the surrounding cells are overstrained.

Angiogenesis is a necessary component of all healing wounds. Tissue hypoxia is a well-known stimulus to angiogenesis [114,115].⁸⁸ Arany et al. 2008 Malignant, ischemic, inflammatory, infectious and immune disorders are characterized by dysregulated formation of new blood vessels [Carmeliet, 2003,92]. There are, however, inconsistent data and widely differing hypotheses concerning how much (what pressure or concentration) is required to support cellular metabolism and how the oxygen levels in cells within tissues are regulated [116].⁸⁹ There is a wide disparity in the published data and in the hypotheses used to interpret the cell and tissue responses to changes in oxygen pressure [117,118].^{90,91}




Figure 3. Scheme showing decreased enzyme activities in tumor cells as compared with those of normal cells, whereas increased enzyme activities are measured around the tumor. Cytochrome c oxidase activity (expressed as micromoles oxidized cytochrome c/mg protein/min) of normal mitochondria, host liver mitochondria, and ascitic mitochondria is different under various conditions (Fresh and aged

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Formatted: Justified, Indent: First line: 0", Space Before: 0 pt, After: 0 pt

mitochondria for 15 min at 22 °C in air or nitrogen-based anaerobiosis), which brings convincing evidence for RpHI hypothesis. Please, see the papers by C. Mihai for experimental details [84,110].

Cytochrome c oxidase activity (expressed as micromoles oxidized cytochrome c/mg protein/min) of normal mitochondria (NM), host liver mitochondria (HM) and ascitic mitochondria (AM) under different conditions (Fresh and aged mitochondria for 15 min at 22 °C in air or nitrogen-based anaerobiosis. See the experiments by C. Mihai for details.⁸⁴

Life style and cancer

The American Cancer Society has recently published the Nutrition and Physical Activity Guidelines, which are developed by a national panel of experts in cancer research, prevention, epidemiology, public health, and policy. They reflect the current scientific evidence related to dietary and activity patterns and cancer risk [119].⁹² Moderate to vigorous levels of physical activity lower the risk of developing several cancers.

Although the lung is a well oxygenated environment, hypoxia has been demonstrated in the center of an invasive lung tumor [Serfin, 2011,104]. The molecular markers of hypoxia like HIF-1 α and CA9 suggest that carbon dioxide removal rather than oxygen supply is a benefit of exercise expected from the physical exercises. However, we may suspect the uncoupling activity of some carcinogens in lung cancer such as tar from smoking, which may inhibit oxidative phosphorylation even in the presence of oxygen. Recent studies referred to the mitochondrial uncoupling in leukaemia cells, with energy dissipation and low ATP synthesis, as a potential mechanism of impairment of mitochondria in Warburg phenomenon [Samudio et al., 2008; Samudio et al., 2009,120,121]. Moreover, uncoupling proteins are over-expressed in breast cancer, supporting the link between mitochondrial uncoupling and the Warburg effect [Upadhyay et al. 2013; Ayyasamy et al., 2011,126,122]. Besides, overexpression of *E2f1* and *c-Myc* was identified at an early dysplastic stage of hepatocarcinogenesis [Coulouarn et al. 2006,123]; the combined activity of *c-Myc* and *E2f1* enhanced the expression of genes involved in mitochondrial metabolism, which results in

685 [increased ATP synthesis by the respiratory chain](#). Overweight and obesity are clearly
686 associated with an increased risk of developing many cancers. ~~Overeating seems to be the~~
687 ~~main avoidable cause of cancer in nonsmokers [Calle & Thun, 2004]124~~. The excess body
688 ~~weight is associated with risk of cancer at several organ sites, including colon, breast,~~
689 ~~endometrium, oesophagus, and kidney [Bianchini et al. 2002]125~~. ~~m~~Maintaining a healthy
690 weight, staying physically active, and consuming a healthy diet can substantially reduce the
691 risk of developing, or dying from cancer. These same behaviors are also associated with a
692 decreased risk of developing cardiovascular disease and diabetes. All these
693 recommendations are concordant with the RpHI hypothesis. Moreover, anxiety, rather than
694 depression, seems to be a problem at long-term cancer survivors and their spouses,
695 compared with healthy controls [126].⁹³ ~~Cancer and depression commonly co-occur, while~~
696 ~~chronic and severe depression may be associated with elevated cancer risk [Spiegel &~~
697 ~~Giese-Davis, 2003]127~~. Some evidence shows that providing psychosocial support to
698 ~~reduce depression, anxiety, and pain may increase survival time with cancer. Depression~~
699 ~~and emotional disorders may reduce treatment success in cancer [Gripp et al. 2007]128~~. A
700 ~~bidirectional relationship between cancer and depression was thus found. Women suffering~~
701 ~~from depression were reported to be more vulnerable to breast cancer [Wellisch &~~
702 ~~Lindberg, 2001]129~~. The complex relationship between the psyche and the nervous,
703 ~~immune, and endocrine systems seems to be implicated in both cancer risk and survival~~
704 ~~[Fife et al. 1996]130~~.

705
706 It was suggested that cancer cells have dysfunctional mitochondria, which prevent
707 them to use the citric acid or other compounds in the Krebs cycle [45,131].^{42,94} ~~We also~~
708 ~~consider that T~~the mitochondrial function ~~can seems to~~ be altered in the precancerous stage.
709 However, if cell content becomes more alkaline due to the continuous work under stress
710 conditions, it will retain carbon dioxide and will get less oxygen regardless of its
711 concentration. The RpHI approach is in full compliance with the findings on cancer
712 occurrence reported by other authors [132].⁹⁵ A more comprehensive theory on cancer
713 etiology to highlight not only the molecular aspects but also the biostructural ones, and

Formatted: Indent: First line: 0.5", Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

possibly psychic-related processes as Macovschi stated, is expected from a more integrative life science [133].⁹⁶

Inflammation and cancer

Chronic inflammatory conditions predispose susceptible cells to neoplastic transformation [134]. In general, the longer the inflammation persists, the higher the risk of cancer. Inflammatory processes have been implicated in the etiology of inflammation-associated cancers [135]; they may induce DNA mutations in cells *via* oxidative/nitrosative stress. However, inflammatory cells and cancer cells themselves produce free radicals and soluble mediators. Reactive intermediates of oxygen and nitrogen may directly oxidize DNA, or may interfere with mechanisms of DNA repair. Free radicals may also rapidly react with proteins, carbohydrates and lipids, and the derivative products may induce a perturbation in the intracellular homeostasis, until DNA mutation. The effectors are represented by an imbalance between pro-oxidant and antioxidant enzyme activities [134].

Viral infection-induced chronic inflammation significantly increases the likelihood of cancer development [136]. Many viruses have already confirmed roles in cancer development and progression. Proteins involved in ROS generation are upregulated during chronic inflammation and cancer [135]. Gastrointestinal cancers are strongly linked with chronic inflammation [137]. Some authors have demonstrated a higher risk of hepatocellular carcinoma in obese individuals. Since obesity is a low-grade inflammatory disease, a direct role of visceral adipose tissue in carcinogenesis was investigated [138]. Cytokines such as IL-6 and TNF- α secreted from the adipose tissue are able to induce a chronic inflammatory condition predisposing the liver to a preneoplastic stage. Chronic inflammation promotes also prostate cancer formation [139]. Cellular glycolytic and oxidative metabolism is significantly influenced by low-level inflammation in prostate epithelial cells. Thus, the micro-environmental inflammatory cytokine TNF α induces aerobic glycolysis while inhibiting oxidative metabolism (Warburg effect). Nevertheless, although the inflammatory cytokine TNF- α can directly induce metabolic changes, similar to those produced by oncogene expression, this effect can simply be blocked by the natural dietary compound curcumin [140].

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

The Warburg effect is an anomalous characteristic of glucose metabolism in cancer cells [141]. Cancer cells are characterized by increased aerobic glycolysis. Moreover, the pro-inflammatory cytokines promote glycolysis in breast cancer cells, where the inflammation-induced miR-155 function is an important mediator. The miR-155/miR-143/HK2 cascade may be involved in the mechanism linking inflammation to the metabolism of cancer cells [141]. The Warburg phenotype may serve as a main causative mechanism for cellular hyperproliferation and resistance to apoptosis [142]. Indeed, hypoxia and inflammation alter cellular metabolism.

According to Warburg, cell respiration is damaged or insufficient in cancer cells. However, oxygen consumption and carbon dioxide production seem to be high in many cancerous tissues and cells, suggesting that their respiration is not irreversibly damaged [24]. The Pasteur effect also appears to be normal in many cancerous tissues since oxygen inhibits lactate production. Many tumor cells show high rates of lactate production in the presence of oxygen, a phenomenon known as the Warburg effect, which has diagnostic and possibly therapeutic implications. The basal lactate production is 3-5 fold higher in T98G glioma cells than in normal astrocytes [143]. On the contrary, the rate of lactate accumulation in response to mitochondrial inhibition with sodium azide is 10 times lower in glioma than in astrocytes, consistent with defective tumor metabolism. Indeed, RPHI hypothesis clarifies this paradox showing that cancer cells are in fact normal cells, which apparently behave differently due to hypoxic conditions. If a cell is in trouble, neighboring cells will help or destroy it. Such phenomenon can not take place in cell culture, and cancer cells thus multiply incessantly. The biostructure breaking down is a gradual process, which may have many stages such as the adaptive one, followed by others like the preneoplastic, neoplastic, or apoptotic stages. The efficacy of cancer therapies, especially for late-stage disease, remains poor overall [144]. Focusing on cancer prevention requires novel targeted

therapies. During the process of identifying preventive agents, the task now is to understand how these agents interact with their target molecular compounds or biostructure components.

A detailed understanding of preneoplastic phenomena and cellular mechanisms of tumor formation can lead to new therapeutic approaches for improvement of clinical outcome.

~~Hypoxia prevents normal ATP production by oxidative phosphorylation.⁹⁷ Hypoxic cells swell, whereas the swollen mitochondria produce less energy. Chronic inflammation begins following continuous exposure to noxious exogenous influences.⁹⁷ Most cancers are mainly attributable to long term exposure to environmental factors, via a multi-year, multi-step, and multi-path process of tumorigenesis.¹¹ Despite ATP synthesis rate is reduced when oxygen availability decreases, the exposure of cells to low oxygen level do not significantly affect the total ATP content, which is quite the same level of normoxia.⁹⁸ These data suggest that cells respond to lowered oxygen tension by increasing the activity of glycolytic enzymes and glucose transporters via the hypoxia inducible factor 1 (HIF).⁹⁸ HIF upregulates up regulates pyruvate dehydrogenase kinase 1, that inhibits the pyruvate dehydrogenase decreasing TCA cycle and consequently oxidative phosphorylation in mitochondria. HIF, which mediates the transcriptional response to hypoxia, is a strong promoter of tumor growth and invasion.⁹⁹⁻¹⁰¹~~

Concluding remarks

Literature findings reviewed here support the hypothesis ~~that on the so-called RpHI, which respiratory and pH imbalance previously described to may~~ explain the long interval between exposure of a certain cell to a mutagen and the expression of the resulting mutations, as well as carcinogenesis without any apparent genetic alteration. Mutations of certain genes have been identified as risk factors for development of certain cases of

Formatted: Indent: First line: 0.5"

Formatted: Indent: First line: 0.5"

Formatted: Space Before: 0 pt, After: 0 pt

cancer. A large body of information suggests that the lethal mutations play an important role, yet not a decisive one. However, gradual progress in understanding the origins of malignant neoplasms should also be considered. The hypothesis discussed here takes into consideration the long standing hypoxia within the whole organism, characteristic to the precancerous stage. Moreover, under hypoxic conditions, the overbusy cells, getting less oxygen than needed, will lead to anaerobic adenosine triphosphate (ATP) production, excessive multiplication and, finally, tumor development. Such assertion is consistent with most literature data. Consequently, new strategies of cancer prevention should be drafted, based on a physiological approach of precancerous stage. Cancer prevention may be associated with carbon dioxide removal from blood in order to stimulate blood and cell oxygenation and oxygen delivery to exchange the in cells, carbon dioxide removal as well as. Besides, pH and glucose concentration control is important to prevent cachexia. Natural mechanisms protecting against cancer should be taken into consideration. Future medicine can also be seen as caring for physiological balances and not only as an emergency surgery.

Further studies are required to highlight the essential mechanisms characteristic to the preneoplastic precancerous stage hidden in the extremely complex oncology. At least, our theoretical approach suggests that oxygen tension, hypoxia, and carbon dioxide loading are key parameters that may positively influence cancer development in some tissues.

References

1. Bode, A.M., Dong Z. (2009) Cancer prevention research - then and now. *Nat Rev Cancer*, **9**, 508-516.
2. Brash, D., Cairns, J. (2009) The mysterious steps in carcinogenesis. *Br J Cancer*, **101**, 379-80.
3. Le Floch, R., Chiche, J., Marchiq, I., Naiken, T., Ilk, K., Murray, C.M., Critchlow, S.E., Roux, D., Simon, M.-P., Pouyssegur, J. (2011) CD147 subunit of lactate/H(+) symporters MCT1 and hypoxia-inducible MCT4 is critical for energetics and growth of glycolytic tumors. *Proc Natl Acad Sci USA*, **108**, 16663-16668.

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Not Highlight

Formatted: Not Highlight

- 833 4. Martin, N.K., Gaffney, E.A., Gatenby, R.A., Gillies, R.J., Robey, I.F., Maini, P.K.
834 (2011) A mathematical model of tumor and blood pH regulation: The $\text{HCO}_3^-/\text{CO}_2$
835 buffering system. *Math Biosci*, **230**, 1-11.
- 836 5. Rakoff-Nahoum S. (2006) Why cancer and inflammation? *Yale J Biol Med*, **79**, 123–130.
- 837 6. Ioannou, S., Voulgarelis, M. (2010) Toll-like receptors, tissue injury, and
838 tumorigenesis. *Mediat Inflamm*, ID 581837, doi: org/10.1155/2010/581837
- 839 7. Niwa, T., Tsukamoto, T., Toyoda, T., Mori, A., Tanaka, H., Maekita, T., Ichinose, M.,
840 Tatematsu, M., Ushijima, T. (2010) Inflammatory processes triggered by
841 *Helicobacter pylori* infection cause aberrant DNA methylation in gastric
842 epithelial cells. *Cancer Res.*, **70**, 1430-1440.
- 843 8. Mantovani, A, Allavena P, Sica A, Balkwill F. (2008) Cancer-related inflammation.
844 *Nature*, **454**, 436-444.
- 845 9. Solinas, G., Germano, G., Mantovani, A., Allavena, P. (2009) Tumor-associated
846 macrophages (TAM) as major players of the cancer-related inflammation. *J*
847 *Leukoc Biol*, **86**, 1065-1073.
- 848 10. Ricciotti, E., FitzGerald, G. A. (2011) Prostaglandins and inflammation. *Arterioscl*
849 *Throm Vasc*, **31**, 986-1000.
- 850 11. Rathore, K. (2011) Suppression of chronically induced breast carcinogenesis and role of
851 mesenchymal stem-like cells. PhD thesis, University of Tennessee-, [pp. 180-182](#).
- 852 12. Rathore, K., Wang, H. C. (2012) Green tea catechin extract in intervention of chronic
853 breast cell carcinogenesis induced by environmental carcinogens. *Mol Carcinogen*,
854 **51**, 280-289.
- 855 13. Rakoff-Nahoum, S., Medzhitov, R. (2009) Toll-like receptors and cancer. *Nat Rev*
856 *Cancer*, **9**, 57-63.
- 857 14. Mattarei, A., Azzolini, M., Carraro, M., Sassi, N., Zoratti, M., Paradisi, C., Biasutto, L.
858 (2013) Acetal derivatives as prodrugs of resveratrol. *Mol Pharmaceut*, **10**, 2781–
859 2792.
- 860 15. Weng, C.-J., Yen, G.-C. (2012) Chemopreventive effects of dietary phytochemicals
861 against cancer invasion and metastasis: Phenolic acids, monophenol, polyphenol,
862 and their derivatives. *Cancer Treat Rev*, **38**, 76-87.

Formatted: Italian (Italy)

- 863 16. Wang, P., Liu, H.-J., Mei, X.-Y., Nakajima, M., Yin, L.-J. (2012) Preliminary study
864 into the factors modulating beta-carotene micelle formation in dispersions using an
865 in vitro digestion model. *Food Hydrocolloid*, **26**, 427-433.
- 866 17. Manson, M. M., Linseisen, J., Rohrmann, S., Sotiroudis, T. G., Kyrtopoulos, S. A.,
867 Hayes, J. D., Kelleher, M. O., Eggleston, I. M., de Kok, T. M., van Breda, S. G.,
868 Rafter, J. (2007) Bioactive components in foods. In: Dietary vitamins,
869 polyphenols, selenium and probiotics: biomarkers of exposure and mechanism of
870 anticarcinogenic action. Ed. Bjorn Akesson, Per Mercke, Lodz, pp. 115-177.
- 871 18. Grill, A. E. (2002) Examining and improving the chemopreventive efficacy of
872 curcumin. PhD Thesis, University of Minnesota, [pp. 14-64](#).
- 873 19. Benamouzig, R., Uzzan, B., Deyra, J., Martin, A., Girard, B., Little, J., Chaussade, S.
874 (2012) Prevention by daily soluble aspirin of colorectal adenoma recurrence: 4-year
875 results of the APACC randomised trial. *Gut*, **61**, 255-261.
- 876 20. Cairns, R. A., Harris, I. S., Mak, T. W. (2011) Regulation of cancer cell metabolism.
877 *Nat Rev Cancer*, **11**, 85-95.
- 878 21. [Seyfried TN, Cancer as a metabolic disease: on the origin, management, and prevention](#)
879 [of cancer. John Wiley and Sons ed. Hoboken, New Jersey; 2012; Cap. 4.3-4.8; 5.1-](#)
880 [5.10; 6.1-6.3; 7.1-7.5; 9.1; 9.2; 10.2-10.4 Cairns, R.A., Harris, I.S., Mak, T.W. \(2011\)](#)
881 [Regulation of cancer cell metabolism. *Nat Rev Cancer*, **11**, 85-95.](#)
- 882 22. Santos de Souza A. C., Zenker J. G., Ribeiro de Araújo D., Martins C. A.D. (2011)
883 Defining the molecular basis of tumor metabolism: a continuing challenge since
884 Warburg's discovery. *Cell Physiol Biochem*, **28**, 771-792.
- 885 23. Bissell, M. J., Hines, W. C. (2011) Why don't we get more cancer? A proposed role of
886 the microenvironment in restraining cancer progression. *Nat Med*, **17**, 320-329.
- 887 24. Drochioiu, G. (1987) Considerations on biochemical mechanism of malignant
888 transformation. *Proc Symp Comp Oncol*, Timișoara, România, **1**, 116-117.
- 889 25. Drochioiu G. (2006) Eugen Macovschi's concept of biostructure and its current
890 development. In *Life and mind. In search of the physical basis*. (S. Savva ed.)
891 Trafford Publ., Canada, USA, Ireland & UK, pp. 43-60.

Formatted: Portuguese (Brazil)

- 892 26. Drochioiu, G. (2008) Chronic metabolic acidosis may be the cause of cachexia: Body
893 fluid pH correction may be an effective therapy. *Med Hypotheses*, **70**, 1167- 1173.
- 894 27. Drochioiu, G. (2008) *Biological systems: a structural-phenomenological approach*, in
895 *Cybernetics and Systems*, Proceedings of the Nineteenth Meeting on Cybernetics and
896 Systems Research, Trappl R. (ed.) Austrian Society for Cybernetic Studies, Vienna, **1**,
897 203-208.
- 898 28. Drochioiu, G. (1985) Respiratory and pH imbalance in carcinogenicity (Romanian).
899 *Știință și tehnică* (Science and Technology-Bucharest), **9**, 38-39.
- 900 29. Drochioiu, G. (1987) The carcinogenic action of some chemicals. *Proc Symp Comp*
901 *Oncol* Timișoara, România, **1**, 115-116.
- 902 30. Douglas, R. G., Shaw, J. H. F. (1990) Metabolic effects of cancer. *Br J Surg*, **77**, 246-
903 254.
- 904 31. Aldwin, C. M. (2007) *Stress, coping, and development: an integrative perspective*.
905 Guilford Press, pp. 37-54.
- 906 32. Lovallo, W. R. (2005) Stress and health: biological and psychological interactions. ^{2nd}
907 Ed., Sage Publ. California, London, New Dehli, pp. 11-22.
- 908 33. Stranks, J. (2013) *Stress at work*. Jeremy Stranks Editor, Routledge Burlington, pp.
909 4-21; 41-44.
- 910 34. Bendelow, G. (2009) Health, emotion and the body. G Bendelow, Edit. -Polity Press,
911 UK Cambridge, UK, Malden, USA, pp. 80-105.
- 912 35. Spiegel D. Cancer. in: Silverthorn, D.U., Ober, W.C., Garrison, C.W., Silverthorn, A.C.,
913 Johnson, B.R. (2009) *Human physiology*. Encyclopedia of stress. Vol. 1 (Fink G,
914 Editor), Person, Benjamin Cummings, USA Academic Press, 2000, pp. 368-375.
- 915 36. Piaget, J. (2013) *Behaviour and evolution*. Routledge. Fawzy FI, Canada AL, Fawzy
916 NW. Cancer treatment. in *Encyclopedia of stress*. Vol. 1 (Fink G, Editor), Academic
917 Press, 2000, pp. 376-380.
- 918 37. Csepregi, G. (2006) *The clever body*. Calgary Press, Canada. Kraemer WJ, Fleck SJ,
919 Deschenes MR. Exercise physiology: integrating theory and application. Kluwer,
920 Philadelphia, Baltimore, New York, London, Buenos Aires, Hong Kong, Sydney, Tokyo,
921 2012, pp. 27-66; 103-134; 167-196.

Formatted: Font: Not Italic

Formatted: Superscript

Formatted: Font: Not Italic

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

- 922 |
- 923 38. Shiraiwa, M., Selzle, K., Pöschl, U. (2012) Hazardous components and health effects of
- 924 atmospheric aerosol particles: reactive oxygen species, soot, polycyclic aromatic
- 925 compounds and allergenic proteins. *Free Radical Res*, **46**, 927-939.
- 926 39. Gong, B., Boor, P. J. (2006) The role of amine oxidases in xenobiotic metabolism.
- 927 *Expert Opin Drug Met*, **2**, 559-571.
- 928 40. Taylor, C. T. (2008) Mitochondria and cellular oxygen sensing in the HIF pathway.
- 929 *Biochem J*, **409**, 19-26.
- 930 41. Erecińska, M., Silver, I. A. (2001) Tissue oxygen tension and brain sensitivity to
- 931 hypoxia. *Resp Physiol*, **128**, 263-276.
- 932 42. King, A., Selak, M. A., Gottlieb, E. (2006) Succinate dehydrogenase and fumarate
- 933 hydratase: linking mitochondrial dysfunction and cancer. *Oncogene*, **25**, 4675-
- 934 4682.
- 935 43. Wiklund, L. (1996) Carbon dioxide formation and elimination in man. *Upsala J Med*
- 936 *Sci*, **101**, 35-68.
- 937 44. Williamson, J. R., Jákob, A., Scholz, R. (1971) Energy cost of gluconeogenesis in rat
- 938 liver. *Metabolism*, **20**, 13-26.
- 939 45. Robin, E. D. (1963) Abnormalities of acid-base regulation in chronic pulmonary
- 940 disease, with special reference to hypercapnia and extracellular alkalosis. *New Engl J*
- 941 *Med*, **268**, 917-922.
- 942 46. West JB. Respiratory physiology: the essentials and pathophysiology: an
- 943 integrated, case-based approach, 2nd ed. Wolters Kluwer Health/Lippincott Williams
- 944 & Wilkins, Baltimore & Philadelphia, 2008; pp. 15-24.
- 945 West, J. B. (2012) Respiratory physiology: the essentials. Kluwer, Canada.
- 946 JB West JB. Respiratory physiology: the essentials. Wolters Kluwer Health/Lippincott Williams &
- 947 Wilkins, Philadelphia, 2008.
- 948 Respiratory physiology: the essentials
- 949 Author: John B West
- 950 Publisher: Philadelphia : Wolters Kluwer Health/Lippincott Williams & Wilkins, ©2008.
- 951 West JB. Respiratory physiology—the essentials. 6th ed. Baltimore (MD), London (U.K.), and Los Angeles
- 952 (CA): Williams & Wilkins; 1999.

Formatted: Indent: Left: 0", Hanging: 0.37"

Formatted: Font: 12 pt, Not Highlight

Formatted: Font: 12 pt, Not Highlight

Formatted: Font: 12 pt, Not Highlight

Formatted: Superscript

Formatted: Font: 12 pt, Not Highlight

Formatted: Font: 12 pt, Not Highlight

Formatted: Font: 12 pt, Not Highlight

- 953 |
- 954 47. Das, M., Bouchev, D. M., Moore, M. J., Hopkins, D. C., Nemenoff, R. A., Stenmark,
955 K. R. (2001) Hypoxia-induced proliferative response of vascular adventitial
956 fibroblasts is dependent on G protein-mediated activation of mitogen-activated
957 protein kinases. *J Biol Chem*, **276**, 15631-15640.
- 958 48. Hall-Stoodley, L., Costerton, J. W., Stoodley, P. (2004) Bacterial biofilms: from the
959 natural environment to infectious diseases. *Nat Rev Microbiol*, **2**, 95-108.
- 960 49. Boutilier, R. G. (2001) Mechanisms of cell survival in hypoxia and hypothermia. *J Exp*
961 *Biol*, **204**, 3171-3181.
- 962 50. Bergers, G., Benjamin, L. E. (2003) Angiogenesis: tumorigenesis and the angiogenic
963 switch. *Nat Rev Cancer*, **3**, 401-410.
- 964 51. Martinou, J. C., Green, D. R. (2001) Breaking the mitochondrial barrier. *Nat Rev Mol*
965 *Cell Bio*, **2**, 63-67.
- 966 52. Morowitz, H., Smith, E. (2007) Energy flow and the organization of life. *Complexity*,
967 **13**, 51-59.
- 968 53. Embley, T. M., Martin, W. (2006) Eukaryotic evolution, changes and challenges.
969 *Nature*, **440**, 623-630.
- 970 54. Schmitt, C. A. (2007) Cellular senescence and cancer treatment. *BBA - Rev Cancer*,
971 **1775**, 5-20.
- 972 55. Nyakas, C., Buwald, B., Luiten, P. G. M. (1996) Hypoxia and brain development. *Prog*
973 *Neurobiol*, **49**, 25-27.
- 974 56. Damjanov, I. (2009) *Pathology secrets*. 3rd Ed., Mosby, Inc., an affiliate of Elsevier Inc,
975 USA.
- 976 57. Hall, J. E. (2010) [Guyton and Hall textbook of medical physiology: enhanced E-book.](#)
977 [12 Edition.](#)
978 [—Elsevier Health Sciences, Philadelphia, Cap. 19, 29, 30, and 39.](#)
- 979 58. Reglin, B., Secomb, T. W., Pries, A. R. (2009) Structural adaptation of microvessel
980 diameters in response to metabolic stimuli: where are the oxygen sensors? *Am J*
981 *Physiol*, **297**, H2206-H2219.

Formatted: German (Germany)

Formatted: Font: Not Italic

- 982 59. Ershov, E., Shlosberg, A., Hanji, V., Kagan, J. (1994) The role of microsomal mixed
983 function oxidases in the metabolism and mechanism of action of certain
984 organophosphorus insecticides. *Drug Metabol Drug Interact*, **11**, 139-152.
- 985 60. Cockbain, A.J., Toogood, G.J., Hull, M.A. (2012) Omega-3 polyunsaturated fatty acids
986 for the treatment and prevention of colorectal cancer. *Gut*, **61**, 135-149.
- 987 61. Kim, J., Namkung, W., Yoon, J.S., Jo, M.J., Lee, S.H., Kim, K.H., Kim, J.Y., Lee,
988 M.G. (2009) The role of translation elongation factor eEF1A in intracellular
989 alkalization-induced tumor cell growth. *Lab Invest*, **89**, 867-74.
- 990 62. Schwartzman, J.M., Sotillo, R., Benezra, R. (2010) Mitotic chromosomal instability and
991 cancer: mouse modeling of the human disease. *Nat Rev Cancer*, **10**, 102-115.
- 992 63. Jakóbsiak, M., Lasek, W., Gołaż, J. (2003) Natural mechanisms protecting against
993 cancer. *Immunol Lett*, **90**, 103-122.
- 994 64. Laird, P.W., Jaenisch, R. (1994) DNA methylation and cancer. *Hum Mol Genet*, **3**,
995 1487-1495.
- 996 65. Arai, E., Kanai, Y. (2011) Genetic and epigenetic alterations during renal
997 carcinogenesis. *Int J Clin Experim Pathol*, **4**, 58-73.
- 998 66. Finnegan, D. J. (2011) Oogenesis: active heterochromatin. *Curr Biol*, **21**, R630- R632.
- 999 67. Vincent, J.P., Kolahgar, G., Gagliardi, M., Piddini, E. (2011) Steep differences in
1000 wingless signaling trigger myc-independent competitive cell interactions. *Dev Cell*,
1001 **21**, 366-374.
- 1002 68. Kim, I., Xu, W., Reed, J. C. (2008) Cell death and endoplasmic reticulum stress:
1003 disease relevance and therapeutic opportunities. *Nat Rev Drug Disc*, **7**, 1013-1030.
- 1004 69. Macovschi, E. (1984) The biostructural theory of carcinogenesis. *Rev Roum Biochim*,
1005 **21**, 3-11.
- 1006 70. Hossmann, K. A., Mies, G., Paschen, W., Szabo, L., Dolan, E., Wechsler, W. (1986)
1007 Regional metabolism of experimental brain tumors. *Acta Neuropathol*, **69**, 139-147.
- 1008 71. Rhodes, C. G., Wise, R. J. S., Gibbs, J. M., Frackowiak, R. S. J., Hatazawa, J., Palmer,
1009 A. J., Thomas, D. G. T., Jones, T. (1983) In vivo disturbance of oxidative metabolism
1010 of glucose in human cerebral gliomas. *Ann Neurol*, **14**, 614-626.

- 1011 72. Gillies, R. J., Robey, I., Gatenby, R. A. (2008) Causes and consequences of increased
1012 glucose metabolism of cancers. *J Nucl Med*, **49**, 24S-42S.
- 1013 73. Hickey, M. M., Simon, M. C. (2006) Regulation of angiogenesis by hypoxia and
1014 hypoxia-inducible factors. *Curr Topics Develop Biol*, **76**, 217-257.
- 1015 74. Taubes G. (2012) Cancer research. Cancer prevention with a diabetes pill? *Science*,
1016 [2012;335\(6064\):29](https://doi.org/10.1126/science.1231231). doi: [10.1126/science.1231231](https://doi.org/10.1126/science.1231231).
- 1017
- 1018 75. Chiche, J., Le Fur, Y., Vilmen, C., Frassinetti, F., Daniel, L., Halestrap, A., Cozzone, P.
1019 J., Pouyssegur, J., Lutz, N. W. (2012) In vivo pH in metabolic-defective Ras-
1020 transformed fibroblast tumors: Key role of the monocarboxylate transporter, MCT4,
1021 for inducing an alkaline intracellular pH. *Int J Cancer*, **130**, 1511-1520.
- 1022 76. Vohwinkel, C.U., Lecuona, E., Sun, H., Sommer, N., Vadász, I., Chandel, N.S.,
1023 Sznajder, J.I. (2011) Elevated CO₂ levels cause mitochondrial dysfunction and impair
1024 cell proliferation. *J Biol Chem*, **286**, 37067-37076.
- 1025 77. Paschen, W., Hossmann, K. A., van den Kerckhoff, W. (1983) Regional assessment of
1026 energy-producing metabolism following prolonged complete ischemia of cat brain. *J*
1027 *Cereb Blood Flow Metab*, **3**, 321-329.
- 1028 78. Junck, L., Blasberg, R., Rottenberg, D. A. (1981) Brain and tumor pH in experimental
1029 leptomeningeal carcinomatosis. *Trans Am Neurol Assoc*, **106**, 298- 301.
- 1030 79. Rottenberg, H. (1984) Membrane potential and surface potential in mitochondria:
1031 uptake and binding of lipophilic cations. *J Membr Biol*, **81**, 127-38.
- 1032 80. Mies, G., Paschen, W., Ebhardt, G., Hossmann, K. A. (1990) Relationship between of
1033 blood flow, glucose metabolism, protein synthesis, glucose and ATP content in
1034 experimentally-induced glioma (RG12.2) of rat brain. *J Neuro-Oncol*, **9**, 17-28.
- 1035 81. Casciari, J. J., Sotirchos, S. V., Sutherland, R. M. (1992) Variations in tumor cell
1036 growth rates and metabolism with oxygen concentration, glucose concentration, and
1037 extracellular pH. *J Cell Physiol*, **151**, 386-94.
- 1038 82. Fais S. (2010) Proton pump inhibitor-induced tumour cell death by inhibition of a
1039 detoxification mechanism. *J Intern Med*, **267**, 515-525.

Formatted: Indent: Left: 0", Hanging: 0.37",
Space Before: 0 pt, After: 0 pt

Formatted: Font: Not Italic

- 1040 83. Packer, L., Fuehr, K. (1977) Low oxygen concentration extends the lifespan of cultured
1041 human diploid cells. *Nature*, **267**, 423-425.
- 1042 84. Mihai, C., Dinescu-Romalo, G. (1977) Anaerobiosis-induced modifications of normal
1043 rat liver mitochondria. *Rev roum Biochim*, **14**, 31-34.
- 1044 85. Higuti, T., Sugiyama, Y., Sato, M., Muraoka, S. (1975) *Biochem Bioph Res Co*, **62**,
1045 602-607.
- 1046 86. Mihai, C., Dinescu-Romalo, G. (1979) The influence of some factors upon cytochrome
1047 c oxidase activity of ascites and Guérin T8 ascites host rat liver mitochondria. *Rev*
1048 *Roum Biochim*, **16**, 49-53.
- 1049 87. Kwast, K. E., Hand, S. C. (1996) Oxygen and pH regulation of protein synthesis in
1050 mitochondria from *Artemia franciscana* embryos. *Biochem J*, **313**, 207-213.
- 1051 88. Knighton, D.R., Silver, I.A., Hunt, T.K. (1981) Regulation of wound-healing
1052 angiogenesis-effect of oxygen gradients and inspired oxygen concentration. *Surgery*,
1053 **90**, 262-270.
- 1054 89. Wilson, D.F. (2008) Quantifying the role of oxygen pressure in tissue function. *Am J*
1055 *Physiol*, **294**, H11-H13.
- 1056 90. [Papkovsky DB, Dmitriev RI. Biological detection by optical oxygen sensing. Chem Soc](#)
1057 [Rev. 2013;42\(22\):8700-32.](#) [Wilson, D. F. \(2008\) Quantifying the role of oxygen](#)
1058 [pressure in tissue function. Am J](#)
1059 [Physiol, 294, H11-H13.](#)
- 1060 91. Cameron, N. E., Cotter, M. A. (1994) The relationship of vascular changes to metabolic
1061 factors in diabetes mellitus and their role in the development of peripheral nerve
1062 complications. *Diabetes Metab Rev*, **10**, 189-224.
- 1063 92. Kushi, L.H., Doyle, C., McCullough, M., Rock, C.L., Demark-Wahnefried, W.,
1064 [Bandera, E.V., Gapstur, S., Patel, A.V., Andrews, K., Gansler, T. \(2012\) Reducing](#)
1065 [the risk of cancer with healthy food choices and physical activity. CA Cancer J](#)
1066 [Clin, 62, 30-67.](#)
- 1067 93. Mitchell, A. J., Ferguson, D. W., Gill, J., Paul, J., Symonds, P. (2013) Depression and
1068 anxiety in long-term cancer survivors compared with spouses and healthy controls: a
1069 systematic review and meta-analysis. *Lancet Oncol*, **14**, 721-732.

Formatted: Swedish (Sweden)

- 1070 94. John, A. P. (2001) Dysfunctional mitochondria, not oxygen insufficiency, cause cancer
1071 cells to produce inordinate amounts of lactic acid: the impact of this on the
1072 treatment of cancer. *Med Hypotheses*, **57**, 429–431.
- 1073 95. Nijsten, M. W. N., van Dam, G. M. (2009) Using the Warburg effect against cancer by
1074 reducing glucose and providing lactate. *Med Hypotheses* **73**, 48–51.
- 1075 96. Murariu, M., Drochioiu, G. (2012) Biostructural theory of the living systems.
1076 *BioSystems*, **109**, 126–132.
- 1077 97. [Geppetti P, Patacchini R, Nassini R. Transient receptor potential channels and](#)
1078 [occupational exposure. Curr Opin Allergy Cl. 2014;14\(2\):77-83.](#) ~~[Damjanov, I. Pathology](#)~~
1079 ~~[secrets. 3rd ed., Mosby, Inc., 2009.](#)~~
- 1080 98. Del Sole, M. *Effect of hypoxia and hyperglycemia on cell bioenergetics*. PhD Thesis,
1081 University of Bologna, 2012, [pp. 31-35.](#)
- 1082 99. Kim, Y., Lin, Q., Glazer, P. M., Yun, Z. (2009) Hypoxic tumor microenvironment and
1083 cancer cell differentiation. *Curr Mol Med*, 9(4), 425-434.
- 1084 100. ~~[Kunz, M., Ibrahim, S. M. \(2003\) Molecular responses to hypoxia in tumor cells. Mol](#)~~
1085 ~~[Cancer](#) **2**, 23, doi:10.1186/1476-4598-2-23~~
- 1086 ~~[101.](#)~~ Wisea, D. R., Warda, P. S., Shayb, J. E. S., Crossa, J. R., Gruberb, J. J., Sachdevab, U.
1087 M., Plattb, J. M., DeMatteoa, R. G., Celeste Simonb, M., Thompson, C. B. (2011)
1088 Hypoxia promotes isocitrate dehydrogenase-dependent carboxylation of α -
1089 ketoglutarate to citrate to support cell growth and viability. *Proc Nat Acad Sci USA*,
1090 108, 19611–19616.
- 1091
- 1092 [Cancer prevention](#)
- 1093 [Nature Reviews Cancer](#) **11**, 211–218 (March 2011) | doi:10.1038/nrc3017
- 1094 **Opinion: Molecular targets of phytochemicals for cancer**
1095 **prevention**
- 1096 [Ki Won Lee^{1,3}, Ann M. Bode^{2,3} & Zigang Dong²—About the authors](#)

Formatted: Indent: First line: 0"

Formatted: Indent: Left: 0"

1097 [Top of page](#)

1098 **Abstract**

1099 [Although successful for a limited number of tumour types, the efficacy of cancer therapies,](#)
1100 [especially for late stage disease, remains poor overall. Many have argued that this could be](#)
1101 [avoided by focusing on cancer prevention, which has now entered the arena of targeted](#)
1102 [therapies. During the process of identifying preventive agents, dietary phytochemicals,](#)
1103 [which are thought to be safe for human use, have emerged as modulators of key cellular](#)
1104 [signalling pathways. The task now is to understand how these chemicals perturb these](#)
1105 [pathways by modelling their interactions with their target proteins.](#)

1106

1107 **Hypoxia**

1108 [Zeng W, Liu P, Pan W, Singh SR, Wei Y. Hypoxia and hypoxia inducible factors in tumor](#)
1109 [metabolism. Cancer Lett. 2014; \(*In press*\). DOI 10.1016/j.canlet.2014.01.032.](#)

1110

1111 [20. Schulze A, Harris AL. How cancer metabolism is tuned for proliferation and vulnerable](#)
1112 [to disruption. Nature. 2012;491:364–73.](#)

1113 [21. Warburg O. On the origin of cancer cells. Science. 1956;123:309–14.](#)

1114 [22. Wu W, Zhao S. Metabolic changes in cancer: beyond the Warburg effect. Acta Biochim](#)
1115 [Biophys Sin. 2013;45\(1\):18-26.](#)

1116 [26. Upadhyay M, Samal J, Kandpal M, Singh OV, Vivekanandan P. The Warburg effect:](#)
1117 [insights from the past decade. Pharmacol Therapeut. 2013;137\(3\):318-30.](#)

1118
1119 [27. Gogvadze V, Zhivotovsky B, Orrenius S. The Warburg effect and mitochondrial](#)
1120 [stability in cancer cells. Mol Aspects Med. 2010;31\(1\):60-74.](#)

1121 [28. Ferreira LMR. Cancer metabolism: the Warburg effect today. Exp Mol Pathol.](#)
1122 [2010;89\(3\):372-80.](#)

1123 [30. Hitosugi T, Chen J. Post-translational modifications and the Warburg effect.](#)

1124 [Oncogene. 2014; \(*In press*\). DOI:10.1038/onc.2013.406](#)

Formatted: Indent: Left: 0"

Formatted: Font: (Default) Times New Roman, 12 pt

- 1125 [66. Loeb LA, Harris CC. Advances in chemical carcinogenesis: a historical review and](#)
 1126 [prospective. Cancer Res. 2008;68:6863-72. DOI: 10.1158/0008-5472](#)
- 1127 [67. Abel EL, Angel JM, Kiguchi K, DiGiovanni J. Multi-stage chemical carcinogenesis in](#)
 1128 [mouse skin: fundamentals and applications. Nat Protoc. 2009;4:1350-62.](#)
 1129 [DOI:10.1038/nprot.2009.120](#)
- 1130 [68. Kaminsky LS, Fasco MJ. Small intestinal cytochromes P450. Crit Rev Toxicol.](#)
 1131 [1992;21\(6\):407-22. DOI:10.3109/10408449209089881](#)
- 1132 [74. Weisenberger DJ. Characterizing DNA methylation alterations from The Cancer](#)
 1133 [Genome Atlas. J Clin Invest. 2014;124\(1\):17-23. DOI: 10.1172/JCI69740](#)
- 1134
- 1135 [75. Jones PA. At the tipping point for epigenetic therapies in cancer. J Clin Invest.](#)
 1136 [2014;124\(1\):14-16. DOI: 10.1172/JCI74145](#)
- 1137 [76. The Cancer Genome Atlas Research Network. Comprehensive molecular](#)
 1138 [characterization of clear cell renal cell carcinoma. Nature. 2013;499\(7456\):43-49.](#)
- 1139 [77. Baylin SB. DNA methylation and gene silencing in cancer. Nat Clin Pract Oncol.](#)
 1140 [2005;2:S4-S11. doi:10.1038/ncponc0354](#)
- 1141 [78. Christensen BC, Houseman EA, Marsit CJ, Zheng S, Wrensch MR, Wiemels JL, et al.](#)
 1142 [Aging and environmental exposures alter tissue-specific DNA methylation dependent upon](#)
 1143 [CpG island context. Plos Genet. 2009; DOI: 10.1371/journal.pgen.1000602](#)
- 1144 [80. Teschendorff AE, Menon U, Gentry-Maharaj A, Ramus2 SJ, Weisenberger DJ, Shen H,](#)
 1145 [et al. Age-dependent DNA methylation of genes that are suppressed in stem cells is a](#)
 1146 [hallmark of cancer. Genome Res. 2010;20:440-46. DOI: 10.1101/gr.103606.109](#)
- 1147 [84. Macovschi E. Biostructure. Romanian Academy Publ. House, Bucharest, 1969, pp. 57,](#)
 1148 [188-202. Romanian](#)
- 1149 [86. Macovschi E. The confirmation of the biostructural theory by the high-voltage electron](#)
 1150 [microscopy. Scientific and Encyclopedic Publ. House, Bucharest, 1981, pp. 15-44.](#)
- 1151 [92. Carmeliet, P. Angiogenesis in health and disease. Nat Med. 2003;9\(6\):653-60.](#)

Formatted: German (Germany)

Formatted: German (Germany)

Formatted: Space Before: 0 pt, After: 0 pt

1152 [94. Leone A, Di Gennaro E, Bruzzese F, Avallone A, Budillon A. New perspective for an](#)
 1153 [old antidiabetic drug: metformin as anticancer agent. Cancer Treat Res. 2014;159:355-76.](#)

1154 [104. Serfin J, Carragher J, Groman A, Dexter EU, Yendamuri S. Outcome prediction using](#)
 1155 [markers of aerobic glycolysis \(the Warburg effect\) varies between tumor regions in stage I](#)
 1156 [non-small cell lung cancer. J Mol Biomark Diagn. 2011;2\(116\). DOI:10.4172/2155-](#)
 1157 [9929.1000116](#)

1158 [105. Zeng W, Liu P, Pan W, Singh SR, Wei Y. Hypoxia and hypoxia inducible factors in](#)
 1159 [tumor metabolism. Cancer Lett. 2014; \(*In press*\). DOI 10.1016/j.canlet.2014.01.032.](#)

1160 [115. Arany Z, Foo SY, Ma Y, Ruas JL, Bommi-Reddy A, Girnun G, et al. HIF-independent](#)
 1161 [regulation of VEGF and angiogenesis by the transcriptional coactivator PGC-1alpha.](#)
 1162 [Nature. 2008;451:1008-12. doi:10.1038/nature06613](#)

1163

1164 [120. Samudio I, Fiegl M, McQueen T, Clise-Dwyer K, Andreeff M. The Warburg effect in](#)
 1165 [leukemia-stroma cocultures is mediated by mitochondrial uncoupling associated with](#)
 1166 [uncoupling protein 2 activation. Cancer Res. 2008;68:5198-205.](#)

1167 [121. Samudio I, Fiegl M, Andreeff M. Mitochondrial uncoupling and the Warburg effect:](#)
 1168 [molecular basis for the reprogramming of cancer cell metabolism. Cancer Res.](#)
 1169 [2009;69:2163-66.](#)

1170 [122. Ayyasamy V, Owens KM, Desouki MM, Liang P, Bakin A, Thangaraj K, et al.](#)
 1171 [Cellular model of Warburg effect identifies tumor promoting function of UCP2 in breast](#)
 1172 [cancer and its suppression by genipin. PLoS One. 2011;6\(9\):e24792. DOI:](#)
 1173 [10.1371/journal.pone.0024792](#)

1174 [123. Coulouarn, C., Gomez-Quiroz, L. E., Lee, J.-S., Kaposi-Novak, P., Conner, E. A.,](#)
 1175 [Goldina, T. A., et al. Oncogene-specific gene expression signatures at preneoplastic](#)
 1176 [stage in mice define distinct mechanisms of hepatocarcinogenesis. Hepatology.](#)
 1177 [2006;44:1003-11. DOI: 10.1002/hep.21293](#)

1178 [124. Calle EE, Thun MJ. Obesity and cancer. Oncogene. 2004;23:6365-78.](#)
 1179 [doi:10.1038/sj.onc.1207751](#)

Formatted: Not Highlight

Formatted: English (United States)

Formatted: Indent: First line: 0.39"

- 1180 [125. Bianchini F, Kaaks R, Vainio H. Overweight, obesity, and cancer risk. *Lancet Oncol*. 2002;3\(9\):565-74.](#) Formatted: Indent: Left: 0", Hanging: 0.39"
- 1181
- 1182 [127. Spiegel D, Giese-Davis J. Depression and cancer: mechanisms and disease](#)
- 1183 [progression. *Biol Psychiat*. 2003;54\(3\):269-82. DOI: 10.1016/S0006-](#)
- 1184 [3223\(03\)00566-3](#)
- 1185 [128. Gripp S, Moeller S, Bölke E, Schmitt G, Matuschek C, Asgari S, et al. Survival](#) Formatted: German (Germany)
- 1186 [prediction in terminally ill cancer patients by clinical estimates, laboratory tests, and](#) Formatted: Indent: Left: 0", Hanging: 0.42"
- 1187 [self-rated anxiety and depression. *J Clin Oncol*. 2007;25\(22\):3313-20. DOI:](#)
- 1188 [10.1200/JCO.2006.10.5411](#)
- 1189 [129. Wellisch DK, Lindberg NM. A psychological profile of depressed and nondepressed](#) Formatted: Indent: Left: 0", Hanging: 0.39"
- 1190 [women at high risk for breast cancer. *Psychosomatics*.2001;42\(4\):330-36. DOI:](#)
- 1191 [10.1176/appi.psy.42.4.330](#)
- 1192 [130. Fife A, Beasley PJ, Fertig DL. Psychoneuroimmunology and cancer: historical](#)
- 1193 [perspectives and current research. *Adv Neuroimmun*. 1996;6\(2\):179-90. DOI:](#)
- 1194 [10.1016/0960-5428\(96\)00016-2](#)
- 1195 [134. Federico A, Morgillo F, Tuccillo C, Ciardiello F, Loguercio C. Chronic inflammation](#)
- 1196 [and oxidative stress in human carcinogenesis. *Int J Cancer*. 2007;121\(11\):2381-86.](#)
- 1197 [DOI: 10.1002/ijc.23192](#)
- 1198 [135. Wu Y, Antony S, Meitzler JL, Doroshow JH. Molecular mechanisms underlying](#)
- 1199 [chronic inflammation-associated cancers. *Cancer Lett*. 2014;345\(2\):164-73.](#)
- 1200 [136. Read SA, Douglas MW. Virus induced inflammation and cancer development. *Cancer*](#)
- 1201 [Lett. 2014;345\(2\):174-81.](#)
- 1202 [137. Marusawa H, Jenkins BJ. Inflammation and gastrointestinal cancer: an overview.](#)
- 1203 [Cancer Lett. 2014;345\(2\):153-56.](#)
- 1204 [138. Zhao J, Lawless MW. Stop feeding cancer: Pro-inflammatory role of visceral](#)
- 1205 [adiposity in liver cancer. *Cytokine*. 2013;64\(3\):626-37.](#)
- 1206 [139. Vaughan RA, Garcia-Smith R, Trujillo KA, Bisoffi M. Tumor necrosis factor alpha](#)
- 1207 [increases aerobic glycolysis and reduces oxidative metabolism in prostate epithelial](#)
- 1208 [cells. *Prostate*. 2013;73\(14\): 1538-46.](#)

1209 [140. Vaughan RA, Garcia-Smith R, Dorsey J, Griffith JK, Bisoffi M, Trujillo KA. Tumor](#)
 1210 [necrosis factor alpha induces Warburg-like metabolism and is reversed by anti-](#)
 1211 [inflammatory curcumin in breast epithelial cells. Int J Cancer. 2013;133\(10\): 2504-](#)
 1212 [10.](#)

1213 [141. Jiang S, Zhang LF, Zhang HW, Hu S, Lu MH, Liang S, et al. A novel miR-155/miR-](#)
 1214 [143 cascade controls glycolysis by regulating hexokinase 2 in breast cancer cells.](#)
 1215 [EMBO J. 2012;31\(8\):1985-98.](#)

1216 [142. Cottrill KA, Chan SY. Metabolic dysfunction in pulmonary hypertension: the](#)
 1217 [expanding relevance of the Warburg effect. Eur J Clin Invest. 2013;43\(8\): 855-65.](#)

1218 [143. San Martín A, Ceballo S, Ruminot I, Lerchundi R, Frommer WB, Barros LF. A](#)
 1219 [genetically encoded FRET lactate sensor and its use to detect the Warburg effect in](#)
 1220 [single cancer cells. PLoS ONE. 2013;8\(2\):e57712.](#)

1221 [144. ~~134. Federico A, Morgillo F, Tuccillo C, Ciardiello F, Loguercio C. Chronic~~](#)
 1222 [inflammation and oxidative stress in human carcinogenesis. Int J Cancer.
 1223 \[2007;121\\(11\\):2381-86. DOI: 10.1002/ijc.23192\]\(#\)](#)

1224 [Lee KW, Bode AM, Dong Z. Molecular targets of phytochemicals for cancer prevention.](#)
 1225 [Nat Rev Cancer. 2011;11:211-18. DOI:10.1038/nrc3017](#)

Formatted: English (United States)

Formatted: Font: Not Bold

Formatted: Indent: Left: 0", Hanging: 0.39"

Formatted: Normal, Justified, Indent: Left: 0",
Hanging: 0.42", Don't adjust space between
Latin and Asian text, Don't adjust space
between Asian text and numbers

1239
1240
1241
1242
1243
1244
1245
1246
1247

Formatted: Normal, Justified, Indent: Left: 0",
Don't adjust space between Latin and Asian
text, Don't adjust space between Asian text and
numbers