

Original Research Article

Anti-venom Activity of *Mucuna pruriens* Leaves Extract Against Cobra Snake

(*Naja hannah*) Venom

Okere O. Shekins¹, Gabriel O. Anyanwu^{1*}, Philip. M. Nmadu², Olufunsho D. Olowoniyi³

¹Department of Biochemistry, Faculty of Science and Technology, Bingham University, Nasarawa State, Nigeria.

²Department of Biological Sciences, Faculty of Science and Technology, Bingham University, Nasarawa State, Nigeria.

³Department of Science Laboratory Technology, Federal Polytechnic, Nasarawa State, Nigeria.

gabrielanyanwu@yahoo.com, gabrielanyanwu@yahoo.com,

ABSTRACT

Aim: The study was done to investigate the anti-venom activity of *Mucuna pruriens* leaves extract against cobra snake (*Naja hannah*) venom.

Study Design: The mice were randomly grouped into six groups (A, B, C, D, E, and F) of five rats each. Group A served as the normal control (no induction), and the mice in the group were given normal saline (1ml/kg/body weight). Group B served as the test control (snake venom was induced but no treatment administered), Group C served as the standard control (snake venom was induced and treated with antivenin, a standard drug), Group D, E and F were all induced with the cobra snake venom and treated with ethanolic extracts of the leaves of *M. pruriens* for 14days.

Methodology: The induction with cobra snake venom was done with 0.075mg/kg b.w of venom and thereafter the treatment with *M. pruriens* extract for Group D, E and F were done with 40 mg/ kg, 60 mg/ kg and 80 mg/ kg respectively intraperitoneally in the mice. Serum blood of the animals was used to assay for total cholesterol, bilirubin, AST, ALT, GSH and catalase levels after 14days.

Result: The injection of crude venom of cobra snake (*Naja hannah*) caused an increase in cholesterol, AST, ALT, bilirubin, catalase and glutathione in envenomated mice which

significantly reduced ($p < 0.05$) compared to all the controls after 14 days of treatment with the extract.

Conclusion: The results suggests that 80 mg/ kg of the plant extract is more effective than the standard drug, therefore *M. pruriens* leaves has a greater anti-venom potential for curing snake bite, than antivenin.

Keywords: Anti-venom, *Mucuna pruriens*, Antivenin, Cobra snake, Haemorrhage

1.0 INTRODUCTION

Each year in the world a lot of people receive venomous bites by snake and about 40,000 of them die [1]. *Echis carinatus* and *Naja hannah* are the major causes of snakebite deaths in Plateau State, Nigeria. Snake venoms of Viperidae and Elapidae are known to consist of a complex mixtures of toxins and enzymes which are responsible for haemorrhage, myonecrosis, neurotoxicity and alteration of blood coagulation [2,3].

The only effective treatment of the modern medicine for serious snakebites is the use of the antidote (antivenin), derived from antibodies, produced in horse's blood serum after injecting the animal with snake venom. In humans, antivenin is administered either through the veins or injected into muscle and acts by neutralizing the snake venom which has entered the body. Because antivenin is obtained from horses, snakebite victims sensitive to horse products must be very carefully treated. The danger is that they could develop an adverse reaction or even an anaphylactic shock. Moreover the efficacy of the serums is dependent on the rapidity with which the specificity treatment is

started and it is only specific for the venom used for the immunization. These represent an important limitation of this kind of therapy.

The genus *Mucuna*, belonging to the Fabaceae family, sub-family Papilionaceae, includes approximately 150 species of annual and perennial legumes. Among the various under-utilized wild legumes, the velvet bean *Mucuna pruriens* (“Cowitch” and “cowhage” are the common English names) is widespread in tropical and sub-tropical regions of the world. It is considered a viable source of dietary proteins [4,5] due to its high protein concentration (23–35%) in addition its digestibility, which is comparable to that of other pulses such as soybean, rice bean, and lima bean [6]. It is therefore regarded as a good source of food.

M. pruriens is a popular Indian medicinal plant, which has long been used in traditional Ayurvedic Indian medicine, for diseases including Parkinsonism [7]. The beans have also been employed as a powerful aphrodisiac [8] and have been used to treat nervous disorders and arthritis [9]. The bean, if applied as a paste on scorpion stings, is thought to absorb the poison [9]. The non-protein amino acid-derived L-dopa (3, 4-dihydroxy phenylalanine) found in this underutilized legume leaves resists attack from insects, and thus controls biological infestation during storage. According to D’Mello, all anti-nutritional compounds confer insect and disease resistance to plants [10]. Further, L-dopa has been extracted from the leaves to provide commercial drugs for the treatment of Parkinson’s disease. L-Dopa is a potent neurotransmitter precursor that is believed, in part, to be responsible for the toxicity of the *Mucuna* leaves [11]. Antiepileptic and anti-neoplastic activity of ethanolic extract from *M. pruriens* had been reported [12].

An ethanolic extract of *M. pruriens* leaves has demonstrated significant in vitro anti-oxidant activity, and there are also indications that ethanolic extracts of *Mucuna pruriens* may be a potential source of natural anti-oxidants and anti-microbial agents [13]. All parts of *M. pruriens* possess valuable medicinal properties and it has been investigated in various contexts, for its anti-diabetic, aphrodisiac, anti-neoplastic, anti-epileptic, and anti-microbial activities [7]. Its anti-venom activities have been investigated and its anti-helminthic activity has been demonstrated [14,15]. *M. pruriens* has also been shown to be neuroprotective [16], and has demonstrated analgesic and anti-inflammatory activity [17]. *M. pruriens* has been reported to be a potential natural source of antioxidants with greater importance as a therapeutic agent in preventing or slowing oxidative stress related degenerative diseases [18]. In Nigeria, where the *M. pruriens* seeds are locally prescribed as an oral prophylactic for snake bite, it is claimed that when two seeds are swallowed, they protect the individual for a year against snake bites [19]. Therefore, the primary objective of the present study is to confirm the bioactivity of *M. pruriens* leaves extract against cobra snake (*Naja hannah*) venom.

2.0 METHODOLOGY

2.1 Collection and Preparation of the Plant Sample

The fresh leaves of *M. pruriens* were collected from Kogi State, Nigeria. It was identified by Dr. S. M. Ayodele (Dept. of Botany, Kogi State University). The leaves were dried under room temperature for 5days. The dried sample was ground powdery form, using the warring commercial blender.

2.2 Extraction of *M. pruriens* leaves

The plant material (500g) was defatted with 400ml hexane (C₆ H₁₄) by using a Soxhlet apparatus for 5h. The defatted powder plant material was air-dried. The air-dried defatted powdered plant material was then extracted with 400 ml ethanol (C₂H₅OH) by using a Soxhlet apparatus for 8h. The residue was dried over night and extracted with 250 ml water (H₂O) by using a shaking water bath at 70⁰C for 2h. The extraction with water was repeated three time. The water filtrates were mixed together. The ethanol and water extract were filtered and evaporated by using a rotary evaporator and freeze dryer to give the crude-dried extract. The dried extracts were stored at -20⁰C until used.

Calculation:

$$\text{Percentage yield (\%)} = \frac{\text{weight of the plant extract}}{\text{Weight of the dried plant used}} \times 100$$

2.3 Proximate Analysis af *M. pruriens*

The moisture content, ash content, carbohydrate content, crude fibre and crude protein were determined using methods as described by AOAC [20].

2.4 Phytochemical Screening

The ethanolic extract of *M. pruriens* leaves were screened for the presence of phytochemical compound as described by Treatise and Evans [21] and Sofowora [22].

2.5 Animals

Experimental mice were purchased from the animal house of NIPRID. The animals were housed in steel cages and kept at room temperature. The mice had no history of

drug consumption that is; they had not been used for any investigation. The mice were put on standard mice pellet (feed) and pure drinking water and allowed to get acclimatized for 21 days before the start of the experiment. The study was done in accordance with the guidelines for animal use of Faculty of Sciences, Bingham University and had the project ID: BCH/02/012.

2.6 Induction of Anti-Snake Venom

The cobra snake venom was procured in a sample bottle from National Veterinary Research Institute, Vom, Plateau State, Nigeria. The cobra snake venom native preparations were given intraperitoneally (i.p) to the mice at a dose which were proportional to the weight of the animals. The volumes of preparation were identical and the same amounts were injected. The venom dose used in this study was estimated according to our preliminary experiments.

2.7 Experimental Design

White male albino mice of wistar strain of body weight ranging between 15-30g were used for the research study. The mice were randomly grouped into six groups (A, B, C, D, E, and F) of five rats each. Group A served as the normal control (no induction), and the mice in the group were given normal saline (1ml/kg body weight). Group B served as the test control (snake venom was induced but no treatment administered), Group C served as the standard control (snake venom was induced and treated with antivenin, a standard drug), Group D, E and F were all induced with the cobra snake venom and treated with ethanolic extracts of the leaves of *M. pruriens* (Table 1).

Table 1: Experimental Design

Group	Name	Treatment for 14 days
A	Normal Control	No induction, No treatment
B	Test Control	Induced with 0.075mg/kg b.w of venom, but no treatment
C	Standard Control	Induced with 0.075mg/kg b.w of venom and treated with antivenin (a standard drug).
D	Venom + 40mg/kg	Induced with 0.075mg/kg b.w of venom and treated with 40mg/kg extract
E	Venom + 60mg/kg	Induced with 0.075mg/kg b.w of venom and treated with 60mg/kg extract
F	Venom + 80mg/kg	Induced with 0.075mg/kg b.w of venom and treated with 80mg/kg extract

2.8 Determination of Serum Cholesterol Level

Total cholesterol was determined by the enzymatic endpoint method as described by Trinder [23]. In this method, cholesterol was determined after enzymatic hydrolysis and oxidation in a series of reaction. The indicator quinoneimine used in this method was formed from hydrogen peroxide and 4-aminoantipyrine in the presence of phenol and peroxidase.

2.9 Determination of Serum Bilirubin Level (Vander Bergh's Reaction)

The total bilirubin in serum or plasma is determined using the method of Jendrassik and Gróf [24] by coupling with diazotized sulfanilic acid after the addition of caffeine, sodium benzoate and sodium acetate. A blue azobilirubin is formed in alkaline Fehling solution II. This blue compound can also be determined selectively in the presence of yellow by products (green mixed coloration) by photometry at 578 nm.

2.10 Determination of Transaminases (ALT and AST)

Alanine Aminotransferase (ALT) was determined in a method as described by Reitman and Frankel [25]. In this method, ALT was measured by monitoring the concentration of pyruvate hydrazone formed with 2,4-dinitrophenylhydrazine in a reaction. Aspartate Aminotransferase (AST) was determined in a method as described by Reitman and Frankel [25]. In this method, AST is measured by monitoring the concentration of oxaloacetate hydrazone formed with 2,4-dinitrophenylhydrazine in a reaction.

2.11 Determination of Serum Glutathione level

The principle was based on the reduction of 5,5 dithiobis (2-nitrobenzoic acid) (DTNB) with reduced glutathione (GSH) to produce a yellow compound. The reduced chromogen was directly proportional to GSH concentration and its absorbance was measured at 405 nm [26].

2.12 Determination of Serum Catalase Level

Catalase was determined by the method described by Cohen *et al.* [27]. In this method, catalase catalysed the conversion of hydrogen peroxide to oxygen and water in a reaction.

2.13 Statistical analysis

Values are expressed as mean + S.E.M randomized complete block design analysis of variance was used for statistical analysis. P values less than 0.05 was considered significant.

174

175 **3.0 RESULTS**176 **3.1 Percentage Yield of Extract**

177 The extract was thick and greenish in colour, with an ethanolic extraction of
 178 *M. pruriens* leaves which indicated the yield of 6.73%.

179 **3.2 Phytochemical Screening**

180 The preliminary phytochemicals test reveals that the major phytochemical constituents
 181 in *M. pruriens* leaves are alkaloids, flavonoids, tannins, saponins, steroids, **terpenoids**,
 182 cardiac glycosides and anthraquinones (Table 2).

183

184 Table 2: Phytochemical composition of *M. pruriens* leaves

Phytochemicals	Presence in <i>M. pruriens</i> leaves
Alkaloids	++
Flavonoids	++
Tannins	+
Saponins	++
Steroids	++
Terpenoids	++
Cardiac glycosides	+
Anthraquinones	+

185 (+) indicates weak presence; (++) indicates strong presence

186 **3.3 Proximate Analysis**

187 The proximate analysis of the leaves in Table 3 has a moisture content of 11.37%,
 188 crude fiber 31.91%, crude fat 2.97%, carbohydrate 45.65%, Ash content 3.00%.

189

Table 3: Proximate composition of *M. pruriens* leaves

PARAMETER	COMPOSITION
Moisture content	11.37 %
Crude protein	31.91%
Crude fat	2.97 %
Carbohydrate	45.65 %
Ash	3.00 %

3.4 Effect of Venom Induction and Extract on Biochemical Parameters

The results revealed in Table 4a and 4b that, the injection of crude venom of cobra snake (*Naja hannah*) caused an increase in cholesterol, AST, ALT, bilirubin, catalase and glutathione in envenomated mice compared to the normal control mice. There was no significant difference in the levels of cholesterol, AST, ALT, bilirubin, catalase and glutathione of the treated groups when compared to the test control group.

Table 4a: Result obtained on the effect of extract on Cholesterol, AST and ALT at day one after venom induction

Treatment	Cholesterol (mg/dl)	AST (U/l)	ALT (U/l)
Normal control	55.94 ± 0.07	28.93 ± 0.14	22.60 ± 0.49
Test control	127.41 ± 0.13 ^a	74.34 ± 2.24 ^a	36.64 ± 0.64 ^a
Standard control	125.91 ± 0.06 ^a	74.96 ± 0.18 ^a	34.96 ± 0.35 ^a
Venom + 40mg/kg	127.48 ± 0.05 ^a	75.67 ± 0.21 ^a	36.20 ± 0.42 ^a
Venom + 60mg/kg	128.30 ± 0.04 ^a	73.61 ± 0.14 ^a	37.10 ± 0.21 ^a
Venom + 80mg/kg	128.11 ± 0.03 ^a	73.93 ± 0.07 ^a	37.04 ± 0.14 ^a

Values are expressed as means ± SEM. ^a indicate values that are significantly different when compared to the normal control at (p < 0.05).

Table 4b: Result obtained on the effect of extract on Bilirubin, Catalase and Glutathione at day one after venom induction

Treatment	Bilirubin (mg/dl)	Catalase (mg/dl)	Glutathione (U/L)
Normal control	0.42 ± 0.04	17.19 ± 0.05	4.69 ± 0.51
Test control	0.95 ± 0.07 ^a	25.45 ± 0.93 ^a	8.47 ± 1.21 ^a
Standard control	0.90 ± 0.02 ^a	26.06 ± 0.06 ^a	8.21 ± 0.68 ^a
Venom + 40mg/kg	0.94 ± 0.05 ^a	25.82 ± 0.74 ^a	8.37 ± 0.74 ^a
Venom + 60mg/kg	0.91 ± 0.03 ^a	27.48 ± 0.62 ^a	7.95 ± 0.55 ^a
Venom + 80mg/kg	0.95 ± 0.02 ^a	25.59 ± 0.50 ^a	8.11 ± 0.48 ^a

Values are expressed as means ± SEM. ^a indicate values that are significantly different when compared to the normal control at (p < 0.05).

The levels of cholesterol, AST, ALT, bilirubin, catalase and glutathione were significantly reduced when compared to the test control after 14 days of treatment with the extract (Table 5a and 5b).

Table 5a: Result obtained on the effect of extract on Cholesterol, AST and ALT at day fourteen

Treatment	Cholesterol (mg/dl)	AST (U/l)	ALT (U/l)
Normal control	50.00 ± 0.07	25.54 ± 0.07	17.46 ± 0.20
Test control	113.45 ± 0.17	62.25 ± 1.24	38.88 ± 0.44
Standard control	60.00 ± 0.06 ^{ab}	30.37 ± 0.19 ^{ab}	19.71 ± 0.28 ^{ab}
Venom + 40mg/kg	96.67 ± 0.05 ^{abc}	40.45 ± 0.21 ^{abc}	26.24 ± 0.35 ^{abc}
Venom + 60mg/kg	83.33 ± 0.04 ^{abc}	36.39 ± 0.14 ^{abc}	21.46 ± 0.21 ^{ab}
Venom + 80mg/kg	55.67 ± 0.03 ^{abc}	28.82 ± 0.09 ^{abc}	18.41 ± 0.17 ^b

Values are expressed as means ± SEM. ^a indicate values that are significantly different when compared to the normal control at (p < 0.05), ^b indicate values that are significantly different when compared to the test control at (p < 0.05) and ^c indicate values that are significantly different when compared to the standard control at (p < 0.05).

Table 5b: Result obtained on the effect of extract on Bilirubin, Catalase and Glutathione at day fourteen

Treatment	Bilirubin (mg/dl)	Catalase (mg/dl)	Glutathione (U/L)
Normal control	0.34 ± 0.02	14.16 ± 0.32	2.59 ± 0.31
Test control	0.98 ± 0.07	20.45 ± 1.30	6.47 ± 1.21
Standard control	0.47 ± 0.03 ^{ab}	13.06 ± 0.56 ^b	1.83 ± 0.33 ^{ab}
Venom + 40mg/kg	0.76 ± 0.04 ^{abc}	16.67 ± 0.95 ^{abc}	4.32 ± 0.49 ^{abc}
Venom + 60mg/kg	0.54 ± 0.03 ^{abc}	15.89 ± 0.84 ^{bc}	3.97 ± 0.22 ^{abc}
Venom + 80mg/kg	0.41 ± 0.01 ^{abc}	12.95 ± 0.72 ^{abc}	1.82 ± 0.12 ^{ab}

Values are expressed as means ± SEM. ^a indicate values that are significantly different when compared to the normal control at (p < 0.05), ^b indicate values that are significantly different when compared to the test control at (p < 0.05) and ^c indicate values that are significantly different when compared to the standard control at (p < 0.05).

4.0 DISCUSSION

The phytochemical constituents in *M. pruriens* leaves are alkaloids, flavonoids, tannins, saponins, steroids, terpenoids, cardiac glycosides and anthraquinones (Table 2). The anti-venom activity observed in the mice treated with *M. pruriens* extract may be attributed to the presence of any of these compounds alkanoids, tannins, flavonoids, steroids and terpenoid [28]. The proximate analysis of the leaves in Table 3 has a moisture content of 11.37%, crude fiber 31.91%, crude fat 2.97%, carbohydrate 45.65%, Ash content 3.00%. The Ash content is known to enhance digestibility, slow down the release of glucose into the blood stream and reduces blood cholesterol level.

The present study revealed in Table 4a and 4b that, the injection of crude venom of cobra snake (*Naja hannah*) caused an increase in cholesterol, AST, ALT, bilirubin,

catalase and glutathione in envenomated mice which significantly reduced after 14 days of treatment with the extract as shown in Table 5a and 5b. These findings are in agreement with other investigators who reported that the reduction in cholesterol, AST, ALT, bilirubin and catalase in envenomated mice was observed in laboratory animals treated with the extracts of *M. pruriens* leaves [29]. It might be assumed that, the increased levels of these serum constituents could be due to disturbance in renal functions as well as haemorrhages in some internal organs when challenged with a snake venom. In addition, the increasing in vascular permeability and haemorrhages in vital organs due to the toxic action of various snake venoms have been reported [30, 31]. Also, the reduction in serum cholesterol, AST, ALT, catalase, glutathione albumin and total bilirubin levels in the envenomated mice could be attributed to the anti-venom potentials of the extract of *M. pruriens* administered.

Furthermore, acute renal damage together with glomerular, tubular and vascular lesions following various snake bites have been reported [32,33,34] with additional, increased vascular permeability and hemorrhages in various vital organs. Another factor is the increase vascular permeability due to toxic action of the venom which could contribute to the low level of protein from plasma and tissue [35]. An earlier study reported that injection of *M. pruriens* seed extract conferred protection against *C. rhodostoma* venom, as the pretreatment attenuated (albeit only partially) the cardio-respiratory depressant effects of *C. rhodostoma* venoms in anesthetized rats [36].

In Table 4a, elevation of ALT and AST in the mice administered with venom as observed in the serum have serious implication on health of the animals. Such elevations are found in cases of both liver damage and myocardial infarction [37]. The elevation of AST and ALT makes the liver a target of suspicion as this is usual in cases of hepatotoxicity caused by toxic agent (Rosalki, 1974). From the experiment it was observed that some of the mice died after 30 minute of induction with the snake venom except the (Normal control) and the ones treated with different dose of the venom/extract (Groups D-F) which survived till the end of the experiment.

There was significant difference between the time of death in the extract treated group and those treated with venom only showing that the plant extract had effect on the activity of the venom. Thus, it was obvious that *M. pruriens* leaves did show greater anti-venom activity, the extract of *M. pruriens* showed a better anti-snake activity compared with antivenin. This can be attributed to the fact that tannins are able to non-specifically bind to *Naja hannah* venom proteins and precipitate them, thus provoking the anti-lethal effects. However, a report suggested that the protective effect of *M. pruriens* extract pretreatment against cobra venom involved a direct action of *M. pruriens* extract on the heart, and, as such, immunological neutralization is not the only mechanism of protective effect of the *M. pruriens* extract pretreatment [38]. Also, the protective effect of *M. pruriens* seed against snake venom poisoning has been reported [39], which corroborates the results of this study that *M. pruriens* plant has great potential for use in the treatment of common cobra bites from the *Naja* subspecies.

285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304

CONCLUSION

In conclusion, this investigation revealed that the ethanolic extract of *M. pruriens* leaves has the following phytochemical components of saponin, terpenoid, flavonoids, steroids, alkaloids, anthraquinones. It was observed that 80 mg/ kg of the plant extract is more effective than the standard drug, therefore *M. pruriens* leaves has a greater medicinal plant for curing snake bite, than anti-venin.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

CONSENT

Not applicable

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

REFERENCES

1. Chippaux JP. Snake Bite: Appraisal of the Global Situation. Bulletin WHO 1998;76:515-524.

305 2. Markland F. Snake venoms and the hemostatic system. *Toxicon*. 1998;36, 1749–
 306 1800.

307 3. Warrell DA. Snake venoms in science and clinical medicine. Russell's viper: biology,
 308 venom and treatment of bites. *Trans Roy Soc Trop Med Hyg*. 1989;83, 732–740.

309 4. Janardhanan K, Gurumoorthi P, Pugalenth M. Nutritional potential of five accessions
 310 of a South Indian tribal pulse, *Mucuna pruriens* var. utilis. Part I. The effect of
 311 processing methods on the contents of L-Dopa, phytic acid, and oligosaccharides. *J*
 312 *Trop and Subtrop Agro-ecosys*. 2003;1, 141-152.

313 5. Pugalenth M, Vadivel V, Siddhuraju P. Alternative Food/Feed Perspectives of an
 314 Underutilized Legume *Mucuna pruriens* var. Utilis—A Review. *Plant Foods Hum. Nutr.*
 315 2005;60: 201–218.

316 6. Gurumoorthi P, Pugalenth M, Janardhanan K. Nutritional potential of five accessions
 317 of a south Indian tribal pulse *Mucuna pruriens* var. utilis; II. Investigation on total free
 318 phenolics, tannins, trypsin and chymotrypsin inhibitors, phytohaemagglutinins, and in
 319 vitro protein digestibility. *Trop. Subtrop. Agroecosys*. 2003;1,153-158.

320 7. Sathyanarayanan G, Garg PK, Prasad H, Tandon RK. Elevated level of interleukin-6
 321 predicts organ failure and severe disease in patients with acute pancreatitis. *J*
 322 *Gastroenterol Hepatol*. 2007;22:550-4.

323 8. Amin KMY, Khan MN, Zillur-Rehman S, Khan NA. Sexual function *Mucuna pruriens*
 324 roving effect of *Mucuna pruriens* in sexually normal male rats. *Fitoterapia*. 1996; 67, 53-
 325 56.

- 326 9. Jeyaweera, DMA. Madicinal plants used in Ceylon. Colombo, Sri Lanka; National
327 Science Council of Sri Lanka, 1981.
- 328 10. D'Mello JPF. Leguminous leaf meals in non-ruminant nutrition. In: Tropical Legumes
329 in Animal Nutrition. J.P.F. D'Mello and C. Devendra, Eds. CAB International,
330 Wallingford, UK. 1995.
- 331 11. Lorenzetti E, Maclsaac S, Arnason JT, Awang DVC, Buckles D. The
332 phytochemistry, toxicology and food potential of velvet bean (*Mucuna adans* spp.
333 Fabaceae). In D. Buckles, O. Osiname, M. Galiba, & G. Galiano, Cover crops of West
334 Africa; contributing to sustainable agriculture. IDRC, Ottawa, Canada & IITA, Ibadan,
335 Nigeria, 1998.
- 336 12. Gupta M, Mazumder UK, Chakraborti S, Bhattacharya S, Rath N, Bhawal SR.
337 Antiepileptic and anticancer activity of some indigenous plants. Indian J. Physiol. Allied
338 Sci. 1997;51, 53-56.
- 339 13. Rajeshwar, Y., Kumar, S.G.P., Gupta, M., Mazumder, K.U., (2005). Studies on in
340 vitro antioxidant activities of mhetanol extract of *Mucuna pruriens* (Fabaceae) seeds.
341 Euro Bull Drug Res., 13, 31-39.
- 342 14. Guerranti R, Aguiyi JC, Errico E, Pagani R, Marinello E. Effects of *Mucuna pruriens*
343 extract on activation of prothrombin by *Echis carinatus* venom. J Ethnopharmacol.
344 2001;75, 175-180.
- 345 15. Jalalpure SS, Alagawadi KR, Mahajanashell CS. In vitro antihelmintic property of
346 various seed oils against *Pheritima posthuma*. Ind. Pharm. Sci. 2007;69, 158-160.

- 347 16. Misra L, Wagner H. Extraction of bioactive principles from *Mucuna pruriens* seeds.
348 Indian J. Biochem. Biophys. 2007;44, 56-60.
- 349 17. Hishika R, Shastry S, Shinde S, Gupta SS. Preliminary, phytochemical and anti-
350 inflammatory activity of seeds of *Mucuna pruriens*. Indian J. pharmacol. 1981;13 (1), 97-
351 98.
- 352 18. Sonpetkar JM, Nipate SS, Shenoy PA, Chaudhari PD. In vitro antioxidant activity of
353 ethanolic extract of *Mucuna pruriens* (L.) DC. Seeds. J Pharm Res. 2012;5 (5):2769.
- 354 19. Scirè A, Tanfani F, Bertoli E, Furlani E, Nadozie HO, Cerutti H, Cortelazzo A, Bini L,
355 Guerranti R. The belonging of gpMuc, a glycoprotein from *Mucuna pruriens* seeds, to
356 the Kunitz-type trypsin inhibitor family explains its direct anti-snake venom activity.
357 Phytomed. 2011;18(10):887-95.
- 358 20. AOAC. Official Methods of analysis. 15th Edn. Association of Official Agricultural
359 Chemists, Washington, DC. 1990.
- 360 21. Treatise GE, Evans WC. Pharmacognosy. 11th edn. Brailliar Tiridel Can. Macmillan
361 publishers. Ltd, Ibadan, 1989.
- 362 22. Sofowora. A Medicinal Plants and Traditional Medicine in Africa. Spectrum Books
363 Ltd, Ibadan, 1993.
- 364 23. Trinder P. Determination of glucose in blood using glucose oxidase with an
365 alternative oxygen acceptor. Anal Clin Biochem. 1969;6: 24-27.
- 366 24. Jendrassik L, Grof P. Colorimetric Method of Determination of bilirubin. Biochem Z.
367 1938; 297:81-82

- 368 25. Reitman S, Frankel S. A colourimetric method for the determination of serum
369 glutamate-oxaloacetate and pyruvate transaminases. Am J Clin Pathol. 1957;28: 56-63.
- 370 26. Tietz NW. Clinical Guide to Laboratory Tests, Second Edition W. B. Saunders
371 Company, Philadelphia, USA,1990.
- 372 27. Cohen G, Dembiec D, Marcus J. (1970). Measurement of catalase activity in
373 tissues extracts. Anal. Biochem. 1998;34: 30-38.
- 374 28. Rajendran K, Annie S, Maneesh MR, Vijaya B. Invitro and In vivo anti snake venom
375 (Daboia russelli) studies on various leaf extracts of Acalpha indica Linn. Int J Phytomed.
376 2010;2: 217-220
- 377 29. Abdul-Nabi IM, Raafat A, El-Shany H. Biological effect of intraperitoneal injection of
378 rats with the venom of the snake *Echis carnetus*. Egypt J. Zool. 1997;29:195-205.
- 379 30. Meier J, Stocker K. Effect of venoms on homeostasis. Toxicol. 1999;21 (3): 1711-
380 1820.
- 381 31. Meier J, Theakston RDG. Approximate LD50 determinations of snake venoms using
382 eight to ten experimental animals. Toxicon 1986;24, 395 – 401.
- 383 32. Sitprija V, Suvanpha R, Pochanugool C, Chusil S, Tungsanga K. (1982). Acute
384 intestinal nephritis snakebite. Am J Trop Med and Hyg. 2000; 31:408-410.
- 385 33. Sani SM, Purandare NM. Autopsy study of cases of snake bite with special
386 reference to renal lesions. J. Postgrad Med. 1972;18: 181-188.

- 387 34. Aung-Khin M. Histological and ultrastructural changes of the kidney in renal failure
388 after viper envenomation. *Toxicon*. 1978;16: 71-75.
- 389 35. Olajide OA, Makinde JM, Awe SO. Effect of aqueous extract of *Mucuna pruriens* stem
390 bark on corragenan induced oedema and granuloma tissue formation in rats and mice.
391 *J. Ethnopharmacol*. 1999;66(1): 113-177
- 392 36. Fung SY, Tan NH, Sim SM. Protective effects of *Mucuna pruriens* seed extract
393 pretreatment against cardiovascular and respiratory depressant effects of *Calloselasma*
394 *rhodostoma* (Malayan pit viper) venom in rats. *Trop Biomed*. 2010;27(3): 366–372.
- 395 37. Gray C, Howorth PJN. *Clinical Chemical Pathology*. 9th ed. The ELBS, Edward
396 Arnold Publishers Ltd., 1982.
- 397 38. Fung SY, Tan NH, Sim SM, Aguiyi JC. Effect of *Mucuna pruriens* Seed Extract
398 Pretreatment on the Responses of Spontaneously Beating Rat Atria and Aortic Ring to
399 *Naja sputatrix* (Javan Spitting Cobra) Venom. *Evid Based Complement Alternat Med*.
400 2012;486390: 6.
- 401 39. Tan NH, Fung SY, Sim SM, Marinello E, Guerranti R, Aguiyi JC. The protective
402 effect of *Mucuna pruriens* seeds against snake venom poisoning. *J Ethnopharmacol*.
403 2009; 123: 356–358.

404