

IN VITRO ANTIMICROBIAL EFFECT OF CRUDE TANNINS ISOLATED FROM THE LEAF OF *ANNONA SENEGALENSIS*

AUTHORS: MAHMOUD, SULEIMAN JADA, smjada2007@yahoo.com, DEPARTMENT OF BIOCHEMISTRY, MODIBBO ADAMA UNIVERSITY OF TECHNOLOGY, YOLA, ADAMAWA STATE, NIGERIA.

WUROCHEKKE, ABDULLAHI USMAN, wchekke@yahoo.co.uk, DEPARTMENT OF BIOCHEMISTRY, MODIBBO ADAMA UNIVERSITY OF TECHNOLOGY, YOLA, ADAMAWA STATE, NIGERIA.

YA'U, ADAMU, adamuyau@rocketmail.com, DEPARTMENT OF BIOCHEMISTRY, MODIBBO ADAMA UNIVERSITY OF TECHNOLOGY, YOLA, ADAMAWA STATE, NIGERIA.

ABSTRACT

This study was aimed at evaluating the antimicrobial effect of tannin isolated from the leaf of *Annona senegalensis* on *Shigella dysenteriae*, *Escherichia coli* and *Salmonella typhi*. The study was carried out in the laboratory of Biochemistry department, Modibbo Adama University of Technology, Yola, Adamawa State, Nigeria, between July and October, 2013. The antimicrobial activity was carried out using agar disc diffusion method. The phytochemical screening of the leaf of the plant revealed the presence of tannins, alkaloids, saponins, flavonoids, terpenoids, cardiac glycosides and anthraquinones while phenols and steroids were absent. Quantitatively, tannins were found to be 24% in the leaf of the plant. Isolated tannin exhibited antibacterial activities against all the tested bacteria; *Shigella dysenteriae* gave the zone of inhibition (18.6mm), *Escherichia coli* (13.9mm) and *Salmonella typhi* (10.6mm). Minimum Inhibitory concentration of the crude tannins on the test organisms ranged from 6.25mg/ml to 12.5mg/ml. These findings confirmed the basis of traditional use of the leaf of *Annona senegalensis* in the treatment of diseases such as dysentery and diarrhea.

KEY WORDS: Antimicrobial, Phytochemicals, Zone of inhibition, and Crude tannins.

INTRODUCTION

Medicinal plants are cheap starting materials for the synthesis of new drugs and the chances of the body accepting preparations from plants are more than that prepared from laboratory [1]. Also, Fansoworth [2] has reported that aqueous decoction of drug has a greater bioavailability to the body than synthetic formulation today in the market. This is due to the fact that presentation of the active constituent of the plant material in the solubilized form brings about increase absorption of the extract by the body.

In recent years, the interest to evaluate plants possessing antibacterial activity for various diseases is still growing [3]. Herbal Medicine is still the main stream of about 75 to 80% of the whole

population and the major part of traditional therapy involves the use of plant extract and their active constituents [4]. Some plant has also been reported to exhibit antimicrobial activities against various pathogens [5].

Annona senegalensis commonly known as African custard apple is an important plant that is used in traditional medicine to cure some diseases like toothache and diarrhea. The leaf of the plant is used for treatment of cancerous tumor, intestinal trouble and venereal disease. It is used for astringent and for treating dysentery and malaria [6]. Locally, the plant is known as “Gwandar daaji” in Hausa, “Uburu Ocha” in Igbo and Abo in Yoruba [7]. The part of the plant that is used for therapeutic purposes include root, bark, leaf, flower and seed.

Tannins are polyphenols that are obtained from various parts of different plants [8]. In addition to use in leather processing industries, tannin have shown potential antiviral [9], anti bacterial [10][11] and anti parasitic effects [12]. In the past few years tannins have been studied for this effect against cancer through different mechanisms [13].

This work is designed to extract crude tannin from the leaf of *Annona senegalensis* and to evaluate its efficacy or otherwise on causative organisms of the dysentery..

MATERIALS AND METHODS

Collection of Plant Material

Fresh leaves of *Annona senegalensis* was collected in Sangere under Girei Local government Area of Adamawa state, Nigeria, in the month of July, 2013. It was identified and authenticated By Mr. Briston Pola of the department of Plant Science, Modibbo Adama University of Technology, Yola, Adamawa State, Nigeria.

Sources of Test Organisms

The microorganisms *Shigella dysenteriae*, *Salmonella typhi* and *Escherichia coli* were obtained from the Laboratory of Microbiology Department, Modibbo Adama University of Technology, Yola, Adamawa State, Nigeria. All the microorganisms are clinical isolates identified in the mentioned laboratory.

Preparation of the Extracts For Phytochemical Analysis

Exactly 50g of the powdered sample was weighed and transferred into 500ml of distilled water in a conical flask. This was allowed to stand for 24 hours at room temperature. The suspension was filtered using sterile Whatman No. 1 filter paper to obtain crude extract. A rotary evaporator was used to concentrate the extract. The concentrated extract obtained was stored in a refrigerator in a screw-cap bottle until when it was required.

Phytochemical Analysis

Chemical tests were carried out on the aqueous extracts to identify the constituents as described by Sofowora and Adewumi [14], Trease and Evans [15].

Extraction and Quantitative Determination of Tannins

The dry leaf was ground and 100mg of the fine powdered portion was dissolved with 1ml of concentrated HCl in methanol (1:100, v/v) in a water bath at 30°C for 30mins and was shaken for every 5 minutes. The sample was centrifuged at 1200g and the supernatant was collected. Extraction was repeated twice; the combined supernatant was then filtered, evaporated to dryness and the weight was taken. The extract was kept for the antimicrobial test[16].

Preparation of Culture Medium

Nutrient agar used for the study was manufactured by Titan Biotech. Ltd, Rajasthan, India and was prepared according to manufacturer's instructions.

Antimicrobial Activity Testing

The method described by Emeruwa [17] was used for the study. Wells were made using a cork borer on the surface of 20ml nutrient agar plates that was previously streaked with test organisms using 1.0McFarland standard (3×10^8 organisms). Extracted tannin of 0.2ml was aseptically introduced into the wells made; 100mg of ciprofloxacin was used as positive control which was added into a separate well. The plates were allowed to stand on the working bench for 30minutes and were incubated for 24hours at 37°C. The presence of zone of inhibition was regarded as the presence of antimicrobial activity and this was expressed as the length of the diameter of the zones of inhibition.

Determination of Minimum Inhibitory Concentration (MIC)

Different concentrations of tannins extracted from the leaf of *Annon asenegalensis* ranging between 1.5 and 50 mg/ml were introduced into different test tubes; each tube was inoculated with an overnight culture of *Salmonella typhi*, *Shigella dysenteriae* and *Escherichia coli*. The tubes were incubated at 37°C for 24 hours. The least concentration of tannin that did not permit any visible growth of the incubated test organism in broth culture was regarded as the Minimum Inhibitory Concentration (MIC) in each case[18].

RESULTS

The leaf of *Annona senegalensis* was found to contain tannins, alkaloids, saponins, flavonoids, terpenoids, anthraquinone and cardiac glycosides. Phenols and steroids were absent (Table 1). Quantitative determination of tannins yielded 24% in the fresh leaf of the plant.

The crude tannin exhibited antibacterial activities against all the tested bacteria; *Shigella dysenteriae* gave the zone of inhibition (18.6mm), *Escherichia coli* (13.9mm) and *Salmonella typhi* (10.6mm) as shown in table 2.

Determination Minimum Inhibitory concentration (MIC) of the crude tannins gave 6.25mg/ml on *Shigella dysenteriae* and 12.5mg/ml on both *Escherichia coli* and *Salmonella typhi*. This is presented in table 3.

Table 1: Phytochemical Screening Of Leaf of *Annona senegalensis*

Phytochemicals	Results
Tannins	++
Alkaloids	+
Saponins	+
Terpenoids	+
Phenols	-
Steroids	-
Anthraquinones	+
Flavonoids	++
Cardiac glycosides	++

Key: + = present, - = absent and ++ = highly present

Table 2: Antimicrobial activity of tannin isolated from the leaf of *Annona senegalensis*

Test Organisms	Zones of inhibition (mm)	
	Crude Tanins	Ciprofloxacin
<i>E.coli</i>	13.90	34.00
<i>S. typhi</i>	10.60	21.00
<i>S. dysenteriae</i>	18.60	26.00

Table 3: Minimum Inhibitory Concentration of Tannins Isolated from the Leaf of *Annona senegalensis*

Organisms	Concentrations (mg/ml)					
	50	25	12.5	6.25	3.0	1.5
<i>E. coli</i>	-	-	-	+	+	+
<i>S. typhi</i>	-	-	-	+	+	+
<i>S. dysenteriae</i>	-	-	-	-	+	+

Keys: + = indicates growth and - = indicates no growth

DISCUSSION

Phytochemical analysis of the extract of leaf of *Annona senegalensis* revealed the presence of chemical constituents namely tannins, alkaloid, saponin, flavonoid, terpenoid, cardiac glycoside and anthraquinone. However, the presence of alkaloid and anthraquinone was in contrast with the work of Tijjani *et al* [19], who reported that alkaloid and anthraquinone were absent in the leaf of the plant. The observed differences may be due to the different solvent used or where the plant is harvested or could be due to storage.

The various phytochemical compound detected are known to have beneficial medicinal value. There are established records that showed the benefits of these compounds detected in *Annona senegalensis*, for instance, tannins are reported to exhibit antiviral, antibacterial, antitumor activities. It was also reported that certain tannins are able to inhibit HIV replication selectively and is also used as diuretic[20].

Saponin in medicine, it is used in hypercholesterolaemia, hyperglycemia, antioxidant, anti-cancer, anti-inflammatory and weight loss etc. it is also known to have anti-fungi properties[21].

Flavonoids have been referred to as nature's biological response modifiers because of strong experimental evidence of their inherent ability to modify the body's reaction to allergic and carcinogens. They show anti-allergic, anti-inflammatory, anti-microbial and anti-cancer activity[22]. Anthraquinones are generally used as dyes and are also known as antibacterial agent[23].

Cardiac glycosides are used in the treatment of congestive heart failure. They are also used to strengthen a weakened heart and allow it to function more efficiently though the dosage must be controlled carefully, since the therapeutic is close to toxic dose[21]. Terpenoid are active against bacteria, fungi and viruses [24] and protozoa.

Moreover, these chemical constituent were reported to exhibit prominent effect on human and other animal system through different mechanisms of action which make them useful for medicinal purposes. For instance anthraquinone are known to complex irreversibly with nucleophilic amino acid in protein [25]. Often leading to inactivation of the protein and loss of function. For that reason, the potential range of quinone antimicrobial effect is great. Probable targets in the microbial cell are surface – exposed adhesions, cell wall polypeptides and membrane bound enzymes. Quinone may also render substrate unavailable to the microorganism[25].

Mechanism of action of alkaloids such as berberine and harmaline is attributed to their ability to intercalate with DNA / cell wall of an organism thereby disrupting the activity of the microorganism[26].

However, the mechanism of action of terpenoids is not fully understood but is speculated to involve in membrane disruption by the lipophilic compound. According to Mendez *et al*, [27] found that increasing the hydrophilicity of kaurenediterpenoids by addition of a methyl group drastically reduced their antimicrobial activity[21].

Cardiac glycosides are known to work by inhibiting the Na^+/K^+ pump. This causes an increase in the level of sodium ion in the myocyte which then lead to a rise in the level of Ca^{+} . This inhibition

increases the amount of Ca^{+} ion available for contraction of the heart, muscles which improve cardiac output and reduce distension of the heart [21].

Crude tannins from the leaf of the plant were found to be 24% and this is in agreement with the earlier study carried out by Simon [28] who obtained the same value. However, Bakar [29] reported that tannin is very rich in fruit of *Annona senegalensis* than in the leaf even though he has not mentioned the values.

The antimicrobial activity of tannins isolated from the leaf showed highest activity against *Shigella dysenteriae* and moderate in *Escherichia coli* and *Salmonella typhi* and ciprofloxacin was used as standard antibiotic as shown in table 2 .

The sensitivity observed in this study is in agreement with earlier investigation by Doss *et al.*, [30] and Akiyama *et al.*, [10]. It was reported by Jone *et al.* [31] that tannins has the ability to interfere with the bacterial cell wall synthesis where they formed complex with polysaccharide through so called non-specific forces such as hydrogen bonding and hydrogen effects as well as by covalent bond formation. Thus, tannin irreversibly inhibits the enzyme trans peptidase by reacting with serine residue in the trans peptidase. This reaction is irreversible and so the growth of the bacteria cell is inhibited. Also reported by Viljoen *et al.*, [32], the ability for the tannin compounds to cause the bacterial colonies to disintegrate probably results from their interference with the bacterial cell wall, thereby inhibiting the microbial growth.

The antimicrobial effect exhibited by ciprofloxacin gave higher zones of inhibition on the test organisms than the extracted tannin and this may be attributed to the concentration of the drug used.

The minimum inhibitory concentration (MIC) of tannin isolated in this study against the test organism ranged between 1.5 and 50mg/ml as shown in table 3. Antimicrobial agent with low activity against an organism had a high MIC while a highly active antimicrobial agent gave low MIC. For instance *Shigella dysenteriae* has the highest zone of inhibition at the same concentration with other organism, but gave the low MIC. The antibacterial effectiveness with the increasing concentration of tannins observed in the present study is in agreement with the earlier investigation as reported by Kurosaki and Nishi [33] and Banso and Adeyemo [34]. However, a minimum inhibitory concentration (MIC) is generally regarded as the most basic laboratory measurement of activity of an antimicrobial agent against an organism.

The clinical significance of Minimum Inhibitory Concentration (MIC) are used not only to determine the amount of antibiotic that the patient will receive but also the type of antibiotic used, which in turn lowers the opportunity for microbial agents. Applying MIC testing to number of bacterial strains in the same species provides an estimate of the concentration that inhibits 50% (MIC_{50}) and 90% (MIC_{90}) of bacterial isolates and can indicate shifts in the susceptibility of bacterial population to antibiotics [35].

CONCLUSION

In conclusion, the result of this study supports the traditional use of the *Annona senegalensis* for the treatment of diseases such as dysentery, diarrhea and typhoid. It also suggests that the tannin isolated from the test plant could serve as a potential source of antibiotics.

ACKNOWLEDGEMENTS.

The authors are very grateful to the department of Biochemistry of Modibbo Adama University of Technology for giving them the privilege to use their laboratory during the study. The authors are also indebted to the Chief Laboratory Technologist of the department for his assistance and valuable contributions.

REFERENCES

1. Sofowora, E.A. (1988). Medicinal Plants and Traditional Medicine in Africa. John Wiley and Sons Limited, Chechester.
2. Fansworth, N.R. (2001). The Present and Future of Pharmacognosy. News Letter of the Americans Society for Pharmacognosy, pp. 4 – 12.
3. Clark, A.M. and Hufford, C.D. (1993). Discco and Development of Novel Prototype Antibiotic for Opportunistic Infections related to the Acquired Immunodeficiency Syndrome. In human medical agents from plant American Chemical Society (ACS Symposium Series 534), Washington, DC., pp. 228 – 241.
4. Acharya, C.A. Deepak, K. and Shrivastava, A. (2008). Indigenous Herbal Medicine: Tribal Formulations and Traditional. Herbal Practices, Aavishkar Publisher Distributor, Jaipur – India. pp. 440.
5. Oladunmoye, M.K. (2006). Comparative Evaluation of Antimicrobial Activities and Phytochemical Screening of Two of Acalypha Wilkesiana. *Trends Applied Sci. Res.* Vo. 1, No. 1, p. 538 – 541.
6. Diallo, D. and Paulsen, B.S. (2000). Pharmaceutical Research and Traditional Practitioners in Mali: Experiences with Benefit sharing. In responding to bioprospecting (from biodiversity in the South to medicines in the North). Edited by: Svarstad H, Dhillion S.S., SpartacusForlagi. pp. 133 – 144.
7. Alqasim, A.M. (2013). *Annona senegalensis* pers on: A Multiposes Shrub, its science and technology. Volume 2, No. ISSN 2049- 7318.
8. Gajendiran, N. and Mahadevan, A. (1990). Utilization of Catechin by Rhizobium sp. *Plant Soil.* 108, 263 – 266.
9. Lin, L.U., Shuwen, L., Shibo, J. and Shuguang, W. (2004). Tannin inhibits HIV-1 entry by targeting pg. 41 .*Acta Pharmacol Sin.* 25(2):213 – 218. *Lioydia.* 39:375 – 377. (Pub Med).
10. Akiyama, H., Kazuyasu, F., Yamasaki, O., Oono, T. and Iwatsuki, K. (2001). Antibacterial action of several tannins against *Staphylococcus aureus*. *J. Antimicrobial Chemotherapy.* 48(4):487 – 491.
11. Funatogawa, K., Hayashi, S., Shimomura, H., Yoshida, T., Hatano, T., Ito, H. and Iria, Y. (2004). Antibacterial activity of hydrolysable tannins derived from medicinal plants against *Helicobacter pylori*. *Microbiol. Immunol.* 48(4):251 – 261.

12. Bhagavalli, A., Narayanana, Otto Geoffroya, Mark, C., Willingham, Gian, G., Reb and Daniel, W. Nixon (1999). Expression and its possible role in G1 arrest and apoptosis in ellagic acid treated cancer cells. *Cancer Letters*. 136, 215 – 221.
13. Sowemimo, A.A., Fakoya, F.A., Awopetu, I., Omobuwajo, O.R. and Adesanya, S.A. (2007). Toxicity and Mutagenic Activity of some selected Nigerian Plants. *J. Ethnopharmacol.* 113:427 – 432.
14. Sofowora, E.A. and Adewunmi, C.O. (1980). Preliminary screening of some plant extracts for molluscidal activity. *Plantamedica*. 39:57-65.
15. Trease, G.E. and Evans, I.C. (1989). *Pharmacognosy* (12th Edn) BailliereTindall London. pp. 21 – 22.
16. Morrison, I. M., Asiedu, E. A., Stuchbury, T., and Powell, A. A. (1995), Determination of Liginin and Tannin Contents of Cowpea Seed Coats. *Annals of Botany* 76: pp. 287 – 290.
17. Emeruwa, A.C. (1982). Antibacterial substances from *Carica papaya* fruit extract. *Lloydia*. 45(2):123 – 127.
18. Collins, G.H., Lynes, P.M. and Grange, J.M. (1995). *Microbiological Methods* (7th Edition). Butterworth Heinemann Ltd, Britain. Pp. 175 – 190.
19. Tijjani, M.A., Abdurahaman, F.I., Abba, A., Idris, Y.S., Baburo, M., Mala, B.S.I., Mala, G.A., Dungus, M.H.M., Aji, B.M. and Abubakar, K.I. (2013). Evaluation of Proximate and Phytochemical Composition of leaves of *Annona senegalensis*. *Journal of Pharmaceutical & Scientific Innovation*. Vol. 2, Issue 1, p. 7.
20. Nonaka, G.I., Nishioka, I., Nishizawa, M., Yamagishi, T., Kashiwada, Y., Dutschman, G.E., Bodner, A.J., Kilkuskie, R.E., Cheng, Y.C. and Lee, K.H. (1990). Anti-AIDS Agents 2. Inhibitory Effect of Tannins on HIV Reverse Transcriptase and HIV replication in H9 lymphocyte cells. *J. Nat. Prod.* 53:387 – 595 (Pub Med).
21. Denwick, P.M. (2002). *Natural Products: A Biosynthetic Approach*. 2nd Edition; John Wiley and Sons Ltd., England. Pp. 241 – 243.
22. Pengsuparp, T., Cail, Constant, H., Fong, H.H., Lin, L.Z., Kinghorn, A.D., Ingoltsdottir, K., Wagner, H. (1995). Mechanistic evolution of new plant derived compounds that inhibit HIV-1 reverse transcriptase. *J. Nat. Prod.* 58:1024 – 1031. (Pub Med).
23. Hemen, D. and Lalita, L. (2012). A Review on Anthraquinone isolated from Cassie species and their applications. *Indian Journal of Natural Products and Resources*. Vol. 3(3):291 – 319.
24. Fujjoka, T. and Kashwada Y. (1996). Anti-AIDS agents. 11. Betulinc acid and platonic acid as anti-HIV principles form syzigiumClaviflorium and the anti-HIV activity of structurally related triterpenoids. *J. Nat. Prod.* 57:243 – 247.
25. Steinberg, P.D., Hagerman, A.E., Stern, J.L., Manson, P.K. (1996). Phloro tannin-protein interactions. *J. Chem. Ecol.* 22:1887 – 1899.
26. Hopp, K.H., Cunningham, L.V.J., Wahba, Khalil, K. (2003). *In vitro* antitrypanosomal activity of certain alkaloids against *Trypanosomalewisi*.
27. Mendez, E., Moreno, A., Colilla, F. Pelaez, F. Limas, G.G., Mendez, R., Soxiano, F., Salinas, M., DeHaro, C. primary structure and inhibition of protein synthesis in eukaryotic cell

- fresystem of a novel thionin, gamma horothionin from barely endosperm Eur (Biochem, 2000; 194:333339. [pub med]).
28. Simon, M. (1993). The Systematic Distribution of tannins in the leaves of Angiosperms: A tool for Ecological Studies. *Biochemical Systematics and Ecology*. 21(8):833 – 846. Doi:10.1016/0308-1978. (93)90096-.
 29. Bakar, K. (2007). *In vitro* antioxidant studies in leaves of *Anonnaspecies*. *Indian Journal of Experiment Biology*. Vol. 45, pp. 480 – 485.
 30. Doss, A., Mohammed, H.M. and Dhanabalan, R. (2009). Antibacterial activity of tannins from the leaves of *Solanum trilobatum* Linn. *Indian J. Sci. Technology*. Vol. 2, No. 2. ISSN:0974 – 6848.
 31. Jones, G.A., McAllister, T.A., Mur, A.D. and Cheng, K.J. (2006). Effect of Saifonin (*Onobrychis viciolia* Scop). Condensed tannins on growth and proteolysis by four strain of terminal bacteria. *Appl. Environ. microbial*. 60: 1374 – 1378.
 32. Viljoen, A., Van Vuuren, S., Ernest, E., Klepser, M., Demiri, B., Bassar, H. and Van Wyk, B.E. (2003). *Osmitopsis asteriscoides* (Asteraceae)- the antimicrobial and essential oil composition of Cape – Dutch remedy. J. *Enthnopharmacol*. 88, 137 – 143.
 33. Kurosaki F. and Nishi A (1983) isolation and Antimicrobial activity of the Phytoalexin-6-methoxymellein from cultured carrots cells. *Phytochemistry*. 22(3), 669-672.
 34. Banso, A. and Adeyemo, S.O. (2007). Evaluation of antibacterial properties of tannins isolated from *Dichrostachys cinerea*. *Afr. J. Biotechnol*. 6(15):1785– 1787.
 35. Davison, H.C., Low, J.C., Woodhouse, M.E. (2000). “What is antibiotic resistance and how can we measure it? *Trend in Microbiology*. 8(112):554 – 559. PMID: 11115751.